



**HAL**  
open science

## Supporting the mental health of adults bereaved by suicide: A systematic review of protective factors

Clémence Jacquet, Geoffrey Gauvin, Edouard Leaune, Jules Mercier, Damien Fouques

### ► To cite this version:

Clémence Jacquet, Geoffrey Gauvin, Edouard Leaune, Jules Mercier, Damien Fouques. Supporting the mental health of adults bereaved by suicide: A systematic review of protective factors. 20th European Symposium on Suicide and Suicidal Behaviour, Aug 2024, Rome, Italy. hal-04742327

**HAL Id: hal-04742327**

**<https://hal.science/hal-04742327v1>**

Submitted on 17 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Supporting the mental health of adults bereaved by suicide : A systematic review of protective factors

Clémence Jacquet<sup>1,2</sup>, Geoffrey Gauvin<sup>1,2</sup>, Edouard Leune<sup>3,4</sup>, Jules Mercier<sup>4</sup> and Damien Fouques<sup>1</sup>

<sup>1</sup> Laboratoire Psychopathologie et Processus de Changement (LPPC) - UR 2027 ; Université Paris 8  
<sup>2</sup> Centre de recherche et d'intervention sur le suicide, enjeux éthiques et pratiques de fin de vie (CRISE)  
<sup>3</sup> RESHAPE (RESearch on HealthAre PErformance) - INSERM U1290 - Université Claude Bernard Lyon 1  
<sup>4</sup> Centre de Prévention du Suicide - Centre Hospitalier Le Vinatier

## BACKGROUND

- Exposure to the suicide of a loved one is associated with poor mental health outcomes, including an increased risk of **depression**<sup>1</sup>, **post-traumatic stress disorder**<sup>2</sup>, **prolonged grief disorder**<sup>2</sup> and **suicidal behavior**<sup>3</sup>.
- Thus, understanding the characteristics and mechanisms that promote mental health in suicide-loss survivors is essential for developing appropriate interventions and support for this vulnerable population.

➔ We therefore sought to systematically synthesize the evidence on protective factors in adults bereaved by suicide.

## METHOD

- Following guidelines for systematic reviews, searches were performed using **PubMed, PsycINFO, Scopus, Web of Science** and **Embase** by two independent reviewers using the same criteria and search procedures.
- Studies were included if they **1)** were original research published in peer-reviewed journals in English or French prior to May 2024, and **2)** focused on protective factors that moderate the risk of developing psychopathologies or support good mental health in adults bereaved by the suicide of a close relative.

## FINDINGS

- After screening (see Figure 1), **31 articles** were included, including 4 longitudinal, 17 cross-sectional and 10 qualitative studies.





Fig.1. PRISMA Flow Chart

- Results revealed several **interpersonal**, **intrapersonal** and **sociodemographic** factors that can prevent the development of mental disorders in adults bereaved by suicide.

## Mapping Protective Factors and Mental Health in the Bereaved by Suicide



## DISCUSSION & CONCLUSION

- Despite the major impact of suicide on loved ones, various factors can support the mental health of the bereaved. These factors need to be understood from a **biopsychosocial perspective**, involving individual, interpersonal, and sociodemographic characteristics.
- However, most of the included studies were cross-sectional, which limits the ability to make causal inferences between protective factors and outcomes. Additionally, some studies involved the same participants, reducing the scope of their results, and most had limited sample sizes.
- To better identify the elements and underlying mechanisms of resilience in suicide loss survivors, **further large-scale longitudinal studies are needed.**

### Références

- Bolton JM, Au W, Leslie WD, Martens PJ, Enns MW, Roos LL, Katz LY, Wilcox HC, Erlangsen A, Chateau D, Walld R, Spiwak R, Seguin M, Shear K, Sareen J. Parents bereaved by offspring suicide: a population-based longitudinal case-control study. *JAMA Psychiatry*. 2013 Feb;70(2):158-67. doi: 10.1001/jamapsychiatry.2013.275. PMID: 232298802
- Grafiadeli R, Glaesmer H, Wagner B. Loss-Related Characteristics and Symptoms of Depression, Prolonged Grief, and Posttraumatic Stress Following Suicide Bereavement. *Int J Environ Res Public Health*. 2022 Aug 18;19(16):10277. doi: 10.3390/ijerph191610277. PMID: 36011928; PMCID: PMC9408305
- Pitman A, Osborn D, King M, Erlangsen A. Effects of suicide bereavement on mental health and suicide risk. *Lancet Psychiatry*. 2014 Jun;1(1):86-94. doi: 10.1016/S2215-0366(14)70224-X. Epub 2014 Jun 4. PMID: 26360405.