



**HAL**  
open science

# On the connection between the Beurling-Malliavin density and the asymptotic density

Rita Giuliano, Georges Grekos

► **To cite this version:**

Rita Giuliano, Georges Grekos. On the connection between the Beurling-Malliavin density and the asymptotic density. *Uniform Distribution Theory*, 2024, 19 (1), pp.67-86. hal-04742362

**HAL Id: hal-04742362**

**<https://hal.science/hal-04742362v1>**

Submitted on 23 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On the connection between the Beurling-Malliavin density and the asymptotic density

Rita Giuliano\*

Georges Grekos†‡

## Abstract

We study the notion of Beurling-Malliavin density from the point of view of Number Theory. We prove a general relation between the Beurling-Malliavin density and the upper asymptotic density; we identify a class of sequences for which the two densities coincide; this class contains the arithmetic progressions. Last, by means of an alternative definition of Beurling-Malliavin density, we study the connection with the asymptotic density for another kind of sequences that again generalizes the arithmetic progressions.

*Keywords:* Beurling-Malliavin density, asymptotic density, increasing sequence, substantial interval, complete system, first digit set, regularly varying function, set of powers

*2020 Mathematical Subject Classification:* 11B25, 11B05

## 1 Introduction

Let  $\Lambda = (\lambda_n)_{n \in \mathbb{Z}}$  be a sequence of real numbers. In order to solve the problem of finding the radius of completeness of  $\Lambda$  (denoted by  $\mathcal{R}(\Lambda)$ ), A. Beurling and P. Malliavin introduced for the first time in the paper [2] the quantity  $b(\Lambda)$ , defined as follows: if  $(I_n)_{n \in \mathbb{Z}}$  is a sequence of disjoint intervals on  $\mathbb{R}$ , call it *short* if

$$\sum_{n \in \mathbb{Z}} \frac{|I_n|^2}{1 + \text{dist}^2(0, I_n)} < \infty$$

and *long* otherwise; then define

$$b(\Lambda) := \sup\{d : \exists \text{ long } (I_n)_{n \in \mathbb{Z}} \text{ such that } \#(\Lambda \cap I_n) > d|I_n|, \forall n \in \mathbb{Z}\}$$

(the particular formulation used above comes from [12] and [14]). The celebrated Theorem of [2] states that the radius of completeness of  $\Lambda$  is connected with  $b(\Lambda)$  by the formula

$$\mathcal{R}(\Lambda) = 2\pi b(\Lambda).$$

Later the same quantity  $b(\Lambda)$  was studied by other authors, for instance [8], [9], and used in several papers; we cite [7], [10] and [15]; for an extensive study and generalizations see [13], [12], [1] (see [14])

---

\*Dipartimento di Matematica, Università di Pisa, Largo Bruno Pontecorvo 5, I-56127 Pisa, Italy (email: [rita.giuliano@unipi.it](mailto:rita.giuliano@unipi.it), ORCID: 0000-0002-1638-2651)

†Faculté des Sciences et Techniques, Université Jean Monnet, 23 rue Dr Paul Michelon. 42023 Saint-Etienne Cedex 2, France (email: [grekos@univ-st-etienne.fr](mailto:grekos@univ-st-etienne.fr), ORCID: 0000-0002-3681-8336).

‡The authors are grateful to the CIRM of Marseille for the hospitality offered to them in 2019 (under the Program *Research in pairs* <https://conferences.cirm-math.fr/2019-calendar.html>), when this research was at the beginning.

for a complete list of references); in particular the author of [15] discovers an alternative equivalent definition, which will be stated and used later (see Definition 2.1).

In the whole literature on the subject the quantity  $b(\Lambda)$  is called the “density” of  $\Lambda$ ; in the sequel we call it “Beurling-Malliavin density”(or “BM-density” for short); anyway, to the best of our knowledge, no paper addresses the problem of investigating whether it “deserves”(so to say) the name of “density”, in the sense in which this term is used in Number Theory. Just to give a trivial example, we have not found any paper stating that the BM-density of the set of multiples of the positive integer  $p$  is  $\frac{1}{p}$ .

Thus, the most natural notion of density of a sequence  $\Lambda$  of positive integers used in Number Theory being the asymptotic density (denoted by the symbol  $d(\Lambda)$ ), the aim of the present investigation is to study the connection between  $b(\Lambda)$  and  $d(\Lambda)$ .

The Beurling-Malliavin density is defined in [2] for general sequences  $\Lambda = (\lambda_n)_{n \in \mathbb{Z}}$  of real numbers; on the other hand, if  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  is a sequence of integers, we may be interested in calculating its asymptotic density. Thus, in order to compare these two concepts, in what follows we shall confine ourselves to sequences of real numbers indexed by  $n \in \mathbb{N}^*$ , positive, ultimately strictly increasing and such that  $\lim_{n \rightarrow \infty} \lambda_n = +\infty$ . In the sequel, by the term *sequence* we always mean a sequence with such properties, unless otherwise specified.

**Remark 1.1.** *Notice that sequences with a finite number of elements are excluded from this discussion, but this is a trivial case (both the BM-density and asymptotic density vanish). Similarly, we are not dealing with sequences with repetitions (see Remark 6.2).*

The paper is structured as follows. Section 2 contains two equivalent definitions of the BM-density; similarly, Section 3 discusses two equivalent definitions of the asymptotic density (upper and lower); in Section 4 we prove a general inequality relating the upper asymptotic density and the BM-density; in Section 5 we prove the coincidence of the asymptotic density and the BM-density for a particular class of sequences, to which arithmetic progressions belong. Last, in Section 6 (i) we show how to use the alternative definition 2.1 for estimating the BM-density of some kind of sequences that again generalizes arithmetic progressions; (ii) we find a nice characterization of the BM-density (see Theorem 6.2 and Corollary 6.2).

We shall use the standard symbols  $[x]$  and  $\lceil x \rceil$  to mean respectively the greatest integer less than or equal to  $x$  (*integer part of  $x$* ) and the least integer greater than or equal to  $x$ .

The *counting function* of  $\Lambda = \{\lambda_n\}_{n \in \mathbb{N}^*}$  is defined as

$$F_\Lambda(t) = \begin{cases} 0 & t = 0 \\ \#\{k \in \mathbb{N}^* : \lambda_k \leq t\} & t > 0. \end{cases}$$

For  $a < b$  we have clearly

$$F_\Lambda(b) - F_\Lambda(a) = \#\{k \in \mathbb{N}^* : \lambda_k \in (a, b]\}.$$

## 2 Various equivalent definitions of the Beurling-Malliavin density

The Beurling-Malliavin density  $b(\Lambda)$ , firstly defined in [2], has been studied later in [9] and in [15] (among others); both the definitions of [9] and [15] are different but equivalent to the original one. In the very recent paper [6] a further equivalent definition has been given. In the present paper we shall use first the approach of [15] and later the one of [6].

## 2.1 The definition of [15]

In [15] it is proved that an equivalent form of Beurling–Malliavin density is given by the following

**Definition 2.1.** *The Beurling Malliavin density  $b(\Lambda)$  of the sequence  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  is the infimum of the numbers  $a \geq 0$  for which there exists a sequence of distinct positive integers  $N = (n_k)_{k \in \mathbb{N}^*}$  such that*

$$\sum_{k=1}^{\infty} \left| \frac{1}{\lambda_k} - \frac{a}{n_k} \right| < \infty.$$

In [15] it is shown that the set  $\mathcal{S}$  of such  $a$  is a right infinite interval in  $\mathbb{R}^+$  (i.e. if  $a$  belongs to  $\mathcal{S}$  and  $b > a$ , then  $b$  belongs to  $\mathcal{S}$  as well). If  $\mathcal{S}$  is empty, then  $b(\Lambda) = +\infty$ .

**Remark 2.1.** *If  $\Lambda$  is a sequence of integers, we have obviously  $b(\Lambda) \leq 1$  (take  $a = 1$  and  $N = \Lambda$  in the above definition).*

## 2.2 The definition of [6]

Let  $\mathfrak{C}$  be the family of all sequences

$$\mathcal{I} = ((a_n, b_n])_{n \in \mathbb{N}^*}$$

of intervals in  $(0, +\infty)$  such that  $a_n < b_n \leq a_{n+1}$  for all  $n \in \mathbb{N}^*$  and

$$\sum_{n=1}^{\infty} \left( \frac{b_n}{a_n} - 1 \right)^2 = +\infty.$$

**Remark 2.2.** *These systems of intervals are referred to as long in [12] and [14] (see the Introduction). In [9] they are called substantial.*

For every  $\mathcal{I} = ((a_n, b_n])_{n \in \mathbb{N}^*} \in \mathfrak{C}$ , let

$$\ell_{\mathcal{I}} = \liminf_{n \rightarrow \infty} \frac{F_{\Lambda}(b_n) - F_{\Lambda}(a_n)}{b_n - a_n}. \quad (1)$$

In [6] it is proved that

**Proposition 2.1.** *The following relation holds true*

$$b(\Lambda) = \sup\{\ell_{\mathcal{I}}, \mathcal{I} \in \mathfrak{C}\}.$$

We shall be interested in the subset of  $\mathfrak{C}$  defined as

$$\mathfrak{C}_{>1} = \left\{ \mathcal{I} = ((a_n, b_n])_{n \in \mathbb{N}^*} \in \mathfrak{C} : \limsup_{n \rightarrow \infty} \frac{b_n}{a_n} > 1 \right\}.$$

Accordingly, we shall denote

$$b_{>1}(\Lambda) = \sup\{\ell_{\mathcal{I}}, \mathcal{I} \in \mathfrak{C}_{>1}\}. \quad (2)$$

From Proposition 2.1 it follows that

$$b_{>1}(\Lambda) \leq b(\Lambda). \quad (3)$$

### 3 Two equivalent definitions of the asymptotic density

Let  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  be a sequence of positive numbers.

**Definition 3.1.** (i) The lower (resp. upper) asymptotic density of  $\Lambda$  is

$$\underline{d}(\Lambda) = \liminf_{n \rightarrow \infty} \frac{F_\Lambda(n)}{n}, \quad (\text{resp. } \bar{d}(\Lambda) = \limsup_{n \rightarrow \infty} \frac{F_\Lambda(n)}{n}).$$

(ii) We say that  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  has asymptotic density  $d(\Lambda)$  if

$$\underline{d}(\Lambda) = \bar{d}(\Lambda) (= d(\Lambda)).$$

It is well known that, besides the definition, there is another way for calculating the upper and lower asymptotic densities of  $\Lambda$ . Precisely we have

**Proposition 3.1.** The set of limit points of the sequence  $(\frac{k}{\lambda_k})_{k \in \mathbb{N}^*}$  coincides with the set of limit points of the sequence  $(\frac{F_\Lambda(n)}{n})_{n \in \mathbb{N}^*}$ .

In particular the following relations hold true

$$\underline{d}(\Lambda) = \liminf_{k \rightarrow \infty} \frac{k}{\lambda_k}; \quad \bar{d}(\Lambda) = \limsup_{k \rightarrow \infty} \frac{k}{\lambda_k}.$$

Moreover  $d(\Lambda)$  exists if either of the two limits

$$\lim_{n \rightarrow \infty} \frac{F_\Lambda(n)}{n}, \quad \lim_{k \rightarrow \infty} \frac{k}{\lambda_k}$$

exists; in this case

$$d(\Lambda) = \lim_{n \rightarrow \infty} \frac{F_\Lambda(n)}{n} = \lim_{k \rightarrow \infty} \frac{k}{\lambda_k}.$$

*Proof.* In fact, if  $\lambda_{k_n} \leq n < \lambda_{k_n+1}$ , we have  $F_\Lambda(n) = k_n$ , and

$$\frac{k_n}{\lambda_{k_n+1}} < \frac{F_\Lambda(n)}{n} \leq \frac{k_n}{\lambda_{k_n}}. \quad (4)$$

Notice that  $\lim_{n \rightarrow \infty} k_n = \infty$  and let  $\ell$  be a limit point for the sequence  $(\frac{F_\Lambda(n)}{n})_{n \in \mathbb{N}^*}$ . Then there exists a subsequence  $(n_r)_{r \in \mathbb{N}^*}$  converging to  $\infty$  such that  $\frac{F_\Lambda(n_r)}{n_r} \rightarrow \ell$  as  $r \rightarrow \infty$ . Relation (4) implies that

$$\bar{\kappa} := \limsup_{r \rightarrow \infty} \frac{k_{n_r}}{\lambda_{k_{n_r}+1}} = \limsup_{r \rightarrow \infty} \frac{k_{n_r}}{\lambda_{k_{n_r}+1}} \leq \ell \leq \liminf_{r \rightarrow \infty} \frac{k_{n_r}}{\lambda_{k_{n_r}}} =: \underline{\kappa}.$$

In other words every limit point for  $(\frac{F_\Lambda(n)}{n})_{n \in \mathbb{N}^*}$  is a limit point for  $(\frac{k}{\lambda_k})_{k \in \mathbb{N}^*}$ .

To go the inverse direction, notice that  $F_\Lambda(\lambda_k) = k$ ; thus, for  $n_k \leq \lambda_k < n_k + 1$ ,

$$\frac{F_\Lambda(n_k)}{n_k + 1} \leq \frac{k}{\lambda_k} \leq \frac{F_\Lambda(n_k + 1)}{n_k};$$

now argue as in the first part of this proof, using the fact that  $\lim_{k \rightarrow \infty} n_k = \infty$ .

□

## 4 The general inequality

**Theorem 4.1.** *For every sequence  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$ , the relation*

$$\bar{d}(\Lambda) \leq b(\Lambda) \tag{5}$$

holds true.

*Proof.* Recall that the Beurling–Malliavin Theorem states that, putting

$$\mathcal{E}_\Lambda = \{e^{\pm i\lambda_n x}\}, \quad \mathcal{R}(\Lambda) = \sup\{a \in \mathbb{R}^+ : \mathcal{E}_\Lambda \text{ is complete in } L^2(0, a)\},$$

we have

$$\mathcal{R}(\Lambda) = 2\pi b(\Lambda).$$

Hence it suffices to prove that, for every  $\epsilon > 0$ ,  $\mathcal{E}_\Lambda$  is complete in  $L^2(0, 2\pi(\bar{d}(\Lambda) - \epsilon))$ . Let  $f$  be a function in  $L^2(0, 2\pi(\bar{d}(\Lambda) - \epsilon))$  and assume that

$$\int_0^{2\pi(\bar{d}(\Lambda) - \epsilon)} e^{\pm i\lambda_n x} f(x) dx = 0, \quad n = 1, 2, 3, \dots$$

By the change of variable  $y = \frac{x}{\bar{d}(\Lambda) - \epsilon} - \pi$ , the above integral becomes

$$(\bar{d}(\Lambda) - \epsilon) e^{\pm i\lambda_n (\bar{d}(\Lambda) - \epsilon)\pi} \int_{-\pi}^{\pi} e^{\pm i\lambda_n (\bar{d}(\Lambda) - \epsilon)y} f((\bar{d}(\Lambda) - \epsilon)(y + \pi)) dy.$$

In [3] the following result is proved (see Theorem XXVIII, p. 84):

**Theorem 4.2.** *Let  $0 < m_1 < m_2 < \dots$  and let*

$$\limsup_{n \rightarrow \infty} \frac{n}{m_n} > 1.$$

*Then, if  $f \in L^2(-\pi, \pi)$  and*

$$\int_{-\pi}^{\pi} e^{\pm im_n x} f(x) dx = 0, \quad n = 1, 2, 3, \dots,$$

*then  $f(x) = 0$  for almost every  $x \in (-\pi, \pi)$ .*

Setting  $m_n = (\bar{d}(\Lambda) - \epsilon)\lambda_n$ , we obtain that

$$\int_{-\pi}^{\pi} e^{\pm im_n y} g(y) dy = 0,$$

where  $g$  is the function  $y \in (-\pi, \pi) \mapsto f((\bar{d}(\Lambda) - \epsilon)(y + \pi))$ .

Observe that

$$\limsup_{n \rightarrow \infty} \frac{n}{m_n} = \limsup_{n \rightarrow \infty} \frac{n}{(\bar{d}(\Lambda) - \epsilon)\lambda_n} = \frac{\bar{d}(\Lambda)}{\bar{d}(\Lambda) - \epsilon} > 1.$$

Hence Theorem 4.2 is in force and we deduce that  $g$  vanishes almost everywhere on  $(-\pi, \pi)$ . This is obviously equivalent to saying that  $f$  vanishes almost everywhere on  $2\pi(\bar{d}(\Lambda) - \epsilon)$ , i.e. that  $\mathcal{E}_\Lambda$  is complete in  $L^2(0, 2\pi(\bar{d}(\Lambda) - \epsilon))$ . □

**Remark 4.1.** *Theorem 4.1 is stated but not proved in [13]. In any case, [13] doesn't formulate it in terms of densities; see slide 3 in [13].*

The following example shows that the inequality (5) may be strict.

**Example 4.1.** *Let  $p$  be an integer with  $1 \leq p \leq 9$  and denote by  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  the - strictly increasing - sequence formed by the positive integers having first digit equal to  $p$ . It is well known (see for instance [5]) that  $\bar{d}(\Lambda) = \frac{10}{9(p+1)} < 1$ , and now we prove that  $b(\Lambda) = 1$ . Recall that the BM-density of any set of integers is  $\leq 1$ . By formula (2) and relation (3), it suffices to prove that there exists a family of intervals  $\mathcal{I} = ((a_n, b_n])_{n \in \mathbb{N}^*}$  such that  $\limsup_{n \rightarrow \infty} \frac{b_n}{a_n} > 1$  and*

$$\ell_{\mathcal{I}} = \liminf_{n \rightarrow \infty} \frac{F_{\Lambda}(b_n) - F_{\Lambda}(a_n)}{b_n - a_n} \geq 1.$$

Take

$$a_n = p10^n, \quad b_n = (p+1)10^n - 1.$$

We have  $b_n - a_n = 10^n - 1$  and

$$F_{\Lambda}(b_n) = \sum_{k=1}^n \sum_{j=p10^k}^{(p+1)10^k-1} 1 = \sum_{k=1}^n \{(p+1)10^k - p10^k\} = \sum_{k=1}^n 10^k = \frac{10}{9}(10^n - 1);$$

$$F_{\Lambda}(a_n) = F_{\Lambda}(b_{n-1}) + 1 = \frac{10}{9}(10^{n-1} - 1) + 1 = \frac{1}{9}(10^n - 1).$$

Thus

$$\ell_{\mathcal{I}} = \liminf_{n \rightarrow \infty} \frac{F_{\Lambda}(b_n) - F_{\Lambda}(a_n)}{b_n - a_n} = \liminf_{n \rightarrow \infty} \frac{\frac{10}{9}(10^n - 1) - \frac{1}{9}(10^n - 1)}{10^n - 1} = 1.$$

**Remark 4.2.** *It is easy to check that for  $a_n = (p+1)10^n - 1$  and  $b_n = a_{n+1} = (p+1)10^{n+1} - 1$  we have*

$$\ell_{\mathcal{I}} = \bar{d}(\Lambda) = \frac{10}{9(p+1)}.$$

## 5 A general result concerning the equality

The aim of this Section is to prove the following result, in which a class of sequences  $\Lambda$  is identified for which  $d(\Lambda) = b(\Lambda)$ .

**Theorem 5.1.** *Assume that  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  is a sequence with the following property: there exists an integer  $p \geq 1$  and non-negative constants  $a_0, a_1, \dots, a_{p-1}$  such that*

$$\lim_{k \rightarrow \infty} \{\lambda_{kp+j+1} - \lambda_{kp+j}\} = a_j, \quad \forall j = 0, 1, 2, \dots, p-1.$$

Assume in addition that  $\sum_{j=0}^{p-1} a_j > 0$ . Then

$$d(\Lambda) = b(\Lambda) = \frac{p}{\sum_{j=0}^{p-1} a_j}.$$

For the proof of this result, we need some lemmas.

**Lemma 5.1.** *Assume that  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  verifies the conditions of Theorem 5.1. Then*

$$\lim_{n \rightarrow \infty} \frac{\lambda_n}{n} = \frac{\sum_{j=0}^{p-1} a_j}{p}.$$

*Proof.* We shall prove that, for every sequence  $(\lambda_{kp+j})_{k \in \mathbb{N}^*}$ ,  $j = 0, 1, \dots, p-1$ , we have

$$\lim_{k \rightarrow \infty} \frac{\lambda_{kp+j}}{kp+j} = \frac{\sum_{j=0}^{p-1} a_j}{p}.$$

First, by Cesaro Theorem,

$$\lim_{k \rightarrow \infty} \frac{\lambda_{kp+j}}{kp+j} = \lim_{k \rightarrow \infty} \frac{\lambda_{(k+1)p+j} - \lambda_{kp+j}}{(k+1)p+j - (kp+j)} = \lim_{k \rightarrow \infty} \frac{\lambda_{(k+1)p+j} - \lambda_{kp+j}}{p}.$$

Now

$$\begin{aligned} \lim_{k \rightarrow \infty} \{\lambda_{(k+1)p+j} - \lambda_{kp+j}\} &= \sum_{r=j}^{p+j-1} \lim_{k \rightarrow \infty} \{\lambda_{kp+r+1} - \lambda_{kp+r}\} \\ &= \sum_{r=j}^{p-1} \lim_{k \rightarrow \infty} \{\lambda_{kp+r+1} - \lambda_{kp+r}\} + \sum_{r=p}^{p+j-1} \lim_{k \rightarrow \infty} \{\lambda_{kp+r+1} - \lambda_{kp+r}\} \\ &= \sum_{r=j}^{p-1} a_r + \sum_{s=0}^{j-1} \lim_{k \rightarrow \infty} \{\lambda_{(k+1)p+s+1} - \lambda_{(k+1)p+s}\} \\ &= \sum_{r=j}^{p-1} a_r + \sum_{s=0}^{j-1} \lim_{k \rightarrow \infty} \{\lambda_{kp+s+1} - \lambda_{kp+s}\} \\ &= \sum_{r=j}^{p-1} a_r + \sum_{s=0}^{j-1} a_s = \sum_{r=0}^{p-1} a_r. \end{aligned}$$

□

**Remark 5.1.** Lemma 5.1 implies that any sequence  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  satisfying the assumptions of Theorem 5.1 has the form

$$\lambda_n = \frac{\sum_{j=0}^{p-1} a_j}{p} n + \psi_n,$$

where

$$\lim_{n \rightarrow \infty} \frac{\psi_n}{n} = 0. \quad (6)$$

Besides (6), a necessary condition for  $\Lambda$  to satisfy the assumptions of Theorem 5.1 is that

$$\lim_{n \rightarrow \infty} \psi_{n+p} - \psi_n = 0.$$

This is easily seen:

$$\psi_{n+p} - \psi_n = \lambda_{n+p} - \frac{\sum_{j=0}^{p-1} a_j}{p} (n+p) - \lambda_n + \frac{\sum_{j=0}^{p-1} a_j}{p} n = \lambda_{n+p} - \lambda_n - \sum_{j=0}^{p-1} a_j = 0$$

since, for  $n = kp + j$ ,  $j = 0, \dots, p-1$ , we have

$$\lim_{k \rightarrow \infty} \{\lambda_{kp+j+p} - \lambda_{kp+j}\} = \sum_{j=0}^{p-1} a_j,$$

as has been proven in Lemma 5.1.



**Lemma 5.2.** Let  $(x_n)_{n \in \mathbb{N}^*}$  and  $(y_n)_{n \in \mathbb{N}^*}$  be two sequences of numbers, with

$$\lim_{n \rightarrow \infty} (y_n - x_n) = +\infty. \quad (7)$$

Then

$$\lim_{n \rightarrow \infty} \frac{\lfloor y_n \rfloor - \lfloor x_n \rfloor}{y_n - x_n} = 1.$$

*Proof.* Let  $\{x\} = x - \lfloor x \rfloor$  be the fractional part of the number  $x$  and write

$$\frac{\lfloor y_n \rfloor - \lfloor x_n \rfloor}{y_n - x_n} = \frac{1}{1 + \frac{\{y_n\} - \{x_n\}}{\lfloor y_n \rfloor - \lfloor x_n \rfloor}}.$$

Observe that  $|\{y_n\} - \{x_n\}| < 1$  (from  $0 \leq \{x\} < 1$ ) and that  $\lim_{n \rightarrow \infty} (\lfloor y_n \rfloor - \lfloor x_n \rfloor) = +\infty$  (since  $\lfloor y_n \rfloor - \lfloor x_n \rfloor \geq y_n - x_n - 1$ ). The statement follows.  $\square$

Denote by  $\bar{\lambda}$  the interpolated linear function relative to  $\Lambda$ , i.e. the function defined by

$$\bar{\lambda}(x) = (\lambda_{k+1} - \lambda_k)(x - k) + \lambda_k, \quad k \leq x < k + 1, \quad k = 1, 2, \dots$$

or equivalently

$$\bar{\lambda}(x) = (\lambda_{\lceil x \rceil} - \lambda_{\lfloor x \rfloor})(x - \lfloor x \rfloor) + \lambda_{\lfloor x \rfloor}.$$

From [4] we recall the

**Definition 5.1.** The function  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is regularly varying with exponent  $\rho > 0$  if, for every  $t > 0$ , we have

$$\lim_{x \rightarrow +\infty} \frac{f(tx)}{f(x)} = t^\rho.$$

**Lemma 5.3.** Assume that  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  is such that

$$\lim_{n \rightarrow \infty} \frac{\lambda_n}{n} = \ell > 0. \quad (8)$$

Then

(i)  $\lim_{x \rightarrow +\infty} \frac{\bar{\lambda}(x)}{x} = \ell;$

(ii)  $\bar{\lambda}$  is regularly varying with exponent 1;

(iii) for every  $\beta > 1$  we have

$$\lim_{\substack{x \rightarrow +\infty \\ \beta x < y \rightarrow +\infty}} \frac{\bar{\lambda}(y) - \bar{\lambda}(x)}{y - x} = \ell.$$

*Proof.* First, from the assumption (8) and the relations

$$\frac{\lfloor x \rfloor}{\lceil x \rceil} \cdot \frac{\bar{\lambda}(\lfloor x \rfloor)}{\lfloor x \rfloor} = \frac{\bar{\lambda}(\lfloor x \rfloor)}{\lceil x \rceil} \leq \frac{\bar{\lambda}(\lfloor x \rfloor)}{x} \leq \frac{\bar{\lambda}(\lceil x \rceil)}{\lfloor x \rfloor}$$

and

$$\frac{\bar{\lambda}(\lceil x \rceil)}{\lceil x \rceil} \leq \frac{\bar{\lambda}(\lceil x \rceil)}{x} \leq \frac{\bar{\lambda}(\lceil x \rceil)}{\lfloor x \rfloor} = \frac{\bar{\lambda}(\lceil x \rceil)}{\lceil x \rceil} \cdot \frac{\lceil x \rceil}{\lfloor x \rfloor},$$

(where we have used also the fact that  $\bar{\lambda}$  is non-decreasing) and by the relation  $\lim_{x \rightarrow +\infty} \frac{[x]}{x} = 1$ , we obtain

$$\lim_{x \rightarrow +\infty} \frac{\bar{\lambda}(\lfloor x \rfloor)}{x} = \lim_{x \rightarrow +\infty} \frac{\bar{\lambda}(\lceil x \rceil)}{x} = \ell, \quad (9)$$

and (i) is obvious, since

$$\frac{\bar{\lambda}(\lfloor x \rfloor)}{x} \leq \frac{\bar{\lambda}(x)}{x} \leq \frac{\bar{\lambda}(\lceil x \rceil)}{x}.$$

For (ii), we have to prove that for every  $t > 0$ ,

$$\lim_{x \rightarrow \infty} \frac{\bar{\lambda}(tx)}{\bar{\lambda}(x)} = t,$$

and this can be obtained by writing

$$\frac{\bar{\lambda}(tx)}{\bar{\lambda}(x)} = t \cdot \frac{\bar{\lambda}(tx)}{tx} \cdot \frac{x}{\bar{\lambda}(x)}$$

and using point (i).

Now we prove point (iii). Write

$$\frac{\bar{\lambda}(y) - \bar{\lambda}(x)}{y - x} = \frac{(\lambda_{\lfloor y \rfloor} - \lambda_{\lfloor y \rfloor})(y - \lfloor y \rfloor)}{y - x} + \frac{(\lambda_{\lceil x \rceil} - \lambda_{\lceil x \rceil})(x - \lceil x \rceil)}{y - x} + \frac{\lambda_{\lfloor y \rfloor} - \lambda_{\lceil x \rceil}}{y - x}.$$

Notice that

$$\limsup_{x \rightarrow +\infty} (\lambda_{\lceil x \rceil} - \lambda_{\lceil x \rceil})(x - \lceil x \rceil) < +\infty.$$

This is because  $x - \lceil x \rceil = \{x\}$  (fractional part of  $x$ ) is between 0 and 1; furthermore

$$\limsup_{x \rightarrow +\infty} (\lambda_{\lceil x \rceil} - \lambda_{\lceil x \rceil}) = \max_{0 \leq j \leq p-1} a_j.$$

Hence the first two summands above converge to 0, since  $y - x > x(\beta - 1) \rightarrow +\infty$  as  $x \rightarrow +\infty$ .

Concerning the last summand, we write it as

$$\frac{\lambda_{\lfloor y \rfloor}}{y} + \left( \frac{\lambda_{\lfloor y \rfloor}}{y} - \frac{\lambda_{\lceil x \rceil}}{x} \right) \cdot \frac{x}{y - x}$$

which allows to obtain the claimed result by (9) and the fact that

$$0 \leq \frac{x}{y - x} \leq \frac{1}{\beta - 1}.$$

□

*Proof of Theorem 5.1.*

Recall that the asymptotic density  $d(\Lambda)$  of the sequence  $\Lambda$  is the  $\lim_{n \rightarrow \infty} \frac{n}{\lambda_n}$ , if this limit exists (see Proposition 3.1). Hence by Lemma 5.1 we have  $d(\Lambda) = \frac{p}{\sum_{j=0}^{p-1} a_j}$ , and now we shall prove that

$$\frac{p}{\sum_{j=0}^{p-1} a_j} \leq b_{>1}(\Lambda). \quad (10)$$

Recalling (2) and the formula preceding (2), let  $(a_n)_{n \in \mathbb{N}^*}$  and  $(b_n)_{n \in \mathbb{N}^*}$  be such that  $\lim_{n \rightarrow \infty} a_n = +\infty$  and

$$\limsup_{n \rightarrow \infty} \frac{b_n}{a_n} \geq \liminf_{n \rightarrow \infty} \frac{b_n}{a_n} = \alpha > 1;$$

we are interested in calculating  $\frac{F_\Lambda(b_n) - F_\Lambda(a_n)}{b_n - a_n}$  (see relation (1)).

For every  $x \in \mathbb{R}^+$  with  $\lambda_k \leq x < \lambda_{k+1}$  we have  $F_\Lambda(x) = k$ ; since  $k \leq (\bar{\lambda})^{-1}(x) < k + 1$ , we can write  $F_\Lambda(x) = \lfloor (\bar{\lambda})^{-1}(x) \rfloor$ .

Thus

$$\frac{F_\Lambda(b_n) - F_\Lambda(a_n)}{b_n - a_n} = \frac{\lfloor (\bar{\lambda})^{-1}(b_n) \rfloor - \lfloor (\bar{\lambda})^{-1}(a_n) \rfloor}{b_n - a_n} = \frac{\lfloor y_n \rfloor - \lfloor x_n \rfloor}{\bar{\lambda}(y_n) - \bar{\lambda}(x_n)} = \frac{y_n - x_n}{\bar{\lambda}(y_n) - \bar{\lambda}(x_n)} \cdot \frac{\lfloor y_n \rfloor - \lfloor x_n \rfloor}{y_n - x_n},$$

where

$$x_n = (\bar{\lambda})^{-1}(a_n), \quad y_n = (\bar{\lambda})^{-1}(b_n).$$

Notice that  $x_n \rightarrow +\infty$ . Then (10) follows immediately from (3), Lemma 5.3 (iii) and Lemma 5.2 (see (7)) if we prove that there exists  $\beta > 1$  such that  $y_n > \beta x_n$  for every sufficiently large  $n$ .

Let  $\epsilon > 0$  be fixed with  $\epsilon < \alpha - 1$ . There exists  $n_0$  such that, for  $n > n_0$ ,  $(\alpha - \epsilon)a_n \leq b_n$ , which yields

$$(\bar{\lambda})^{-1}((\alpha - \epsilon)a_n) \leq (\bar{\lambda})^{-1}(b_n).$$

By Lemma 5.3 (ii) and Theorem 1.5.12 in [4],  $(\bar{\lambda})^{-1}$  is regularly varying with exponent 1, hence

$$\frac{(\bar{\lambda})^{-1}((\alpha - \epsilon)a_n)}{(\bar{\lambda})^{-1}(a_n)} \rightarrow \alpha - \epsilon, \quad n \rightarrow \infty.$$

It follows that for every  $\delta > 0$  with  $\delta < \alpha - \epsilon - 1$ , we have

$$(\bar{\lambda})^{-1}(a_n)(\alpha - \epsilon - \delta) \leq (\bar{\lambda})^{-1}((\alpha - \epsilon)a_n) \leq (\bar{\lambda})^{-1}(b_n)$$

or  $y_n \geq \beta x_n$  with  $\beta = \alpha - \epsilon - \delta > 1$ .

Now an application of Theorem 4.1 concludes the proof. □

**Example 5.1.** *Arithmetic progressions satisfy the assumption of Theorem 5.1. A less trivial example is*

$$\lambda_n = \frac{3}{2}n + \frac{3 + (-1)^n}{4};$$

*notice that the values of this sequence are all integers. Hence*

$$b(\Lambda) = d(\Lambda) = \frac{2}{3}.$$

## 6 Some results that follow from the definition of [15]

In this Section we show how to use Definition 2.1 to get information about  $b(\Lambda)$  and its connection with  $d(\Lambda)$ .

**Proposition 6.1.** *Assume that there exist  $\ell > 0$  and a sequence  $(\psi_n)_{n \in \mathbb{N}^*}$  such that*

$$\lambda_n = \ell n + \psi_n.$$

*If either*

*(i)  $(\psi_n)_{n \in \mathbb{N}^*}$  is non-decreasing (ultimately)*

or

$$(ii) \ \kappa := \liminf_{n \rightarrow \infty} \left| \ell + \frac{\psi_n}{n} \right| > 0 \text{ and } \sum_n \frac{|\psi_n|}{n^2} < +\infty,$$

then

$$b(\Lambda) \leq \frac{1}{\ell}. \quad (11)$$

Moreover, in both cases, we have  $\alpha := \liminf_{n \rightarrow \infty} \frac{\psi_n}{n} \geq 0$ . In particular, if  $\alpha = 0$ , then

$$b(\Lambda) = \bar{d}(\Lambda) = \frac{1}{\ell}.$$

*Proof.* We prove that, in both cases (i) and (ii), the number  $\frac{1}{\ell}$  verifies the property stated in Definition 2.1.

(i) Let  $n_k = k + \lfloor \frac{\psi_k}{\ell} \rfloor$ . This sequence of integers is made of distinct integers since

$$n_{k+1} - n_k = \left\lfloor \frac{\psi_{k+1}}{\ell} \right\rfloor - \left\lfloor \frac{\psi_k}{\ell} \right\rfloor + 1 \geq 1.$$

Moreover

$$\begin{aligned} \sum_k \left| \frac{1}{\lambda_k} - \frac{1}{n_k} \right| &= \frac{1}{\ell} \sum_k \left| \frac{1}{k + \frac{\psi_k}{\ell}} - \frac{1}{k + \lfloor \frac{\psi_k}{\ell} \rfloor} \right| = \frac{1}{\ell} \sum_k \frac{|\frac{\psi_k}{\ell} - \lfloor \frac{\psi_k}{\ell} \rfloor|}{(k + \frac{\psi_k}{\ell})(k + \lfloor \frac{\psi_k}{\ell} \rfloor)} \\ &\leq \frac{1}{\ell} \sum_k \frac{1}{k^2 (1 + \frac{\psi_k}{\ell k})(1 + \frac{1}{k} \lfloor \frac{\psi_k}{\ell} \rfloor)}. \end{aligned}$$

Since  $(\psi_n)_{n \in \mathbb{N}^*}$  is non-decreasing, there exists an integer  $k_0$  and a constant  $\alpha$  such that for  $k \geq k_0$ ,

$$1 + \frac{\alpha}{k} > 0, \quad 1 + \frac{\psi_k}{\ell k} \geq 1 + \frac{\alpha}{k}, \quad 1 + \frac{1}{k} \lfloor \frac{\psi_k}{\ell} \rfloor \geq 1 + \frac{\alpha}{k}.$$

Thus the series  $\sum_k \frac{1}{k^2 (1 + \frac{\psi_k}{\ell k})(1 + \frac{1}{k} \lfloor \frac{\psi_k}{\ell} \rfloor)}$  is majorized by

$$\text{const.} + \sum_k \frac{1}{k^2 (1 + \frac{\alpha}{k})^2},$$

which behaves like  $\sum_k \frac{1}{k^2} < +\infty$ .

(ii) Take  $n_k = k$ . We have

$$\sum_k \left| \frac{1}{\lambda_k} - \frac{1}{k} \right| = \frac{1}{\ell^2} \sum_k \frac{|\psi_k|}{k^2 |1 + \frac{\psi_k}{\ell k}|}$$

which, by the assumption  $\kappa > 0$ , behaves like  $\sum_k \frac{|\psi_k|}{k^2} < +\infty$ .

Concerning  $\bar{d}(\Lambda)$ , we have

$$\bar{d}(\Lambda) = \limsup_{n \rightarrow \infty} \frac{n}{\lambda_n} = \limsup_{n \rightarrow \infty} \frac{1}{\ell + \frac{\psi_n}{n}} = \frac{1}{\ell + \limsup_{n \rightarrow \infty} \frac{\psi_n}{n}} = \frac{1}{\ell + \alpha}.$$

The inequality  $\alpha \geq 0$  is obvious in case (i); in case (ii) it follows from the basic inequality (5). If  $\alpha = 0$ , the above discussion implies that

$$b(\Lambda) \leq \frac{1}{\ell} = \bar{d}(\Lambda),$$

and we get the equality once more by (5). □

**Example 6.1.** Any arithmetic progression  $\lambda_n = cn + a$  verifies both the assumptions (i) and (ii) of Proposition 6.1; in particular the set of multiples of the integer  $p$  has Beurling-Malliavin density equal to  $\frac{1}{p}$ .

The sequence of Example 5.1 verifies (ii) of Proposition 6.1.

**Remark 6.1.** There are sequences which satisfy the assumptions of Proposition 6.1 but not those of Theorem 5.1. For instance take

$$\lambda_n = \ell n + \sin n.$$

By Proposition 6.1 (ii)

$$b(\Lambda) = d(\Lambda) = \frac{1}{\ell}$$

( $d(\Lambda)$  exists since  $\lim_{n \rightarrow \infty} \frac{\sin n}{n} = 0$ ). Theorem 5.1 doesn't hold since, for every  $p$ , the sequence  $(\sin(n+p) - \sin n)_{n \in \mathbb{N}}$  doesn't have any limit (see Remark 5.1).

**Proposition 6.2.** Assume that  $\liminf_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n) =: \ell > 0$ . Then

$$b(\Lambda) \leq \frac{1}{\ell}.$$

*Proof.* Recall that  $\Lambda$  is strictly increasing, hence  $\ell \geq 0$ . Now let  $\ell > 0$ . It suffices to prove that

$$b(\Lambda) \leq \frac{1}{\ell - \epsilon}$$

for every  $\epsilon$  with  $0 < \epsilon < \ell$ .

Take  $n_k = \lfloor \frac{\lambda_k}{\ell - \epsilon} \rfloor$ . Then

$$n_{k+1} - n_k = \left\lfloor \frac{\lambda_{k+1}}{\ell - \epsilon} \right\rfloor - \left\lfloor \frac{\lambda_k}{\ell - \epsilon} \right\rfloor \geq \frac{\lambda_{k+1}}{\ell - \epsilon} - 1 - \frac{\lambda_k}{\ell - \epsilon} > 0$$

ultimately, since  $\lambda_{k+1} - \lambda_k > \ell - \epsilon$  ultimately. Hence the sequence  $(n_k)_{k \in \mathbb{N}^*}$  is made of distinct integers (ultimately).

Moreover

$$\sum_k \left| \frac{1}{\lambda_k} - \frac{1}{n_k} \right| = \sum_k \left| \frac{1}{\lambda_k} - \frac{1}{\lfloor \frac{\lambda_k}{\ell - \epsilon} \rfloor} \right| = \sum_k \frac{|\lfloor \frac{\lambda_k}{\ell - \epsilon} \rfloor - \frac{\lambda_k}{\ell - \epsilon}|}{\lambda_k \lfloor \frac{\lambda_k}{\ell - \epsilon} \rfloor} \leq \sum_k \frac{1}{\lambda_k \lfloor \frac{\lambda_k}{\ell - \epsilon} \rfloor},$$

and the last series has the same behaviour as  $\sum_k \frac{1}{\lambda_k^2}$  (since  $\lambda_k \rightarrow \infty$ ). It remains to prove that

$$\sum_k \frac{1}{\lambda_k^2} < +\infty.$$

Putting  $\psi_n = \lambda_{n+1} - \lambda_n$ , it is easy to see by induction that

$$\lambda_n = \lambda_1 + \sum_{k=1}^{n-1} \psi_k.$$

As stated above, there exists  $n_0$  such that  $\psi_n > \ell - \epsilon$  for  $n > n_0$ . Hence, for  $n > n_0 + 1$ , we have

$$\lambda_n = \lambda_1 + \sum_{k=1}^{n_0} \psi_k + \sum_{k=n_0+1}^{n-1} \psi_k > \lambda_1 + \sum_{k=1}^{n_0} \psi_k + (\ell - \epsilon)(n - 1 - n_0) = (\ell - \epsilon)n + \text{const.},$$

which implies that  $\sum_k \frac{1}{\lambda_k^2}$  is  $\leq \text{const} \cdot \sum_k \frac{1}{k^2} < +\infty$ . □

The preceding Proposition and Theorem 4.1 yield immediately the following result, which is a particular case of (actually it has been a motivation for) Theorem 5.1:

**Corollary 6.1.** *If the limit*

$$\ell := \lim_{n \rightarrow \infty} (\lambda_{n+1} - \lambda_n)$$

*exists (as an extended number) and is strictly positive, then*

$$d(\Lambda) = b(\Lambda) = \frac{1}{\ell},$$

where we adopt the convention  $\frac{1}{0} = +\infty$ .

**Remark 6.2.** *For the sequence  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  with  $\lambda_n = \log n$  we have  $\ell = 0$  and as a consequence  $b(\Lambda) = +\infty$ . We conjecture that the same happens for  $M = (\mu_n)_{n \in \mathbb{N}^*}$  with  $\mu_n = \lfloor \log n \rfloor$ , as well as for sequences with “too many” repetitions; but we have no proof at present for this.*

**Proposition 6.3.** *Assume that  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  is a sequence such that*

$$\sum_k \frac{1}{\lambda_k} < +\infty.$$

*Then  $d(\Lambda) = b(\Lambda) = 0$ .*

**Remark 6.3.** *This Proposition was proved in [16], but only in the part concerning  $d(\Lambda)$ .*

*Proof.* Recall Olivier’s Theorem (also known as as Abel’s or Pringsheim’s Theorem), proved in [11]:

**Theorem 6.1.** *Let  $(a_n)_{n \in \mathbb{N}^*}$  be a non-increasing sequence of positive numbers such that the corresponding series  $\sum_{n=1}^{\infty} a_n$  is convergent. Then  $\lim_{n \rightarrow \infty} n a_n = 0$ .*

Applying it with  $a_n = \frac{1}{\lambda_n}$  we have

$$d(\Lambda) = \lim_{n \rightarrow \infty} \frac{n}{\lambda_n} = 0.$$

Moreover, by Definition 2.1,  $b(\Lambda) = 0$ . □

**Example 6.2.** *This result applies for instance to the set of powers  $(k^p)_{k \in \mathbb{N}^*}$  ( $p \in \mathbb{N}^*$ ,  $p \geq 2$ ). More generally, it applies to any increasing sequence of the form  $(L(k)k^p)_{k \in \mathbb{N}^*}$ , where  $(L_k)_{k \in \mathbb{N}^*}$  is a slowly varying sequence: recall that a slowly varying sequence  $(L_k)_{k \in \mathbb{N}^*}$  has the property that, for any  $\epsilon > 0$ , ultimately we have  $k^{-\epsilon} < L_k$  (this is an easy consequence of Theorem 1.3.1 in [4]), which implies that  $\sum_k \frac{1}{\lambda_k} < \sum_k \frac{1}{k^{p-\epsilon}} < +\infty$  if  $\epsilon < p - 1$ .*

Last, here below is the announced characterization of the BM-density:

**Theorem 6.2.** *Let  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  be a sequence with  $b(\Lambda) \in (0, +\infty)$ ; assume that there exist a strictly positive number  $\ell$  and another sequence  $M = (\mu_n)_{n \in \mathbb{N}^*}$  with  $b(M) \in (0, +\infty)$  such that*

$$\sum_k \left| \frac{1}{\lambda_k} - \frac{1}{\ell \mu_k} \right| < +\infty.$$

*Then  $\ell$  is unique; furthermore*

$$b(M) = \ell b(\Lambda). \tag{12}$$

*Proof.* Denote by

$$\mathcal{A} = \left\{ \ell > 0 : \sum_k \left| \frac{1}{\lambda_k} - \frac{1}{\ell \mu_k} \right| < +\infty \right\}$$

and

$$\sigma = \inf \mathcal{A}, \quad \tau = \sup \mathcal{A}.$$

Let  $\epsilon > 0$  and  $N = (n_k)_{k \in \mathbb{N}^*}$  a sequence of distinct integers such that

$$\sum_k \left| \frac{1}{\mu_k} - \frac{b(M) + \epsilon}{n_k} \right| < +\infty. \quad (13)$$

Then, for every  $\ell \in \mathcal{A}$ ,

$$\sum_k \left| \frac{1}{\lambda_k} - \frac{b(M) + \epsilon}{\ell n_k} \right| \leq \sum_k \left| \frac{1}{\lambda_k} - \frac{1}{\ell \mu_k} \right| + \frac{1}{\ell} \sum_k \left| \frac{1}{\mu_k} - \frac{b(M) + \epsilon}{n_k} \right| < +\infty.$$

This yields

$$b(\Lambda) \leq \frac{b(M) + \epsilon}{\ell}$$

and by optimizing in  $\ell$  and  $\epsilon$  we conclude that

$$b(\Lambda) \leq \frac{b(M)}{\tau}. \quad (14)$$

Relation (14), together with the assumption  $b(\Lambda) > 0$ , implies that  $\tau < \infty$ .

Now interchange the roles of  $\Lambda$  and  $M$ ; we obtain that

$$b(M) \leq \ell(b(\Lambda) + \epsilon),$$

and optimizing

$$b(M) \leq \sigma b(\Lambda). \quad (15)$$

From relation (15) and the assumption  $b(M) > 0$  we deduce that  $\sigma > 0$ ; and now, putting together (14) and (15), we get

$$b(\Lambda) \leq \frac{b(M)}{\tau} \leq \frac{\sigma}{\tau} b(\Lambda);$$

this proves the unicity of  $\ell$  and the relation (12). □

We put in evidence the following particular case (see also Proposition 6.3 for the case  $\ell = 0$ ):

**Corollary 6.2.** *Let  $\Lambda = (\lambda_n)_{n \in \mathbb{N}^*}$  be a sequence with  $b(\Lambda) < +\infty$ ; if there exists a positive number  $\ell$  such that*

$$\sum_k \left| \frac{1}{\lambda_k} - \frac{\ell}{k} \right| < \infty.$$

*then  $\ell = b(\Lambda)$ .*

## References

- [1] Baranov, A., Belov, Y., Ulanovskii, A., (2017), Gap Problem for Separated Sequences without Pain *J. Fourier Anal. Appl.* 23, 877–885, DOI:10.1007/s00041-016-9495-5.
- [2] A. Beurling and P. Malliavin, (1967), On the closure of characters and the zeros of entire functions, *Acta Math.*, 118, 79–93.
- [3] Wiener, N., Paley, R. C., (1934), *Fourier Transforms in the Complex Domain* Am. Math. Soc. Colloquium Publications, Vol. 19.
- [4] Bingham, N. H., Goldie, C. M., Teugels, J. L., (1989), *Regular Variation*, Cambridge University Press.
- [5] Fuchs, A., Giuliano Antonini, R., (1990), Théorie générale des densités. *Rend. Accad. Naz. Sci. XL Mem. Mat.* (5) 14 , no. 1, 253–294.
- [6] Giuliano, R., Grekos, G., Mišik, L., (2023), On the Beurling–Malliavin density and its connection with Pólya density, to appear.
- [7] Khabibullin, B.N., (1995), Nonconstructive Proofs of the Beurling–Malliavin Theorem on the Radius of Completeness, and Nonuniqueness Theorems for Entire Functions *Izv. Math.* 45, 125–149.
- [8] Kahane, J.-P., (1962), Travaux de Beurling et Malliavin, *Séminaire Bourbaki : année 1961/62, exposés 223-240, Séminaire Bourbaki*, 7, Talk no. 225, 13 p.
- [9] P. Koosis, (1992) *The Logarithmic Integral II*, Cambridge University Press.
- [10] Krasichkov-Ternovskii, I. F., (1990), An interpretation of the Beurling–Malliavin theory on the radius of completeness, *Math. USSR Sb.*, 66, 405–429.
- [11] Olivier, L., (1827) Remarques sur les séries infinies et leur convergence. *J. reine angew. Math.*, 2:31–44.
- [12] Poltoratski, A., (2012), Spectral gaps for sets and measures, *Acta Math.* 208, 1, 151–209.
- [13] Poltoratski, A., (2018), Completeness of exponentials. Beurling–Malliavin and Type Problems, <https://eta.impa.br/icm-files/invited/section-8/IL.8.18.pdf>
- [14] Poltoratski, A., (2018), Toeplitz Methods in Completeness and Spectral Problems, *Proc. Int. Cong. of Math. – 2018* Rio de Janeiro, Vol. 3 (1771–1796).
- [15] Redheffer, R., (1972), Two consequences of the Beurling–Malliavin Theory, *Proc. Am. Math. Soc.*, 36, 1, 116–122.
- [16] Salát, T., (1964), On subseries, *Math. Z.*, 85, 209–225.