



HAL
open science

Effects of one session of theta or high alpha neurofeedback on EEG activity and working memory

Samy Chikhi, Nadine Matton, Marie Sanna, Sophie Blanchet

► To cite this version:

Samy Chikhi, Nadine Matton, Marie Sanna, Sophie Blanchet. Effects of one session of theta or high alpha neurofeedback on EEG activity and working memory. *Cognitive, Affective, and Behavioral Neuroscience*, 2024, 10.3758/s13415-024-01218-4. hal-04742953

HAL Id: hal-04742953

<https://hal.science/hal-04742953v1>

Submitted on 18 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Effects of one session of theta or high alpha neurofeedback on EEG activity and working
memory**

Samy Chikhi^{1,4*}, Nadine Matton^{2,3}, Marie Sanna¹, Sophie Blanchet¹

¹ Université Paris Cité, Laboratoire Mémoire, Cerveau et Cognition, F-92100 Boulogne-
Billancourt, France.

² CLLE - Cognition, Langues, Langage, Ergonomie, Université de Toulouse, Toulouse, France.

³ Fédération ENAC ISAE-SUPAERO ONERA, Université de Toulouse, France.

⁴ Integrative Neuroscience and Cognition Center, Université Paris Cité, F-75006, Paris, France.

Author Note

Correspondence concerning this article should be addressed to Samy Chikhi, Integrative Neuroscience and Cognition Center, UMR 8002, Université Paris Cité, 45 rue des Saints-Pères, 75006 Paris, France. Email: sychikhi@gmail.com

Abstract

Neurofeedback techniques provide participants immediate feedback on neuronal signals, enabling them to modulate their brain activity. This technique holds promise in unveiling brain-behavior relationship, and offers opportunities for neuroenhancement. Establishing causal relationships between modulated brain activity and behavioral improvements requires rigorous experimental designs, including appropriate control groups and large samples. Our primary objective was to examine whether a single neurofeedback session, designed to enhance working memory through the modulation of theta or high-alpha frequencies, elicits specific changes in electrophysiological and cognitive outcomes. Additionally, we explored predictors of successful neuromodulation. One hundred and one healthy adults were assigned to groups trained to increase frontal theta, parietal high alpha, or random frequencies (active control group). We measured resting-state EEG, working memory performance, and self-reported psychological states before and after one neurofeedback session. Although our analyses revealed improvements in electrophysiological and behavioral outcomes, these gains were not specific to the experimental groups. An increase in the frequency targeted by the training has been observed for the theta and high alpha groups, but training aimed at increasing randomly selected frequencies appears to induce more generalized neuromodulation compared to targeting a specific frequency. Among all the predictors of neuromodulation examined, resting theta and high alpha amplitudes predicted specifically the increase of those frequencies during the training. These results highlight the challenge of integrating a control group based on enhancing randomly selected frequency bands and suggest potential avenues for optimizing interventions (e.g., by including a control group trained in both up- and down-regulation).

Keywords: neurofeedback, working memory, theta, high alpha, active control group

Introduction

The brain consists of numerous neurons forming intricate networks. Within these networks, information transmission occurs through rhythmic electrical activity known as neural oscillations (Buzsaki & Draguhn, 2004; Engel et al., 2001; Weisz & Keil, 2022). Neural oscillations play a crucial role in various cognitive processes (Başar et al., 2001), including sensory processing (Haegens & Golumbic, 2018), attention (Clayton et al., 2015) and working memory (Chikhi et al., 2022; Pavlov & Kotchoubey, 2022). Given the foundational role of neural oscillations in cognitive processes, modulating specific brain activity parameters holds promise for improving cognition (Chiasson et al., 2023; Gruzelier, 2014; Viviani & Vallesi, 2021) and elucidating the causal relationship between these parameters and cognitive processes (Kvamme et al., 2022a; Ramot & Martin, 2022).

Neurofeedback is a brain-computer interface that trains participants to voluntarily modulate a specific parameter of their brain activity through real-time feedback on their activity (Enriquez-Geppert et al., 2017; Sitaram et al., 2017). Participants engage in a closed loop where their mental actions (i.e., changing mental states) modify the feedback signal, subsequently influencing their forthcoming mental actions (Birbaumer et al., 2013; Lubianiker et al., 2022). Through repetition and trial-and-error learning, positive feedbacks that reward the mental state corresponding to the targeted neuronal state (e.g., increase of theta frequency above a certain threshold) will promote the reoccurrence of this mental state (Birbaumer et al., 2013; Gaume et al., 2016; Sherlin et al., 2011). Repeated production of the rewarded neuronal state may lead to neuroplasticity, strengthening synaptic connections within the underlying neural network (Davelaar, 2018; Shibata et al., 2019). Thus, the neurofeedback technique could enable individuals to learn to self-regulate their brain activity autonomously, without the aid of an external device (e.g., neurostimulation techniques; Herrmann et al., 2016). This can be achieved not only within controlled laboratory or clinical settings but also extended to real-world environments (Bassett & Khambhati, 2017; Herrmann et al., 2016).

In the field of EEG-neurofeedback, numerous studies have sought to increase the amplitude of two specific brain frequencies to enhance cognition: theta and alpha frequencies (Pfeiffer et al., 2024; Yeh et al., 2021; Yeh et al., 2022). The frontal theta frequency (4-8 Hz), intricately linked with cognitive control (Cavanagh, & Frank, 2014; Senoussi et al., 2022), has been targeted to improve attention (Brandmeyer & Delorme, 2020; Kerick et al., 2023), episodic memory (Eschmann et al., 2020; Tseng et al., 2021), autobiographic memory (Shoji et al., 2017), memory consolidation (Reiner et al., 2014; Rozengurt et al., 2016; Rozengurt et al., 2017; Shtoots et al., 2021), motor performance (Eschmann et al., 2022), working memory (Li et al., 2019; Reis et al., 2016; Wang & Hsieh, 2013) and executive functions (Enriquez-Geppert et al., 2014; Eschmann & Mecklinger, 2022; Smit et al., 2023). The posterior alpha frequency (8-12 Hz) serves as a cortical inhibitor, suppressing irrelevant information (Klimesch et al., 2007; Jensen & Mazaheri, 2010; Van Diepen et al., 2019) and contributing to working memory processes (Chen et al., 2023; Riddle et al., 2020). This frequency, and in particular the higher range of this frequency (referred to as high alpha, 10-12 Hz), has been a target of neurofeedback training aiming to enhance mental rotation (Hanslmayr et al., 2005; Zoefel et al., 2011), episodic memory (Guez et al., 2015), attention (Navarro Gil et al., 2018), executive control (Nawaz et al., 2022) and working memory processes (Bobby, & Prakash, 2017; Chen & Sui, 2023; Dehghanpour et al., 2018; Escolano et al., 2012; Escolano et al., 2014; Esteves et al., 2021; Hsueh et al., 2016; Naas et al., 2019; Shen et al., 2023; Takabatake et al., 2021; Wei et al., 2017). While the training duration varies across studies (from one to twenty), some research has indicated successful neuromodulation of the trained frequencies within a single session (Eschmann et al., 2022; Escolano et al., 2012; Escolano et al., 2014; Reiner et al., 2014; Rozengurt et al., 2016; Rozengurt et al., 2017; Shtoots et al., 2021). Although single neurofeedback sessions may not suffice for achieving mastery over one's brain activity, they do induce notable modifications in cortical excitability (Ros et al., 2010) as well as neuroplastic changes in structural and functional connectivity (Marins et al., 2019; Ros et al., 2013;

Sampaio-Baptista et al., 2021). These observations suggest that brief neurofeedback training can transiently affect brain activity and behavioral outcomes.

These effects are particularly interesting given the existing dilemma within the field of cognitive training. Indeed, a choice often has to be made between a large sample size or conducting a larger number of training sessions, due to limited resources. The field of neurofeedback is not exempt from these constraints, and numerous protocols suffer from insufficient statistical power. This often results in an inflation of effect sizes and low reproducibility of results (Button et al., 2013; Szucs & Ioannidis, 2017; Thibault & Pedder, 2022). In fact, a recent systematic review highlighted that only 4% of neurofeedback studies had sample sizes exceeding 50 participants, with an average number of participants per group of $n = 16.64$ (Chiasson et al., 2023). Hence, it is crucial for the neurofeedback field to rely on larger samples to rigorously evaluate the effects of this technique. Additional concerns arise regarding the potential influence of non-specific factors on neurofeedback outcomes. Non-specific factors encompass various extraneous variables that are not directly related to the neurofeedback protocol but can significantly contribute to the observed effects (Ros et al., 2020). Such factors include expectancy effects, placebo effects, and unmeasured confounding variables (e.g., La Marca et al., 2018; Thibault et al., 2017). Inadequate control of these non-specific factors may compromise the validity and interpretability of neurofeedback research findings (Thibault et al., 2016). Lastly, the psychological, cognitive, and electrophysiological variables that determine the ability to modulate brain activity remain insufficiently identified (Haugg et al. 2021; Jeunet et al., 2018; Kadosh & Staunton, 2019; Weber et al., 2020). Identifying these variables would optimize participant recruitment, mitigate inefficiency issues (Alkoby et al., 2018), and enable the conduct of more powerful studies without increasing sample size (Thibault & Pedder, 2022).

This study aimed to investigate the specific effects of a single neurofeedback session on brain frequencies amplitude and working memory. We compared the performance of a group that was trained

to increase the amplitude of frontal theta frequency with a group trained to increase the amplitude of posterior high alpha frequency during a single neurofeedback session. Theta oscillations facilitate cognitive control and item organization in working memory (e.g., Cooper et al., 2019; Deiber et al., 2007; Hanslmayr et al., 2019; Hsieh et al., 2011; Itthipuripat et al., 2013; Pesonen et al., 2007; Riddle et al., 2020; Roux & Uhlhaas, 2014; Sauseng et al., 2010, 2009, 2005), while alpha oscillations may inhibit task-irrelevant information (e.g., de Vries et al., 2020; Hsu & Hämäläinen, 2022; Pesonen et al., 2007; Sghirripa et al., 2021). We hypothesize that neurofeedback-based amplitude modulation of these frequencies will improve performance on working memory tasks. To better distinguish between the specific and non-specific effects of the neurofeedback, we included an active control group that was trained to modulate randomly selected frequencies. By selecting random frequency bands, participants engage in a genuine neurofeedback task that targets their own neurophysiological signals. However, because they cannot learn to modulate a specific frequency, this approach minimizes potential interference with their cognitive processes. Additionally, unlike sham control groups (Kvamme et al., 2022b), providing real EEG feedback reduces the risk of inducing a sense of learned helplessness in participants (Aliño et al., 2016). We assessed brain frequency amplitude, performance on working memory tasks (including digit span, *n*-Back, and Corsi blocks), as well as self-reported psychological states related to emotional and attentional aspects before and after the training period. We also aimed to identify factors (psychological, cognitive or electrophysiological) that could predict neuromodulation success during the neurofeedback task.

Method

Sample

G-Power (Faul et al., 2007) was utilized to determine the sample size needed for an analysis of variance with repeated measures focusing on within-between interactions. For the investigation of the impact of training on working memory, where a comparison was made between pre- and post-training performances, we computed the required sample size based on the following parameters: effect size $f =$

.20, power level = .90, number of groups = 3, number of measures = 2, correlation between repeated measures = .50. An estimated total sample size of 84 participants was obtained. 70 participants were randomly assigned to either the Theta group, aiming to increase the amplitude of the theta frequency in the frontal cortex, or to the Control group, aiming to increase randomly selected frequencies within each block. Data from some participants had to be excluded ($N = 8$, 11%) due to low-quality EEG data (i.e., more than 20% of the signal was removed after artifact rejection, $N = 7$) and one participant was too fatigued to complete the entire protocol and chose to discontinue the experiment. The data from the group trained to increase parietal high alpha amplitude were obtained through a prior identical experimental protocol where the research question aimed to investigate the effects of a strategy list on neuromodulation (Chikhi et al., 2023). A total of 101 participants were included in our analyses. The Theta group consisted of 28 participants (Mean age = 19.4 ± 1.81 , 75% woman), the Control group comprised 34 participants (Mean age = 21.41 ± 8.28 , 91% woman) and the High alpha group consisted of 39 participants (Mean age = 23.28 ± 5.96 , 82% woman). The inclusion criteria were normal or corrected-to-normal vision, no psychiatric or neurological disorders, and no usage of medication. All participants were naïve to the neurofeedback technique.

Neurofeedback task and procedure


Each participant underwent a single experimental session lasting approximately 120 minutes (Figure 1) and received course credits for their participation. Prior to and after neurofeedback session, participants completed computerized tasks assessing working memory (digit span, spatial span and n -back task) from the Psychology Experiment Building Language (Mueller & Piper, 2014). Additionally, participants completed a shortened version of the Neurofeedback Evaluation & Training questionnaire (NExT questionnaire; Bismuth et al., 2020; Jaumard-Hakoun et al., 2017; Pillette et al., 2021) to evaluate their subjective state during the neurofeedback protocol. Neurofeedback training began by measuring the dimensions of the participants' heads to determine electrode locations. Next, conductive gel and paste

were applied to prepare the skin and improve signal quality at each electrode location. Once the correct impedance was achieved at each electrode, a baseline was measured by recording the resting-state EEG signal during 3 minutes. Participants were instructed to maintain their gaze on a black screen positioned in front of them. They were required to remain relaxed with their eyes open, refraining from blinking, making any bodily movements, or clenching their jaw.

After recording the baseline, the average amplitude of theta and high alpha frequencies were extracted to determine the threshold for activating positive feedback. In the Control group, participants received feedback based on a threshold that automatically adapted to the average amplitude of the targeted brain signal. The frequency bands used for the participants in the Control group were randomly generated 2 Hz bands for each block and each participant. The randomly selected frequencies included a range of 2Hz, situated between 1 and 30 Hz, except for the frequency bands targeted by the experimental groups (4-8Hz and 10-12Hz). Participants were always required to increase the amplitude of the selected frequency band. Irrespective of the group, all participants were informed that the gauge shown on the left side of the screen depicted the real-time amplitude of a specific brain frequency. When the gauge exceeded the threshold (as denoted by an orange bar), a screen animation (depicting a roller coaster) was triggered, accompanied by enjoyable music. The participants' goal was to 'keep the level of the gauge above the threshold for as long and as frequently as possible'. All participants were informed that the audiovisual feedback would convey information about their brain activity and aid in achieving a mental state that triggers the reward feedback. They were not provided with particular mental strategies for regulating their brain activity. Following the completion of the 10 training blocks, the resting-state EEG signal was once again recorded. Subsequently, the electrodes were removed, the participants' scalps were cleaned, and they proceeded to complete the NExT questionnaire and computerized working memory tasks once more.

Figure 1

Experimental protocol overview



Note. Preceding and subsequent to the neurofeedback session, they engaged in computerized tasks designed to evaluate their working memory (Mueller & Piper, 2014). Additionally, they responded to a concise questionnaire aimed at assessing their psychological state (Bismuth et al., 2020; Jaumard-Hakoun et al., 2017). The neurofeedback protocol encompassed measuring a 3-minute resting-state EEG before

and after the 10 neurofeedback training blocks. For the Theta and High alpha groups, the positive feedback activation threshold was determined based on the mean amplitude of theta and high alpha frequencies, respectively. Both groups were provided real-time feedback, presented as a gauge positioned on the left side of the screen. Exceeding the established threshold led to the activation of animation and music. The primary objective was to sustain the gauge level above this threshold. Following the 10 training blocks and the second resting-state EEG, participants undertook post-session working memory tasks and questionnaire.

EEG acquisition and processing

An encoder (ProComp T740M) was connected to a laptop via a fiber optic cable and USB interface for recording the EEG signal. The BioGraph Infiniti software (version 6.1, Thought Technology Ltd., Montreal, QC, Canada) was used to analyze the signal and convert the targeted brain activity into audio-visual feedback. The High alpha group was trained using an electrode positioned at Pz. For the Theta group, the electrode was positioned at Fz. In the Control group, the electrode was placed at the vertex (Cz), which is equidistant from Fz and Pz (Pimenta et al., 2018). Additionally, a reference electrode and a ground electrode were positioned on the right and left earlobes, respectively. The signal was sampled at 256 Hz, filtered using a bandpass filter (0.1 to 60 Hz), and a notch filter (50 Hz) to eliminate electrical interference. The software extracted the amplitude of the signal trained for each group (theta, high alpha, or random frequency) and converted it into audio-visual feedback. The main brain frequencies were recorded for offline analysis: delta (1-4 Hz), theta (4-8 Hz), low alpha (8-10 Hz), high alpha (10-12 Hz), beta (12-35 Hz), and gamma (35-64 Hz) using fast Fourier transform (FFT). Nuprep Skin Prep Gel (Weaver and Company, Aurora, USA) was used to clean the skin prior to electrode placement. Electrodes were fixed with Ten20 Conductive Paste (Weaver and Company, Aurora, USA). Impedance was assessed using an EEG-Z sensor and kept below 10 k Ω . An automatic artifact rejection threshold was set at 100 μ V before visually inspecting the raw data and manually rejecting artifacts. Additional pre-processing and artifact

correction steps could not be performed due to the software used. The feedback was presented as an animation illustrating a roller coaster, accompanied by calming music (Wang & Hsieh, 2013).

For the groups trained to increase the theta or high alpha amplitude, positive feedback (animation and music) was activated when the average amplitude of the trained frequency exceeded 80% of the baseline's average value. For the Control group, a fixed threshold could not be applied because the trained frequency varied between each block. Therefore, we employed an adaptive threshold based on the amplitude of the selected frequency. This threshold corresponded to +1% of the average signal amplitude, with a delay of 0.2 seconds. This adaptive positive threshold ensures that participants receive feedback that is contingent upon their own neural activity, reducing the risk of frustration and learned helplessness (Sorger et al., 2019).

Working memory tasks

To assess different facets of participants' working memory, three distinct working memory tasks were administered via the Psychology Experiment Building Language (Mueller & Piper, 2014). The tasks included the backward digit span (Wechsler, 1955), the *n*-back task with 1, 2, and 3-back conditions (Owen et al., 2005), and the backward spatial span task, also known as Corsi's blocks task (Corsi, 1972). The backward digit span task required participants to recall a series of randomly presented digits in reverse order (Richardson, 2007). To complete this task, participants utilized the number pad of a standard keyboard. Throughout the task, the length of the digit sequence gradually increased. The dependent variable was defined as the longest successfully recalled sequence. The *n*-back task involves presenting participants with a series of stimuli (e.g., a letter), presented one after another in a sequential manner. The task required participants to identify if the current stimulus matches the one presented '*n*' position earlier in the sequence. For example, during the 2-back condition, if the letter sequence displayed on the screen was F-N-B-N-B-F, participants were instructed to press the 'yes' key when the second 'N' and the second 'B' appeared. They had to press the 'no' key for all other letters. The *n*-back task included three

conditions: 1-back, 2-back, and 3-back. The dependent variable for each condition was the percentage of correct responses. The computerized Corsi block task comprised nine fixed squares displayed on a computer screen in a pseudo-random arrangement. These squares change color individually over a short period. In the backward condition, participants were instructed to click on the squares in the reverse order of their original sequence. As the experiment advanced, the length of the block sequence progressively increased. The dependent variable was the longest accurately recalled sequence's length (Mueller & Piper, 2014). The digit span backward task was used to engage the executive control component of working memory (Lezak et al., 2004). The Corsi blocks backward task was employed to involve visuospatial working memory processing (Kessels et al., 2008). The *n*-back task involves simultaneously retaining previously presented stimuli, processing the currently presented stimulus, and updating the contents of working memory (Lezak et al., 2004; Soveri et al., 2017). By using tasks that target various working memory processes, our aim was to distinguish the cognitive effects of theta and high alpha amplitude modulation.

Self-reported questionnaire

To effectively assess the psychological state of participants before, during, and after neurofeedback, while minimizing the burden of the experimental procedure and cognitive load, we utilized a condensed version of the NEXt questionnaire (Bismuth et al., 2020; Jaumard-Hakoun et al., 2017; Pillette et al., 2021). The questionnaire items were adapted from established and validated questionnaires (see Supplementary material in Chikhi et al., 2022 for item sources). Each participant was asked to rate their experiences using a 5-point Likert scale. The questionnaire covered five distinct dimensions: emotional state ('calm,' 'energetic,' 'happy,' 'relaxed,' and 'satisfied'), attentional state ('receptive' and 'focused'), cognitive load, motivation, and agency ('sense of control over the feedback signal' and 'predictability of the feedback signal'). The questionnaire was administered on two occasions. Firstly, it only consisted of the first two dimensions, assessing emotional and attentional state just before the neurofeedback session. This measurement aimed to evaluate the effect of self-reported psychological

states before training on neuromodulation ability. Secondly, the questionnaire was administered immediately after the neurofeedback session and included all five dimensions. This second administration aimed to assess the impact of neurofeedback on participants' psychological state changes (via the first two dimensions). The remaining three dimensions were used to compare participants' psychological state during the neurofeedback task, including cognitive load, motivation, and sense of agency. This enabled an investigation into the potential influence of these variables on the participants' success in the neurofeedback task. For example, participants in the control group might exhibit a reduced sense of agency towards the neurofeedback device, indicating a weaker perception of control over the feedback signal, in contrast to the experimental groups. Such a disparity could potentially affect their motivation and consequently lead to a decline in performance both during the neurofeedback task and in the subsequent behavioral assessments.

Statistical analysis

Statistical analyses were performed with the RStudio environment (v.4.0.2; R Core Team, 2020) with the following packages: lme4 (v.1.1-27.1; Bates et al., 2015) and lmer (v. 3.1-3; Kuznetsova et al., 2017) for classical and mixed linear regression, multcomp (v. 1.4-17; Hothorn et al., 2008) and modelbased (v. 0.8.0; Makowski et al., 2020a) for post hoc contrasts, and report (v. 0.5.1; Makowski et al., 2020b), ggplot2 (v. 3.3.6; Wickham, 2016) and smplot2 (v. 0.1.0; Min & Zhou, 2021) for formatting and visualization of results. For repeated measures, mixed-effects linear models were estimated using the Restricted Maximum Likelihood (REML) method and fitted with a random intercept (1| Participant). Classical linear regression models were estimated using Ordinary Least Squares (OLS) method. P-values for Type III ANOVA F-tests were obtained using Kenward-Roger approximation for degrees of freedom (Kenward & Roger, 1997; Luke, 2017). In cases where an effect was statistically significant, we applied Tukey's post hoc contrasts with Holm correction for multiple comparisons (Schad et al., 2020). Only significant differences from post hoc comparisons were detailed. The significance threshold for all

analyses was set at $\alpha = 0.05$. Effect sizes were interpreted as follows: $\eta_p^2 = 0.01$ as a small effect, $\eta_p^2 = 0.06$ as a moderate effect, and $\eta_p^2 = 0.14$ as a large effect.

Results

Effects of neurofeedback on targeted brain signal (Figure 2)

Effect of neurofeedback on theta amplitude during training

A mixed linear model was employed to predict the amplitude of theta frequency with Group (Theta, High alpha, Control) and Training Block (1 to 10) as fixed effects, and Participant as a random effect. The ANOVA revealed a significant and strong main effect of Group ($F(2) = 14.14, p < .001; \eta_p^2 = 0.22, 95\% CI [0.11 - 1.00]$). Post hoc analysis indicated that the average theta frequency amplitude was significantly lower in the High alpha group compared to the Theta group ($\beta = -1.37, SE = 0.44, p = .004$). The difference between the Theta group and the Control group was not statistically significant. The main effect of Training Block was also statistically significant and weak ($F(9) = 2.10, p = 0.027; \eta_p^2 = 0.02, 95\% CI [0.001 - 1.00]$), however, post hoc comparisons revealed no significant differences in amplitude between the blocks. The interaction between Group and Training Block was not statistically significant ($p = .053$).

The absence of global interaction among the three groups and the training blocks might mask divergent effects depending on the pair of groups under consideration. Thus, we conducted two complementary analyses investigating the Group x Training Block interaction, specifically comparing (1) Theta versus High alpha group and (2) Theta versus Control group. In the first mixed linear model, we predicted the amplitude of theta frequency with Group (Theta x High alpha) and Training Block as fixed effects, and Participant as a random effect. The ANOVA revealed a significant interaction between Group and Training Block ($F(9) = 2.43, p = .010; \eta_p^2 = 0.04, 95\% CI [0.005, 1.00]$), and subsequent post hoc analysis revealed that the average theta frequency amplitude was significantly lower in the High Alpha group compared to the Theta group across blocks 3 to 10 ($SE = 0.44, all p < .01$). The second mixed linear model, predicting the amplitude of theta frequency with Group (Theta x Control) and Training Block as fixed

effects, and Participant as a random effect, indicated that the interaction between Group and Training Block was not statistically significant ($F(9) = 1.17, p = .313; \eta_p^2 = 0.02, 95\% CI [0.00, 1.00]$).

Effect of neurofeedback on theta amplitude during resting-state

A mixed linear model was performed to predict the amplitude of theta frequency recorded during resting-state EEG, considering Group and Resting-EEG (pre- and post-training EEG recording) as fixed effects and Participant as a random effect. The ANOVA revealed a statistically significant and moderate main effect of Group ($F(2) = 5.64, p = .005; \eta_p^2 = 0.10, 95\% CI [0.02 - 1.00]$). Post hoc analysis indicated that the average theta amplitude was significantly lower in the High alpha group compared to the Control group ($\beta = -1.67, ES = .60, p = .02$), while no significant difference was observed between the Theta group and the Control group ($p = .52$) nor between the Theta group and the High alpha group ($p = .09$). Additionally, the main effect of Resting-EEG was statistically significant and weak ($F(1) = 4.82, p = 0.03; \eta_p^2 = 0.05, 95\% CI [0.002 - 1.00]$), with a slightly higher mean amplitude observed after training ($M = 10.13 \pm 2.86$) compared to before ($M = 9.94 \pm 2.42$). However, neither the overall interaction between the Group and Resting EEG reached significance ($p = .11$), nor did the pairwise comparisons (Theta x High Alpha, $p = .42$; Theta x Control, $p = .24$). To assess the impact of resting EEG activity on the neurofeedback task, we applied a classical linear model to compare the average theta amplitude activity during the first resting-state among the groups. The ANOVA indicated that the main effect of Group was statistically significant and medium ($F(2, 98) = 5.11, p = .008; \eta_p^2 = 0.09, 95\% CI [0.02, 1.00]$). Post-hoc comparisons revealed that the mean theta amplitude was significantly lower in the High alpha group compared to the Control group ($\beta = 1.67, p < .01$). No significant differences were observed between the Theta group and the Control group ($p = .48$) or between the Theta group and the High alpha group ($p = .06$).

Effect of neurofeedback on high alpha amplitude during training

A mixed linear model was performed to predict the amplitude of high alpha frequency with Group and Training Block as fixed effects, and Participant as a random effect. The ANOVA revealed that the main

effect of Training Block was statistically significant yet weak ($F(9) = 5.60$, $p < .001$; $\eta_p^2 = 0.05$, 95% CI [0.02, 1.00]). Post hoc analysis revealed no significant differences in amplitude between the blocks. Moreover, neither the main effect of Group nor the interaction between Group and Training Block reached statistical significance ($p = .21$ and $p = .08$, respectively).

Again, the lack of global interaction between the three groups and the training blocks may mask differential effects depending on the pair of groups considered. We conducted two complementary analyses testing the Group x Block interaction when specifically comparing (1) Theta vs. High alpha group and (2) Theta vs. Control group. Using a mixed linear model, we predicted the amplitude of high alpha frequency with Group (High alpha and Theta) and Training Block as fixed effects, and Participant as a random effect. The ANOVA indicated a significant interaction between Group and Training Block ($F(9) = 2.25$, $p = .018$; $\eta_p^2 = 0.03$, 95% CI [3.02e-03, 1.00]). However, post hoc analysis did not indicate any amplitude difference across blocks. The second mixed linear model, predicting the amplitude of high alpha frequency with Group (High alpha and Control) and Training Block as fixed effects, and Participant as a random effect, indicated that the interaction between Group and Block was not statistically significant ($F(9) = 0.60$, $p = .795$; $\eta_p^2 = 0.008$, 95% CI [0.00, 1.00]).


Effect of neurofeedback on high alpha amplitude during resting-state

A mixed linear model was performed to predict the amplitude of high alpha frequency recorded during resting-state EEG, with Group and Resting-EEG as fixed effects and Participant as a random effect. The ANOVA revealed a statistically significant and robust main effect of Resting-EEG ($F(1) = 22.37$, $p < .001$; $\eta_p^2 = 0.19$, 95% CI [0.08 - 1.00]), indicating a higher average amplitude after training ($M = 6.18 \pm 2.29$) compared to before ($M = 5.83 \pm 2.17$). However, the main effect of Group ($p = .39$) and the interaction between Group and Resting EEG (global: $p = .99$; Theta x High Alpha, $p = .97$; Theta x Control, $p = .89$) was not statistically significant. We also assessed the difference in EEG activity during the first resting-state

using a classical linear model. The ANOVA indicated that the main effect of Group was not statistically significant ($p = .36$).

Figure 2

Evolution of theta and high alpha amplitude during neurofeedback and resting-states



Note. **A.** Changes in theta amplitude during neurofeedback and resting-states. The first line graph shows that, after an initial decrease, there has been a gradual increase of theta amplitude in the Theta and Control group. The second graph represents the mean amplitude and individual data points of theta frequency amplitude during the resting state, measured just before and just after the neurofeedback session. The third and final plot is an interaction plot showing the differential effects of neurofeedback training on the theta amplitude of each group. **B.** Changes in high alpha amplitude during neurofeedback and resting-states. The first line graph shows that there has been an increase of high alpha amplitude in


the three groups. The second graph shows the mean amplitude of the high alpha frequency and the individual data points for each resting-state and the last graph is an interaction plot.

Effects of neurofeedback on all brain frequencies

Figure 3 illustrate the effect of neurofeedback on all frequency bands. Statistical analysis of neurofeedback effects on other frequencies amplitude during training and resting-state are presented in Supplementary material (Section 1.1).

Figure 3

Effect of neurofeedback on EEG frequency bands.



Note. Forest plot illustrating the mean slope of frequency band amplitudes across the training, error bars represent standard errors.

Effect of neurofeedback on working memory performance (Figure 4)

Effect of neurofeedback on backward digit span task performance

We conducted a mixed linear model to predict the score on a backward digit span task, with Group and Cognitive Screening (pre-and post-training working memory measurement) as fixed effects and Participant as a random effect. The ANOVA revealed that the main effect of Cognitive Screening was statistically significant and strong ($F(1) = 29.39$, $p < .001$; $\eta_p^2 = 0.23$, 95% CI [0.12 – 1.00]), with a higher

mean score after training ($M = 6.71 \pm 1.62$) compared to before ($M = 5.80 \pm 1.69$). However, the main effect of Group and the interaction between Group and Cognitive Screening were not statistically significant ($p = .12$ and $p = .18$, respectively).

Effect of neurofeedback on 1-back and 2-back tasks performance

We conducted a mixed linear model to predict the score on the 1-back task, with Group and Cognitive Screening as fixed effects and Participant as a random effect. The ANOVA revealed that the main effect of Cognitive Screening was statistically significant and moderate ($F(1) = 6.69$, $p = .011$; $\eta_p^2 = 0.06$, 95% $CI [0.008 - 1.00]$), with a higher mean score after training ($M = 0.96 \pm 0.08$) compared to before ($M = 0.93 \pm 0.10$). The main effect of Group and the interaction between Group and Cognitive Screening were not statistically significant ($p = .19$ and $p = .16$, respectively). We then conducted a mixed linear model to predict the score on the 2-back task, with Group and Cognitive Screening as fixed effects and Participant as a random effect. The ANOVA revealed a statistically significant and robust main effect of Cognitive Screening ($F(1) = 25.64$, $p < .001$; $\eta_p^2 = 0.21$, 95% $CI [0.10 - 1.00]$). Post-training scores ($M = 0.93 \pm 0.08$) were significantly higher compared to pre-training scores ($M = 0.88 \pm 0.09$). However, the main effect of Group and the interaction between Group and Cognitive Screening were not significant ($p = .66$ and $p = .56$).

Effect of neurofeedback on 3-back and Corsi block tasks performance

Further analyses were conducted to evaluate the effects of Group, Cognitive Screening, and their interaction on 3-back scores and Corsi block performance. Our mixed models and ANOVA results indicated that the main effects of Group ($p = .70$ and $p = .08$) and Cognitive Screening ($p = .08$ and $p = .22$), as well as the interaction between these variables ($p = .53$ and $p = .12$), were not statistically significant.


Effect of brain activity modulation on working memory performance

To assess the link between brain activity and cognitive performance, we evaluated the statistical relationship between the slope of theta and high alpha amplitude during neurofeedback and the changes

in working memory task performance (measured through the difference between post-training and pre-training scores). Five linear models were performed to predict the changes in performance across working memory tasks based on the modulation of the amplitudes of two frequencies targeted during training (slope of the amplitude evolution for theta and high alpha frequencies) and the Group variable as predictor variables. The ANOVA tests showed no significant main effects nor interactions (results are presented in the Supplementary material, Section 1.2).

Figure 4

Changes in working memory tasks performance



Note. Average performance and individual data points on working memory tasks, namely digit span, spatial span and n -back (1, 2 & 3-back). Descriptively, there was an average increase in performance on the digit span, 1-back and 2-back tasks. As this increase was observed in all three groups, we did not observe an interaction effect. The span block task presented less unequivocal results, with an average increase for the Theta group, a decrease for the Control group and stagnation of performance for the High

alpha group. Conversely, on the 3-back task, the Control group showed the greatest improvement in performance, followed by the High alpha group and then the Theta group, whose performance stagnated. However, as presented in the text, we observed no interaction effect between groups and those two behavioral measures.

Effect of neurofeedback on subjective psychological state (Figure 5)

Calm score

We conducted a mixed-effects linear model to predict the calm score using Group and Self-Reported Measure as fixed effects and Participant as a random effect. The ANOVA indicated that the main effect of the Self-Reported Measure was statistically significant and moderate ($F(1) = 9.22, p = .003; \eta_p^2 = 0.09, 95\% CI [0.02 - 1.00]$), with a higher average score after training ($M = 3.19 \pm 0.97$) compared to before ($M = 2.78 \pm 1.05$). The main effect of the Group and the interaction between Group and Self-Reported Measure were not significant ($p = .92$ and $p = .52$, respectively).

Energetic score

We performed a mixed-effects linear model to predict the energy score using Group and Self-Reported Measure as fixed effects and Participant as a random effect. The ANOVA revealed that the main effect of the Self-Reported Measure was statistically significant and strong ($F(1) = 41.51, p < .001; \eta_p^2 = 0.30, 95\% CI [0.18 - 1.00]$), with a lower average score after training ($M = 1.16 \pm 0.97$) compared to before ($M = 1.88 \pm 0.98$). The main effect of the Group and the interaction between Group and Self-Reported Measure were not significant ($p = .19$ and $p = .38$).

Relaxed score

We fitted a mixed-effects linear model to predict the relaxation score using Group and Self-Reported Measure as fixed effects and Participant as a random effect. The ANOVA demonstrated that the main effect of the Self-Reported Measure was statistically significant and strong ($F(1) = 38.52, p < .001; \eta_p^2 = 0.28, 95\% CI [0.16 - 1.00]$), with a higher average score after training ($M = 3.08 \pm 0.94$) compared to

before ($M = 2.25 \pm 1.28$). The main effect of the Group and the interaction between Group and Self-Reported Measure were not significant ($p = .69$ and $p = .78$).

Satisfaction score

We conducted a mixed linear model to predict the satisfaction score, with Group and Self-Reported Measure as fixed effects and Participant as a random effect. The ANOVA revealed a statistically significant and moderate main effect of the Self-Reported Measure ($F(1) = 7.58, p = .007; \eta_p^2 = 0.07, 95\% CI [0.01 - 1.00]$), indicating a higher average score after the training ($M = 2.69 \pm 0.98$) compared to before ($M = 2.42 \pm 1.19$). However, the interaction effect between Group and Self-Reported Measure was statistically significant and moderate ($F(2) = 4.70, p = .011; \eta_p^2 = 0.09, 95\% CI [0.01, 1.00]$). Post hoc analysis revealed that for the Control group, the average score was significantly higher after the training compared to before ($\beta = 0.59, SE = 0.19, p = .03$). The main effect of Group was not significant ($p = .053$).

Happiness, Receptivity, and Concentration scores

We conducted three mixed linear models to predict the happiness score, receptivity score, and concentration score, with Group and Self-Reported Measure as fixed effects and Participant as a random effect. The ANOVA indicated that the main effects of Group, Cognitive Screening, and the interaction between these two variables were not significant for these three items (results are presented in the Supplementary material, Section 1.3).

Cognitive Load score


We performed a linear model to predict the subjective cognitive load score during neurofeedback training, with Group as the predictor variable. The ANOVA revealed a statistically significant and strong main effect of Group ($F(2, 98) = 38.19, p < .001; \eta_p^2 = 0.44, 95\% CI [0.32 - 1.00]$). Post hoc analysis demonstrated that the mean cognitive load score was significantly lower in the High Alpha group compared to the Theta group ($\beta = -1.92, t(98) = 7.78, SE = 0.25, p < .001$) and the Control group ($\beta = -1.63, t(98) = 7.01, SE = 0.23, p < .001$).

Flow, Mastery, and Coherence scores

We employed three linear models to predict the subjective feeling of flow, mastery and coherence of the feedback signal during the neurofeedback training, with Group as the predictive variable. The ANOVAs revealed that there were no significant main effects of Group on any of the scores ($p = .37$, $p = .86$ and $p = .39$, respectively).

Figure 5

Statistically significant changes in self-reported psychological state




Note. Average score and individual data point of questionnaire scores measured before and after neurofeedback. In a similar way for the three groups, there was an increase in the calm and relaxation scores and a decrease in the energy score. In terms of satisfaction, the Control group showed the strongest increase, as evidenced by the significant interaction effect (see Results section), followed by the Theta group. The High alpha group showed a slight decrease. *** $p < .001$.

Predictors of neuromodulation

To better understand the factors involved in neuromodulation, we utilized resting-state EEG activity, pre-training scores on working memory tasks and questionnaires to predict the neuromodulation of targeted signals through neurofeedback. A mixed linear model was employed to predict the slope of theta amplitude during neurofeedback using Group and resting-state brain frequencies (delta, theta, low alpha, high alpha, beta, and gamma) as predictor variables. ANOVA results revealed a statistically significant and strong main effect of resting-state theta amplitude ($F(1, 92) = 14.06, p < .001; \eta_p^2 = 0.13, 95\% CI [0.04 - 1.00]$). The other main effects were not significant (see Supplementary material, Section 1.4.a). A mixed linear model was utilized to predict the slope of high alpha amplitude during neurofeedback using Group and resting-state brain frequencies (delta, theta, low alpha, high alpha, beta, and gamma) as predictor variables. ANOVA indicated a statistically significant and moderate main effect of resting-state low alpha amplitude ($F(1, 92) = 7.80, p = .006; \eta_p^2 = 0.08, 95\% CI [0.01 - 1.00]$) and a strong main effect of resting-state high alpha amplitude ($F(1, 92) = 21.97, p < .001; \eta_p^2 = 0.19, 95\% CI [0.09 - 1.00]$). The other main effects were not significant (see Supplementary material, Section 1.4.b). The performance scores on working memory tasks and the questionnaire measured before neurofeedback did not predict the changes in the amplitude of the signals targeted by the training (results are presented in the Supplementary material, Section 1.4).

Figure 6

Scatter plots of the two predictors of neuromodulation



Note. We predicted the evolution of the amplitude (i.e., slope) of the signals targeted by neurofeedback (i.e., theta and high alpha) via psychological (self-reported questionnaire), cognitive (working memory performance) and electrophysiological (resting-state EEG) variables measured before neurofeedback. Only resting-state EEG of the trained signal predicted the slope of those signals (A: theta slope, B: high alpha slope). The two scatter plots show the slope of the trained signal as a function of the mean amplitude recorded in the resting-state. The points represent individual data, the line represents the fitted linear model and the shaded area the 95% confidence interval of the fitted values.

Discussion

In this study, we investigated the effect of a single neurofeedback session on brain oscillations amplitude, working memory task performance, and self-reported psychological states. We hypothesized that the neurofeedback's effects on brain activity would vary depending on whether the group was trained to enhance the amplitude of the theta frequency, the high alpha frequency, or random frequencies (active control group). This modulation of the targeted brain signal was expected to result in improved

performance in working memory tasks. To gain a deeper understanding of the self-regulation mechanism, we utilized resting-state EEG activity, working memory task performance, and self-reported psychological states to predict the neuromodulation of the trained signal. While our analyses did reveal electrophysiological and behavioral improvements, these gains were not exclusive to the trained signals. An increase in the frequency targeted by the training has been observed for the theta and high alpha groups, but no significant difference has been observed between these groups and the active control group. Among all the predictors of neuromodulation that have been examined, theta and high alpha amplitudes at rest have been shown to predict the ability to modulate the frequencies targeted by neurofeedback. Participants who showed the greatest increase in amplitude for the trained frequency during neurofeedback were those with the highest amplitudes during resting EEG measurements. Overall, these findings contribute to a better understanding of the factors influencing the outcomes of a neurofeedback training and highlight the challenges associated with using random frequencies training as an active control group.

Effect of neurofeedback on the amplitude of trained brain oscillations

The main goal of this study was to investigate the effects of a neurofeedback session designed to enhance the amplitude of theta and high alpha frequencies. The first hypothesis was that engaging in neurofeedback to increase the amplitudes of theta and high alpha frequencies would indeed affect the amplitudes of the targeted frequencies.

Theta frequency

When comparing the three groups together, we observed a main effect of neurofeedback training on the average amplitude of theta frequency but no interaction effect. Nonetheless, the absence of a global interaction effect may mask specific differences among the groups. Indeed, we observed a significant interaction between the Theta and High Alpha groups, indicating an increase in theta amplitude in the former compared to the latter. This interaction effect is consistent with previous research that

indicates an increase in theta amplitude during a single training session (Eschmann et al., 2021; Rozengurt et al., 2016; Rozengurt et al., 2017; Shtoots et al., 2021), and suggests that the theta training had specific effects compared to a group trained to modulate another frequency. However, we did not observe significant differences between the Theta group and our active Control group. In the previously cited studies, the active control group underwent training to enhance the amplitude of the beta frequency, resulting in slight decreases (Rozengurt et al., 2017), no change (Rozengurt et al., 2016), or slight increases in theta amplitude (8%, Shtoots et al., 2021). Another study utilized neurofeedback to increase beta1 frequency (15-18 Hz) amplitude and showed a slight decrease in theta amplitude during the training (Pimenta et al., 2018). The lack of difference could be attributed to our control group, which used random frequencies instead of the beta frequency. According to our findings, training to increase multiple frequencies appears to induce a more generalized neuromodulation than training targeting a specific frequency. Thus, a more appropriate control group would be one trained either to decrease randomly selected frequencies or to increase a different and uncorrelated frequency (e.g., beta). Concerning the effect of training on resting brain activity, our findings revealed an increase in the amplitude of resting-state theta amplitude after the training. However, no significant differences were observed between the groups concerning changes in theta amplitude, whether through assessing the overall interaction among the three groups or through additional pairwise comparisons. This lack of interaction effect cannot be directly compared to the findings of earlier studies (Eschmann et al., 2021; Rozengurt et al., 2016; Rozengurt et al., 2017; Shtoots et al., 2021), as these previous studies solely assessed theta frequency amplitude at rest before the neurofeedback, without considering post-training measurements.

High alpha frequency

Consistent with previous studies (Escolano et al., 2012; Escolano et al., 2014), we observed that a single high alpha neurofeedback training session leads to an increase in the amplitude of the high alpha frequency both during training (compared to the theta group) and in subsequent resting states. However,

we did not observe an interaction effect when comparing the high alpha group with the active control group, as the latter also exhibited an increase in high alpha frequency. Again, this suggests that training to increase different frequencies across neurofeedback training may result in nonspecific, broad effects on frequency amplitude.

Other frequencies

The analysis of the neurofeedback effect on the amplitude of other brain frequencies has shown an increase in delta frequency amplitude within the Theta and Control group, a reduction in beta frequency amplitude among the High alpha group, and a decrease in gamma frequency amplitude across all groups. Thus, our findings corroborate the absence of selectivity resulting from a single neurofeedback session on EEG activity (Dessy et al., 2020; Jurewicz et al., 2018; Pimenta et al., 2018). This outcome was expected within the Control group, which underwent training to regulate various brain frequencies, but not within the Theta and High alpha groups. A unique neurofeedback session does not appear to enable the modulation of brain activity exclusively within the frequency targeted by the neurofeedback device.

Effect of training on working memory performance

The second hypothesis of this study was that enhancing the amplitude of theta or high alpha frequencies through neurofeedback would improve performance on working memory tasks. While significant improvements were observed in performance on the digit span task and in two conditions of the n -back task (1-back and 2-back), these gains were observed across all groups. These improvements in performance across the three groups may be attributed to the neuromodulation observed in each of the groups. To test this hypothesis, we predicted the difference in working memory performance (post-training minus pre-training) using the slope of the evolution of theta and high alpha amplitude during training. Similar to the study conducted by Eschmann and colleagues (2021), which evaluated the link between the modulation of theta amplitude and changes in performance on a 2-back task, no statistically significant relationship was observed. The absence of statistically significant relationships in both studies

underscores the possibility that improvements in behavioral tasks may not directly result from the targeted neurofeedback training, but rather from non-specific factors such as increased task familiarity or increased motivation. It must, however, be emphasized that our ability to make psychophysiological inferences is limited here given that we did not record brain activity during the behavioral tasks. Further research examining cerebral activity during working memory tasks may yield more robust and comprehensive explanations.

Effect of training on self-reported questionnaire

The analysis of the questionnaires assessing participants' emotional state and consciousness revealed significant changes. A substantial decrease in the sense of energy was observed, alongside an increase in feelings of calmness and relaxation, regardless of the experimental group. The absence of differences between the groups suggests that these effects were not limited to the trained frequency and may indicate an increase in participants' levels of fatigue. Despite having short breaks between each block, the 30-minute neurofeedback training demanded continuous vigilance in focusing on the feedback signal. This increase in fatigue could potentially explain the observed modulation of various brain frequencies—namely, delta, theta, alpha, and beta. A recent meta-analysis emphasized the connection between fatigue and changes in amplitude within these frequencies (Tran et al., 2020). Interestingly, self-reported questionnaire responses indicated that both the Theta and Control groups experienced higher cognitive loads during training than the High alpha group. While this may support the established link between theta amplitude and cognitive load (Chikhi et al., 2022), we found no statistically significant correlation between theta activity and perceived cognitive load. Nonetheless, this result, along with similar motivation and agency scores across groups, suggests that participants of the Control group were equally engaged in the task as participants in the Theta and High alpha groups.

Towards specificity of EEG neurofeedback

Here, a single neurofeedback session aimed at increasing one frequency amplitude (theta or high alpha), does not elicit specific modulation of brain activity when compared to a group trained to increase random frequencies. First, this absence of significant differences may be attributed to the brief duration of our training, which was limited to a single session. Although there is no consensus on the time required for learning to occur, theoretical models suggest that the repeated production of the neural state rewarded by feedback induces neuroplasticity, facilitating the re-establishment of the neural state targeted by the training (Birbaumer et al., 2013; Davelaar, 2018; Lubianiker et al., 2022; Niv, 2013; Ros, 2014; Shibata et al., 2019). Thus, a prolonged training duration could strengthen the neural connections associated with the specific targeted frequencies, leading to more durable and robust neural changes compared to a group trained to modulate random signals. Consequently, by prioritizing sample size over the number of training sessions, our findings do not preclude the possibility that mastering the self-regulation of theta or high alpha frequencies over an extended period could produce distinct effects compared to a group trained to increase multiple frequency bands.

However, brain frequencies may be inter-correlated (Canolty & Knight, 2010; Klimesch, 2018), to the extent that EEG-neurofeedback training leads to non-specific modulation in the amplitude of brain frequencies. In this respect, the use of conventional EEG frequency bands could be sub-optimal when targeting a specific parameter of brain activity to positively impact a specific cognitive component. Therefore, it may be crucial to find functionally relevant biomarkers that can be trained independently. Hence, recent studies have used innovative methods for analyzing EEG signals in neurofeedback protocols. For example, studies have used multivariate pattern analysis (Haynes & Rees, 2006; Taschereau-Dumouchel et al., 2021) to "decode" cognitive states using machine learning techniques, thereby targeting specific brain networks involved in the cognitive state to be modified (Bu et al., 2019; Faller et al., 2019; Keynan et al., 2019; Tuckute et al., 2021; Rana et al., 2020; Wang et al., 2021). Other

studies have applied micro-state analysis, which identifies short, stable, distinct patterns of electrical activity measured by EEG across the scalp (Férat et al., 2022; Khanna et al., 2015; Michel & Koenig, 2018), representing various cognitive processes (Tarailis et al., 2024). Currently, few studies have attempted to modulate these neural states via neurofeedback (Asai et al., 2022; Diaz Hernandez et al., 2016), and it is necessary to assess larger samples in well-controlled protocols to highlight the potential of these methods compared to the use of conventional brain frequencies.

Predictive variables of neuromodulation of trained signals

In this research, our second objective was to identify factors that could potentially predict the modulation of brain signals through the use of a neurofeedback. We found that neither working memory performance nor pre-training questionnaire scores could predict the neuromodulation of the trained signals. Instead, the sole predictor for both trained frequencies (theta and high alpha) was the average amplitude of the targeted frequency recorded during rest before the training. This effect, already observed for alpha frequency (Su et al., 2021; Nan et al., 2018; Wan et al., 2014), was replicated here for theta frequency. This effect is particularly intriguing due to the specificity of the result: neuromodulation during training is not predicted by other brain signals or variables recorded in our study. The sole exception to this specificity pertains to the high alpha frequency, where modulation is also predicted by the amplitude of the resting low alpha. This result may suggest that the two sub-bands of the alpha frequency are not totally independent, at least within the temporal scope of our study. This effect could also contribute to an explanation for the observed increase in theta amplitude in the Control group. Indeed, the Control group exhibited higher resting theta activity compared to the High Alpha group and demonstrated similar activity to the Theta group, reflecting the neuromodulation pattern observed during neurofeedback. Our study appears to validate the existing relationship between neuromodulation capacity and resting-state signal amplitude. Thus, it may be relevant to consider using this measure as an inclusion criterion to select a sample with a higher proportion of learners. This approach would enhance

the statistical power of studies without increasing the sample size (Thibault & Pedder, 2022). What remains to be elucidated, however, is the mechanism behind this statistical relationship. In cognitive training, a lower baseline level is usually associated with greater improvement in the trained variable (Roheger et al., 2020; Traut et al., 2021). In the case of neuromodulation, our findings suggest an inverse relationship: higher amplitude of the trained frequency seems to indicate a higher likelihood of neuronal synchronization in the targeted brain region. One hypothesis could be that the oscillatory activity level (i.e., amplitude) recorded at rest reflects the level of neuronal excitability in the targeted frequency (Ogata et al., 2019; Schutter & Hortensius, 2011). To further explore this hypothesis, experimental testing could stimulate brain activity immediately prior to neurofeedback session (Orendáčová & Kvašňák, 2021).

Conclusion

Our neurofeedback training led to electrophysiological and behavioral improvements in all groups, including the active control group. Although these results may suggest a lack of specific neurofeedback effect, identifying the non-specific factors contributing to neuromodulation remains complex due to the sensory, cognitive, and affective factors involved in the self-regulation task (Lubianiker et al., 2019). A factor that appears robustly associated with brain activity neuromodulation is the resting-state activity recorded before training. Still, longer-term protocols with sufficient statistical power are still necessary to draw solid conclusions about the mechanisms involved in EEG activity modulation by neurofeedback.

Declarations

Funding

S.C.'s thesis was funded by the Agence de l'innovation de défense (AID) and the Ecole Nationale de l'Aviation Civile (ENAC). The views expressed are those of the authors and not those of the AID or the ENAC. The authors have no competing interests to declare that are relevant to the content of this article.

Conflicts of interest/Competing interest

The authors have no relevant financial or non-financial interests to disclose.

Ethics approval

This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Ethics Committee of the University Paris Cité (IRB No. = 00012019-41).

Consent to participate/Consent for publication

Informed consent was obtained from all individual participants included in the study.

Availability of data and code (Open Practices Statement)

The data and R script are available on this OSF repository:

https://osf.io/zs97e/?view_only=7d01849a74454cdfa42a09a92938cf64

References

Asai, T., Hamamoto, T., Kashihara, S., & Imamizu, H. (2022). Real-time detection and feedback of canonical electroencephalogram microstates: Validating a neurofeedback system as a function of delay. *Frontiers in Systems Neuroscience*, *16*, 786200.

<https://doi.org/10.3389/fnsys.2022.786200>

- Alkoby, O., Abu-Rmileh, A., Shriki, O., & Todder, D. (2018). Can we predict who will respond to neurofeedback? A review of the inefficacy problem and existing predictors for successful EEG neurofeedback learning. *Neuroscience*, *378*, 155-164.
- Başar, E., Başar-Eroglu, C., Karakaş, S., & Schürmann, M. (2001). Gamma, alpha, delta, and theta oscillations govern cognitive processes. *International journal of psychophysiology*, *39*(2-3), 241-248.
- Bassett, D. S., & Khambhati, A. N. (2017). A network engineering perspective on probing and perturbing cognition with neurofeedback. *Annals of the New York Academy of Sciences*, *1396*(1), 126-143.
- Bates, D., Kliegl, R., Vasishth, S., & Baayen, H. (2015). Parsimonious mixed models. *arXiv preprint arXiv:1506.04967*.
- Birbaumer, N., Ruiz, S., & Sitaram, R. (2013). Learned regulation of brain metabolism. *Trends in cognitive sciences*, *17*(6), 295-302.
- Bismuth, J., Vialatte, F., & Lefaucheur, J. P. (2020). Relieving peripheral neuropathic pain by increasing the power-ratio of low- β over high- β activities in the central cortical region with EEG-based neurofeedback: Study protocol for a controlled pilot trial (SMRPain study). *Neurophysiologie Clinique*, *50*(1), 5-20.
- Bobby, J. S., & Prakash, S. (2017). Upper alpha neurofeedback training enhances working memory performance using LabVIEW. *International Journal of Biomedical Engineering and Technology*, *25*(2-4), 120-132.
- Brandmeyer, T., & Delorme, A. (2020). Closed-loop frontal midline θ neurofeedback: A novel approach for training focused-attention meditation. *Frontiers in Human Neuroscience*, *14*, 246.
- Button, K. S., Ioannidis, J. P., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S., & Munafò, M. R. (2013). Power failure: why small sample size undermines the reliability of neuroscience. *Nature reviews neuroscience*, *14*(5), 365-376.

- Bu, J., Young, K. D., Hong, W., Ma, R., Song, H., Wang, Y., ... & Zhang, X. (2019). Effect of deactivation of activity patterns related to smoking cue reactivity on nicotine addiction. *Brain*, *142*(6), 1827-1841. <https://doi.org/10.1093/brain/awz114>
- Buzsaki, G., & Draguhn, A. (2004). Neuronal oscillations in cortical networks. *Science*, *304*(5679), 1926-1929.
- Canolty, R. T., & Knight, R. T. (2010). The functional role of cross-frequency coupling. *Trends in cognitive sciences*, *14*(11), 506-515.
- Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in cognitive sciences*, *18*(8), 414-421.
- Chen, X., Ma, R., Zhang, W., Zeng, G. Q., Wu, Q., Yimiti, A., ... & Zhang, X. (2023). Alpha oscillatory activity is causally linked to working memory retention. *PLoS Biology*, *21*(2), e3001999.
- Chen, X., & Sui, L. (2023). Alpha band neurofeedback training based on a portable device improves working memory performance of young people. *Biomedical Signal Processing and Control*, *80*, 104308.
- Chiasson, P., Boylan, M. R., Elhamiasl, M., Pruitt, J. M., Ranjan, S., Riels, K., ... & Keil, A. (2023). Effects of Neurofeedback training on performance in laboratory tasks: A systematic review. *International Journal of Psychophysiology*, *189*, 42-56.
- Chikhi, S., Matton, N., & Blanchet, S. (2022). EEG power spectral measures of cognitive workload: A meta-analysis. *Psychophysiology*, *59*(6), e14009.
- Chikhi, S., Matton, N., Sanna, M., & Blanchet, S. (2023). Mental strategies and resting state EEG: Effect on high alpha amplitude modulation by neurofeedback in healthy young adults. *Biological Psychology*, *178*, 108521.
- Clayton, M. S., Yeung, N., & Kadosh, R. C. (2015). The roles of cortical oscillations in sustained attention. *Trends in cognitive sciences*, *19*(4), 188-195.

- Cooper, P. S., Karayanidis, F., McKewen, M., McLellan-Hall, S., Wong, A. S., Skippen, P., & Cavanagh, J. F. (2019). Frontal theta predicts specific cognitive control-induced behavioural changes beyond general reaction time slowing. *Neuroimage*, *189*, 130-140.
<https://doi.org/10.1016/j.neuroimage.2019.01.022>
- Corsi, P. M. (1973). Human memory and the medial temporal region of the brain. *Dissertation Abstracts International*, *34*(2-B), 891.
- Davelaar, E. J. (2018). Mechanisms of neurofeedback: a computation-theoretic approach. *Neuroscience*, *378*, 175-188.
- Dehghanpour, P., Farokhi, F., & Rostami, R. (2018). Improvement of working memory performance by parietal upper alpha neurofeedback training. *International Journal of Smart Electrical Engineering*, *7*(02), 77-81.
- Deiber, M. P., Missonnier, P., Bertrand, O., Gold, G., Fazio-Costa, L., Ibanez, V., & Giannakopoulos, P. (2007). Distinction between perceptual and attentional processing in working memory tasks: a study of phase-locked and induced oscillatory brain dynamics. *Journal of cognitive neuroscience*, *19*(1), 158-172. <https://doi.org/10.1162/jocn.2007.19.1.158>
- Dessy, E., Mairesse, O., Van Puyvelde, M., Cortoos, A., Neyt, X., & Pattyn, N. (2020). Train your brain? Can we really selectively train specific EEG frequencies with neurofeedback training. *Frontiers in Human Neuroscience*, *14*, 22.
- de Vries, I. E., Slagter, H. A., & Olivers, C. N. (2020). Oscillatory control over representational states in working memory. *Trends in cognitive sciences*, *24*(2), 150-162.
<https://doi.org/10.1016/j.tics.2019.11.006>
- Diaz Hernandez, L., Rieger, K., Baenninger, A., Brandeis, D., & Koenig, T. (2016). Towards using microstate-neurofeedback for the treatment of psychotic symptoms in schizophrenia. A feasibility study in healthy participants. *Brain topography*, *29*, 308-321.

- Dobrushina, O. R., Vlasova, R. M., Rumshiskaya, A. D., Litvinova, L. D., Mershina, E. A., Sinitsyn, V. E., & Pechenkova, E. V. (2020). Modulation of intrinsic brain connectivity by implicit electroencephalographic neurofeedback. *Frontiers in human neuroscience*, 192.
- Domingos, C., Peralta, M., Prazeres, P., Nan, W., Rosa, A., & Pereira, J. G. (2021). Session frequency matters in neurofeedback training of athletes. *Applied Psychophysiology and Biofeedback*, 46, 195-204.
- Engel, A. K., Fries, P., & Singer, W. (2001). Dynamic predictions: oscillations and synchrony in top-down processing. *Nature Reviews Neuroscience*, 2(10), 704-716.
- Enriquez-Geppert, S., Huster, R. J., Figge, C., & Herrmann, C. S. (2014). Self-regulation of frontal-midline theta facilitates memory updating and mental set shifting. *Frontiers in behavioral neuroscience*, 8, 420.
- Enriquez-Geppert, S., Huster, R. J., & Herrmann, C. S. (2017). EEG-neurofeedback as a tool to modulate cognition and behavior: a review tutorial. *Frontiers in human neuroscience*, 11, 51.
- Eschmann, K. C., Bader, R., & Mecklinger, A. (2020). Improving episodic memory: Frontal-midline theta neurofeedback training increases source memory performance. *NeuroImage*, 222, 117219.
- Eschmann, K. C., & Mecklinger, A. (2022). Improving cognitive control: Is theta neurofeedback training associated with proactive rather than reactive control enhancement?. *Psychophysiology*, 59(5), e13873.
- Eschmann, K. C., Riedel, L., & Mecklinger, A. (2022). Theta neurofeedback training supports motor performance and flow experience. *Journal of Cognitive Enhancement*, 1-17.
- Escolano, C., Oliván, B., Lopez-del-Hoyo, Y., Garcia-Campayo, J., & Minguez, J. (2012, August). Double-blind single-session neurofeedback training in upper-alpha for cognitive enhancement of healthy subjects. In *2012 Annual International Conference of the IEEE Engineering in Medicine and Biology Society* (pp. 4643-4647). IEEE.

- Escolano, C., Navarro-Gil, M., Garcia-Campayo, J., & Minguéz, J. (2014). The effects of a single session of upper alpha neurofeedback for cognitive enhancement: A sham-controlled study. *Applied psychophysiology and biofeedback, 39*, 227-236.
- Esteves, I., Nan, W., Alves, C., Calapez, A., Melício, F., & Rosa, A. (2021). An exploratory study of training intensity in EEG neurofeedback. *Neural Plasticity, 2021*.
- Faller, J., Cummings, J., Saproo, S., & Sajda, P. (2019). Regulation of arousal via online neurofeedback improves human performance in a demanding sensory-motor task. *Proceedings of the National Academy of Sciences, 116*(13), 6482-6490. <https://doi.org/10.1073/pnas.1817207116>
- Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior research methods, 39*(2), 175-191.
- Férat, V., Seeber, M., Michel, C. M., & Ros, T. (2022). Beyond broadband: Towards a spectral decomposition of electroencephalography microstates. *Human Brain Mapping, 43*(10), 3047-3061.
- Gaume, A., Vialatte, A., Mora-Sánchez, A., Ramdani, C., & Vialatte, F. B. (2016). A psychoengineering paradigm for the neurocognitive mechanisms of biofeedback and neurofeedback. *Neuroscience & Biobehavioral Reviews, 68*, 891-910.
- Gruzelier, J. H. (2014). EEG-neurofeedback for optimising performance. I: A review of cognitive and affective outcome in healthy participants. *Neuroscience & Biobehavioral Reviews, 44*, 124-141.
- Guez, J., Rogel, A., Getter, N., Keha, E., Cohen, T., Amor, T., ... & Todder, D. (2015). Influence of electroencephalography neurofeedback training on episodic memory: A randomized, sham-controlled, double-blind study. *Memory, 23*(5), 683-694.
- Haegens, S., & Golombic, E. Z. (2018). Rhythmic facilitation of sensory processing: A critical review. *Neuroscience & Biobehavioral Reviews, 86*, 150-165.

- Hanslmayr, S., Axmacher, N., & Inman, C. S. (2019). Modulating human memory via entrainment of brain oscillations. *Trends in neurosciences*, 42(7), 485-499. <https://doi.org/10.1016/j.tins.2019.04.004>
- Hanslmayr, S., Sauseng, P., Doppelmayr, M., Schabus, M., & Klimesch, W. (2005). Increasing individual upper alpha power by neurofeedback improves cognitive performance in human subjects. *Applied psychophysiology and biofeedback*, 30, 1-10.
- Haugg, A., Renz, F. M., Nicholson, A. A., Lor, C., Götzendorfer, S. J., Sladky, R., ... & Steyrl, D. (2021). Predictors of real-time fMRI neurofeedback performance and improvement—A machine learning mega-analysis. *Neuroimage*, 237, 118207.
- Haynes, J. D., & Rees, G. (2006). Decoding mental states from brain activity in humans. *Nature reviews neuroscience*, 7(7), 523-534.
- Hothorn, T., Bretz, F., Westfall, P., Heiberger, R. M., Schuetzenmeister, A., Scheibe, S., & Hothorn, M. T. (2016). Package 'multcomp'. *Simultaneous inference in general parametric models. Project for Statistical Computing, Vienna, Austria*. Available from <http://multcomp.r-forge.r-project.org>.
- Hsieh, L. T., Ekstrom, A. D., & Ranganath, C. (2011). Neural oscillations associated with item and temporal order maintenance in working memory. *Journal of Neuroscience*, 31(30), 10803-10810. <https://doi.org/10.1523/JNEUROSCI.0828-11.2011>
- Hsu, Y. F., & Hämäläinen, J. A. (2022). Load-dependent alpha suppression is related to working memory capacity for numbers. *Brain Research*, 1791, 147994. <https://doi.org/10.1016/j.brainres.2022.147994>
- Hsueh, J. J., Chen, T. S., Chen, J. J., & Shaw, F. Z. (2016). Neurofeedback training of EEG alpha rhythm enhances episodic and working memory. *Human brain mapping*, 37(7), 2662-2675.
- Itthipuripat, S., Wessel, J. R., & Aron, A. R. (2013). Frontal theta is a signature of successful working memory manipulation. *Experimental brain research*, 224, 255-262. <https://doi.org/10.1007/s00221-012-3305-3>

- Jaumard-Hakoun, A., Chikhi, S., Medani, T., Nair, A., Dreyfus, G., & Vialatte, F. B. (2017). An apparatus to investigate western opera singing skill learning using performance and result biofeedback, and measuring its neural correlates. *Interspeech*, *55*, 102-111.
- Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: gating by inhibition. *Frontiers in human neuroscience*, *4*, 186.
- Jeunet, C., Lotte, F., Batail, J. M., Philip, P., & Franchi, J. A. M. (2018). Using recent BCI literature to deepen our understanding of clinical neurofeedback: a short review. *Neuroscience*, *378*, 225-233.
- Jurewicz, K., Paluch, K., Kublik, E., Rogala, J., Mikicin, M., & Wróbel, A. (2018). EEG-neurofeedback training of beta band (12–22 Hz) affects alpha and beta frequencies—A controlled study of a healthy population. *Neuropsychologia*, *108*, 13-24.
- Kadosh, K. C., & Staunton, G. (2019). A systematic review of the psychological factors that influence neurofeedback learning outcomes. *Neuroimage*, *185*, 545-555.
- Kenward, M. G., & Roger, J. H. (1997). Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics*, 983-997.
- Kerick, S. E., Asbee, J., Spangler, D. P., Brooks, J. B., Garcia, J. O., Parsons, T. D., ... & Robucci, R. (2023). Neural and behavioral adaptations to frontal theta neurofeedback training: A proof of concept study. *Plos one*, *18*(3), e0283418.
- Kessels, R. P., van Den Berg, E., Ruis, C., & Brands, A. M. (2008). The backward span of the Corsi Block-Tapping Task and its association with the WAIS-III Digit Span. *Assessment*, *15*(4), 426-434.
- Keynan, J. N., Cohen, A., Jackont, G., Green, N., Goldway, N., Davidov, A., ... & Hendler, T. (2019). Electrical fingerprint of the amygdala guides neurofeedback training for stress resilience. *Nature human behaviour*, *3*(1), 63-73. <https://doi.org/10.1038/s41562-018-0484-3>

- Khanna, A., Pascual-Leone, A., Michel, C. M., & Farzan, F. (2015). Microstates in resting-state EEG: current status and future directions. *Neuroscience & Biobehavioral Reviews*, *49*, 105-113.
<https://doi.org/10.1016/j.neubiorev.2014.12.010>
- Klimesch, W. (2018). The frequency architecture of brain and brain body oscillations: an analysis. *European Journal of Neuroscience*, *48*(7), 2431-2453.
- Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: the inhibition–timing hypothesis. *Brain research reviews*, *53*(1), 63-88.
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. (2017). lmerTest package: tests in linear mixed effects models. *Journal of statistical software*, *82*, 1-26.
- Kvamme, T. L., Ros, T., & Overgaard, M. (2022a). Can neurofeedback provide evidence of direct brain-behavior causality?. *Neuroimage*, *258*, 119400.
- Kvamme, T. L., Sarmanlu, M., & Overgaard, M. (2022b). Doubting the double-blind: Introducing a questionnaire for awareness of experimental purposes in neurofeedback studies. *Consciousness and Cognition*, *104*, 103381.
- La Marca, J. P., Cruz, D., Fandino, J., Cacciaguerra, F. R., Fresco, J. J., & Guerra, A. T. (2018). Evaluation of artifact-corrected electroencephalographic (EEG) training: a pilot study. *Journal of Neural Transmission*, *125*, 1087-1097.
- Lezak, M. D., Howieson, D. B., Loring, D. W., & Fischer, J. S. (2004). *Neuropsychological assessment*. Oxford University Press.
- Li, Z., Wang, H., Wu, X., Xu, X., Wei, S., & Yao, L. (2019, February). Working Memory Training Using EEG Neurofeedback Based on Theta Coherence of Brain Regions. In *2019 7th International Winter Conference on Brain-Computer Interface (BCI)* (pp. 1-6). IEEE.
- Lubianiker, N., Paret, C., Dayan, P., & Hendler, T. (2022). Neurofeedback through the lens of reinforcement learning. *Trends in Neurosciences*, *45*(8), 579-593.

Lubianiker, N., Goldway, N., Fruchtman-Steinbok, T., Paret, C., Keynan, J. N., Singer, N., ... & Hendler, T.

(2019). Process-based framework for precise neuromodulation. *Nature human behaviour*, 3(5), 436-445.

Luke, S. G. (2017). Evaluating significance in linear mixed-effects models in R. *Behavior research methods*, 49(4), 1494-1502.

Makowski, D., Lüdecke, D., & Ben-Shachar, M. S. (2020a). Modelbased: Estimation of model-based predictions, contrasts and means. *R package version 0.3.0*. CRAN. Available from <https://github.com/easystats/modelbased>.

Makowski, D., Ben-Shachar, M.S., Patil, I. & Lüdecke, D. (2020b). Automated Results Reporting as a Practical Tool to Improve Reproducibility and Methodological Best Practices Adoption. CRAN. Available from <https://github.com/easystats/report>.

Marins, T., Rodrigues, E. C., Bortolini, T., Melo, B., Moll, J., & Tovar-Moll, F. (2019). Structural and functional connectivity changes in response to short-term neurofeedback training with motor imagery. *Neuroimage*, 194, 283-290.

Michel, C. M., & Koenig, T. (2018). EEG microstates as a tool for studying the temporal dynamics of whole-brain neuronal networks: a review. *Neuroimage*, 180, 577-593.

Min, S. H., & Zhou, J. (2021). Smplot: An R package for easy and elegant data visualization. *Frontiers in Genetics*, 12, 802894.

Mueller, S. T., & Piper, B. J. (2014). The psychology experiment building language (PEBL) and PEBL test battery. *Journal of neuroscience methods*, 222, 250-259.

Muñoz-Moldes, S., & Cleeremans, A. (2020). Delineating implicit and explicit processes in neurofeedback learning. *Neuroscience & Biobehavioral Reviews*, 118, 681-688.

- Naas, A., Rodrigues, J., Knirsch, J. P., & Sonderegger, A. (2019). Neurofeedback training with a low-priced EEG device leads to faster alpha enhancement but shows no effect on cognitive performance: A single-blind, sham-feedback study. *PLoS One*, *14*(9), e0211668.
- Nan, W., Wan, F., Tang, Q., Wong, C. M., Wang, B., & Rosa, A. (2018). Eyes-closed resting EEG predicts the learning of alpha down-regulation in neurofeedback training. *Frontiers in psychology*, *9*, 1607.
- Navarro Gil, M., Escolano Marco, C., Montero-Marín, J., Minguez Zafra, J., Shonin, E., & García Campayo, J. (2018). Efficacy of neurofeedback on the increase of mindfulness-related capacities in healthy individuals: a controlled trial. *Mindfulness*, *9*, 303-311.
- Nawaz, R., Nisar, H., Yap, V. V., & Tsai, C. Y. (2022). The Effect of Alpha Neurofeedback Training on Cognitive Performance in Healthy Adults. *Mathematics*, *10*(7), 1095.
- Ninaus, M., Kober, S. E., Witte, M., Koschutnig, K., Stangl, M., Neuper, C., & Wood, G. (2013). Neural substrates of cognitive control under the belief of getting neurofeedback training. *Frontiers in human neuroscience*, *7*, 914.
- Niv, S. (2013). Clinical efficacy and potential mechanisms of neurofeedback. *Personality and Individual Differences*, *54*(6), 676-686. <https://doi.org/10.1016/j.paid.2012.11.037>
- Ogata, K., Nakazono, H., Uehara, T., & Tobimatsu, S. (2019). Prestimulus cortical EEG oscillations can predict the excitability of the primary motor cortex. *Brain Stimulation*, *12*(6), 1508-1516.
- Orendáčová, M., & Kvašňák, E. (2021). Effects of transcranial alternating current stimulation and neurofeedback on alpha (EEG) dynamics: a review. *Frontiers in Human Neuroscience*, *15*.
- Owen, A. M., McMillan, K. M., Laird, A. R., & Bullmore, E. (2005). N-back working memory paradigm: A meta-analysis of normative functional neuroimaging studies. *Human brain mapping*, *25*(1), 46-59.

- Pavlov, Y. G., & Kotchoubey, B. (2022). Oscillatory brain activity and maintenance of verbal and visual working memory: A systematic review. *Psychophysiology*, *59*(5), e13735.
- Pesonen, M., Hämäläinen, H., & Krause, C. M. (2007). Brain oscillatory 4–30 Hz responses during a visual n-back memory task with varying memory load. *Brain research*, *1138*, 171-177.
<https://doi.org/10.1016/j.brainres.2006.12.076>
- Pfeiffer, M., Kübler, A., & Hilger, K. (2024). Modulation of Human Frontal Midline Theta by Neurofeedback: A Systematic Review and Quantitative Meta-Analysis. *Neuroscience & Biobehavioral Reviews*, 105696.
- Pimenta, M. G., van Run, C., de Fockert, J. W., & Gruzelier, J. H. (2018). Neurofeedback of SMR and beta1 frequencies: an investigation of learning indices and frequency-specific effects. *Neuroscience*, *378*, 211-224.
- Pillette, L., Roc, A., N’Kaoua, B., & Lotte, F. (2021). Experimenters' influence on mental-imagery based brain-computer interface user training. *International Journal of Human-Computer Studies*, *149*, 102603.
- R Core Team. (2020). *R: A language and environment for statistical computing (4.0.3)*. R Foundation for Statistical Computing.
- Ramot, M., & Martin, A. (2022). Closed-loop neuromodulation for studying spontaneous activity and causality. *Trends in cognitive sciences*, *26*, 290-299.
- Ramot, M., Grossman, S., Friedman, D., & Malach, R. (2016). Covert neurofeedback without awareness shapes cortical network spontaneous connectivity. *Proceedings of the National Academy of Sciences*, *113*(17), E2413-E2420.
- Rana, K. D., Khan, S., Hämäläinen, M. S., & Vaina, L. M. (2020). A computational paradigm for real-time MEG neurofeedback for dynamic allocation of spatial attention. *Biomedical engineering online*, *19*, 1-17. <https://doi.org/10.1186/s12938-020-00787-y>

- Reiner, M., Rozengurt, R., & Barnea, A. (2014). Better than sleep: Theta neurofeedback training accelerates memory consolidation. *Biological psychology, 95*, 45-53.
- Reis, J., Portugal, A. M., Fernandes, L., Afonso, N., Pereira, M., Sousa, N., & Dias, N. S. (2016). An alpha and theta intensive and short neurofeedback protocol for healthy aging working-memory training. *Frontiers in aging neuroscience, 157*.
- Richardson, J. T. (2007). Measures of short-term memory: a historical review. *Cortex, 43*(5), 635-650.
- Riddle, J., Scimeca, J. M., Cellier, D., Dhanani, S., & D'Esposito, M. (2020). Causal evidence for a role of theta and alpha oscillations in the control of working memory. *Current Biology, 30*(9), 1748-1754.
- Roheger, M., Meyer, J., Kessler, J., & Kalbe, E. (2020). Predicting short-and long-term cognitive training success in healthy older adults: who benefits?. *Aging, Neuropsychology, and Cognition, 27*(3), 351-369.
- Ros, T., J. Baars, B., Lanius, R. A., & Vuilleumier, P. (2014). Tuning pathological brain oscillations with neurofeedback: a systems neuroscience framework. *Frontiers in human neuroscience, 8*, 1008. <https://doi.org/10.3389/fnhum.2014.01008>
- Ros, T., Enriquez-Geppert, S., Zotev, V., Young, K. D., Wood, G., Whitfield-Gabrieli, S., ... & Thibault, R. T. (2020). Consensus on the reporting and experimental design of clinical and cognitive-behavioural neurofeedback studies (CRED-nf checklist). *Brain, 143*(6), 1674-1685.
- Ros, T., Munneke, M. A., Ruge, D., Gruzelier, J. H., & Rothwell, J. C. (2010). Endogenous control of waking brain rhythms induces neuroplasticity in humans. *European Journal of Neuroscience, 31*(4), 770-778.
- Ros, T., Théberge, J., Frewen, P. A., Kluetsch, R., Densmore, M., Calhoun, V. D., & Lanius, R. A. (2013). Mind over chatter: plastic up-regulation of the fMRI salience network directly after EEG neurofeedback. *Neuroimage, 65*, 324-335.

- Roux, F., & Uhlhaas, P. J. (2014). Working memory and neural oscillations: alpha–gamma versus theta–gamma codes for distinct WM information?. *Trends in cognitive sciences*, *18*(1), 16-25.
<https://doi.org/10.1016/j.tics.2013.10.010>
- Rozengurt, R., Barnea, A., Uchida, S., & Levy, D. A. (2016). Theta EEG neurofeedback benefits early consolidation of motor sequence learning. *Psychophysiology*, *53*(7), 965-973.
- Rozengurt, R., Shtoots, L., Sheriff, A., Sadka, O., & Levy, D. A. (2017). Enhancing early consolidation of human episodic memory by theta EEG neurofeedback. *Neurobiology of learning and memory*, *145*, 165-171.
- Sampaio-Baptista, C., Neyedli, H. F., Sanders, Z. B., Havard, D., Huang, Y., Andersson, J. L., ... & Johansen-Berg, H. (2021). fMRI neurofeedback in the motor system elicits bidirectional changes in activity and in white matter structure in the adult human brain. *Cell Reports*, *37*(4).
- Sauseng, P., Griesmayr, B., Freunberger, R., & Klimesch, W. (2010). Control mechanisms in working memory: a possible function of EEG theta oscillations. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1015-1022. <https://doi.org/10.1016/j.neubiorev.2009.12.006>
- Sauseng, P., Klimesch, W., Heise, K. F., Gruber, W. R., Holz, E., Karim, A. A., ... & Hummel, F. C. (2009). Brain oscillatory substrates of visual short-term memory capacity. *Current biology*, *19*(21), 1846-1852. <https://doi.org/10.1016/j.cub.2009.08.062>
- Sauseng, P., Klimesch, W., Schabus, M., & Doppelmayr, M. (2005). Fronto-parietal EEG coherence in theta and upper alpha reflect central executive functions of working memory. *International journal of Psychophysiology*, *57*(2), 97-103. <https://doi.org/10.1016/j.ijpsycho.2005.03.018>
- Senoussi, M., Verbeke, P., Desender, K., De Loof, E., Talsma, D., & Verguts, T. (2022). Theta oscillations shift towards optimal frequency for cognitive control. *Nature Human Behaviour*, *6*(7), 1000-1013.

- Schad, D. J., Vasishth, S., Hohenstein, S., & Kliegl, R. (2020). How to capitalize on a priori contrasts in linear (mixed) models: A tutorial. *Journal of Memory and Language*, *110*, 104038.
- Schutter, D. J., & Hortensius, R. (2011). Brain oscillations and frequency-dependent modulation of cortical excitability. *Brain stimulation*, *4*(2), 97-103.
- Sghirripa, S., Graetz, L., Merkin, A., Rogasch, N. C., Ridding, M. C., Semmler, J. G., & Goldsworthy, M. R. (2021). The role of alpha power in the suppression of anticipated distractors during verbal working memory. *Brain Topography*, *34*, 102-109. <https://doi.org/10.1007/s10548-020-00810-4>
- Shen, L., Jiang, Y., Wan, F., Ku, Y., & Nan, W. (2023). Successful Alpha Neurofeedback Training Enhances Working Memory Updating and Event-related Potential Activity. *Neurobiology of Learning and Memory*, 107834
- Sherlin, L. H., Arns, M., Lubar, J., Heinrich, H., Kerson, C., Strehl, U., & Serman, M. B. (2011). Neurofeedback and basic learning theory: implications for research and practice. *Journal of Neurotherapy*, *15*(4), 292-304.
- Shibata, K., Lisi, G., Cortese, A., Watanabe, T., Sasaki, Y., & Kawato, M. (2019). Toward a comprehensive understanding of the neural mechanisms of decoded neurofeedback. *Neuroimage*, *188*, 539-556.
- Shibata, K., Watanabe, T., Sasaki, Y., & Kawato, M. (2011). Perceptual learning incepted by decoded fMRI neurofeedback without stimulus presentation. *Science*, *334*(6061), 1413-1415.
- Shoji, Y., Patti, C. R., & Cvetkovic, D. (2017, July). Electroencephalographic neurofeedback to up-regulate frontal theta rhythms: preliminary results. In *2017 39th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)* (pp. 1425-1428). IEEE.
- Shtoots, L., Dagan, T., Levine, J., Rothstein, A., Shati, L., & Levy, D. A. (2021). The Effects of Theta EEG Neurofeedback on the Consolidation of Spatial Memory. *Clinical EEG and Neuroscience*, *52*(5), 338-344.

- Sitaram, R., Ros, T., Stoeckel, L., Haller, S., Scharnowski, F., Lewis-Peacock, J., ... & Sulzer, J. (2017). Closed-loop brain training: the science of neurofeedback. *Nature Reviews Neuroscience*, *18*(2), 86-100.
- Smit, D., Dapor, C., Koerts, J., Tucha, O. M., Huster, R. J., & Enriquez-Geppert, S. (2023). Long-term improvements in executive functions after frontal-midline theta neurofeedback in a (sub) clinical group. *Frontiers in Human Neuroscience*, *17*, 1163380.
- Sorger, B., Scharnowski, F., Linden, D. E., Hampson, M., & Young, K. D. (2019). Control freaks: Towards optimal selection of control conditions for fMRI neurofeedback studies. *Neuroimage*, *186*, 256-265.
- Soveri, A., Antfolk, J., Karlsson, L., Salo, B., & Laine, M. (2017). Working memory training revisited: A multi-level meta-analysis of n-back training studies. *Psychonomic bulletin & review*, *24*(4), 1077-1096.
- Su, K. H., Hsueh, J. J., Chen, T., & Shaw, F. Z. (2021). Validation of eyes-closed resting alpha amplitude predicting neurofeedback learning of upregulation alpha activity. *Scientific reports*, *11*(1), 19615.
- Szucs, D., & Ioannidis, J. P. (2017). Empirical assessment of published effect sizes and power in the recent cognitive neuroscience and psychology literature. *PLoS biology*, *15*(3), e2000797.
- Takabatake, K., Kunii, N., Nakatomi, H., Shimada, S., Yanai, K., Takasago, M., & Saito, N. (2021). Musical auditory alpha wave neurofeedback: Validation and cognitive perspectives. *Applied Psychophysiology and Biofeedback*, *46*(4), 323-334.
- Tarailis, P., Koenig, T., Michel, C. M., & Griškova-Bulanova, I. (2024). The functional aspects of resting EEG microstates: a systematic review. *Brain topography*, *37*(2), 181-217.
<https://doi.org/10.1007/s10548-023-00958-9>
- Taschereau-Dumouchel, V., Cortese, A., Lau, H., & Kawato, M. (2021). Conducting decoded neurofeedback studies. *Social Cognitive and Affective Neuroscience*, *16*(8), 838-848.

- Thibault, R. T., Lifshitz, M., & Raz, A. (2016). The self-regulating brain and neurofeedback: Experimental science and clinical promise. *Cortex*, *74*, 247-261.
- Thibault, R. T., Lifshitz, M., & Raz, A. (2017). Neurofeedback or neuroplacebo?. *Brain*, *140*(4), 862-864.
- Thibault, R. T., & Pedder, H. (2022). Excess significance and power miscalculations in neurofeedback research. *NeuroImage. Clinical*, *35*, 103008.
- Tran, Y., Craig, A., Craig, R., Chai, R., & Nguyen, H. (2020). The influence of mental fatigue on brain activity: Evidence from a systematic review with meta-analyses. *Psychophysiology*, *57*(5), e13554.
- Tseng, Y. H., Tamura, K., & Okamoto, T. (2021). Neurofeedback training improves episodic and semantic long-term memory performance. *Scientific reports*, *11*(1), 17274.
- Tuckute, G., Hansen, S. T., Kjaer, T. W., & Hansen, L. K. (2021). Real-time decoding of attentional states using closed-loop EEG neurofeedback. *Neural Computation*, *33*(4), 967-1004.
- Uslu, S., & Vögele, C. (2023). The more, the better? Learning rate and self-pacing in neurofeedback enhance cognitive performance in healthy adults. *Frontiers in Human Neuroscience*.
- Van Diepen, R. M., Foxe, J. J., & Mazaheri, A. (2019). The functional role of alpha-band activity in attentional processing: the current zeitgeist and future outlook. *Current opinion in psychology*, *29*, 229-238.
- Viviani, G., & Vallesi, A. (2021). EEG-neurofeedback and executive function enhancement in healthy adults: A systematic review. *Psychophysiology*, *58*(9), e13874.
- Wan, F., Nan, W., Vai, M. I., & Rosa, A. (2014). Resting alpha activity predicts learning ability in alpha neurofeedback. *Frontiers in human neuroscience*, *8*, 500.
- Wang, J. R., & Hsieh, S. (2013). Neurofeedback training improves attention and working memory performance. *Clinical Neurophysiology*, *124*(12), 2406-2420.

- Wang, B., Xu, Z., Luo, T., & Pan, J. (2021). EEG-Based Closed-Loop Neurofeedback for Attention Monitoring and Training in Young Adults. *Journal of Healthcare Engineering, 2021*(1), 5535810. <https://doi.org/10.1155/2021/5535810>
- Weber, L. A., Ethofer, T., & Ehlis, A. C. (2020). Predictors of neurofeedback training outcome: A systematic review. *NeuroImage: Clinical, 27*, 102301.
- Wechsler, D. (1955). Wechsler adult intelligence scale. *Archives of Clinical Neuropsychology*.
- Wei, T. Y., Chang, D. W., Liu, Y. D., Liu, C. W., Young, C. P., Liang, S. F., & Shaw, F. Z. (2017). Portable wireless neurofeedback system of EEG alpha rhythm enhances memory. *Biomedical engineering online, 16*(1), 1-18.
- Weisz, N., & Keil, A. (2022). Introduction to the special issue of human oscillatory brain activity: Methods, models, and mechanisms. *Psychophysiology, 59*(5), e14038.
- Wickham, H. (2016). Data analysis. In *ggplot2* (pp. 189-201). Springer, Cham.
- Yeh, W. H., Hsueh, J. J., & Shaw, F. Z. (2021). Neurofeedback of alpha activity on memory in healthy participants: A systematic review and meta-analysis. *Frontiers in Human Neuroscience, 14*, 562360.
- Yeh, W. H., Ju, Y. J., Liu, Y. T., & Wang, T. Y. (2022). Systematic review and meta-analysis on the effects of neurofeedback training of theta activity on working memory and episodic memory in healthy population. *International Journal of Environmental Research and Public Health, 19*(17), 11037.
- Zoefel, B., Huster, R. J., & Herrmann, C. S. (2011). Neurofeedback training of the upper alpha frequency band in EEG improves cognitive performance. *Neuroimage, 54*(2), 1427-1431. ISO 690.