

Addressing the climate adaptation tracking gap: an assessment method and its application to the Caribbean region

Camille Laurent, Virginie K E Duvat

To cite this version:

Camille Laurent, Virginie K E Duvat. Addressing the climate adaptation tracking gap: an assessment method and its application to the Caribbean region. Regional Environmental Change, 2024, 24 (4), pp.147. 10.1007/s10113-024-02301-9. hal-04745306

HAL Id: hal-04745306 <https://hal.science/hal-04745306v1>

Submitted on 20 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Distributed under a Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

ORIGINAL ARTICLE

Addressing the climate adaptation tracking gap: an assessment method and its application to the Caribbean region

Camille Laurent¹ · Virginie K. E. Duvat1,[2](http://orcid.org/0000-0002-9336-3833)

Received: 18 February 2024 / Accepted: 28 August 2024 © The Author(s) 2024

Abstract

This article addresses the climate adaptation tracking gap. Indeed, we still ignore the intensity, nature, spatial distribution, efectiveness, and recent evolution of adaptation eforts at the national, regional, and global scales. We propose a webbased replicable assessment method using key variables to document adaptation eforts: country/territory, location, goal, implementation date, type of action, holder, funding source. Applying it to the Caribbean region, we analyzed 100 coastal adaptation actions. This studies the method while also highlighting the difficulties faced to track adaptation. We found that coastal adaptation eforts are substantial and increasing in the Caribbean, revealing the use of diversifed adaptation actions; prevalence of hard protection (51%); increase use of Nature-based Solutions (22%); limited use of retreat (6%); and accommodation (2%). Combined actions (17%) increased over time, due to the failure of single actions and need to fnd tradeofs between human asset protection encouraging hard protection and the maintenance of attractive tourist beaches encouraging beach nourishment. Puerto Rico and Trinidad and Tobago fall under the engineering-based "one-size-fts-all" adaptation model, whereas Jamaica and Barbados experiment diversifed options and combinations of options. Jamaica, Puerto Rico, Trinidad and Tobago, and Barbados are particularly active in taking adaptation action, while most dependent islands and sub-national island jurisdictions have no adaptation action reported. Considering the advantages and limitations of a web-based method compared to a feld-based approach, we recommend the combined use of these two complementary approaches to support adaptation tracking and help structuring communities of practice to the benefts of decision-makers and practitioners and scholars.

Keywords Coastal adaptation · Adaptation tracking · Assessment method · Tropical islands · Caribbean region · Small islands

Introduction

Tropical Islands (TI) are highly exposed to compound risks and their cascading impacts on coastal areas where most of the population, infrastructure, and economic activities concentrate (Glavovic et al. 2022; Mycoo et al. 2022). They

Communicated by Robbert Biesbroek

 \boxtimes Virginie K. E. Duvat virginie.duvat@univ-lr.fr Camille Laurent

camille.laurentm@gmail.com

¹ UMR LIENSs 7266, La Rochelle University-CNRS, Bâtiment ILE, 2 Rue Olympe de Gouges, 17 000, La Rochelle, France

Institut Universitaire de France (IUF), Paris, France

experience increasing coastal risks due to the combination of climate change impacts, especially sea-level rise (SLR), climate extremes, and anthropogenic disturbances (Magnan et al. 2019). Climate extremes mainly include extra-tropical and tropical cyclones and El Niño Southern Oscillation (ENSO) events. Whereas the strong swells and heavy precipitations generated by cyclones cause extensive fooding and accelerated shoreline retreat that have destructive impacts on human assets, ENSO events are involved in mass coral bleaching that alters the protection provided by coral reefs (Cooper et al. 2013; Duvat et al. 2019; Eddy et al. 2021). Coastal risks increasingly result from the compounding efects of such events (Bevacqua et al. 2020; Ford et al. 2018). The colonization process amplifed these risks through the promotion of highly vulnerable export-driven and climate-sensitive economies and accelerated coastal urbanization (Ferdinand 2019; Pichler and Striessnig 2013).

Sand mining from beach-dune systems and nearshore areas, the disturbance of physical-ecological processes by coastal development, and maladaptive responses, exacerbated coastal risks (Duvat et al. 2019; Jackson et al. 2012; Klöck et al. 2022). Because coastal risks are likely to make some islands and island coastal areas uninhabitable over this century (e.g., Duvat et al. 2021; Le Cozannet et al. 2021), they increasingly trigger climate adaptation in island countries and territories. This creates a scientifc opportunity to better understand coastal adaptation strategies in TI, which this article seizes.

Increasing climate risks call for adaptation which, in human systems, refers to "the process of adjustment to actual or expected climate and its efects, in order to moderate harm or exploit benefcial opportunities" (IPCC 2022). Adaptation aims at reducing current and future risks, as not addressing short- and long-term challenges in a coordinated manner can lead to maladaptation (Magnan et al. 2016). A wide range of coastal adaptation options exist, which include technical, institutional, ecological, and behavioral actions (Oppenheimer et al. 2019). This paper focuses on technical interventions aimed at reducing coastal climate-related risks, namely coastal erosion and marine fooding. In line with previous studies, these technical interventions include no response, hard protection, accommodation, ecosystem-based adaptation (EbA), retreat, advance with land raising, and deengineering, which can be used separately or in combination (Duvat et al. 2020; Oppenheimer et al. 2019). No response or "do-nothing" refers to a voluntary absence of action, maintaining coastal risks at their current level and letting them increase in the future (Hoggart et al. 2014). Hard protection consists in reducing coastal risks through the construction of structures (e.g., seawalls or riprap) that fx the shoreline and/ or reduce wave impact. Accommodation reduces the structural vulnerability of coastal human assets, including human constructions (e.g., food-proof buildings). Ecosystem-based adaptation (EbA), which is part of Nature-based Solutions (NbS), reduces climate risk through the protection, improved management, restoration, or creation of coastal and marine buffering ecosystems (Cohen-Shacham et al. 2016). Retreat reduces coastal risks by relocating exposed people, settlements, and human activities, to safer inland areas. Advance with land raising consists of advancing the shoreline through the creation of new and elevated land by reclaiming shallow water areas (Brown et al. 2019). These diferent actions can be used separately or in combination. For example, the hybrid type combines hard protection and EbA (Nicholls 2018). In line with Duvat et al. (2020: 110), these actions are here referred to as "adaptation-labelled" actions (ALA), since they have not necessarily proven to be efective.

In TI, "there is a consequent need to understand where adaptation takes place and what kinds of interventions are undertaken" (Klöck and Nunn 2019: 3). Whereas numerous

scientifc studies addressed coastal risks by quantifying hazards (e.g., Giardino et al. 2018) or evaluating vulnerability (e.g., Robinson 2017), limited attention was paid to on-theground coastal ALA. Some studies contributed to fll this knowledge gap by providing insights about pilot projects (e.g., Anisimov et al. 2020; Reguero et al. 2018), a given country or territory (e.g., Duvat 2020; Duvat et al. 2020), or a specifc type of action (e.g., Betzold and Mohamed 2017; Barnett et al. 2022). Although these studies have improved our understanding of coastal adaptation efforts in TI, we still do not have a comprehensive picture of the technical responses implemented in these settings (Chausson et al. 2020). We still ignore the intensity, nature, spatial distribution, efectiveness, and recent evolution of coastal adaptation efforts in TI. This situation is due to the lack of adaptation tracking at the global and national scales, which makes assessments relying upon available technical reports (e.g., Ferrario et al. 2014) and scattered feld investigations (e.g., Duvat 2013). This knowledge gap is exacerbated in archipelagos comprising tens to thousands of islands, due to distance, remoteness, and limited scientifc capacities of SIDS compared to continental states (Schultz et al. 2018).

This article thus falls under adaptation tracking research. In line with recent efforts made by scholars (e.g., Magnan et al. 2023), it contributes to the mapping of adaptation efforts in TI through a twofold approach. First, it proposes a methodological protocol aimed at overcoming the limitations of focused studies addressing a specifc location or adaptation response. In line with systematic assessments (e.g., Araos et al. 2016), our protocol uses web-based research, following the assumption that this approach allows to cover an entire island region by investigating with the same level of intensity the islands composing it. Key variables are documented for each ALA, including the country/ territory concerned, its precise location, main goal, implementation date, type (using the above-mentioned classifcation) and sub-type, brief description, holder, funding source, and information source. Second, it presents the results of the application of this protocol to coastal ALA in the Caribbean region. Third, based on the lessons learnt from this assessment, which highlight the benefts and limitations of webbased methods, we call for the combined use of web- and feld-based approaches to track adaptation.

Context and materials and methods

Study area

This study focuses on 26 Caribbean countries and territories composed of one or several islands, which are part of the Greater and Lesser Antilles and the Lucayan Archipelago (Fig. 1). French overseas territories are not included in this

Fig. 1 Study area

study, as they were the focus of another study in preparation. Study islands vary in size, from 13 km^2 for Saba to more than $110,000 \text{ km}^2$ for Cuba. Although most of them are high islands of continental, volcanic, or tectonic origin, several low-lying coral islands are included.

Caribbean TI have a high exposure and vulnerability to climate disasters, especially hurricanes and drought. For example, Hurricane Ivan (2004) affected 79% of Grenada's population (Manuel-Navarrete et al. 2007). Their high vulnerability is driven by a variety of factors, some of which are inherited from the colonial period, including the limited diversifcation of economic activities, marked societal divisions and inequalities, and widespread environmental degradation (Barclay et al. 2019; Popke and Rhiney 2019; Stanciof et al. 2018). In addition, the small size of Caribbean TI limits available natural and human resources and their coping capacity in the face of climate disasters (Medina et al. 2020; Schultz et al. 2018). Since the mid-twentieth century, coastal urbanization driven by external private investors for beach-oriented tourism and real estate further increased economic vulnerability and population exposure to such disasters (Klein 2018; Scott et al. 2012; Seraphin 2018). Under climate change, Caribbean TI experience an increase in their economic and human vulnerability, especially as their high debt levels challenge their fnancial capacity to invest in climate adaptation and address adaptation-related governance challenges. This has contributed to maladaptation by constraining the fnancial resources available at the national level to adapt and encouraging externally driven unsuitable projects (Gheuens et al. 2019; Robinson 2018).

Caribbean TI face increasing coastal erosion and marine fooding. Beach loss was reported in Anguilla, Antigua and Barbuda, St Kitts and Nevis, Montserrat, Dominica, and Grenada, where shoreline retreat reached approximately 0.5 m/year between 1985 and 2000, and it was exacerbated by the 2017 hurricanes (Barreto-Orta et al. 2019; Cambers 2009; Duvat et al. 2019; Pillet et al. 2019). Between 1993 and 2014, relative SLR was greater than average in the Eastern (3–5 mm/year) and North-Western (2.5–3 mm/year) Caribbean (Becker et al. 2019). Hurricanes caused major floods, as highlighted by the 2017 events in Saint-Martin, Saint-Barthelemy, and Cuba (Chew et al. 2020; Rey et al. 2019). In this region, coastal risks are signifcantly exacerbated by the decline of marine and coastal ecosystems. Seagrasses declined in Bermuda, the Cayman Islands, and the US Virgin Islands, whereas regional-wide bleaching

events caused extensive coral mortality, e.g., in 2005 when 70% of corals were afected (Mycoo et al. 2022). Category 4 and 5 hurricanes are a major risk to population and infrastructure, causing injuries, infectious diseases, and fatalities, especially among vulnerable populations, as well as severe economic loss. For example, Hurricane Maria (2017) caused more fatalities in Puerto Rico's municipalities exhibiting the lowest socio-economic development; it coincided with the frst case of *leptospirosis* in the US Virgin Islands; and losses amounted to more than 225% of the annual GDP in Dominica (Eckstein et al. 2018; Mycoo et al. 2022). However, the vulnerability of Caribbean islands varies depending on island size, relative isolation, resources, and socioeconomic features, with Haiti being recognized as one of the most vulnerable TI worldwide (Lam et al. 2014; Pichler and Striessnig 2013; Robinson 2017). Coastal erosion is projected to aggravate under accelerated SLR, especially under the RCP8.5 climate scenario (Le Cozannet et al. 2019). In the Caribbean region, where about 22 million people lived below 6 m in elevation in 2017, hurricanes are expected to remain the main driver of fooding (Cashman and Nagdee 2017; Ranasinghe et al. 2021). Even under the optimistic 1.5 °C global warming scenario, SLR would cause the loss of 3100 km^2 of land and be the dominant driver of financial adaptation needs in the region (Mycoo 2018; Tiggeloven et al. 2020).

Identifcation of coastal adaptation‑labelled actions

This study mainly relies on desktop web-based research conducted primarily in English and Spanish, and secondarily in French. It targeted all relevant data sources available online, including scientifc and grey literature (mainly technical reports), web media, adaptation program websites, adaptation project brochures, government websites, and climate fnance institution websites. We used the search engines *Scopus* and *Google Scholar* to identify relevant scientifc publications and *Google* for other sources. The scientifc literature search was conducted using keywords associated with the name of the ocean basin (i.e., *Caribbean*). The objective was to collect publications providing information on ALA, either through case studies and pilot projects or through overviews of the situation of study countries and territories. However, this search was not satisfactory, as most collected articles proposed an inventory of risks and recommendations to face them without mentioning concrete actions implemented on the ground. To overcome this limitation, we extended the search to the *Google* engine. First, we focused on a selection of international organizations (i.e., Adaptation Fund, World Bank, Organization of Eastern Caribbean States, Inter-American Development Bank, Caribbean Development Bank, United Nations Development Program, and international NGOs) associated with the name of the ocean basin and keywords related to the topic of interest, including *disaster risk management*, *disaster risk reduction*, *coastal risk reduction*, *coastal protection*, *coastal risk protection*, *climate change adaptation*, *coastal risk infrastructure*, *coastal relocation*, *coastal retreat*, *coastal accommodation*, *coastal ecosystem-based adaptation*, and *coastal nature-based solution* (Fig. 2, step 1). This search allowed us to identify international and regional organizations developing or funding large-scale adaptation programs. These programs were further investigated, either through additional queries on search engines or through the links provided on the web pages visited and the grey literature analyzed. This complementary search allowed us to determine whether ALA were conducted in the TI of interest. If so, the program and related actions were retained. Second, a similar search using the same keywords associated with the name of the ocean basin was run, without entering the name of international organizations (Fig. 2, step 2). This search aimed at completing the frst fndings by highlighting actions led by less popular organizations and other entities (e.g., public institutions). Third, every program and project retained was searched on *Google* along with the name of the concerned country (Fig. 2, step 3). This allowed us to fnd concrete cases of ALA. Collectively, these three interrelated steps provided a frst sample of ALA. However, for most of them, this search provided more information on risks or governance than on ALA. It mainly served as a basis for the identifcation of funding programs. A fourth step was then added to the protocol: for each TI, a specifc search using the name of the country or territory, and the above-mentioned keywords was run (Fig. 2, step 4). This search mainly led to government websites, including those of ministries and public agencies in charge of environmental matters, planning, and public works. It also directed us to the websites of associations and NGOs holding or implementing projects, and to local press articles. All identifed programs and projects were then searched, associated with the name of the country or territory or locality (preferably), to obtain as much information as possible on each ALA (Fig. 2, step 5). Finally, this was completed by a test aimed at checking the relevance of a series of keywords that could potentially improve the inventory already carried out, including *seawall*, *dike*, *levee*, *artifcial reef*, *coastal hard defense*, *nature-based solution*, *ecosystem-based adaptation*, *beach nourishment*, *beach restoration*, *coral gardening*, *house on stilts*, *coastal setback*, *coastal managed retreat*, *coastal relocation*, and *coastal managed realignment*. This second list of keywords was used for two territories where no project was found (Anguilla and Curaçao) and for two territories where several projects were identifed (the Dominican Republic and Cuba). This last search allowed for the identifcation of a few additional ALA, but generally it mostly revealed actions that had already been identifed.

Fig. 2 Web-based search method used to assess coastal adaptation-labelled actions. This fowchart describes the fve steps of the search method used to identify and characterize adaptation actions in this study

Database creation

Each inventoried action was entered into an Excel database, with the following attributes flled in: (1) Identifcation number; (2) Country or territory; (3) Location (from a region to a district); (4) Main goal (reduce coastal erosion, reduce coastal fooding, reduce both coastal erosion and fooding, reduce one of these two risks while also pursuing other objectives, e.g., preserve biodiversity or promote tourism); (5) Implementation date; (6) Action type (e.g., hard protection, following the IPCC categorization described above); (7) Action sub-type (e.g., seawall for hard protection, based on the fndings of the search; see sub-types and their defnition in Supplementary Material 2); (8) Brief description of the action (see Supplementary Material 2); (9) Holder type (e.g., national public authorities, local public authorities, and NGOs; see full description in Supplementary Material 2); (10) Holder identity; (11) Funding source (national institutions, local institutions, foreign government, NGO, international/regional organizations, private, co-funding; see full description in Supplementary Material 2); (12, 13) Funder type (e.g., hotel company) and identity; (14) Information sources (see Supplementary Material 1). Some of these attributes have predetermined modalities (see Supplementary Material 2 for the full description of the methodological protocol and the defnition of the abovementioned modalities). Wherever successive actions were implemented at the same site by the same holder, they were considered as distinct actions and documented as distinct lines in the database, using the same identifcation number with letters in alphabetical order (e.g., 1a, 1b). For those projects exhibiting missing information, a complementary search was conducted, using photointerpretation of satellite images freely provided by *Google Earth* and contacts with key informants (e.g., project managers or scientists). Thirty-four resource persons were contacted (Supplementary Material 3). The responses obtained provided additional information on actions (27%) and/or new key contacts (21%), who were solicited to collect complementary data. Sixty-two percent of resource persons had invalid email address, did not respond to our request, or could not provide additional information. Those actions for which we were able to document all attributes were kept in the database for processing and analysis. Data were analyzed based on statistical calculations involving fat sorting for each modality and bivariate analysis. Statistical results were reported on maps.

Advantages and limitations of the assessment method

Compared to field-based methods (e.g., Duvat et al. 2020), this protocol had the main advantage of allowing us to cover the whole Caribbean region and to identify a high number of ALA, 143 in total. Among these, we were able to document in full 100 ALA, which constitute the sample used in this study. These ALA were implemented between 1986 and 2021, and most of them over the past decade. Our assessment was constrained by data availability on the web, causing a "reporting bias" (Araos et al. 2016: 4) related to the communication efforts made by project holders. Indeed, ALA implemented on the ground may not be reported on the web. First, this *reporting bias* impacted the visibility of island countries and territories. Overseas territories were found to have limited visibility, since national communication efforts on adaptation generally focus on the mainland. In addition, because we did not conduct any search in Dutch, the Netherlands overseas territories are under-represented in this study. Second, this *reporting bias* led to the over- or under-representation of some types of ALA. We noted over-communication boosting the representation of EbA-NbS and very limited reporting of "do nothing." Likewise, accommodation is under-represented in this study, as it is generally part of national programs upgrading the standard of buildings or addressing the "build back better" challenge. Third, using a web-based protocol makes international and regional organizations and public institutions more visible than local communities and economic actors, since the former undertake communication campaigns highlighting their adaptation efforts whereas the latter do not.

Results and discussion: coastal adaptation‑labelled actions in the Caribbean region

As a reminder, 100 ALA were integrated into the database and analyzed. The following sections present the main results obtained. They focus on the main goals of ALA, their type and sub-type, their spatial distribution, their funding, and the risk reduction pathways highlighted by detailed case studies.

Predominance of coastal erosion‑oriented actions

Forty-eight percent of ALA aimed at reducing coastal erosion, whereas 28% addressed marine fooding and 21% these two risks (Fig. 3). Although the results may be biased by the inclusion of a high number of erosion-oriented actions implemented in Puerto Rico in our sample, based on the detailed study conducted by Bush et al. (2009), they highlight two key elements. First, the strong economic dependence of Caribbean Islands on beach-side tourism, which pushes public and private actors to take action to maintain or create attractive sandy beaches (Contact No.2, Supplementary Material 3). Second, the role of slow onset erosion accelerated by extreme events in triggering action, compared to rare and temporary fooding (Fabian et al. 2014; Wynne et al. 2016). These results fall in line with observations made in other touristic TI, e.g., in Mauritius (Duvat et al. 2020).

Prevalence of hard protection, Nature‑based Solutions, and combined actions

The results reveal the unequal use of the diferent types of adaptation actions. First, hard protection represents 51% of ALA (Fig. 4a) and is found in most countries and territories. Its use was stable over the past decades. Among hard

Fig. 4 Distribution of coastal adaptation-labelled actions by type and sub-type. **a** ALA distribution by type; **b** hard protection measures distribution by sub-type; **c** NbS distribution by sub-type

structures, seawalls and riprap prevail, representing 23.5% and 13.7% of these structures respectively, and are often used in combination (15.7%), including with other hard structures (7.8%) (Fig. 4b). These fndings are consistent with current knowledge emphasizing the prevalence of hard structures in SIDS, due to the difusion of the "seawall mindset" from developed to developing countries over recent decades (Banton et al. 2015; Jackson et al. 2012; Klöck et al. 2022). The widespread use of seawalls in Caribbean islands is consistent with observations made in the Pacifc Ocean (e.g., Fiji and Kiribati; Duvat 2013; Klöck et al. 2022; Nunn et al. 2021) and Indian Ocean (e.g., Reunion, Comoros, the Maldives;

Betzold and Mohammed 2017; Duvat 2020; Magnan and Duvat 2018; Ratter et al. 2019). This has caused a hard path dependency involving well-established institutional practices, past policies, and former investments, which contributed to its persistence and reinforcement over time, despite the failure and maladaptive character of hard structures at many locations and the emergence of alternative and more promising nature-based options (Morris et al. 2018; Nunn et al. 2021; Parsons et al. 2019; Temmerman et al. 2013).

Second, NbS represent 22% of adaptation actions, 64% of which were implemented over the past decade (Fig. 4a and Supplementary Material 1). The increasing use of NbS worldwide and in TI is due to a variety of factors, including the recognition of the failure of hard protection to reduce coastal risks, the growing international support provided to these alternative options (e.g., by the Convention on Biological Diversity or World Economic Forum), the promotion of community-based adaptation, and the recommendations made by the Intergovernmental Panel on Climate Change (IPCC) and Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) for their wider difusion (Chausson et al. 2020; Cohen-Shacham et al. 2019; Logan et al. 2018; Nesshöver et al. 2017; Reid et al. 2018; Scarano 2017; Wilson and Forsyth 2018). This strong support relies upon their potentially high efectiveness, low-cost compared to hard measures, "no-regrets" character, and the multiple co-benefts they generate, especially in TI (Beck et al. 2022; Ferrario et al. 2014; Manes et al. 2022; Narayan et al. 2016; Ruangpan et al. 2020). The sustained communication efforts made by NbS funders may have boosted NbS in our sample. In addition, our results reveal that the dominant type of NbS in the Caribbean is ecosystem restoration, whereas we only found a few cases of ecosystem protection and no case of ecosystem creation. With 59.4% of NbS, mangrove restoration is the most common NbS in the region. In contrast, seagrass restoration only accounts for 4.5% of NbS (Fig. 4c), which is in line with the observation made by Wilson and Forsyth (2018: 6) that "Restoration of seagrasses have generally received less attention than [restoration of] mangroves and coral reefs." We found two cases of hybrid actions involving coral gardening and hard protection in Antigua and Barbuda and in the Cayman Islands (Fig. 6a). The limited use of reef-based actions in our sample may be due frst, to the high level of degradation of reefs in the Caribbean which negatively affects the protection they can provide, and second, to the challenging character of reef restoration more generally (Duarte et al. 2020; Lirman and Schopmeyer 2016). Cost differences of restoration between marine ecosystems may also explain these results, as seagrass and reef restoration are much more expensive than mangrove restoration (Bayraktarov et al. 2016; Morrison et al. 2019). NbS were mainly found in Jamaica, Barbados, the British

Virgin Islands, Haiti, Grenada, the Dominican Republic, and Puerto Rico (Fig. 5).

Third, our results emphasize the use of other adaptation options, although these latter proved rarer (Fig. 4a). Retreat, which mainly consists in the relocation of people in our sample, represents 6% of ALA, with all cases found in Jamaica (Fig. 5). Accommodation is also used in the Caribbean, but to a lesser extent with only 2% of ALA. We only found one case of de-engineering, in Puerto Rico (Figs. 5 and 6), and one case of "do nothing," in Barbados. Advance with land rising was used as part of mixed actions in Jamaica (Fig. 6d). Thus, the various types of adaptation actions listed in the scientifc literature are used in the Caribbean region, although their distribution is uneven. The above-mentioned *reporting bias* may contribute to the under-representation of accommodation and "do nothing" in our sample. Because accommodation is generally implemented under national programs through building codes or at individual level by inhabitants, it requires feldwork to be adequately reported (López-Marrero 2010). Besides, the "do nothing" strategy is rarely reported, as the public authorities in charge of adaptation are supposed to take action, that is, "to do something" to reduce risk.

Fourth, 17% of ALA consist of combined actions. Among these, hybrid actions (combining hard protection and NbS) and mixed actions (other combinations) respectively represent 10% and 7% of ALA (Fig. 4a). Beach nourishment/ reprofling is involved in 90% of hybrid actions, including in association with other NbS. For example, seawalls, vegetation planting, and beach nourishment/reprofling were implemented together in Anguilla, whereas breakwaters, groins, and beach nourishment/reprofling were used jointly in Santa Lucia and St Kitts. Mixed actions, which generally involve accommodation, combine up to six technical interventions, as in Annotto Bay, Jamaica, where they associate hard protection, advance, NbS, and accommodation (Fig. 6d). We also noted combinations involving various sub-types of hard structures (joint use of seawalls and riprap; Fig. 4b), and various sub-types of NbS, with combinations representing 18% of the total for NbS (Fig. 4c and Fig. 6). The increasing development of combined ALA is in line with a growing assumption that optimal adaptation requires a combination of measures, preferably "green" and "grey" (Reddy et al. 2016).

Finally, we noted the occurrence of uncoordinated actions implemented at a given location by diferent stakeholders over time, sometimes with diferent objectives and without making any reference to former actions (Fig. 6). Six examples were identifed, four of which in Jamaica. In Old Harbor Bay, Jamaica, relocation was implemented by the Office of National Reconstruction and followed by the installation of an artifcial reef by the National Environment and Planning Agency. Likewise, in Rincon, Puerto Rico, hard structures were built by the local public authorities to reduce coastal erosion, which were later complemented by vegetation

planting by residents. In Telescope, Grenada, two projects of mangrove restoration took place almost simultaneously without coordination (Contact No.34, Supplementary Material 3).

Spatial distribution of coastal adaptation actions

ALA are unevenly distributed across the Caribbean (Fig. 5). Out of the 26 countries and territories considered, eight are missing from the inventory, all of which are either dependent islands (DI) or sub-national island jurisdictions (SNIJ), following the typology proposed by Petzold and Magnan (2019). Although the reporting biases may contribute to the low visibility of these territories, this fnding confrms the difficulties for DI and SNIJ to implement ALA due to the lack of local means and human resources and expertise, absence of representation in United Nations negotiations, limited bilateral help, lack of fnancial and legislative support from mainland, and eviction from international and regional programs (Petzold and Magnan 2019). Moreover, some countries and territories are much more active in climate adaptation and communication than others. Puerto Rico (22 actions), Jamaica (18), and Trinidad and Tobago (10) gather 50% of the actions reported. This is due, frst, to the existence of comprehensive data sources inventorying adaptation actions in these countries and territories (e.g., Bush et al. 2009 for Puerto Rico); second, to their strong communication efforts. For example, the Coastal Protection Unit of the Ministry of Work and Transport, Trinidad and Tobago, provides a list of ALA on its website. Likewise, web publications emanating from the government, the local press, and the Adaptation Fund extensively document ALA in Jamaica. The fact that Trinidad and Tobago and Puerto Rico are high-income countries probably contributes to their capacity to both implement and communicate around adaptation actions (Chausson et al. 2020). Among the 18 countries and territories that implemented ALA, 13 implemented less than 5 actions, whereas 3 implemented 6 to 10 actions (Fig. 5).

The unequal spatial distribution of ALA is also obvious at the scale of countries and territories, at which a

Fig. 5 Spatial distribution of coastal adaptation-labelled actions in the Caribbean region. This map emphasizes the uneven distribution of actions between the 26 countries and territories considered. Whereas eight countries and territories exhibit no action, five countries and territories concentrate most of inventoried actions, including Puerto Rico (22), Jamaica (18), Trinidad and Tobago (10), Barbados (10), and the Dominican Republic (7)

Fig. 6 Selected island examples of coastal adaptation-labelled actions in the Caribbean region. This fgure shows the spatial distribution of actions at the island scale. It highlights, frst, the high number of actions implemented in Trinidad and Tobago, Jamaica, and Puerto

Rico; second, the prevalence of hard protection in some countries (e.g., Trinidad and Tobago, Puerto Rico); third, the extended use of combined options, mostly mixed

core-periphery gradient emerges, with capital islands concentrating most actions, e.g., in the Bahamas, St Kitts and Nevis, Antigua and Barbuda, and Trinidad and Tobago (Fig. 6a, b, and c). This fnding is consistent with the assertion made by Klöck and Nunn (2019: 8) that "the core benefts from higher levels of external support and tends to have more and better infrastructure and technical know-how [compared to the periphery]." This statement was since then confrmed by additional studies (Nunn et al. 2021; Klöck et al. 2022). The spatial analysis also revealed the preference of some countries and territories for some ALA. Whereas the coastlines of Trinidad, Puerto Rico, and the Dominican Republic, are largely fxed by hard structures, Jamaica and Barbados show greater diversity of ALA (Fig. 6c, d, and e).

Funding of coastal adaptation actions

First, the major actors involved in coastal adaptation funding are national and local institutions (named public institutions hereafter), international and regional institutions, and the private sector (Fig. 7). In our sample, public institutions funded 40.5% of ALA, with national institutions (23.3%) prevailing over local institutions (17.2%). International organizations (Adaptation Fund, World Bank, and United Nations programs; 10.1%) and regional organizations (Inter-American Development Bank, Caribbean Development Bank; 10.1%), which played an increasing role in coastal adaptation funding since 2010, 20.2% of ALA. This result is in line with the fndings of Robinson and Gilfllan (2017). In addition, 14.1% of ALA were co-funded by public institutions and international and regional organizations. The private sector, mainly represented by hotel companies in our sample, funded 20.2% of ALA. Last, minor funding sources include NGOs (3.0% of actions) and foreign governments $(2.0\%).$

Second, these various funders supported diferent types of ALA (Fig. 7). Public institutions mainly supported hard protection (with 53% of hard protection actions), often used as a "one-size-fts-all" solution in TI because it benefts from an important visibility and provides an immediate sense of protection to residents (Betzold and Mohamed 2017; Logan et al. 2018; Nunn et al. 2021; Ratter et al. 2019). In addition, public institutions supported all retreat projects whereas they were little involved in NbS (10% of NbS funding only). In contrast, international and regional organizations extensively funded NbS, either separately (32% of NbS actions) or through co-funding involving partnership with governments (27%), and most mixed actions (43%). Private actors, especially hotel companies, were mainly involved in hybrid actions aimed at protecting exposed buildings while also supporting the maintenance of threatened beaches (Contact No.2, Supplementary Material 3).

Third, the analysis of ALA funding reveals marked differences between countries and territories. Whereas climate adaptation relies on public institutions in some countries (Jamaica, Barbados, and Trinidad and Tobago), it mainly depends on external funding in others (the British Virgin Islands, Haiti, the Bahamas). For example, in the British Virgin Islands, almost all ALA were funded by international and regional organizations. This situation questions the ability of such countries to bear the cost of these actions over time, which in turn questions the long-term efectiveness of actions requiring maintenance and monitoring, such as hard protection and NbS (Nunn and Kumar 2019).

Fig. 7 Funding of coastal adaptation-labelled actions in the Caribbean region. Public institutions include both national and local institutions. Private actors include hotel companies and residents. Co-funding refers to any combination of diferent funding sources (e.g., regional organization and national government; national government and foreign government)

Past‑to‑present risk reduction pathways

At six locations, various ALA were successively implemented by the same stakeholder over time to reduce coastal risks (Fig. 8). In line with Duvat et al. (2020), this situation allows to highlight risk reduction pathways. Most cases involved the combination of soft and hard measures. At some locations, stakeholders frst used NbS and then shifted to hard or hybrid structures. In Maundays Bay, Anguilla, a hotel company frst implemented repeated beach nourishment/reprofling in the 1990s. Because this action failed in protecting beach villas from wave-driven damages during Hurricane Lenny in 1999, the hotel company installed a seawall while also continuing beach nourishment and replanting vegetation in the 2000s (Wynne et al. 2016; Contact No.2, Supplementary Material 3). Similarly, in Piñones, Puerto Rico, beach nourishment was frst carried out in 1986 to rebuild a dune afected by sand mining that protected the coastal road. The artifcial dune was destroyed by storm waves and eventually replaced by riprap in the 1990s. In these cases, the successive actions undertaken were motivated by the failure of previous actions in stopping beach erosion and associated damage. In Telescope area, Grenada, to reduce marine fooding, a project was implemented which involved frst mangrove restoration and a year later, the installation of four artifcial reefs (Reguero et al. 2018). At other locations, adaptation actions shifted from hard to hybrid measures or NbS. At Gran Dominicus Resort, Dominican Republic, artifcial reefs were frst installed, followed by beach nourishment (Fabian et al. 2014). This second action was motivated by the fact that the reefs did not signifcantly reduce wave energy. This led to their strengthening by the addition of rock units. These artifcial reefs then caused excessive beach accretion and were for this reason moved further offshore (Fabian et al. 2014). The same type of situation occurred at Grand Cayman Marriott Resort,

Fig. 8 Examples of coastal risk reduction pathways in Caribbean Islands. These risk reduction pathways highlight shifts in adaptation actions over time and show, although the sample is small, the increased use of hybrid options

Cayman Islands, where artifcial reefs were frst installed to promote beach accretion in 2002, and complemented a few years later with new artifcial reefs, coral gardening, and beach nourishment (Fabian et al. 2014). In contrast, in one case, the same type of response was maintained over time. This occurred in Rincon, Puerto Rico, where gabions were installed in the early 1990s and replaced by seawalls the next decade to stop beach erosion, with both actions failing. Interestingly, fve out of these six risk reduction pathways underline hybridization of actions. The tourism-oriented function of beaches contributed to the choice of NbS, especially beach nourishment, which reinforces the aesthetic value of beaches (Contact No.2, Supplementary Material 3).

Way forward to improve adaptation tracking

We designed and applied to the Caribbean region a webbased methodological protocol aimed at tracking coastal adaptation. This protocol allowed us to map 100 coastal adaptation actions across the Caribbean region and to document them using nine key variables, including the country/territory where the action was implemented, the precise location of the action, its main goal, implementation date, type (using the IPCC categorization), sub-type, brief description, holder, and funding source. This demonstrates that our web-based method (see the "Identifcation of coastal adaptation-labelled actions" section), which is transferable to non-coastal adaptation actions and to other regions of the world, can provide a valuable support to adaptation tracking and thereby help to build an overall picture of climate adaptation worldwide (Table 1).

However, using the web to document adaptation actions revealed six main limitations. First, whereas the web search allowed us to identify 143 ALA, we were only able to document 100 ALA in full. This is because the actors involved in adaptation actions do not necessarily communicate in detail

about them on their websites and because no inventory effort is made at the country or territory level to report on adaptation. Consequently, the amount and type of information provided on adaptation actions varies signifcantly from one project to the other. To address this limitation and increase both the number of ALA included in our sample and the amount of information provided on each ALA, we used photointerpretation of satellite images and contacts with key informants (e.g., project managers or scientists). Although this allowed us to strengthen our ALA sample, 62% of the informants who were contacted did not provide additional information (invalid email address, no response, no information to share). Second, information gaps prevented us to determine for each ALA the reasons for its implementation, especially if it was deployed to address climate change impacts and/or pursue other goals (e.g., improve beach condition or support tourism revenues). Third, using the web to track adaptation led to the under-representation of some countries (eight out of the 26 countries targeted are missing in this assessment) and territories (i.e., overseas territories). Fourth, the web-based search also led to the under-representation of some types of ALA for which communication efforts are limited, including "do nothing" (as adaptation actors communicate about "the actions taken") and accommodation. The latter is generally supported by the public authorities through sectoral policies (e.g., housing policies aimed at promoting food-proof buildings), making

it difficult to capture localized accommodation actions (e.g., house on stilts) using a risk-oriented web-based protocol. Fifth, our web-based search led to the over-representation of the actions taken by "visible actors"—i.e., having a website and feeding it (i.e., public institutions, international and regional organizations, and NGOs)—and under-representation of the actions taken by actors who are less visible or "invisible" in the web (e.g., local communities and private individuals) or do not communicate about the adaptation actions they take. Sixth, our web-based method made it difficult to reconstruct risk reduction pathways (only six pathways highlighted), although the latter are crucial to learnt from past experiments to improve the effectiveness of adaptation actions.

These methodological limitations can be overcome by combining a web-based method such as the one used in this study with a feld-based approach involving local scientists and adaptation stakeholders at large (Table 2). Local social scientists could, for example, be involved in feld visits and semi-structured interviews and focus groups with public institutions, NGOs, and local associations, to collect detailed information on adaptation actions (e.g., Anisimov et al. 2020; Duvat et al. 2020; López-Marrero 2010). This would allow to describe in detail these actions and detect actions that are "invisible" on the web. Additionally, local ecologists and geomorphologists could provide valuable information on the successes and failures of past

Table 1 Benefts and limitations of a web-based approach for adaptation tracking

Variables Web-based approach

This table is based on the lessons learnt from the present study

Variables	Field-based approach	
	Benefits	Limitations
1. Spatial scale	- Local (site) - National	
2. Temporal scale	tion progress	- Allows repeated assessments aimed at measuring adapta- - Requires a large amount of time (field visits, interviews)
3. Description of adaptation actions	- Provides detailed information on adaptation actions, including the reasons that triggered action, the various types and sub-types of actions, actions taken by "invis- ible" actors - Allows to reconstruct some risk reduction pathways	- Difficulties to reconstruct a large number of risk reduction pathways due to the lack of reporting of actions and high staff turn-over, especially in public institutions
4. Information sources	- Allows to identify some adaptation stakeholders (local communities, private individuals, and companies) - Allows to identify local key informants and collect infor- mation among them	- Some adaptation stakeholders (especially international and regional organizations, NGOs, foreign governments) and the information they have can be missed

Table 2 Benefts and limitations of feld-based approaches for adaptation tracking

This table is based on the lessons learnt from a previous feld-based study (Duvat et al. 2020)

adaptations actions and help reconstruct risk reduction pathways. The comparative analysis of the specifc benefts and advantages of web- and feld-based approaches demonstrates that they are resolutely complementary (Tables 1 and 2). Web-based methods allow to cover entire countries and regions (e.g., the Caribbean region in this study) and to detect a large number of adaptation actions (143 actions in this study). This research effort is needed to develop region-wide assessments which, once aggregated, can feed global databases allowing to track adaptation and adaptation progress (through the reiteration of the approach) globally. Moreover, regional assessments are crucial to foster adaptation progress through experience sharing. In addition, feld-based approaches can help flling the gaps of web-based methods by providing information on "invisible" countries and territories and detailed information on adaptation actions to document (i) the nine key variables identifed to describe actions, (ii) the specifc reasons why they were implemented, (iii) actions that are "invisible" on the web, (iv) actions taken by actors that are "invisible" on the web, and (v) risk reduction pathways. On this latter point, a previous feld-based assessment completed in Mauritius showed that reconstructing risk reduction pathways based on interviews and locally available grey literature is challenging (Duvat et al. 2020), due to the lack of reporting of adaptation actions and a high staff turnover explaining a limited memory of past actions in public institutions. Beyond this challenge, combining web- and feld-based approaches would help creating or structuring communities of adaptation practice through increased networking efforts, which would in turn help designing robust standards for the development of national and global adaptation databases to the advantage of decision makers and practitioners and scholars.

Conclusion

This study provides an overview of coastal adaptationlabelled actions in the Caribbean, based on a sample of 100 technical actions identified using a web-based approach. Although it is not exhaustive and has some limitations that are inherent to the search method used, it demonstrates that desktop research can provide a valuable contribution to adaptation tracking at various spatial scales (including island, national, and regional), and proposes a replicable protocol to address this gap. First, we found that adaptation efforts aimed at reducing coastal erosion (48% of actions) and marine fooding (28%) are substantial in the Caribbean. Second, in line with Pacifc and Indian Ocean studies, this article highlights the prevalence of hard protection (51% of actions), especially seawalls (23.5% of hard structures) and riprap (13.7%), over Nature-based Solutions (22% of actions). The latter were more recently implemented, and they mainly consist of mangrove restoration (59.4% of NbS). Retreat and accommodation only represent 6% and 2% of actions, respectively. Combined actions, including hybrid actions involving hard protection and NbS (90% of which involve beach restoration) and mixed actions (other combinations), represent 17% of actions and increased over time. The hybridization of actions is confrmed by reconstructed risk reduction pathways. Combinations of actions and the accumulation of technical interventions over time result from the failure of single actions and the need to protect efectively human assets while also maintaining attractive beaches for tourism. Whereas some countries fall under the engineering-based "one-size-fts-all" adaptation model (e.g., Puerto Rico and Trinidad and Tobago), others (e.g., Jamaica and Barbados) experiment diversifed options and

combinations of options to adapt. Third, this study highlights the unequal spatial distribution of adaptation efforts, with Jamaica, Puerto Rico, Trinidad and Tobago, and Barbados being particularly active, whereas most dependent islands and sub-national island jurisdictions have no adaptation action reported. This emphasizes signifcant diferences in island countries and territories capacity to take adaptation action. A core-periphery gradient was also noted. Fourth, coastal adaptation funding mainly relies on public institutions, international and regional organizations, and the private sector. These various actors support diferent types of actions. Public actors mainly fund hard protection and retreat, whereas international and regional organizations support NbS and combined actions. The latter also have the preference of hotel companies. Some island countries and territories are highly dependent on external funding, which questions their ability to face adaptation challenges over the long run.

We identifed six limitations that are inherent to webbased methods, including difficulties to (i) document adaptations actions deployed in some countries (eight out of the 26 countries targeted have no action reported) and territories (especially overseas); (ii) document adaptation actions in full using our nine key variables; (iii) determine the reasons that triggered action (climate-related or not); (iv) report on the various types of adaptation actions equally ("do nothing" and accommodation are under-reported in the web); (v) report on the actions taken by local communities, private individuals, and private companies; (vi) reconstruct risk reduction pathways, due to the lack of reporting of past actions and high staff turn-over, especially in public institutions. We argue that these fve limitations could be overcome by combining the web-based method used in this study with a feld-based approach. Each of these two approaches has specifc benefts and limitations. A web-based protocol allows to identify a large number of actions at the national and regional scales and to provide a frst description of these actions. In addition, feld-based methods involving local scientists (from social to physical sciences) and stakeholders (especially public institutions, NGOs and local associations, and private companies) would allow to collect in-depth information on actions and thereby help flling the abovementioned gaps. Furthermore, bridging web- and feld-based approaches would allow creating or structuring communities of adaptation practice, which would in turn help designing robust standards for the development of national to global adaptation databases to the advantage of decision-makers and practitioners and scholars.

Supplementary Information The online version contains supplementary material available at<https://doi.org/10.1007/s10113-024-02301-9>. **Acknowledgements** The authors warmly thank the resource persons who provided information on coastal adaptation actions.

Funding This work was supported by the French National Research Agency under the STORISK (No. ANR- 15-CE03-0003) and the FUTURISKS (No. ANR-22-POCE-0002) projects. VKED also beneftted from the support of the French University Institute (IUF).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit<http://creativecommons.org/licenses/by/4.0/>.

References

- Anisimov A, Magnan A, Duvat V (2020) Learning from risk reduction pilot projects for enhancing long-term adaptation governance: the case of Mauritius Island (Indian Ocean). Environ Sci Policy 108:93–103.<https://doi.org/10.1016/j.envsci.2020.03.016>
- Araos M, Berrang-Ford L, Ford JD, Austin SE, Biesbroek R et al (2016) Climate change adaptation planning in large cities: a systematic global assessment. Environ Sci Policy 66:375–382. <https://doi.org/10.1016/j.envsci.2016.06.009>
- Banton J, Morin V, Warner P, Smith D (2015) Selection of appropriate coastal protection strategies for Caribbean Coastlines. In: Wallendorf L, Cox D.T. (eds). Coastal Structures and Solutions to Coastal Disasters 2015: Resilient Coastal Communities American Society of Civil Engineers, pp 570–581. ISBN: 978–0–7844–8030–4
- Barclay J, Wilkinson E, White CS, Shelton C, Forster J, et al. (2019) Historical trajectories of disaster risk in Dominica. Int J Disaster Risk Sci 10:149–165.<https://doi.org/10.1007/s13753-019-0215-z>
- Barnett J, Jarillo S, Swearer SE, Lovelock CE, Pomeroy A et al (2022) Nature-based solutions for atoll habitability. Phil Trans R Soc B 377:20210124.<https://doi.org/10.1098/rstb.2021.0124>
- Barreto-Orta M, Méndez-Tejeda R, Rodriguez E, Cabrera N, Diaz E, et al. (2019) State of the beaches in Puerto Rico after Hurricane Maria (2017). Shore & Beach 87(1):16–23
- Bayraktarov ES, Saunders MI, Abdullah S, Mills M, Beher J et al (2016) The cost and feasibility of marine coastal restoration. Ecol Appl 26(4):1055–1074.<https://doi.org/10.1890/15-1077>
- Beck MW, Heck N, Narayan S, Menénedez P, Reguero BG et al (2022) Return on investment for mangrove and reef flood protection. Ecosyst Serv 56:101440. [https://doi.org/10.1016/j.ecoser.2022.](https://doi.org/10.1016/j.ecoser.2022.101440) [101440](https://doi.org/10.1016/j.ecoser.2022.101440)
- Becker M, Karpytchev M, Papa F (2019) Hotspots of relative sea-level rise in the tropics. Tropical extremes: natural variability and trends 203–262.<https://doi.org/10.1016/B978-0-12-809248-4.00007-8>
- Betzold C, Mohamed I (2017) Seawalls as a response to coastal erosion and fooding: a case study from Grande Comore, Comoros (West Indian Ocean). Reg EnvironChange 17(4):1077–1087. [https://doi.](https://doi.org/10.1007/s10113-016-1044-x) [org/10.1007/s10113-016-1044-x](https://doi.org/10.1007/s10113-016-1044-x)
- Bevacqua E, Vousdoukas MI, Zappa G, Hodges K, Shpeherd TG et al (2020) More meteorological events that drive compound coastal

fooding are projected under climate change. Commun Earth Environ 1(1):47.<https://doi.org/10.1038/s43247-020-00044-z>

- Brown S, Wadey MP, Nicholls RJ, Shareef A, Khaleel Z et al (2019) Land raising as a solution to sea-level rise: an analysis of coastal fooding on an artifcial island in the Maldives. Flood Risk Management 13(Suppl. 1):e12567.<https://doi.org/10.1111/jfr3.12567>
- Bush DM, Neal WJ, Jackson CW (2009) Summary of Puerto Rico's vulnerability to coastal hazards: risk, mitigation, and management with examples. Geol Soc Am Spec Pap 460:149–165. [https://doi.](https://doi.org/10.1130/2009.2460(11)) [org/10.1130/2009.2460\(11\)](https://doi.org/10.1130/2009.2460(11))
- Cambers G (2009) Caribbean beach changes and climate change adaptation. Aquat Ecosyst Health Manage 12(2):168–176. <https://doi.org/10.1080/14634980902907987>
- Cashman A, Nagdee M (2017). Impacts of climate change on settlements and infrastructure in the coastal and marine environments of Caribbean Small Island Developing States (SIDS). Caribb Mar Clim Chang Rep Card Sci Rev 155–173. [https://assets.publi](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/605066/11._Settlements_and_Infrastructure_combined.docx.pdf) [shing.service.gov.uk/government/uploads/system/uploads/attac](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/605066/11._Settlements_and_Infrastructure_combined.docx.pdf) [hment_data/fle/605066/11._Settlements_and_Infrastructure_](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/605066/11._Settlements_and_Infrastructure_combined.docx.pdf) [combined.docx.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/605066/11._Settlements_and_Infrastructure_combined.docx.pdf)
- Chausson A, Turner B, Seddon D, Chabaneix N, Girardin CJ et al (2020) Mapping the efectiveness of nature-based solutions for climate change adaptation. Glob Change Biol 26:6134–6155. <https://doi.org/10.1111/gcb.15310>
- Chew C, Reager JT, Small E (2020) CYGNSS data map flood inundation during the 2017 Atlantic hurricane season. Sci Rep 8:9336.<https://doi.org/10.1038/s41598-018-27673-x>
- Cohen-Shacham E, Andrade A, Dalton J, Dudley N, Jones M et al (2019) Core principles for successfully implementing and upscaling Nature-based Solutions. Environ Sci Policy 98:20–29. <https://doi.org/10.1016/j.envsci.2019.04.014>
- Cohen-Shacham E, Walters G, Janzen C. Maginnis S (eds) (2016). Nature-based Solutions to address global societal challenges. Gland, Switzerland: IUCN. xiii + 97
- Cooper JAG, Jackson DWT, Gore S (2013) A groundswell event on the coast of the British Virgin Islands: spatial variability in morphological impact. J Coast Res SI 65.<https://doi.org/10.2112/SI65-118.1>
- Duarte CM, Agusti S, Barbier E, Britten GL, Castilla JC et al (2020) Rebuilding marine life. Nature 580:39. [https://doi.org/10.1038/](https://doi.org/10.1038/s41586-020-2146-7) [s41586-020-2146-7](https://doi.org/10.1038/s41586-020-2146-7)
- Duvat V (2013) Coastal protection structures in Tarawa Atoll. Republic of Kiribati Sustain Sci 8(3):363–379. [https://doi.org/](https://doi.org/10.1007/s11625-013-0205-9) [10.1007/s11625-013-0205-9](https://doi.org/10.1007/s11625-013-0205-9)
- Duvat VKE (2020) Human-driven atoll island expansion in the Maldives. Anthropocene 32:100265. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.ancene.2020.100265) [ancene.2020.100265](https://doi.org/10.1016/j.ancene.2020.100265)
- Duvat VKE, Pillet V, Volto N, Krien Y, Cece R et al (2019) High human infuence on beach response to tropical cyclones in small islands: Saint-Martin Island, Lesser Antilles. Geomorphology 325:70–91.<https://doi.org/10.1016/j.geomorph.2018.09.029>
- Duvat VKE, Anisimov A, Magnan AK (2020) Assessment of coastal risk reduction and adaptation-labelled responses in Mauritius Island (Indian Ocean). Region Environ Change 20(4). [https://](https://doi.org/10.1007/s10113-020-01699-2) doi.org/10.1007/s10113-020-01699-2
- Duvat VKE, Magnan AK, Perry CT, Spencer T, Bell JD et al (2021) Risks to future atoll habitability from climate-driven environmental changes. WIREs Clim Change 12;e700. [https://doi.org/](https://doi.org/10.1002/wcc.700) [10.1002/wcc.700](https://doi.org/10.1002/wcc.700)
- Eckstein D, Hutfls M-L, Winges M (2018) Global climate risk index 2019. Who suffers most from extreme weather events? Weatherrelated loss events in 2017 and 1998 to 2017. Think tank. Germanwatch, pp 36
- Eddy TD, Lam VWY, Reygondeau G, Cisneros-Montemayor AM, Greer K et al (2021) Global decline in capacity of coral reefs to provide ecosystem services. One Earth 4:1278–1285. [https://](https://doi.org/10.1016/j.oneear.2021.08.016) doi.org/10.1016/j.oneear.2021.08.016
- Fabian R, Beck M, Potts D (2014) Reef restoration for coastal defense: a review. University of California, Santa Cruz, p 59
- Ferdinand M (2019) Une écologie décoloniale. Le Seuil, Paris, p 464 Ferrario F, Beck MW, Storlazzi CD, Micheli F, Shepard CC et al (2014) The efectiveness of coral reefs for coastal hazard risk reduction and adaptation. Nat Commun 5:3794. [https://doi.org/](https://doi.org/10.1038/ncomms4794) [10.1038/ncomms4794](https://doi.org/10.1038/ncomms4794)
- Ford M, Merrifeld M, Becker J (2018) Inundation of a low-lying urban atoll island: Majuro. Marshall Islands Nat Hazards 91(3):1273– 1297. <https://doi.org/10.1007/s11069-018-3183-5>
- Gheuens J, Nagabhatla N, Pradeep Perera ED (2019) Disaster-risk, water security challenges and strategies in Small Island Developing States (SIDS). Water 11:637. [https://doi.org/10.3390/w1104](https://doi.org/10.3390/w11040637) [0637](https://doi.org/10.3390/w11040637)
- Giardino A, Nederhoff K, Vousdoukas M (2018) Coastal hazard risk assessment for small islands: assessing the impact of climate change and disaster reduction measures on Ebeye (Marshall Islands). Reg Environ Change 18:2237–2248. [https://doi.org/10.](https://doi.org/10.1007/s10113-018-1353-3) [1007/s10113-018-1353-3](https://doi.org/10.1007/s10113-018-1353-3)
- Glavovic BC, Dawson R, Chow W, Garschagen M, Haasnoot M et al (2022) Cross-chapter paper 2: Cities and settlements by the sea. In: *Climate change 2022: impacts, adaptation and vulnerability.* Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Pörtner H-O, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegría A, Craig M, Langsdorf S, Löschke S, Möller V, Okem A, Rama B (eds)]. Cambridge University Press, Cambridge, UK and New York, pp 2163–2194,<https://doi.org/10.1017/9781009325844.019>
- Hoggart SPG, Hanley ME, Parker DJ, Simmonds DJ, Bilton DT et al (2014) The consequences of doing nothing: the effects of seawater fooding on coastal zones. Coast Eng 87:169–182. [https://doi.org/](https://doi.org/10.1016/j.coastaleng.2013.12.001) [10.1016/j.coastaleng.2013.12.001](https://doi.org/10.1016/j.coastaleng.2013.12.001)
- IPCC (2022) Annex II: Glossary. Möller V, van Diemen R, Matthews JBR, Méndez C, Semenov S et al (eds.). In: Climate change 2022: impacts, adaptation and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK and New York, pp. 2897–2930. [https://doi.org/10.](https://doi.org/10.1017/9781009325844.029) [1017/9781009325844.029](https://doi.org/10.1017/9781009325844.029)
- Jackson CW, Bush DM, Neal WJ (2012) Documenting beach loss in front of seawalls in Puerto Rico: pitfalls of engineering a small island nation shore. In: Cooper JAG, Pilkey OH (eds). Pitfalls of shoreline stabilisation: selected case studies. Springer, Dordrecht, pp 53–71.<https://doi.org/10.1007/978-94-007-4123-2>
- Klein N (2018) The battle for paradise. Haymarket Books, Chicago, p 96
- Klöck C, Nunn PD (2019) Adaptation to climate change in Small Island Developing States: a systematic literature review of academic research. J Environ Dev 28(2):196–218. [https://doi.org/10.1177/](https://doi.org/10.1177/1070496519835895) [1070496519835895](https://doi.org/10.1177/1070496519835895)
- Klöck C, Duvat VKE, Nunn PD (2022) Maladaptive difusion? The spread of hard protection to adapt to coastal erosion and fooding along island coasts in the Pacifc and Indian Ocean. Reg Environ Change 22:136.<https://doi.org/10.1007/s10113-022-01989-x>
- Lam NSN, Arenas H, Brito PL, Liu KB (2014) Assessment of vulnerability and adaptive capacity to coastal hazards in the Caribbean Region. J Coast Res 70(10070):473–478. [https://doi.org/10.2112/](https://doi.org/10.2112/SI70-080.1) [SI70-080.1](https://doi.org/10.2112/SI70-080.1)
- Le Cozannet G, Idier D, de Michele M, Legendre Y, Moisan M, et al. (2021) Timescales of emergence of chronic fooding in the major economic center of Guadeloupe. Nat Hazards Earth Syst Sci 21:703–722.<https://doi.org/10.5194/nhess-21-703-2021>
- Le Cozannet G, Bulteau T, Castelle B, Ranasinghe R, Wöppelmann et al (2019) Quantifying uncertainties of sandy shoreline change projections as sea level rises. Sci Rep 9: 42. [https://doi.org/10.](https://doi.org/10.1038/s41598-018-37017-4) [1038/s41598-018-37017-4](https://doi.org/10.1038/s41598-018-37017-4)
- Lirman D, Schopmeyer S (2016) Ecological solutions to reef degradation: optimizing coral reef restoration in the Caribbean and Western Atlantic. PeerJ 4:e2597.<https://doi.org/10.7717/peerj.2597>
- Logan TM, Guikema SD, Bricker JD (2018) Hard-adaptive measures can increase vulnerability to storm surge and tsunami hazards over time. Nat Sustai 1(9):526–530. [https://doi.org/10.1038/](https://doi.org/10.1038/s41893-018-0137-6) [s41893-018-0137-6](https://doi.org/10.1038/s41893-018-0137-6)
- López-Marrero T (2010) An integrative approach to study and promote natural hazards adaptive capacity: a case study of two flood-prone communities in Puerto Rico. Geogr J176(2):150– 163. <https://doi.org/10.1111/j.1475-4959.2010.00353.x>
- Magnan AK, Duvat VKE (2018) Unavoidable solutions for coastal adaptation in Reunion Island (Indian Ocean). Environ Sci Policy 89:393–400. <https://doi.org/10.1016/j.envsci.2018.09.002>
- Magnan AK, Schipper ELF, Burkett M, Bharwani S, Burton I et al (2016) Addressing the risk of maladaptation to climate change. Wiley Interdiscip Rev Clim Change 7(5):646–665. [https://doi.](https://doi.org/10.1002/wcc.409) [org/10.1002/wcc.409](https://doi.org/10.1002/wcc.409)
- Magnan AK, Bell R, Duvat VKE, Ford JD, Garschagen M et al (2023) Status of global coastal adaptation. Nat Clim Change. <https://doi.org/10.1038/s41558-023-01834-x>
- Magnan AK, Garschagen M, Gattuso J-P, Hay JE, Hilmi N et al (2019) Cross-chapter box 9: Integrative cross-chapter box on low-lying islands and coasts. In: Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M et al (eds). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate. Cambridge University Press, Cambridge, UK, and New York, pp 657–674.<https://doi.org/10.1017/9781009157964.009>
- Manes S, Vale MM, Malecha A, Pires APF (2022) Nature-based solutions promote climate change adaptation safeguarding ecosystem services. Ecosyst Serv 55:101439. [https://doi.org/](https://doi.org/10.1016/j.ecoser.2022.101439) [10.1016/j.ecoser.2022.101439](https://doi.org/10.1016/j.ecoser.2022.101439)
- Manuel-Navarrete D, Gomez JJ, Gallopin G (2007) Syndromes of sustainability development for assessing the vulnerability of coupled human-environmental systems. The case of hydrometeorological disasters in Central America and the Caribbean. Global Environ. Change 17(2):207–217. [https://doi.org/10.](https://doi.org/10.1016/j.gloenvcha.2006.07.002) [1016/j.gloenvcha.2006.07.002](https://doi.org/10.1016/j.gloenvcha.2006.07.002)
- Medina N, Abebe YA, Sanchez A, Voljinovic Z (2020) Assessing socioeconomic vulnerability after a hurricane: a combined use of an index-based approach and principal component analysis. Sustainability 12:1452. <https://doi.org/10.3390/su12041452>
- Morris RL, Konlechner TM, Ghisalberti M, Swearer SE (2018) From grey to green: efficacy of eco-engineering solutions for naturebased coastal defence. Glob Change Biol 24:1827–1842. [https://](https://doi.org/10.1111/gcb.14063) doi.org/10.1111/gcb.14063
- Morrison TH, Hughes TP, Adger WN, Brown K, Barnett J, et al. (2019) Save reefs to rescue all ecosystems. Nature 573(7774):333–336. [https://](https://doi.org/10.1038/d41586-019-02737-8) doi. org/ 10. 1038/ [d41586-019-02737-8](https://doi.org/10.1038/d41586-019-02737-8)
- Mycoo M, Wairiu M, Campbell D, Duvat V, Golbuu Y et al (2022) Small islands. In: Pörtner H-O, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K et al (eds). Climate change 2022: impacts, adaptation, and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, UK, and New York, pp 2043–2121.<https://doi.org/10.1017/9781009325844.017>
- Mycoo M (2018) Beyond 1.5°C: vulnerabilities and adaptation strategies for Caribbean Small Island Developing States. Reg. Environ Change 18: 2341–2353. [https://doi.org/10.10007/](https://doi.org/10.10007/s10113-017-1248-8) [s10113-017-1248-8](https://doi.org/10.10007/s10113-017-1248-8)
- Narayan S, Beck MW, Reguero BFG, Losada IJ, van Wesenbeeck B, et al. (2016) The efectiveness, costs and coastal protection benefts of natural and nature-based defences. PLoS ONE 11(5):e0154735. <https://doi.org/10.1371/journal.pone.0154735>
- Nesshöver C, Assmuth T, Irvine KN, Rusch GM, Waylen KA et al (2017) The science, policy and practice of nature-based solutons: an interdisciplinary perspective. Sci Total Environ 579:1215–1227. <https://doi.org/10.1016/j.scitotenv.2016.11.106>
- Nicholls RJ (2018) Adapting to sea-level rise. In: Zommers Z, Alverson K (eds) Resilience: the science of adaptation to climate change. Elsevier, Oxford, pp 13–29
- Nunn PD, Kumar R (2019) Cashless adaptation to climate change: unwelcome yet unavoidable? One Earth 1(1):31–34. [https://doi.](https://doi.org/10.1016/j.oneear.2019.08.004) [org/10.1016/j.oneear.2019.08.004](https://doi.org/10.1016/j.oneear.2019.08.004)
- Nunn PD, Klöck C, Duvat V (2021) Seawalls as maladaptations along island coasts. Ocean Coast Manage 205:105554. [https://doi.org/](https://doi.org/10.1016/j.ocecoaman.2021.105554) [10.1016/j.ocecoaman.2021.105554](https://doi.org/10.1016/j.ocecoaman.2021.105554)
- Oppenheimer M, Glavovic BC, Hinkel J, van de Wal R, Magnan AK et al (2019) Sea level rise and implications for low-lying islands, coasts and communities. In: Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M et al (eds.). IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, Cambridge University Press, Cambridge, United Kingdom and New York,pp 321-445
- Parsons M, Nalau J, Fisher K, Brown C (2019) Disrupting path dependency: making room for Indigenous knowledge in river management. Glob Environ Change 56:95–113. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.gloenvcha.2019.03.008) [gloenvcha.2019.03.008](https://doi.org/10.1016/j.gloenvcha.2019.03.008)
- Petzold J, Magnan AK (2019) Climate change: thinking small islands beyond Small Island Developing States (SIDS). Clim Change 152:145–165. <https://doi.org/10.1007/s10584-018-2363-3>
- Pichler A, Striessnig E (2013) Diferential vulnerability to hurricanes in Cuba, Haiti, and the Dominican Republic: the contribution of education. Ecol Soc 18(3).<https://doi.org/10.5751/ES-05774-180331>
- Pillet V, Duvat VKE, Krien Y, Cécé R, Arnaud G et al (2019) Assessing the impacts of shoreline hardening on beach response to hurricanes: Saint-Barthélemy, Lesser Antilles. Ocean Coast Manage 174:71–91.<https://doi.org/10.1016/j.ocecoaman.2019.03.021>
- Popke J, Rhiney K (2019) Introduction to special issue on the Caribbean after Irma and Maria: climate, development & the posthurricane context. J Extreme Events 6(1):1902001. [https://doi.org/](https://doi.org/10.1142/S2345737619020019) [10.1142/S2345737619020019](https://doi.org/10.1142/S2345737619020019)
- Ranasinghe R, Ruane AC, Vautard R, Arnell N, Coppola E et al (2021) Climate change information for regional impact and for risk assessment. In: Masson-Delmotte V, Zhai P, Pirani A, Connors SL, Péan C et al (eds). Climate change 2021: the physical science basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK and New York, pp 1767–1926.<https://doi.org/10.1017/9781009157896.014>
- Ratter B, Hennig A, Zahid (2019) Challenges for shared responsibility: political and social framing of coastal protection transformation in the Maldives. J Geogr Soc Berlin 3(150):169–183. [https://doi.](https://doi.org/10.12854/erde-2019-426) [org/10.12854/erde-2019-426](https://doi.org/10.12854/erde-2019-426)
- Reddy SM, Guannel G, Griffin R, Faries J, Boucher T et al (2016) Evaluating the role of coastal habitats and sea-level rise in hurricane risk mitigation: an ecological economic assessment method and application to a business decision. Integr Environ Assess Manag 12(2):328–344.<https://doi.org/10.1002/ieam.1678>
- Reguero BG, Beck MW, Agostini VN, Kramer P, Hancock B (2018) Coral reefs for coastal protection: a new methodological approach and engineering case study in Grenada. J Environ Manage 210:146–161. <https://doi.org/10.1016/jenvman.2018.01.024>
- Reid H, Bourne A, Muller H, Podvin K, Scorgie S, Orindi V (2018) A framework for assessing the efectiveness of ecosystem-based approaches to adaptation, pp 207–216 In Part 5: Tools and methods for resilience. London, Elsevier.
- Rey T, Leone F, Candela T, Belmadani A, Palany P et al (2019) Coastal processes and infuence on damage to urban structures during

Hurricane Irma (St-Martin & St-Barthélemy, French West Indies). J Mar Sci Eng 7(7):215.<https://doi.org/10.3390/jmse7070215>

- Robinson S (2017) Climate change adaptation trends in Small Island Developing States. Mitig Adapt Strateg Glob Chang 22(4):669– 691.<https://doi.org/10.1007/s11027-015-9693-5>
- Robinson S, Gilfillan D (2017) Regional organizations and climate change adaptation in Small Island Developing States. Reg Environ Change 17(4):989–1004. [https://doi.org/10.1007/](https://doi.org/10.1007/s10113-016-0991-6) [s10113-016-0991-6](https://doi.org/10.1007/s10113-016-0991-6)
- Robinson S (2018) Adapting to climate change at the national level in Caribbean Small Island Developing States. Isl Stud J 13(1):79– 100.<https://doi.org/10.24043/isj.59>
- Ruangpan L, Vojinovic Z, Di Sabatino S, Leo LS, Capobianco V et al (2020) Nature-based solutions for hydro-meteorological risk reduction: a state-of-the-art review of the research area. Nat Hazards Earth Syst Sci 20:243–270. [https://doi.org/10.5194/](https://doi.org/10.5194/nhess-20-243-2020) [nhess-20-243-2020](https://doi.org/10.5194/nhess-20-243-2020)
- Scarano FR (2017) Ecosystem-based adaptation to climate change: concept, scalability and a role for conservation science. PECON 16:65–73. <https://doi.org/10.1016/j.pecon.2017.05.003>
- Schultz JM, Kossin JP, Shepherd JM, Ransdell JM, Walshe R, et al. (2018) Risks, health consequences, and response challenges for small-island-based populations: observations from the 2017 Atlantic Hurricane season. Disaster Med Public Health Prep. <https://doi.org/10.1017/dmp.2018.28>
- Scott D, Simpson MC, Sim R (2012) The vulnerability of Caribbean coastal tourism to scenarios of climate change related sea level rise. J Sustain Tour 20(6):883–898. [https://doi.org/10.1080/09669](https://doi.org/10.1080/09669532.2012.699063) [532.2012.699063](https://doi.org/10.1080/09669532.2012.699063)
- Seraphin H (2018) Natural disasters and destination management: the case of the Caribbean and hurricane Irma. Curr Issues Tourism. <https://doi.org/10.1080/13683500.2017.1422483>
- Stanciof CE, Stojanov R, Kelman I, Nemec D, Landa J et al. (2018) Local perception of climate change impacts in St. Kitts (Caribbean Sea) and Malé, Maldives (Indian Ocean). Atmosphere 9:459. <https://doi.org/10.3390/atmos9120459>
- Temmerman S, Meire P, Bouma TJ, Herman PMJ, Ysebaert T et al (2013) Ecosystem-based coastal defence in the face of global change. Nature 504(7478):79–83. [https://doi.org/10.1038/natur](https://doi.org/10.1038/nature12859) [e12859](https://doi.org/10.1038/nature12859)
- Tiggeloven T, De Moel H, Winsemius HC, Eilander D, Erkens G et al (2020) Global scale beneft-cost analysis of coastal food adaptation to diferent food risk drivers. Nat Hazards Earth Sys 20:1025–1044.<https://doi.org/10.5194/nhess-2019-330>
- Wilson AMW, Forsyth C (2018) Restoring near-shore marine ecosystems to enhance climate security for island ocean states: aligning international processes and local practices. Mar Policy 93:284– 294.<https://doi.org/10.1016/marpol.2018.01.018>
- Wynne S, Cambers G, Richardson R, Salisbury O (2016) The Anguilla beach monitoring program report 1992–2014: temporal analysis of twenty-two years of monitoring data. Anguilla Fisheries and Marine Resources Research Bulletin No.1.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.