

A genomic duplication spanning multiple P450s contributes to insecticide resistance in the dengue mosquito Aedes aegypti

Tiphaine Bacot, Chloé Haberkorn, Joseph Guilliet, Julien Cattel, Mary Kefi, Louis Nadalin, Jonathan Filee, Frederic Boyer, Thierry Gaude, Frederic Laporte, et al.

▶ To cite this version:

Tiphaine Bacot, Chloé Haberkorn, Joseph Guilliet, Julien Cattel, Mary Kefi, et al.. A genomic duplication spanning multiple P450s contributes to insecticide resistance in the dengue mosquito Aedes aegypti. 2024. hal-04745935

HAL Id: hal-04745935 https://hal.science/hal-04745935v1

Preprint submitted on 21 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

A genomic duplication spanning multiple P450s contributes to insecticide resistance in the dengue mosquito *Aedes aegypti*

Tiphaine Bacot¹, Chloé Haberkorn^{§, 1}, Joseph Guilliet^{§, 1, 2}, Julien Cattel^{§, 1}, Mary Kefi^{§, 3}, Louis Nadalin¹, Jonathan Filee², Frederic Boyer¹, Thierry Gaude¹, Frederic Laporte¹, Jordan Tutagata¹, John Vontas^{3,4}, Isabelle Dusfour⁵, Jean-Marc Bonneville^{#, 1}, Jean-Philippe David^{#, 1, *}

equal contribution

§ equal contribution

* corresponding author

¹ Laboratoire d'Ecologie Alpine (LECA, UMR 5553), Université Grenoble-Alpes (UGA), Université Savoie Mont-Blanc (USMB), CNRS, 38041 Grenoble, France.

² Université Paris-Saclay, CNRS, IRD, UMR Évolution, Génomes, Comportement et Écologie, 91198 Gif-sur-Yvette, France.

³ Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology Hellas, Heraklion, Greece.

⁴ Pesticide Science Laboratory, Department of Crop Science, Agricultural University of Athens, Athens, Greece.

⁵ Vectopôle Amazonien Emile Abonnenc, Institut Pasteur de la Guyane, Cayenne, France ; Global Health Department, Institut Pasteur, Paris, France.

* jean-philippe.david@univ-grenoble-alpes.fr

Abstract

1 Resistance of mosquitoes to insecticides is one example of rapid adaptation to anthropogenic 2 selection pressures having a strong impact on human health and activities. Target-site modification 3 and increased insecticide detoxification are the two main mechanisms underlying insecticide 4 resistance in mosquitoes. While target-sites mutations are well characterised and often used to track 5 resistance in the field, the genomic events associated with insecticide detoxification remain partially 6 characterised. Recent studies evidenced the key role of gene duplications in the over-expression of 7 detoxification enzymes and their potential use to track metabolic resistance alleles in the field. 8 However, such genomic events remain difficult to characterise due to their complex genomic 9 architecture and their co-occurrence with other resistance alleles. In this concern, the present work 10 investigated the role of a large genomic duplication affecting a cluster of detoxification enzymes in 11 conferring resistance to the pyrethroid insecticide deltamethrin in the mosquito Aedes aegypti. 12 Two isofemale lines originating from French Guiana and being deprived from major target-site 13 mutations showed distinct insecticide resistance levels. Combining RNA-seq and whole genome pool-14 seq identified a 220 Kb genomic duplication enhancing the expression of multiple contiguous 15 cytochrome P450s in the resistant line. The genomic architecture of the duplicated loci was 16 elucidated through long read sequencing, evidencing its transposon-mediated evolutionary origin. 17 The involvement of this P450 duplication in deltamethrin survival was supported by a significant 18 phenotypic response to the P450 inhibitor piperonyl butoxide together with genotype-phenotype 19 association and RNA interference. Experimental evolution suggested that this P450 duplication is 20 associated with a significant fitness cost, potentially affecting its adaptive value in presence of other 21 resistance alleles. 22 Overall, this study supports the importance of genomic duplications affecting detoxification enzymes 23 in the rapid adaptation of mosquitoes to insecticides. Deciphering their genomic architecture

provides new insights into the evolutionary processes underlying such rapid adaptation. Such

25 findings and provides new tools for the surveillance and management of resistance in the field.

26 Key words: Mosquito; Insecticide resistance; P450; Gene duplication; Kdr mutation; Aedes aegypti

27

28 Introduction

29 Natural populations experience a wide range of selective pressures, leading to the accumulation of 30 locally adaptive features and the expression of complex phenotypes (Orr, 2005). Among them, those 31 driven by human activities promote novel and strong selective pressures. Understanding the genetic 32 mechanisms allowing populations to quickly respond to rapid environmental changes has become a 33 major goal (Hendry et al., 2008, 2017; Palumbi, 2001). Resistance of insects to insecticides is a key 34 example of rapid evolution under strong anthropogenic selective pressures. This adaptation has 35 occurred quickly and independently in a large number of taxa with frequent parallel trajectories 36 (Ffrench-Constant et al., 2004; Liu, 2015). As a consequence of local and temporal variations of the 37 selection pressures, the genetic modifications affecting resistant populations are often complex and 38 combine several resistance mechanisms that are potentially additive, each of them bringing variable 39 fitness costs. In addition, the very strong selection pressure exerted by insecticides is prone to 40 produce bottleneck effects. Such complexity makes the identification of resistance alleles and the 41 assessment of their respective contribution challenging (Ffrench-Constant et al., 2004; Li et al., 2007). 42 Besides the understanding of the genetics of rapid adaptation and the origins of complex traits, 43 deciphering the molecular bases of insecticide resistance is also essential for improving pest 44 management strategies (Hawkins et al., 2018). Among taxa of major economic and medical 45 importance, mosquitoes represent a major threat for public health worldwide because of their ability 46 to transmit human viruses and pathogens (Lounibos, 2002). Among them, Aedes aegypti is of 47 particular importance because of its wide distribution and its capacity to transmit Yellow fever, 48 Dengue, Zika and Chikungunya viruses (Brown et al., 2014). These arboviruses are now re-emerging 49 worldwide following the expansion of mosquitoes' distribution area as a consequence of global 50 warming, global transportation network and land perturbations (Kraemer et al., 2015). Although 51 efforts are invested in improving vaccine-based prevention strategies (Carvalho & Long, 2021; Garg 52 et al., 2020; Silva et al., 2018), vector control remains the cornerstone of arboviral diseases control. 53 However, decades of insecticide usage have led to the selection and spread of insecticide resistance 54 in this mosquito species, affecting all public health insecticides, including the most-used pyrethroids 55 (Moyes et al., 2017). High pyrethroid resistance has been shown to reduce vector control efficacy 56 (Dusfour et al., 2011; Marcombe, Carron, et al., 2009; Marcombe et al., 2011; Valle et al., 2019). 57 Though greener vector control strategies are being developed, insecticides will likely remain a key 58 component of integrated vector control in high transmission areas for the next decades (Achee et al., 59 2019). In this concern, identifying the genetic factors underlying resistance is crucial for tracking 60 resistance alleles in the field and making a better use of the few authorised public health insecticides 61 through resistance management actions (Cattel et al., 2020; Corbel & N'Guessan, 2013; Dusfour et 62 al., 2019). 63 In mosquitoes, resistance to chemical insecticides is mainly caused by genetic changes decreasing the

64 affinity of the insecticide for its target (target-site resistance), decreasing its penetration (cuticular 65 resistance), or increasing its detoxification through complex biochemical pathways (metabolic 66 resistance) (Li et al., 2007). Pyrethroid insecticides target the neuronal voltage-gated sodium channel 67 (VGSC gene), and the selection of knockdown resistance (kdr) mutations affecting this protein can 68 lead to resistance. Multiple kdr mutations, often combined as haplotypes, have been identified in Ae. 69 *aegypti* with the following ones playing a major role in pyrethroid resistance in South America: 70 Val410Leu, Val1016Ile and Phe1534Cys (Brengues et al., 2003; Haddi et al., 2017; Hirata et al., 2014; 71 Kasai et al., 2022; Saavedra-Rodriguez et al., 2007; Smith et al., 2016; Yanola et al., 2011). Today, 72 these mutations can be tracked in the field using PCR-based diagnostic assays or mass sequencing 73 approaches, providing essential information for resistance management programmes (Melo Costa et 74 al., 2020).

Conversely, the genetic bases of metabolic resistance are far less understood, though it often accounts for a significant part of the phenotype (David et al., 2013). Indeed, the complexity and 77 redundancy of xenobiotic biodegradation pathways often lead to multigenic adaptations that can

78 differ according to the nature and intensity of the selection pressure together with the demographic

and ecological context (Feyereisen et al., 2015; Li et al., 2007). Metabolic resistance to pyrethroids

80 usually results from an increased activity of detoxification enzymes such as cytochrome P450

81 monooxygenases (P450s or *CYPs* for genes), glutathione S-transferases (GSTs)

82 carboxy/cholinesterases (CCEs) and UDP-glycosyl-transferases (UDPGTs) (David et al., 2013;

83 Hemingway et al., 2004; Smith et al., 2016). At the genetic level, this can result from the selection of

84 enzyme variants showing a higher insecticide metabolism rate, or their overexpression through cis-

85 or trans- regulation. Genomic duplications can also contribute to overexpression, and duplications

86 affecting detoxification genes appear frequently associated with insecticide resistance in mosquitoes

87 (Cattel et al., 2020, 2021; Faucon et al., 2015; Weetman et al., 2018).

88 In addition to the genetic complexity of metabolic resistance alleles, their frequent co-occurrence

89 with other resistance mechanisms in natural mosquito populations makes them difficult to

90 characterise. This is the case in French Guiana, where the use of various insecticides, including the

pyrethroid deltamethrin, for decades has led to the selection of highly resistant *Ae. aegypti* populations combining metabolic resistance alleles and multiple *kdr* mutations such as Val410Leu,

Val1016lle or Phe1534Cys (Dusfour et al., 2015; Haddi et al., 2017). Although the over-expression of

several detoxification enzymes was supported by both transcriptomics and proteomics (Dusfour et

al., 2015; Epelboin et al., 2021; Faucon et al., 2017), those most contributing to deltamethrin

96 resistance and the underlying genetic events remain to be identified. A recent study using a

97 composite population from French Guiana resistant to multiple insecticides identified gene copy

98 number variations (CNV) as a probable cause of detoxification enzymes overexpression in this region

99 (Cattel et al., 2020). Through a pool-seq approach targeting >300 candidate genes, this study

identified multiple contiguous P450s from the CYP6 family on chromosome 1 showing an apparent

101 elevated CNV in association with deltamethrin survival. However, despite the use of controlled

102 crosses, the multigenic resistance phenotype of this population (carrying the three above-mentioned

103 Kdr mutations at high frequency together with multiple metabolic resistance alleles) did not allow to 104 conclude about their role in deltamethrin resistance. In addition, the exon-based nature of the

104 conclude about their role in deltamethrin resistance. In addition, the exon-based nature of the 105 sequencing data generated together with the complexity of *Ae. aegypti* genome did not allow

106 resolving the genomic architecture of the duplicated locus.

107 In this context, the present study aimed at confirming the contribution of this P450 duplication in the 108 resistance of Ae. aegypti to deltamethrin and at deciphering its genomic architecture. In order to 109 limit confounding effects from other resistance alleles, two isofemale lines originating from French 110 Guiana and showing contrasted pyrethroid resistance levels but no apparent resistance to 111 organophosphate and carbamate insecticides were used (Epelboin et al., 2021). In addition, both 112 lines lacked the two major kdr mutations Val1016Ile and Val410Leu occurring in South America 113 (Epelboin et al., 2021). Controlled crosses were then used to remove the Phe1534Cys kdr mutation from the resistant line, which retained a P450-mediated resistance phenotype. Then, RNA-seq and 114 115 whole genome pool-seq confirmed the presence of a ~200 Kb genomic duplication enhancing the 116 expression of a cluster of multiple P450 genes in the resistant line. The genomic architecture of the 117 duplicated loci was elucidated through long read sequencing, providing clues about its evolutionary 118 origin. Association studies, experimental evolution and reverse genetic were then used to further

119 investigate the contribution of this duplicated allele to deltamethrin resistance.

120

121 **Results**

122 Deltamethrin resistance is associated with P450 activity and is autosomal

123 Two isofemale lines from Ile Royale island (French Guiana) showing contrasted resistance

124 phenotypes, namely IR03 and IR13 lines, were used as starting material (Table 1). Comparative

deltamethrin bioassays confirmed that the IR13 line can be categorized as susceptible to

deltamethrin as previously shown (Epelboin et al. 2021), though its susceptibility was slightly lower

127 than the laboratory strain Bora-Bora that was maintained for decades in insectary. Mating IR03

128 genotyped individuals allowed producing the IROF line deprived of the three kdr mutations majorly

associated with pyrethroid resistance in South America (*i.e.* Val410Leu, Val1016lle, Phe1534Cys).

130 However, our attempts to remove the only *kdr* mutation remaining, Ile1011Met were not successful

131 (see below). Bioassays showed that the IROF line remained highly resistant to deltamethrin with only

132 44% mortality to a high dose of insecticide. Mortality levels remains similar between the IR03 and the

133 IROF lines suggesting that the removal of the Phe1534Cys *kdr* mutation did not affect the resistance

134 phenotype, or was compensated by the retaining of the Ile1011Met *kdr* mutation.

135

136 Table 1. Origin, resistance status and *kdr* mutations frequencies of the studied lines

Line	Origin, type	Deltamethrin mortality ¹	Resistance status	f(410Leu) 2	f(1016 e) 2	f(1534Cys) 2	f(1011Met) 3
Bora-Bora	Fr. Polynesia, Lab	100	Susceptible	0	0	0	0
R13	Fr. Guiana, isofemale	95 (91-99)	Susceptible	0	0	0.26	0
IR03	Fr. Guiana, isofemale	43 (26-60)	Resistant	0	0	0.26	0.33
ROF	IR03 cross	44 (34-54)	Resistant	0	0	0	0.50

137 ¹ Mortality rates (mean ±SD) previously obtained on 3 days-old females using 0.05% deltamethrin for 40 min (N=10).

138 ² Kdr mutation frequencies estimated by qPCR individual genotyping (N=30) and later confirmed by pool-seq whole genome data.

139 ³ Kdr mutation frequencies estimated from pool-seq whole genome data.

141 The involvement of P450s in the resistance phenotype of the IROF line was then confirmed by

sequentially exposing adult mosquitoes to the P450 inhibitor piperonyl butoxide (PBO) and to the

insecticide. Such PBO pre-exposure did not significantly affect deltamethrin survival in the

susceptible IR13 line, but increased mortality by 20% in the IR0F line (Wilcoxon test P value = 0.009),

supporting the contribution of P450-mediated detoxification in the resistance of the IROF line (Figure

146 **1**).





157

As the duplicated P450 locus is located on chromosome 1, which also carries the sex determining
locus, deltamethrin bioassays were performed on F1 and F2 males and females obtained from both *Bora-Bora x IROF'* reciprocal crosses in order to investigate the mode of transmission of resistance.
Such comparative bioassays did not support any significant bias affecting the transmission of
resistance (Supplementary file 1). For both F1 males and females, mortality levels upon deltamethrin
exposure were similar in the two reciprocal crosses and intermediate to those of the parental lines,
indicating the absence of a maternal effect and a semi-dominant phenotype in heterozygotes. Similar

and intermediate mortality levels were again observed in F2 males and females, suggesting the lack

- 166 of a sex transmission bias.
- 167

168 Overexpression of the duplicated CYP6 gene cluster

169 RNA-seq was used to identify differentially transcribed genes between the two resistant lines (IR03 170 and IROF), and the two susceptible lines (IR13 and Bora-Bora). Among the 11268 protein-coding genes passing our coverage filter (82% of all protein coding genes), 84 genes showed a significant and 171 consistent differential transcription level in the four 'resistant Vs susceptible' pairwise comparisons 172 173 (see methods, expression fold change \geq 1.5 and adjusted P value \leq 0.0005). The 39 genes under-174 transcribed in resistant lines did not include any gene belonging to any family known to be associated 175 with insecticide resistance. Conversely, the 45 genes over-transcribed in resistant lines included eight 176 genes belonging to gene families associated with metabolic resistance (Figure 2 and Supplementary file 2). These included five P450s (CYP6BB2 AAEL014893, CYP6-like AAEL026852, CYP6-like 177 178 AAEL017061, CYP6P12 AAEL014891 and CYP9F-like AAEL028635). The four CYP6 genes showed mean 179 fold changes up to 4.0 folds and belong to a cluster of five consecutive CYP6 genes located at ~271

- 180 Mb on chromosome 1. Among them, *CYP6P12* (AAEL014891) was annotated as a single accession in
- 181 the reference genome while both protein sequence and RNA-seq data support the presence of two
- distinct genes hereafter referred to as *CYP6P12v1* and *CYP6P12v2* when appropriate.



183

184 Figure 2. Genes over transcribed in resistant lines and their associated copy number variations. Only the 45 genes found 185 significantly over transcribed from RNA-seq are shown (FC \ge 1.5 and corrected P value \le 0.0005 in all resistant versus 186 susceptible pairwise comparisons). Genes also showing an elevated copy number from WG DNA-seq (FC ≥ 1.5) in all 187 pairwise comparisons are indicated with stars. Genomic coordinates (chromosome and start position), accession number 188 and description are indicated for all genes. Gene descriptions in grey were obtained from a BlastP against NCBI Refseq. 189 Genes in red belong to a single duplicated region on chromosome 1. Pool-seg read coverage profiles of the susceptible IR13 190 and the resistant IROF lines around the duplicated locus are shown. Pie charts show the proportion of P450s in each 191 dataset.

192

193 Whole genome Pool-seq short read data were then used to identify genes affected by copy number 194 variations in association with resistance base on their normalised exonic coverage. This analysis 195 identified 22 genes showing a congruent elevated copy number (CN) in resistant Vs susceptible lines 196 (FC \ge 1.5 in all *'resistant Vs susceptible'* pairwise comparisons). These included seven P450s. One of 197 them (AAEL009018) was located at ~58.8 Mb on chromosome 1 and was not detected as over-198 transcribed from RNA-seq. The six others were located at ~271 Mb on chromosome 1 and included 199 the five CYP6s found over-transcribed in resistant lines (see above) and another CYP6 (CYP6CC1) from 200 the same gene cluster that was filtered out from RNA-seq analysis due to its low transcription level 201 (Figure 2 and Supplementary file 2). The over-transcribed CYP9F-like located on chromosome 3 202 showed a slight increased CN in the IROF line Vs the Bora-Bora line (1.6 fold) but such increased CN 203 was not confirmed for the three other 'resistant Vs susceptible comparisons' (1.07, 1.4 and 1.2 fold 204 for IR03/IR13, IR0F/IR13 and IR03/Bora respectively). A closer look at the genomic coverage at the 205 CYP6 locus on chromosome 1 revealed a ~2 folds increased coverage affecting a region of ~200 Kb 206 spanning the entire CYP6 cluster. Such duplication was observed in all resistant Vs susceptible 207 comparisons. Altogether, these genomic data showed that the deltamethrin resistance phenotype 208 observed in the resistant lines is associated with the presence of a large duplication on chromosome 209 1 affecting the transcription level of six clustered CYP6 genes.

Whole genome short reads data confirmed the absence of the three major *kdr* mutations Val410Leu,
Val1016Ile and Phe1534Cys in the IROF resistant line, while another *kdr* mutation (Ile1011Met) was
identified at an exact 0.5 frequency (141/282 short reads supporting the 1011Met allele). Indeed,

both short read and long read data confirmed that this lle1011Met *kdr* mutation is affected by an

214 heterogenous genomic duplication (see below and in Martins et al., 2013). Therefore, a 0.5

215 frequency indicates that the Ile-Met/Ile-Met genotype is fixed in the IROF line.

216

217 Genomic architecture of the duplicated loci

218 Though not the major aim of the present study, our genomic data allowed characterizing the 219 architecture of the duplication affecting the Ile1011Met Kdr mutation in the IROF line. Both short 220 read and long read topologies confirmed that the Ile1011Met kdr mutation is part of 125 Kb genomic 221 duplication covering the 21 last coding exons of the VGSC gene AAEL023266 (Supplementary file 3). 222 The breakpoints of this partial gene duplication are located at 315,905,811 bp and 316,030,250 bp on 223 the reference genome AaegL5. The partial 3' copy carries the wild type allele (1011lle) while the full-224 length copy carries the resistant allele (1011Met). The two copies also differed by their intronic 225 sequence with the partial copy bearing a B type intron and the full-length copy bearing a A type 226 intron as previously described (Martins et al., 2013). As expected from an incomplete gene 227 duplication, RNA-seq data supported the sole expression of the 1011Met allele meaning that the 228 IROF line (genotype Ile-Met/Ile-Met) express a Met phenotype. 229 The genomic structure of the P450 duplicated region identified on chromosome 1 was further

230 investigated in light of long reads data obtained from the IR13 susceptible line and the IR0F resistant 231 line. This revealed the presence of a 220.4 Kb duplication in the IROF line, with the two tandem 232 repeats separated by a 6 Kb insertion (Figure 3). The right breakpoint of the duplication (RB) was 233 found ~50 Kb right to CYP6CC1 at position 271,472,082 bp, and the left breakpoint (LB) ~77 Kb left to 234 CYP6BB2 at 271,251,705 bp. These breakpoints were supported by split reads from both long and 235 short read data (Supplementary file 4). The duplication is flanked by two distinct transposons, each 236 having its own terminal inverted repeats (TIR): a 2.4 Kb PiggyBac-like transposon on the right (PYL), 237 and 9 Kb hAT-related transposon on the left (hAT-CYP6). These transposons are described in more

- 238 detail in **Supplementary file 5**.
- 239



Figure 3. Genomic architecture of the duplication. The topology was deduced from both short and long reads sequencing
 data. The six CYP6 genes carried by the duplication in the resistant lines are represented by red arrows. Regions containing
 full or partial transposable elements are shown in greater detail. Orange and yellow triangles denote the inverted terminal
 repeats from hAT (17 pb) and PiggyBac-like (18 pb) elements respectively. Diagram is not to scale.

246 The central sequence joining the tandem copies is chimeric, with 60 bp from the left end of PYL 247 transposon followed by 6001 bp from the right end of the hAT transposon. This duplication is absent 248 from the IR13 susceptible line (no read joining RB and LB sequences and no PYL transposon 249 sequence, neither full nor partial at RB). However, the hAT-CYP6 transposon was identified at the LB 250 in one IR13 haplotype, much like in the IROF line, while a second haplotype was devoid of hAT 251 insertion, much like the reference genome. An excess of tri-allelic SNPs is expected in the duplicated 252 region if the two copies have diverged. Pool-seq short read data were then used to compare the 253 frequency of tri-allelic loci within the duplication and 100 Kb upward and downward. Among the 254 1401 substitutions sites identified within the duplication, five were tri-allelic whereas no such tri-255 allelic variant was identified among the 1334 substitutions identified in the flanking regions. This 256 represents a slight but significant enrichment in the duplicated region (Fisher test p value = 0.031). 257 These tri-allelic sites were all located within a 11.2 Kb intergenic region between the two CYP6-like 258 genes AAEL017061 and AAEL014891v1.

259

240

The P450 duplication and the Kdr 1011 mutation are both associated with deltamethrin
 survival

262 The association of the P450 duplication and the kdr lle1011Met mutation with deltamethrin survival 263 were investigated in F2 individuals obtained from both 'Bora-Bora x IROF' reciprocal crosses. F2 264 females were then exposed to a high dose of deltamethrin (85.5% mortality) before quantifying 265 *CYP6BB2* gene copy number and the frequency of the 1011Met *kdr* allele in dead and survivors using 266 ddPCR (Figure 4). As expected in the presence of a fixed duplication, a 2-fold increase of CYP6BB2 267 copy number was observed in the IROF line $(4.31 \pm 0.21 \text{ copies})$ as compared to Bora-Bora line (2.07 ± 1.01) 268 0.10 copies) while F1 individuals showed an intermediate copy number (3.58 ± 0.14 copies). In F2 269 individuals, CYP6BB2 copy number was higher in survivors than in dead individuals (4.00 ± 0.16) 270 copies versus 3.31 ± 0.15 copies) supporting a genetic linkage between the P450 duplication and 271 deltamethrin survival. Insecticide survival was also linked to the kdr mutation lle1011Met on 272 chromosome 3. In line with an heterogenous kdr duplication, the frequency of the 1011Met allele 273 was close to 0.5 in the IROF resistant line (genotype Ile-Met/Ile-Met) and close to 0.33 in F1



individuals (genotype Ile-Met/Ile). In deltamethrin-exposed F2 individuals, the Met allele frequency was higher in survivors than in dead individuals (0.41 ± 0.02 versus 0.30 ± 0.02).

Figure 4. Genotype-phenotype association study. Top left panel shows the crossings underlying this experiment. Top right panel shows mortality data obtained by exposing adult females from the IROF and Bora-Bora lines and their F1 and F2 progeny to 0.03% deltamethrin for 1 hour. Mean mortalities ± SD 24h after insecticide exposure are shown and distinct letters indicate significant differences (Kruskal-Wallistest followed by post hoc Wilcoxon test with Bonferroni-Holm correction, $N \ge 5$, $p \le 1$ 0.05). Bottom left and right panels show the copy number of CYP6BB2 and the frequency of the kdr 1011Met allele respectively as inferred by ddPCR from pools of individuals (mean ± 95% Cl, Bora:



300

301 Multiple CYP genes carried by the P450 duplication may contribute to resistance

302 The relative importance of the P450s carried by the duplication in deltamethrin survival was then 303 examined by RNA interference. Direct dsRNA injections in adult females of the IROF line were used to 304 specifically knock down the five CYP6 genes showing sufficient expression to be detected by RNA-seq 305 (CYP6BB2, CYP6-like AAEL026582, CYP6-like AAEL017061, CYP6P12v1 and CYP6P12v2). Such 306 approach allowed reaching an acceptable silencing specificity and a moderate silencing efficiency 307 (from 42.2% for AAEL017061 to 69.7% for AAEL026582). Comparative deltamethrin bioassays 308 performed on dsCYP6-injected mosquitoes and dsGFP-injected controls suggested that the two first 309 CYP gene of the duplicated cluster (CYP6BB2 and CYP6-like AAEL026582) contribute to deltamethrin 310 survival (see methods and results in Supplementary file 6). However, data generated across multiple 311 injection experiments were not fully conclusive because a high mortality was frequently observed in 312 dsGFP-injected controls and because the increased mortalities observed in dsCYP-injected 313 mosquitoes were relatively low. Such low mortality variations upon dsCYP6 injection were likely due 314 to the moderate silencing efficiency and the presence of the kdr 1011Met allele in the IROF line.

315

316 The P450 duplication is hardly retained by selection in presence of the Kdr 1011 allele

The co-occurence of the P450 duplication and the Ile1011Met *kdr* mutation in the IROF resistant line was taken as an opportunity to compare their responses to deltamethrin selection. Four independent lines obtained from 'Bora-Bora x IROF' F3 offspring and representing two replicated selection regimes (selected lines: SelA and SelB, non-selected lines: NSA and NSB) were compared for their resistance level, *CYP6BB2* copy number and *kdr* 1011Met allele frequency (**Figure 5**).

322





348 An increase of deltamethrin resistance was observed for both Sel lines, with mortality dropping from 349 ~80% in F4 to less than 20% in F9. Relaxing deltamethrin selection for three successive generations 350 (F9 to F11) lead to an increased mortality in both Sel lines, suggesting that the resistance phenotype 351 is costly. Though subjected to sampling effects, ddPCR data showed that the kdr 1011Met allele was 352 rapidly selected by deltamethrin. Its frequency was close to 0.5 in F9 of both Sel lines, supporting the 353 near fixation of the Ile-Met duplicated allele. Such increased frequency was not seen in NS lines 354 where Met allele frequency decreased to less than 10% in one line and fluctuated under 0.25 in the 355 other line. No such strong response to selection was observed for the P450 duplication, with 356 CYP6BB2 copy number showing a gradual decrease through generations in both Sel and NS lines. 357 However, CYP6BB2 mean copy number were significantly higher in Sel lines than in NS lines across 358 generations, supporting a higher retention rate of the duplicated allele upon deltamethrin selection 359 (Linear Mixed Effect Model, p value = 0.0088).

360

361 **Discussion**

362 In line with their high probability of occurrence, genomic duplications have been shown to frequently 363 contribute to short-term adaptation in various organisms (Kondrashov, 2012). In arthropods facing 364 strong selection pressures from insecticides, duplications are known to be adaptive via two distinct 365 mechanisms (Bass & Field, 2011). First, heterogeneous duplications affecting key insecticide target 366 genes and tethering both susceptible and resistant alleles have been associated with a reduction of 367 fitness costs (Labbe et al., 2007). Second, duplications of detoxification genes can contribute 368 quantitatively to their over-expression, leading to resistance through increased insecticide 369 metabolism (Faucon et al., 2015; Weetman et al., 2018). As opposed to target-site resistance 370 mutations that can be easily tracked by PCR in natural populations, the paucity of DNA markers 371 underlying metabolic resistance hinders their tracking in the field. In this concern, identifying gene 372 duplications associated with metabolic resistance can provide new tools to monitor the dynamics of 373 resistance alleles (Cattel et al., 2021).

374

375 The P450 duplication affects a cluster of genes previously associated with resistance

376 The present study confirmed the occurrence of a large genomic duplication affecting a cluster of six 377 P450s from the CYP6 family in Ae. aegypti populations from French Guiana. The presence of such a 378 duplication was previously suspected though the structure of the duplicated loci could not be 379 resolved from targeted-sequencing data (Faucon et al., 2015). The association of this P450 380 duplication with deltamethrin resistance was supported by controlled crosses, although the presence 381 of other resistance alleles limited the association power (Cattel et al., 2020). A duplication 382 encompassing the CYP6BB2 gene was also associated with pyrethroid resistance in Lao PDR 383 (Marcombe et al., 2019). Though gene duplication was not looked for, the over-transcription of CYP6 384 genes from this cluster (e.g. CYP6BB2, CYP6P12 and CYP6CC1) was frequently associated with 385 pyrethroid resistance worldwide (Bariami et al., 2012; Dusfour et al., 2015; Goindin et al., 2017; Kasai 386 et al., 2014; Marcombe, Poupardin, et al., 2009; Moyes et al., 2017; Reid et al., 2014; Saavedra-387 Rodriguez et al., 2012; Seixas et al., 2017). These genes were also found under directional selection in 388 association with pyrethroid survival in a Mexican resistant line from whole exome SNP data 389 (Saavedra-Rodriguez et al., 2021). In addition, CYP6BB2 was also shown capable of metabolising the 390 pyrethroid permethrin (Kasai et al., 2014). By comparing its copy number and transcriptional level 391 across different lines, Faucon et al. (2017) suggested that both transcriptional regulation and 392 genomic duplication contribute to its over-expression in South America. Interestingly, this gene also 393 responded to selection with the neonicotinoid imidacloprid, suggesting that P450s from this cluster 394 can be selected by and metabolise other insecticides (Riaz et al., 2013; Zoh et al., 2021).

395 Our attempt to isolate this P450 duplication from other resistance alleles occurring in French Guiana 396 was partially successful. Indeed, both isofemale lines were naturally deprived from the major Val1016Iso and Val410Leu kdr mutations, and controlled crosses allowed removing the Phe1534Cys 397 398 mutation. However, the resulting IROF resistant line carrying the P450 duplication still carried the 399 Ile1011Met kdr mutation as a fixed heterogenous duplication (Ile-Met/Ile-Met genotype, Met/Met 400 phenotype). There is evidence that this mutation confers some resistance to pyrethroids in the field 401 (Brengues et al., 2003; Brito et al., 2018; Martins et al., 2009). Nevertheless, the use of the P450 402 inhibitor PBO showed that the resistance phenotype was still associated with P450-mediated 403 metabolism. Considering that four of the five P450s over-transcribed in resistant lines belong to the 404 duplicated locus, the contribution of this P450 duplication to the resistance phenotype is likely. 405 Studying the mode of transmission of resistance did not allow evidencing any significant sexual 406 transmission bias potentially associated with resistance. In line with this, this P450 duplication is 407 located outside of a 210 Mb region of chromosome 1 showing a low recombination rate and a high 408 genetic differentiation between Ae. aegypti males and females (Fontaine et al., 2017).

409

410 Contrasted architectures of the duplications affecting the P450 and the Kdr loci

411 The presence of two distinct transposons at the P450 duplication breakpoints implies that they have 412 played an active role in the generation of this genomic event. Transposon-mediated duplication 413 often arises from ectopic recombination between two homologous transposon copies inserted in the 414 same orientation at distinct positions of a same chromosome (Baker et al., 1996). However, such an 415 event would leave a single full-length transposon copy between the two duplicated genomic regions 416 with no transposon at other breakpoints (Remnant et al., 2013). Here, a more complex scenario must 417 have taken place because two distinct transposons are flanking the duplication, and because the 418 sequence joining the two copies is chimeric. A possible scenario involves independent insertion of 419 the two distinct transposons, then transposition of the hAT into the PiggyBac-like transposon 420 followed by a deletion event, and finally an ectopic homologous recombination event between the 421 two duplicated hAT sequences (Supplementary File 7). One transposon involved is related to the hAT 422 superfamily while the other one is distantly related to PiggyBac (see Supplementary file 5 for more 423 details), both being cut and paste elements having multiple copies and being active in the Ae. aegypti 424 genome (Nene et al., 2007). The divergence of the two tandem copies was supported by the 425 identification of a few tri-allelic substitution sites within the duplicated region, most likely implying 426 mutations occurring post-duplication. The few tri-allelic sites observed represent only a fraction of 427 those, because they imply a pre-duplication mutation at the same position, and because mutations 428 reverting to the reference allele are not detected as such. Nevertheless, such low divergence rate

429 between the two copies supports the recent age of this duplication event.

In contrast, the genomic duplication affecting the VGSC gene at the kdr Ile1011Met locus shows no
remnant of any transposon at breakpoints, suggesting a different genesis than the P450 duplication.
It is likely to be an earlier event, since the copy divergence is much more pronounced, particularly in
intronic regions (Martins et al, 2013). Also, it differs from other heterogenous duplications implicated
so far in insecticide resistance because the incomplete copy carrying the susceptible Ile1011

- 435 pseudoallele, does not contribute to gene expression. It therefore cannot modulate the resistance
- 436 level provided by the 1011Met allele carried by the full copy, nor its fitness. This contrasts with the
- 437 heterogenous duplications affecting the Ace1 target-site mutation observed in *Culex pipiens sp.* and
- 438 Anopheles gambiae, where both copies can contribute to the phenotype, thus allowing a reduction
- 439 of the fitness cost associated with the resistant allele in absence of insecticide (Djogbenou et al.,
- 440 2009; Labbe et al., 2007). In this regard, further investigating the evolutionary origin and dynamics of
- 441 this *kdr* heterogenous duplication deserves further attention.
- 442

443 The P450 duplication has a limited adaptive value in presence of the Kdr 1011 mutation

444 The association study performed on F2 individuals supported the association of the P450 duplication 445 with deltamethrin survival together with the IIe1011Met kdr mutation. The association of the 446 Ile1011Met kdr mutation with deltamethrin survival was expected. As the VGSC gene carrying kdr 447 mutations is located on chromosome 3, a genetic linkage with the P450 duplicated loci is excluded. 448 Therefore, the association of the P450 duplication in F2, following limited recombination events, 449 implies a resistance locus located on chromosome 1, either the P450 duplication itself or another 450 genetically linked locus. However, no other known resistance gene was identified in this genomic 451 region. The relative importance of the different CYP6 genes carried by the P450 duplication in 452 deltamethrin survival was investigated through RNA interference. Despite a limited statistical power, 453 likely due to the presence of the kdr lle1011Met resistance allele and moderate knock down 454 efficiencies, these data suggest that at least two CYP6 genes (CYP6BB2 and AAEL026582) carried by 455 the duplication may contribute to deltamethrin detoxification.

456 Experimental evolution confirmed the strong positive response of the kdr 1011Met allele to selection 457 with a high dose of deltamethrin. Such experiment also evidenced a gradual decrease of this allele in 458 absence of insecticide selection, supporting a significant fitness cost as previously shown for other 459 kdr mutations (Rigby et al., 2020; Uemura et al., 2023). The response of the P450 gene duplication to 460 deltamethrin selection was less clear. A gradual decrease of CYP6BB2 copy number was observed in 461 both selected and non-selected lines, even though such decay was slower in the selected lines. Such 462 result supports a lower adaptive value of the P450 duplication in our experimental conditions. Such 463 unexpected response may indicate the presence of a significant fitness cost associated with this large 464 genomic duplication. Indeed, genomic duplications can be detrimental for various reasons (Schrider 465 et al., 2013): first, gene duplications can alter the gene dosage balance which can lead to metabolic 466 cost. Second, genomic duplications allow the accumulation of deleterious mutations that are less 467 cleared off by background selection because of the functional redundancy of duplicated copies. 468 Third, large duplications can impair recombination locally. In the present case, metabolic costs due to 469 gene dosage balance and/or a low recombination rate at this locus between the Bora-Bora line (well 470 adapted to laboratory conditions) and the IROF line (less fitted to our laboratory conditions) may have impaired the selection of the P450 duplication and favoured the selection of other resistance 471

alleles such as the *kdr* 1011Met mutation. Finally, considering the fitness costs associated with this

473 *kdr* mutation, its additive cost with the P450 duplication may have prevented their concomitant

selection with the use of a high insecticide dose favouring the selection of the former. Such fitness-

475 cost balance at two distinct resistance loci may reflect the complex interactions occurring between

476 resistance alleles *in natura*, leading to distinct adaptive trajectories depending on selection pressures
477 and demographic effects.

478

479 Conclusions

480 Following previous work, the present study supports the contribution of this P450 gene duplication in 481 pyrethroid resistance. However, deciphering its adaptive value versus other resistance alleles and its 482 dynamics in natural mosquito populations deserves further work. The relative importance of the 483 different CYP6 genes carried by this duplication in resistance to pyrethroids (and possibly to other 484 insecticides) also deserves further validation. Despite this incomplete picture, the nature of the 485 P450s affected by this large genomic duplication and their frequent association with insecticide 486 resistance call for further studying its evolutionary origin and dynamics in response to xenobiotics. In 487 this context, though the flanking of the duplication by repeated transposable elements prevented 488 the design of a specific PCR diagnostic assay targeting the breakpoints, the dual-colour quantitative 489 TaqMan ddPCR assay we developed (see **Supplementary file 8**) represents a good alternative to track 490 this resistance allele in natural mosquito populations.

491

492 Methods

493 Mosquitoes

494 The Ae. aegypti laboratory strain Bora-Bora, fully susceptible to insecticides was used as reference in 495 the present study. The isofemale lines used in the present study were derived from a field population 496 collected in 2015 in the Ile Royale (IR) island (5.287° N; 52.590° W) off the coast of French Guiana as 497 described in Epelboin et al. (2021). Among the isofemale lines isolated from this population, the lines 498 IR13 and IR03 showed contrasted pyrethroid resistance levels with the IR03 line being highly 499 resistant and the IR13 line being susceptible as defined by WHO diagnostic dose bioassay (Table 1) 500 and Epelboin et al. (2021). Both lines were shown to lack the two major voltage-gated sodium 501 channel kdr mutations occurring in this geographical area (Val410Leu and Val1016Ile), yet both 502 carried the Phe1534Cys mutation at a moderate frequency (Epelboin et al., 2021). The proteomic 503 profiling of these lines identified multiple cytochrome P450s enriched in the IR03 resistant line as 504 compared to a susceptible line, supporting the presence of metabolic resistance alleles (Epelboin et 505 al., 2021). All mosquito lines were maintained under standard laboratory conditions (27 ± 2 °C, 70 \pm 506 10 % relative humidity, light/dark cycle 14:10h) and large population size to limit drift effects 507 (N>1000). Larvae were reared in tap water and fed with hay pellets. Adults were maintained in mesh 508 cages and fed with a 10% honey solution. Blood feedings were performed on mice. 509

510 Removal of the Phe1534Cys mutation from the IR03 line

511 In an attempt to isolate metabolic resistance alleles, inter-crosses assisted by genotyping were 512 performed on the IR03 resistant line to create the IR0F line depleted from the Phe1534Cys kdr 513 mutation. A total of 245 IR03 couples were individualised at the pupal stage in eppendorf tubes 514 placed in plastic cups covered by a nylon mesh and allowed to emerge and reproduce. The 515 Phe1534Cys Kdr mutation was first searched for from male exuvia for all couples and then from 516 female exuvia of the 58 couples from which males were negatives for the 1534Cys allele. Genotyping 517 was performed on qPCR using the High Resolution Melt Curve Analysis method as described in 518 Saavedra-Rodriguez et al. (2007). The six couples in which both parents were homozygous wild type

(Phe-Phe) were transferred to a mesh cage, blood fed and allowed to lay eggs. The absence of the
1534Cys allele in the offspring was confirmed by genotyping 40 individuals of both sexes. Adults were
then exposed to 0.05% deltamethrin and survivors were allowed to freely reproduce to generate the

522 IROF resistant line. The IROF line was then maintained in standard insectary conditions under

523 moderate deltamethrin selection pressure (*i.e.* 50% mortality every two generations).

524

525 Deltamethrin bioassays

526 All bioassays were performed by exposing replicates of 20-25 three days-old non-blood-fed adults to 527 deltamethrin impregnated filter papers following the standard WHO procedure, with mortality being 528 recorded 24h after insecticide exposure (WHO, 2022). Different insecticide doses and exposure times 529 were used according to the nature of the experiment and the mosquito tested (susceptibility status, 530 sex, use of enzymatic inhibitor or not) in order to keep mortality in the dynamic range and maximise 531 mortality variations between conditions. The bioassays used for assessing the resistance level of the 532 different lines presented in Table 1 were performed on females using 10 replicates per line and a 533 dose of 0.05% deltamethrin for 40 min. The involvement of P450s in the resistance phenotype 534 presented in Figure 1 was assessed by bioassays on 5 replicates of females exposed or not to 4% 535 piperonyl butoxide (PBO) for 1h prior to deltamethrin exposure using a dose of 0.03% deltamethrin 536 for 30 min. Bioassays used to investigate the mode of inheritance of resistance presented in 537 Supplementary Figure 1 were performed on males and females obtained from the susceptible Bora-538 Bora line and the resistant IROF line together with F1 and F2 individuals obtained from each 539 reciprocal crosses (see below). As males and females naturally show different tolerance to 540 insecticides and F1/F2 individuals were expected to be more susceptible than IROF individuals, 541 deltamethrin exposure conditions were accordingly (0.03% deltamethrin for 30 min for females and 542 0.015% deltamethrin for 25 min). Finally, bioassays used for monitoring the resistance level of the 543 selected and non-selected lines through the experimental evolution experiment presented in Figure 544 5 were performed on 5 to 7 replicates per line per generation using a dose of 0.03% deltamethrin for 545 1h.

546

547 Controlled crosses and association studies

548 In order to investigate the mode of inheritance of resistance, reciprocal crosses between the fully 549 susceptible Bora-Bora line and the resistant IROF line were performed (cross A: 'Bora-Bora females x 550 IROF males' and cross B: 'IROF females x Bora-Bora males'). Crosses were performed with 200 virgin 551 females and males of each line. The susceptibility of F1 and F2 males and females from each cross were then compared using deltamethrin bioassays as described above. The association of the P450 552 553 duplication with deltamethrin survival was studied in F2 females obtained from both reciprocal 554 crosses pooled in equal quantities. F2 females were exposed to 0.03% deltamethrin for 1h leading to 555 88.5% mortality after a 24h recovery time. Dead and surviving females were sampled and stored at -556 20°C until molecular analyses. The response of the P450 duplication and the Ile1011Met kdr 557 mutation were further studied across multiple generations using experimental selection. F3 eggs 558 obtained from both reciprocal crosses were pooled in equal guantity and then randomly split in 4 559 lines: the first two lines (NSA and NSB) were maintained without insecticide selection, while the two 560 other lines (SelA and SelB) were selected with deltamethrin. This duplicated line setup was used in 561 order to control for genetic drift that may occur across generations. Deltamethrin mass selection was 562 performed prior mating on both virgin males and virgin females (N>1000 for each line) at generations 563 F4, F6, F8, F12 and F13. A constant dose of deltamethrin was used through the selection process 564 leading to ~70% mortality in each sex at generation F3 (females: 0.03% for 1h; males: 0.015% for 30 565 min). The resistance level of each line was monitored at generations F5, F9, F12, F13 and F14 on 566 females prior to insecticide selection using standard bioassays (see above). Unexposed adult females 567 from each line were sampled at the same generations and stored at -20°C until molecular analysis. 568

569 RNA-sequencing

570 Gene transcription levels were compared across the four lines (Bora-Bora, IR13, IR03 and IROF) using 571 RNA-seq. For each line, four RNA-seq libraries were prepared from distinct batches of 25 calibrated 572 three-day-old non-blood-fed females not exposed to insecticide. Total RNA was extracted using 573 TRIzol (Thermo Fisher Scientific) following manufacturer's instructions. RNA samples were then 574 treated with RNase-free DNase set (Qiagen) to remove gDNA contamination, purified on RNeasy mini 575 columns (Qiagen) and QC checked using Qubit (Thermo Fisher Scientific) and bioanalyzer (Agilent). 576 RNA-seq libraries were prepared from 500 ng total RNA using the NEBNext[®] Ultra™ II directional RNA 577 library Prep Kit for Illumina (New England Biolabs) following manufacturer's instructions. Briefly, 578 mRNAs were captured using oligodT magnetic beads and fragmented before being reverse 579 transcribed using random primers. Double-stranded cDNAs were synthesised, end-repaired, and 580 adaptors were incorporated at both ends. Libraries were then amplified by PCR for 10 cycles and 581 purified before QC check using Qubit and Bioanalyzer. Libraries were then sequenced in multiplex as 582 single 75 bp reads using a NextSeq500 sequencer (Illumina). An average of 51.4 M reads was 583 generated per library. After unplexing and removing adaptors, sequenced reads from each library 584 were loaded into Strand NGS V3.2 (Strand Life Science) and mapped to the latest Ae. aegypti genome 585 assembly (Aaeg L5) using the following parameters: min identity = 90%, max gaps = 5%, min aligned 586 read length = 25, ignore reads with > 5 matches, 3^{\circ} 2 end read trimming if quality < 20, mismatch 587 penalty = 4, gap opening penalty = 6, gap extension penalty = 1. Mapped reads were then filtered 588 based on their quality and alignment score as follows: mean read quality > 25, max N allowed per 589 read = 5, alignment score \geq 90, mapping quality \geq 120, no multiple match allowed, read length \geq 35. 590 These filtering steps allowed retaining ~75% of reads. Quantification of transcription levels was 591 performed on the 13614 protein-coding genes using the DESeq2 method with 1000 iterations 592 (Anders & Huber, 2010, https://bioconductor.org/packages/release/bioc/html/DESeq2.html). Only 593 the 11268 protein-coding genes showing a normalised expression level ≥ 0.5 (~0.05 RPKM) in all 594 replicates for all lines were retained for further analysis. Differential gene transcription levels 595 between each line across all replicates were then computed using a one-way ANOVA followed by a 596 Tukey post hoc test and p values were corrected using the Benjamini and Hochberg multiple testing 597 correction (Benjamini & Hochberg, 1995). Genes showing a transcription variation \geq 1.5 fold (in 598 either direction) and an adjusted p value ≤ 0.0005 in the four pairwise comparisons between 599 resistant and susceptible lines (i.e. IR03 Vs Bora-Bora, IR03 Vs IR13, IR0F Vs Bora-Bora and IR0F Vs 600 IR13), were considered differentially transcribed in association with deltamethrin resistance.

601

602 Whole genome Pool-sequencing

603 The genome of the four lines were compared using short read whole genome pool-seq. For each line, 604 genomic DNA was extracted from two batches of 50 adult females using the PureGene Core Kit A 605 (Qiagen) and the two gDNA extracts were then pooled in equal proportion into a single sequencing 606 library. Whole genome sequencing was performed from 200 ng gDNA. Sequencing libraries were 607 prepared according to the TruSeg DNA Nano Reference guide for Illumina Paired-end Indexed 608 sequencing (version oct. 2017) with an insert size of 550 bp. Sequencing was performed on a 609 NextSeq 550 (Illumina) as 150 bp paired-reads. Sequencing depth was adjusted to reach an average 610 coverage \geq 80X leading to the sequencing of ~713 M reads per library. Reads obtained from each line 611 were loaded into Strand NGS V3.2 and mapped to Ae. aegypti genome (Aaeg L5) using the following 612 parameters: min identity = 90%, max gaps = 5%, min aligned read length = 25, ignore reads with > 5613 matches, 32 end read trimming if quality < 15, mismatch penalty = 4, gap opening penalty = 6, gap 614 extension penalty = 1. Mapped reads were then filtered based on their quality and alignment score 615 as follows: mean read quality > 25, max N allowed per read = 5, alignment score \geq 90, mapping 616 quality \geq 60, no multiple match allowed, read length \geq 100. Finally, interchromosomal split reads and 617 PCR duplicates were removed. These filtering steps allowed retaining ~53% of reads.

618 In order to focus on duplication impacting protein expression, gene copy number variation (CNV) 619 analysis was performed on exons of all protein-coding genes. For each gene, raw coverages were 620 obtained by dividing the number of reads mapped to all (non-overlapping) exons by total exon 621 length. Normalised gene coverages were then obtained by dividing raw gene coverages by the total 622 number of reads passing mapping and quality filters from each sequenced library. In order to limit 623 false positives in low coverage regions, genes showing a normalised coverage value < 1E-8 in at least 624 one line were filtered out (12145 genes retained). As for RNA-seq, genes showing a CNV \geq 1.5 fold (in 625 either direction) in the four pairwise 'resistant vs susceptible' comparisons were considered affected 626 by CNV in association with resistance.

627 The divergence between the two copies of the P450 duplication was investigated through the

628 detection of tri-allelic substitution sites in the IROF resistant line. Substitution were called from reads

629 mapped within a region encompassing the P450 duplication extended by 100 kb on both sides

630 (Chr1:271151705 - Chr1:271572082). Only substitutions showing a coverage >40 reads and minimum

allele frequency = 10% were called, allowing the detection of 1401 and 1334 substitution sites within

and outside the duplicated region respectively. Substitution sites were then considered as tri-allelic

633 if they carried the reference allele together with two distinct variant alleles at the same position,

- 634 each showing a frequency > 20%.
- 635

636 Long read sequencing and de novo assembly of the P450 duplication

637 The genomic architecture of the duplicated locus was further examined in the duplicated IROF line 638 versus the non-duplicated IR13 line. Both lines were submitted to long read sequencing using Oxford 639 Nanopore technology. Genomic DNA was extracted from pools of 75 adult females using the Gentra 640 Puregene Tissue kit (Qiagen) with 3 mL cell lysis buffer. Genomic DNA was quantified using Qubit 641 DNA Broad Range assay (Qiagen) and quality-checked by agarose gel electrophoresis before adjusting 642 concentration to 150 ng/µl. Long genomic DNA fragments were selected using the Short Read 643 Eliminator kit (Circulomics). DNA libraries were prepared using 1 µg gDNA with the ligation 644 sequencing kit SQK-LSK109 kit (Oxford Nanopore) following manufacturer's instructions. Finally, 50 645 fmoles of the resulting DNA library were loaded on a R9.4.1 MinIon flow cell. Long read sequences 646 were collected from 2 independent runs of 48 h for each line. Sequencing generated a total of 647 515K/1017K reads with an average length of 13.5/9.8 Kb, representing 3.4/4.8-fold genome 648 coverage, for IROF and IR13 lines respectively. Reads were mapped on the AaegL5 reference genome 649 using Winnowmap (Jain et al., 2022) and alignments were visualised using IGV (Robinson et al., 650 2011). Finally, de novo genomes of the IROF and IR13 lines were assembled using both short and long 651 reads using MaSuRCA (Zimin et al., 2013). This de novo assembly resulted into 17346/13160 contigs, 652 with a median size of 40/55 kb for each line respectively. Transposable elements were identified 653 using BLAST against the RepBase library (10/12/2021 version).

654

655 Quantification of the P450 duplication

656 The detection of the P450 duplication was achieved through the quantification of CYP6BB2 copy 657 number by digital droplet PCR (ddPCR). Genomic DNA was extracted from pools of mosquitoes using 658 the cetyltrimethylammonium bromide (CTAB) method (Collins et al 1987), resuspended in nuclease-659 free water and quantified using Qubit DNA Broad Range assay (Qiagen). The duplex ddPCR 660 quantification assay was based on the co-amplification of two target genes each detected with 661 specific Taqman probes: the gene CYP6BB2 (AAEL014893, Fam) was used to quantify the P450 662 duplication copy number while the gene CYP4D39 (AAEL007808, Hex) always found as a single copy 663 in various Ae. aegypti lines was used as control for normalisation of gDNA quantities (Faucon et al., 664 2015). Before amplification, gDNA samples were digested with Xhol for 15 min at 37°C directly within 665 the ddPCR reaction mixture. This reaction mixture was partitioned into up to 20,000 nanoliter-sized

droplets using the QX200 droplet generator (Bio-Rad) by mixing 70 μ l of synthetic oil with 20 μ l PCR

mix. Then, $40\mu L$ of the partitioned reaction was amplified for 40 cycles with an annealing

temperature of 60°C (see **Supplementary file 8** for protocol details and primers/probes). After

amplification, the number of positive and negative droplets was quantified for both Fam and Hex

670 channels using the QX200 droplet reader (Bio-Rad) and the positive/negative ratio was used to

estimate the initial gDNA concentration of each gene assuming a Poisson distribution. After

normalisation for gDNA quantity, *CYP6BB2* copy number were expressed as mean copy number (±
 95% CI).

674

675 *Quantification of the lle1011Met kdr mutation frequency*

The frequency of the Ile1011Met kdr mutation was quantified on pools of mosquitoes using ddPCR.

677 Genomic DNA was extracted as described above and digested with Xhol for 15 min at 37°C before

678 partitioning. The duplex TaqMan reaction mixture contained two amplification primers together with

two Taqman probes labelled with Fam and Hex fluorophores corresponding to the Met and Iso alleles
 respectively (see Supplementary file 8 for protocol and primer/probes details). The duplex reaction

681 mixture was emulsioned as described above and amplified for 40 cycles with an annealing

682 temperature of 59°C. After droplet reading, the positive/negative ratio obtained for each channel

- 683 was visualised as a scatterplot and was used to estimate the frequency of each allele assuming a 684 Poisson distribution with 95% CI.
- 685

686 Aknowledgments

687 We thank the Institut Pasteur de La Guyane for providing the isofemale lines IR03 and IR13. We also

- thank Dr. Ademir Jesus Martins from the Fundación Oswaldo Cruz for a critical reading of themanuscript.
- 690

691 Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest inrelation to the content of the article.

694

695 Authors's contributions

596 JMB and JPD designed the study. TB, CH, JG, JC, MK, LN, TG, FL and JT performed experiments and 597 analysed data. JF, FB, JV, JMB and JPD supervised data analysis. ID provided biological materials and 598 data. JMB, JF and JPD acquired the funding. TB, JMB and JPD wrote and revised the paper.

699

700 Data availability

- 701 RNA-seq data have been deposited at EBI short read archive (SRA) under accession number E-MTAB-
- 13325. DNA-seq short and long reads have been deposited at EBI short read archive (SRA) under
 accession number E-MTAB-13310.
- 704

705 Ethical aspects

- 706 Mice were kept in the animal facility of the Biology department of the University of Grenoble-Alpes,
- approved by the French Ministry of Animal Welfare (agreement no. B 38 421 10 001) and used in
- accordance with the laws of the European Union (directive 2010/63/EU). The use of animals for this

study was approved by the ComEth Grenoble-C2EA-12 ethics committee mandated by the French

- 710 Ministry of Higher Education and Research (MENESR). The study was conducted in accordance with
- 711 the ARRIVE guidelines.
- 712

713 Funding

- This study was funded by the European Union's Horizon 2020 research and innovation program
- vnder the ZIKAlliance project [grant agreement no. 734548] and the Infravec2 project [grant
- agreement no. 731060]. MK and TB were respectively supported by mobility fellowships funded by
- the European Union's Horizon 2020 research and innovation under the AIM COST action [grant
- 718 agreement no. CA17108] and the MSCA /NOVEC project [grant agreement no. 101086257]. Views
- and opinions expressed are however those of the authors only and do not necessarily reflect those of
- the European Union. Neither the European Union nor the granting authority can be held responsible
- 721 for them.

722 Supplementary materials

724 Supplementary file 1 (.tiff image). Bioassays data from F1 and F2 offspring used to investigate the

725 mode of inheritance of the resistance phenotype. The reciprocal controlled crosses used to

726 generate F1 and F2 individuals used for bioasays are indicated. Deltamethrin exposure conditions

were 0.03% deltamethrin for 30 min for females and 0.015% deltamethrin for 25 min for males in

order to account for their inherent lower insecticide tolerance. Mortality rates are indicated as mean

229 ± SD. Distinct letters indicate significantly distinct means (Kruskal-Wallis test followed by post hoc

pairwise Wilcoxon test with Bonferroni-Holm correction, N=5, $p \le 0.05$).

731 Supplementary file 2 (.xlsx file). RNA-seq and CNV data sets. All protein coding genes showing a

r32 significant differential transcription (sheet 'RNA-seq') or a significant differential exonic coverage

733 (sheet 'CNV') are shown. See methods for filtering conditions and thresholds. Gene descriptions

beginning with a # were manually inferred from Blast X result.

735 Supplementary File 3 (.tiff image). Genomic architecture of the *kdr* locus in the IROF resistant line.

On chromosome 3, the 3' part of the VGSC gene, including the 21 last exons, is duplicated. Black bars

- indicate the position of the coding exons. The duplication is 125 Kb long, and is imperfect: the two
- copies, R and R' mostly differing in their intronic sequences. The duplication breakpoints, left (LB)

and right (RB) are located at 315,905,811 bp and 316,030,250 bp on the reference genome AaegL5.

The 1011Met codon is located on the upstream copy (R) and followed by a B type intron. The 1011Ile

codon is on the downstream copy and followed by an A type intron (Martins et al, 2013). RNA-seq

- 742 data supported the sole expression of the 1011Met allele.
- 743 Supplementary file 4 (.xlsx file). Long reads overlapping the P450 duplication breakpoints.

Supplementary file 5 (.docx file). Data supporting the identification of the transposable elements
 flanking the P450 duplication.

- Supplementary file 6 (.docx file). RNA interference experiment. This document includes detailed
 results and methods.
- 548 Supplementary file 7 (.tiff image). Proposed evolutionary steps leading to the observed duplication

749 **architecture.** Theoretical structures and those inferred from short and long reads data are indicated.

750 The hAt-like and PiggyBac-like transposons are shown in purple and green respectively. Transposition

751 events are shown as dashed lines. The second hAt-like transposition event would have created an

incomplete copy of the hAT transposon, either directly or through a posterior deletion.

- Recombination events are shown as blue crossed lines. Stars indicates junctions that were confirmedby short reads data.
- 755 Supplementary File 8 (.docx file). Digital droplet PCR assays used to quantify the P450 duplication

756 and the frequency of the Ile1011Met kdr mutation. This document provides all technical details

related these ddPCR assays including amplification primers, TaqMan probes, PCR conditions and datainterpretation.

759 References

- Achee, N. L., Grieco, J. P., Vatandoost, H., Seixas, G., Pinto, J., Ching-Ng, L., Martins, A. J.,
- Juntarajumnong, W., Corbel, V., Gouagna, C., David, J.-P., Logan, J. G., Orsborne, J., Marois, E.,
- 762 Devine, G. J., & Vontas, J. (2019). Alternative strategies for mosquito-borne arbovirus control.
- 763 PLoS Negl Trop Dis, 13(1). https://doi.org/10.1371/journal.pntd.0006822
- Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count data. *Genome Biology*, *11*(10), R106. https://doi.org/10.1186/gb-2010-11-10-r106
- Baker, M. D., Read, L. R., Beatty, B. G., & Ng, P. (1996). Requirements for Ectopic Homologous
 Recombination in Mammalian Somatic Cells. *Molecular and Cellular Biology*, 16(12), 712227132.
 https://doi.org/10.1128/MCB.16.12.7122
- Bariami, V., Jones, C. M., Poupardin, R., Vontas, J., & Ranson, H. (2012). Gene Amplification, ABC
 Transporters and Cytochrome P450s^D: Unraveling the Molecular Basis of Pyrethroid Resistance in
 the Dengue Vector, Aedes aegypti. *PLoS Negl Trop Dis*, 6(6).
 https://doi.org/10.1271/journal.netd.0001692
- https://doi.org/10.1371/journal.pntd.0001692
- Bass, C., & Field, L. M. (2011). Gene amplification and insecticide resistance. *Pest Management Science*, 67(8), 886^[2]890. https://doi.org/10.1002/ps.2189
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate^D: A Practical and Powerful
 Approach to Multiple Testing. *J Royal Stat Soc B*, *57*, 289^D 300. https://doi.org/10.1111/j.25176161.1995.tb02031.x
- Brengues, C., Hawkes, N. J., Chandre, F., McCarroll, L., Duchon, S., Guillet, P., Manguin, S., Morgan, J.
 C., & Hemingway, J. (2003). Pyrethroid and DDT cross-resistance in *Aedes aegypti* is correlated
 with novel mutations in the voltage-gated sodium channel gene. *Med Vet Entomol*, *17*(1), 87^[2]94.
 https://doi.org/DOI: 10.1046/j.1365-2915.2003.00412.x
- Brito, L. P., Carrara, L., Freitas, R. M. D., Lima, J. B. P., & Martins, A. J. (2018). Levels of Resistance to
 Pyrethroid among Distinct *kdr* Alleles in *Aedes aegypti* Laboratory Lines and Frequency of *kdr*Alleles in 27 Natural Populations from Rio de Janeiro, Brazil. *BioMed Research International, 2018*,
 110. https://doi.org/10.1155/2018/2410819
- Brown, J. E., Evans, B. R., Zheng, W., Obas, V., Barrera-Martinez, L., Egizi, A., Zhao, H., Caccone, A., &
 Powell, J. R. (2014). Human impacts have shaped historical and recent evolution in *Aedes aegypti*,
 the dengue and yellow fever mosquito. *Evolution*, 68(2), 5142525.
 https://doi.org/10.1111/evo.12281
- Carvalho, V. L., & Long, M. T. (2021). Perspectives on New Vaccines against Arboviruses Using Insect Specific Viruses as Platforms. *Vaccines*, *9*(3), 263. https://doi.org/10.3390/vaccines9030263
- Cattel, J., Faucon, F., Le Peron, B., Sherpa, S., Monchal, M., Grillet, L., Gaude, T., Laporte, F., Dusfour,
 I., Reynaud, S., & David, J. P. (2020). Combining genetic crosses and pool targeted DNA-seq for
 untangling genomic variations associated with resistance to multiple insecticides in the mosquito
 Aedes aegypti. *Evolutionary Applications*, 3032 317. https://doi.org/10.1111/eva.12867
- Cattel, J., Haberkorn, C., Laporte, F., Gaude, T., Cumer, T., Renaud, J., Sutherland, I. W., Hertz, J. C.,
 Bonneville, J., Arnaud, V., Fustec, B., Boyer, S., Marcombe, S., & David, J. (2021). A genomic
 amplification affecting a carboxylesterase gene cluster confers organophosphate resistance in the
 mosquito Aedes aegypt¹/₂: From genomic characterization to high-throughput field detection.
 Evolutionary Applications, 14(4), 1009¹/₂1022. https://doi.org/10.1111/eva.13177
- Corbel, V., & N'Guessan, R. (2013). Distribution, Mechanisms, Impact and Management of Insecticide
 Resistance in Malaria Vectors^D: A Pragmatic Review, Anopheles mosquitoes -. *New insights into malaria vectors*. https://doi.org/10.5772/56117

 David, J. P., Ismail, H. M., Chandor-Proust, A., & Paine, M. J. I. (2013). Role of cytochrome P450s in insecticide resistance^[]: Impact on the control of mosquito-borne diseases and use of insecticides on Earth. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 368(1612).
 https://doi.org/10.1098/rstb.2012.0429

Djogbenou, L., Labbe, P., Chandre, F., Pasteur, N., & Weill, M. (2009). Ace-1 duplication in Anopheles
gambiae^[2]: A challenge for malaria control. *Malar J, 8*, 70. https://doi.org/10.1186/1475-2875-870

Busfour, I., Thalmensy, V., Gaborit, P., Issaly, J., Carinci, R., & Girod, R. (2011). Multiple insecticide
resistance in *Aedes aegypti* (Diptera[®]: Culicidae) populations compromises the effectiveness of
dengue vector control in French Guiana. *Mem Inst Oswaldo Cruz*, *106*(3), 346[®]352.

814 https://doi.org/DOI: 10.1590/s0074-02762011000300015

Busfour, I., Vontas, J., David, J. P., Weetman, D., Fonseca, D. M., Raghavendra, K., Corbel, V.,
Coulibaly, M., Martins, A. J., & Chandre, F. (2019). Management of insecticide resistance in the
major Aedes vectors of arboviruses^D: Advances and challenges. *Plos Neg Trop Dis*, *13*(10),
e0007615. https://doi.org/10.1371/journal.pntd.0007615

Busfour, I., Zorrilla, P., Guidez, A., Issaly, J., Girod, R., Guillaumot, L., Robello, C., & Strode, C. (2015).
 Deltamethrin Resistance Mechanisms in Aedes aegypti Populations from Three French Overseas
 Territories Worldwide. *PLoS Negl Trop Dis*, *9*(11), e0004226.
 https://doi.org/10.1371/journal.patd.0004226.

822 https://doi.org/10.1371/journal.pntd.0004226

Epelboin, Y., Wang, L., Giai Gianetto, Q., Choumet, V., Gaborit, P., Issaly, J., Guidez, A., Douché, T.,
Chaze, T., Matondo, M., & Dusfour, I. (2021). CYP450 core involvement in multiple resistance
strains of Aedes aegypti from French Guiana highlighted by proteomics, molecular and
biochemical studies. *PLOS ONE*, *16*(1), e0243992. https://doi.org/10.1371/journal.pone.0243992

Faucon, F., Dusfour, I., Gaude, T., Navratil, V., Boyer, F., Chandre, F., Sirisopa, P., Thanispong, K.,
Juntarajumnong, W., Poupardin, R., Chareonviriyaphap, T., Girod, R., Corbel, V., Reynaud, S., &
David, J. P. (2015). Identifying genomic changes associated with insecticide resistance in the
dengue mosquito Aedes aegypti by deep targeted sequencing. *Genome research*, 25(9),
134721359. https://doi.org/10.1101/gr.189225.115

Faucon, F., Gaude, T., Dusfour, I., Navratil, V., Corbel, V., Juntarajumnong, W., Girod, R., Poupardin,
R., Boyer, F., Reynaud, S., & David, J. P. (2017). In the hunt for genomic markers of metabolic
resistance to pyrethroids in the mosquito *Aedes aegypti*^[D]: An integrated next-generation
sequencing approach. *PLoS Negl Trop Dis*, *11*(4), e0005526.

836 https://doi.org/10.1371/journal.pntd.0005526

Feyereisen, R., Dermauw, W., & Van Leeuwen, T. (2015). Genotype to phenotype, the molecular and
 physiological dimensions of resistance in arthropods. *Pesticide Biochemistry and Physiology, 121,* 61¹/₂77. https://doi.org/10.1016/j.pestbp.2015.01.004

Ffrench-Constant, R. H., Daborn, P. J., & Le Goff, G. (2004). The genetics and genomics of insecticide
resistance. *Trends Genet*, 20(3), 1632170. https://doi.org/10.1016/j.tig.2004.01.003

842 Fontaine, A., Filipović, I., Fansiri, T., Hoffmann, A. A., Cheng, C., Kirkpatrick, M., Rašić, G., &

843 Lambrechts, L. (2017). Extensive Genetic Differentiation between Homomorphic Sex

844 Chromosomes in the Mosquito Vector, Aedes aegypti. *Genome Biology and Evolution*, 9(9),
845 232222335. https://doi.org/10.1093/gbe/evx171

Garg, H., Mehmetoglu-Gurbuz, T., & Joshi, A. (2020). Virus Like Particles (VLP) as multivalent vaccine
 candidate against Chikungunya, Japanese Encephalitis, Yellow Fever and Zika Virus. *Scientific Reports*, 10(1), 4017. https://doi.org/10.1038/s41598-020-61103-1

649 Goindin, D., Delannay, C., Gelasse, A., Ramdini, C., Gaude, T., Faucon, F., David, J. P., Gustave, J.,

- 850 Vega-Rua, A., & Fouque, F. (2017). Levels of insecticide resistance to deltamethrin, malathion, and
- temephos, and associated mechanisms in Aedes aegypti mosquitoes from the Guadeloupe and
- 852 Saint Martin islands (French West Indies). *Infect Dis Poverty*, *6*(1), 38.
- 853 https://doi.org/10.1186/s40249-017-0254-x
- Haddi, K., Tome, H. V. V., Du, Y., Valbon, W. R., Nomura, Y., Martins, G. F., Dong, K., & Oliveira, E. E.
 (2017). Detection of a new pyrethroid resistance mutation (V410L) in the sodium channel of *Aedes aegypti*: A potential challenge for mosquito control. *Scientific Reports*, *7*, 46549.
 https://doi.org/10.1038/srep46549
- Hawkins, N. J., Bass, C., Dixon, A., & Neve, P. (2018). The evolutionary origins of pesticide resistance. *Biol Rev Camb Philos Soc*. https://doi.org/10.1111/brv.12440
- Hemingway, J., Hawkes, N. J., McCarroll, L., & Ranson, H. (2004). The molecular basis of insecticide
 resistance in mosquitoes. *Insect Biochemistry and Molecular Biology*, *34*(7), 6532665.
 https://doi.org/10.1016/j.ibmb.2004.03.018
- Hendry, A. P., Farrugia, T. J., & Kinnison, M. T. (2008). Human influences on rates of phenotypic
 change in wild animal populations. *Molecular Ecology*, 17(1), 20229.
 https://doi.org/10.1111/j.1365-294X.2007.03428.x
- Hendry, A. P., Gotanda, K. M., & Svensson, E. I. (2017). Human influences on evolution, and the
 ecological and societal consequences. *Philos Trans R Soc Lond B Biol Sci*, 372(1712).
 https://doi.org/10.1098/rstb.2016.0028
- Hirata, K., Komagata, O., Itokawa, K., Yamamoto, A., Tomita, T., & Kasai, S. (2014). A single crossingover event in voltage-sensitive Na+ channel genes may cause critical failure of dengue mosquito
 control by insecticides. *PLoS Negl Trop Dis*, *8*(8), e3085.
- 872 https://doi.org/10.1371/journal.pntd.0003085
- Jain, C., Rhie, A., Hansen, N. F., Koren, S., & Phillippy, A. M. (2022). Long-read mapping to repetitive
 reference sequences using Winnowmap2. *Nature Methods*, *19*(6), 705¹/₂710.
 https://doi.org/10.1038/s41592-022-01457-8
- Kasai, S., Itokawa, K., Uemura, N., Takaoka, A., Furutani, S., Maekawa, Y., Kobayashi, D., ImanishiKobayashi, N., Amoa-Bosompem, M., Murota, K., Higa, Y., Kawada, H., Minakawa, N., Cuong, T. C.,
 Yen, N. T., Phong, T. V., Keo, S., Kang, K., Miura, K., ... Komagata, O. (2022). Discovery of superinsecticide-resistant dengue mosquitoes in Asia[®]: Threats of concomitant knockdown resistance
 mutations. *Science Advances*, 8(51), eabq7345. https://doi.org/10.1126/sciadv.abq7345
- Kasai, S., Komagata, O., Itokawa, K., Shono, T., Ng, L. C., Kobayashi, M., & Tomita, T. (2014).
 Mechanisms of pyrethroid resistance in the dengue mosquito vector, Aedes aegypti^[2]: Target site
 insensitivity, penetration, and metabolism. *PLoS Negl Trop Dis*, 8(6), e2948.
 https://doi.org/10.1271/journal.aptd.0002048
- 884 https://doi.org/10.1371/journal.pntd.0002948
- Kondrashov, F. A. (2012). Gene duplication as a mechanism of genomic adaptation to a changing
 environment. *Proceedings. Biological Sciences / The Royal Society*, 279(1749), 504825057.
 https://doi.org/10.1098/rspb.2012.1108
- Kraemer, M. U., Sinka, M. E., Duda, K. A., Mylne, A. Q., Shearer, F. M., Barker, C. M., Moore, C. G.,
 Carvalho, R. G., Coelho, G. E., Van Bortel, W., Hendrickx, G., Schaffner, F., Elyazar, I. R., Teng, H.-J.,
 Brady, O. J., Messina, J. P., Pigott, D. M., Scott, T. W., Smith, D. L., ... Hay, S. I. (2015). The global
 distribution of the arbovirus vectors Aedes aegypti and Ae. Albopictus. *eLife*, *4*, e08347.
- 892 https://doi.org/10.7554/eLife.08347
- Labbe, P., Berthomieu, A., Berticat, C., Alout, H., Raymond, M., Lenormand, T., & Weill, M. (2007).
 Independent duplications of the acetylcholinesterase gene conferring insecticide resistance in the

- mosquito Culex pipiens. *Mol Biol Evol*, 24(4), 1056 21067.
- 896 https://doi.org/10.1093/molbev/msm025
- Li, X. C., Schuler, M. A., & Berenbaum, M. R. (2007). Molecular mechanisms of metabolic resistance
 to synthetic and natural xenobiotics. *Annual Review of Entomology*, *52*, 2312253.
 https://doi.org/10.1146/annurev.ento.51.110104.151104
- Liu, N. (2015). Insecticide Resistance in Mosquitoes¹: Impact, Mechanisms, and Research Directions.
 Annual Review of Entomology, 60(1), 537¹/₂559. https://doi.org/10.1146/annurev-ento-010814 020828
- Lounibos, L. P. (2002). Invasions by insect vectors of human disease. *Annu Rev Entomol*, 47, 2332266.
 https://doi.org/10.1146/annurev.ento.47.091201.145206
- Marcombe, S., Carron, A., Darriet, F., Etienne, M., Agnew, P., Tolosa, M., Yp-Tcha, M. M., Lagneau, C.,
 Yebakima, A., & Corbel, V. (2009). Reduced efficacy of pyrethroid space sprays for dengue control
 in an area of Martinique with pyrethroid resistance. *Am J Trop Med Hyg*, *80*(5), 7452751.
 https://doi.org/10.4269/ajtmh.2009.80.745
- Marcombe, S., Darriet, F., Tolosa, M., Agnew, P., Duchon, S., Etienne, M., Yp Tcha, M. M., Chandre, F.,
 Corbel, V., & Yebakima, A. (2011). Pyrethroid resistance reduces the efficacy of space sprays for
 dengue control on the island of Martinique (Caribbean). *PLoS Negl Trop Dis*, 5(6), e1202.
 https://doi.org/10.1371/journal.pntd.0001202
- Marcombe, S., Fustec, B., Cattel, J., Chonephetsarath, S., Thammavong, P., Phommavanh, N., David,
 J.-P., Corbel, V., Sutherland, I. W., Hertz, J. C., & Brey, P. T. (2019). Distribution of insecticide
 resistance and mechanisms involved in the arbovirus vector Aedes aegypti in Laos and implication
 for vector control. *PLOS Neglected Tropical Diseases*, *13*(12), e0007852.
- 917 https://doi.org/10.1371/journal.pntd.0007852
- Marcombe, S., Poupardin, R., Darriet, F., Reynaud, S., Bonnet, J., Strode, C., Brengues, C., Yebakima,
 A., Ranson, H., Corbel, V., & David, J.-P. (2009). Exploring the molecular basis of insecticide
 resistance in the dengue vector Aedes aegypti^D: A case study in Martinique Island (French West
 Indies). *BMC Genomics*, 10. https://doi.org/494 10.1186/1471-2164-10-494
- Martins, A. J., Brito, L. P., Linss, J. G., Rivas, G. B., Machado, R., Bruno, R. V., Lima, J. B., Valle, D., &
 Peixoto, A. A. (2013). Evidence for gene duplication in the voltage-gated sodium channel gene of
 Aedes aegypti. *Evolution, Medicine, and Public Health, 2013*(1), 1482160.
 https://doi.org/10.1093/emph/eot012
- Martins, A. J., Lins, R. M., Linss, J. G., Peixoto, A. A., & Valle, D. (2009). Voltage-gated sodium channel
 polymorphism and metabolic resistance in pyrethroid-resistant Aedes aegypti from Brazil. *Am J Trop Med Hyg*, *81*(1), 1082115. https://doi.org/10.4269/ajtmh.2009.81.108
- Melo Costa, M., Campos, K. B., Brito, L. P., Roux, E., Melo Rodovalho, C., Bellinato, D. F., Lima, J. B. P.,
 & Martins, A. J. (2020). Kdr genotyping in Aedes aegypti from Brazil on a nation-wide scale from
 2017 to 2018. *Scientific Reports, 10*(1), 13267. https://doi.org/10.1038/s41598-020-70029-7
- Moyes, C. L., Vontas, J., Martins, A. J., Ng, L. C., Koou, S. Y., Dusfour, I., Raghavendra, K., Pinto, J.,
 Corbel, V., David, J. P., & Weetman, D. (2017). Contemporary status of insecticide resistance in
 the major Aedes vectors of arboviruses infecting humans. *PLoS Negl Trop Dis*, *11*(7), e0005625.
 https://doi.org/10.1371/journal.pntd.0005625
- 936 Nene, V., Wortman, J. R., Lawson, D., Haas, B., Kodira, C., Tu, Z. J., Loftus, B., Xi, Z., Megy, K.,
- 937 Grabherr, M., Ren, Q., Zdobnov, E. M., Lobo, N. F., Campbell, K. S., Brown, S. E., Bonaldo, M. F.,
- 938 Zhu, J., Sinkins, S. P., Hogenkamp, D. G., ... Severson, D. W. (2007). Genome sequence of Aedes
- 939 *aegypti*, a major arbovirus vector. *Science*, *316*(5832), 1718^[2]1723.
- 940 https://doi.org/10.1126/science.1138878

- Orr, H. A. (2005). The genetic theory of adaptation¹: A brief history. *Nat Rev Genet*, 6(2), 119¹/2127.
 https://doi.org/10.1038/nrg1523
- Palumbi, S. R. (2001). Humans as the world's greatest evolutionary force. *Science*, 293(5536),
 1786¹/₂1790. https://doi.org/10.1126/science.293.5536.1786
- Reid, W. R., Thornton, A., Pridgeon, J. W., Becnel, J. J., Tang, F., Estep, A., Clark, G. G., Allan, S., & Liu,
 N. (2014). Transcriptional Analysis of Four Family 4 P450s in a Puerto Rico Strain of *Aedes aegypti*
- 947 (Diptera[®]: Culicidae) Compared With an Orlando Strain and Their Possible Functional Roles in
- 948 Permethrin Resistance. *Journal of Medical Entomology*, *51*(3), 6052615.
- 949 https://doi.org/10.1603/ME13228
- Remnant, E. J., Good, R. T., Schmidt, J. M., Lumb, C., Robin, C., Daborn, P. J., & Batterham, P. (2013).
 Gene duplication in the major insecticide target site, *Rdl*, in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*, *110*(36), 14705214710.
 https://doi.org/10.1073/pnas.1311341110
- Riaz, M. A., Chandor-Proust, A., Dauphin-Villemant, C., Poupardin, R., Jones, C. M., Strode, C., Regent Kloeckner, M., David, J.-P., & Reynaud, S. (2013). Molecular mechanisms associated with
 increased tolerance to the neonicotinoid insecticide imidacloprid in the dengue vector Aedes
 aegypti. Aquatic Toxicology, 126, 326237. https://doi.org/10.1016/j.aquatox.2012.09.010
- Rigby, L. M., Rašić, G., Peatey, C. L., Hugo, L. E., Beebe, N. W., & Devine, G. J. (2020). Identifying the
 fitness costs of a pyrethroid-resistant genotype in the major arboviral vector Aedes aegypti. *Parasites & Vectors*, 13(1), 358. https://doi.org/10.1186/s13071-020-04238-4
- Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., & Mesirov, J. P.
 (2011). Integrative genomics viewer. *Nature Biotechnology*, 29(1), 24226.
 https://doi.org/10.1038/nbt.1754
- Saavedra-Rodriguez, K., Campbell, C. L., Lozano, S., Penilla-Navarro, P., Lopez-Solis, A., Solis-Santoyo,
 F., Rodriguez, A. D., Perera, R., & Black Iv, W. C. (2021). Permethrin resistance in Aedes aegyptill:
 Genomic variants that confer knockdown resistance, recovery, and death. *PLOS Genetics*, *17*(6),
 e1009606. https://doi.org/10.1371/journal.pgen.1009606
- Saavedra-Rodriguez, K., Suarez, A. F., Salas, I. F., Strode, C., Ranson, H., Hemingway, J., & Black, W. C.
 th. (2012). Transcription of detoxification genes after permethrin selection in the mosquito Aedes
 aegypti. *Insect Mol Biol*, 21(1), 61277. https://doi.org/10.1111/j.1365-2583.2011.01113.x
- Saavedra-Rodriguez, K., Urdaneta-Marquez, L., Rajatileka, S., Moulton, M., Flores, A. E., FernandezSalas, I., Bisset, J., Rodriguez, M., McCall, P. J., Donnelly, M. J., Ranson, H., Hemingway, J., & Black,
 W. C. (2007). A mutation in the voltage-gated sodium channel gene associated with pyrethroid
 resistance in Latin American *Aedes aegypti. Insect Molecular Biology*, *16*(6), 785^[2]798.
 https://doi.org/doi: 10.1111/j.1365-2583.2007.00774.x.
- 975 https://doi.org/doi: 10.1111/j.1365-2583.2007.00774.x.
- Schrider, D. R., Houle, D., Lynch, M., & Hahn, M. W. (2013). Rates and Genomic Consequences of
 Spontaneous Mutational Events in *Drosophila melanogaster*. *Genetics*, 194(4), 9372954.
 https://doi.org/10.1534/genetics.113.151670
- 979 Seixas, G., Grigoraki, L., Weetman, D., Vicente, J. L., Silva, A. C., Pinto, J., Vontas, J., & Sousa, C. A.
 980 (2017). Insecticide resistance is mediated by multiple mechanisms in recently introduced Aedes
 981 aegypti from Madeira Island (Portugal). *PLoS Negl Trop Dis*, *11*(7), e0005799.
 982 https://doi.org/10.1371/journal.pntd.0005799
- 983 Silva, J. V. J., Lopes, T. R. R., Oliveira-Filho, E. F. D., Oliveira, R. A. S., Durães-Carvalho, R., & Gil, L. H. V. 984 G. (2018). Current status, challenges and perspectives in the development of vaccines against
- 985 yellow fever, dengue, Zika and chikungunya viruses. *Acta Tropica*, *182*, 2572263.
- 986 https://doi.org/10.1016/j.actatropica.2018.03.009

987	Smith, L. B., Kasai, S., & Scott, J. G. (2016). Pyrethroid resistance in Aedes aegypti and Aedes
988	albopictus [®] : Important mosquito vectors of human diseases. Pesticide Biochemistry and
989	<i>Physiology, 133,</i> 1212. https://doi.org/10.1016/j.pestbp.2016.03.005

- Uemura, N., Furutani, S., Tomita, T., Itokawa, K., Komagata, O., & Kasai, S. (2023). Concomitant
 knockdown resistance allele, L982W + F1534C, in Aedes aegypti has the potential to impose
 fitness costs without selection pressure. *Pesticide Biochemistry and Physiology*, 193, 105422.
 https://doi.org/10.1016/j.pestbp.2023.105422
- Valle, D., Bellinato, D. F., Viana-Medeiros, P. F., Lima, J. B. P., & Martins Junior, A. D. J. (2019).
 Resistance to temephos and deltamethrin in Aedes aegypti from Brazil between 1985 and 2017. *Memórias Do Instituto Oswaldo Cruz, 114*, e180544. https://doi.org/10.1590/0074-02760180544
- Weetman, D., Djogbenou, L. S., & Lucas, E. (2018). Copy number variation (CNV) and insecticide
 resistance in mosquitoes^[2]: Evolving knowledge or an evolving problem? *Curr Opin Insect Sci, 27*,
 82^[2]88. https://doi.org/10.1016/j.cois.2018.04.005
- WHO. (2022). Standard operating procedure for testing insecticide susceptibility of adult mosquitoes
 in WHO tube tests. SOP version 2: WHO Tube test/01/14. World Health Organization.
 https://ivia.who.ipt/bitstraam/bandle/10665/252216/0720240042821.ang.ndf2caguages=1
- 1002 https://iris.who.int/bitstream/handle/10665/352316/9789240043831-eng.pdf?sequence=1
- Yanola, J., Somboon, P., Walton, C., Nachaiwieng, W., Somwang, P., & Prapanthadara, L. A. (2011).
 High-throughput assays for detection of the F1534C mutation in the voltage-gated sodium
 channel gene in permethrin-resistant *Aedes aegypti* and the distribution of this mutation
 throughout Thailand. *Trop Med Int Health*, *16*(4), 5012509. https://doi.org/10.1111/j.13653156.2011.02725.x
- Zimin, A. V., Marçais, G., Puiu, D., Roberts, M., Salzberg, S. L., & Yorke, J. A. (2013). The MaSuRCA
 genome assembler. *Bioinformatics*, 29(21), 2669^[2]2677.
 https://doi.org/10.1093/bioinformatics/btt476
- 1010 https://doi.org/10.1093/bioinformatics/btt476
- Zoh, M. G., Gaude, T., Prud'homme, S. M., Riaz, M. A., David, J.-P., & Reynaud, S. (2021). Molecular
 bases of P450-mediated resistance to the neonicotinoid insecticide imidacloprid in the mosquito
 Ae. Aegypti. Aquatic Toxicology, 236, 105860. https://doi.org/10.1016/j.aquatox.2021.105860
- 1014