



HAL
open science

Federated Causal Inference: Multi-Centric ATE Estimation beyond Meta-Analysis

Rémi Khellaf, Aurélien Bellet, Julie Josse

► **To cite this version:**

Rémi Khellaf, Aurélien Bellet, Julie Josse. Federated Causal Inference: Multi-Centric ATE Estimation beyond Meta-Analysis. 2024. hal-04747923

HAL Id: hal-04747923

<https://hal.science/hal-04747923v1>

Preprint submitted on 22 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Federated Causal Inference: Multi-Centric ATE Estimation beyond Meta-Analysis

Rémi Khellaf¹, Aurélien Bellet¹, Julie Josse¹
¹INRIA, Université de Montpellier, IDESP

Abstract

We study Federated Causal Inference, an approach to estimate treatment effects from decentralized data across centers. We compare three classes of Average Treatment Effect (ATE) estimators derived from the Plug-in G-Formula, ranging from simple meta-analysis to one-shot and multi-shot federated learning, the latter leveraging the full data to learn the outcome model (albeit requiring more communication). Focusing on Randomized Controlled Trials (RCTs), we derive the asymptotic variance of these estimators for linear models. Our results provide practical guidance on selecting the appropriate estimator for various scenarios, including heterogeneity in sample sizes, covariate distributions, treatment assignment schemes, and center effects. We validate these findings with a simulation study.

1 INTRODUCTION

In modern evidence-based medicine, Randomized Controlled Trials (RCT) are considered the gold standard for estimating the Average Treatment Effect (ATE) because they effectively isolate the treatment effect from confounding factors (Guyatt et al., 2015). The most widely used estimator of the ATE, when expressed as a risk difference, is the difference-in-means (DM) estimator. Recently, however, the U.S. Food and Drug Administration (2023) has recommended to adjust for covariates using linear models for the outcome, as this approach consistently yields more precise ATE estimates than the DM estimator (European Medicines Agency, 2024; Tsiatis et al., 2008; Benkeser et al., 2021) even when the assumption of linearity does not hold (Lin, 2013; Wager, 2020; Lei and Ding, 2021; Van Lancker et al., 2024).

Nevertheless, concerns have been raised about the limited scope of RCTs, including their stringent eligibility criteria, short timeframes, limited sample size, etc. Consequently, regulatory agencies tasked with making high-stakes decisions on drug approvals—decisions that directly impact public health and for which the reimbursement of the drug is often tied to its efficacy (French Health Authority, 2024)—frequently turn to meta-analysis to guide their choices. Meta-analysis, which aggregates estimated effects from multiple studies conducted across various centers (Hunter and Schmidt, 2004; Borenstein et al., 2021), represents the

pinnacle of evidence in clinical research (Blunt, 2015). They can lead to increased statistical power and more precise estimates, while also offering valuable insights into rare adverse events.

Despite extensive guidelines on conducting meta-analyses (Moher et al., 1999; Liberati et al., 2009; Higgins et al., 2019), multi-centric approaches still face significant challenges. These primarily arise from heterogeneity caused by imbalances in datasets, variations in populations across studies, and center effects due to differing practices across institutions (Berlin and Golub, 2014). Moreover, simply aggregating local estimates is not the only approach to conducting meta-analyses. However, implementing “one-stage” meta-analyses (Morris et al., 2018) that pool individual patient data from all centers is practically challenging due to data silos and personal data regulations.

Federated causal inference, an emerging field combining federated learning (Kairouz et al., 2021) and causal inference (Imbens and Rubin, 2015; Hernan, 2020) to estimate causal effects from decentralized data sources, offers a compelling alternative to traditional meta-analysis. Federated Learning (FL) enables multiple centers to collaboratively train a model without sharing their raw data, instead exchanging only model updates that are iteratively aggregated by an orchestrating server. This decentralized approach is especially valuable in fields like medicine (Sheller et al., 2020; Che et al., 2022; Prospero et al., 2020), where strong incentives exist to keep data on-site—whether to comply with data protection regulations, maintain ownership and control over the data, or avoid unwanted knowledge transfer. However, traditional FL algorithms are designed to learn predictive models (Kairouz et al., 2021), rather than estimating causal effects.

Contributions. In this paper, we aim to estimate the ATE for a population represented by multiple RCTs conducted across different centers, accounting for potential heterogeneity, using a federated approach. We study and compare three classes of federated estimators of the ATE: (i) *meta-analysis estimators*, which aggregate ATE estimates computed independently at each center; (ii) *one-shot federated estimators*, where outcome model parameters are estimated at each center, aggregated, and shared back for centers to compute and aggregate ATE estimates; and (iii) *gradient-based federated estimators*, where outcome model parameters are learned on joint data using federated gradient descent before each center computes and aggregates its ATE estimate. These estimators entail different communication costs, which often act as the bottleneck in real-world systems (Kairouz et al., 2021).

Our primary contribution is the derivation of asymptotic variances for these estimators under a linear outcome model. This modeling choice is known to provide a variance reduction compared to the classic DM estimator, even when the underlying model is not linear (Lin, 2013; Wager, 2020; Lei and Ding, 2021; Van Lancker et al., 2024), and aligns with recent recommendations from regulatory agencies (U.S. Food and Drug Administration, 2023). We specifically address scenarios involving heterogeneity, including distributional shifts (varying covariate distributions across centers) and center effects on the outcome. Our results shed light on the trade-offs between the statistical efficiency, communication costs, and underlying modeling assumptions of the considered estimators. We find that, despite their simplicity, low communication overhead and minimal assumptions, meta-analysis estimators can achieve statistical efficiency comparable to pooled data analysis when sufficient data is available at

each center, while naturally accommodating center effects. In contrast, when local datasets are small, gradient-based federated estimators stand out as the only viable option. One-shot estimators offer an interesting middle ground in some cases: they can recover the same ATE estimate as pooled data analysis while being robust to distributional shifts in covariates and differences in treatment assignments, but suffer from increased variance when center effects are present. These conclusions are supported by a simulation study that illustrates the behavior of the estimators under the different scenarios. Ultimately, our work provides clear guidelines on selecting the most suitable estimator for different scenarios, as summarized by a decision diagram designed for practitioners (Figure 6 in Appendix A).

Related work. The work closest to ours is Xiong et al. (2023), which adapts estimators of the ATE through a one-shot federated estimation of the outcome/propensity score model parameters. However, their work does not compare the efficiency of these one-shot federated estimators with traditional meta-analysis and pooled dataset estimators, nor does it consider gradient-based federated alternatives. Our results offer clear guidelines on when the One-Shot estimators proposed by Xiong et al. (2023) should be preferred over other methods.

Other work on federated causal inference (see Brantner et al. (2023) and Edmondson et al. (2023) for an overview) consider different settings and objectives than the ones considered in our work. Vo et al. (2022b) employ a Bayesian framework using Gaussian processes to estimate the ATE under uniform data distributions across centers. Terrail et al. (2023) focus on federating an external control arm for time-to-event outcomes, adapting a gradient-based algorithm for Cox hazard model parameters. While our work aims to estimate the causal effect of treatment across the joint population of centers, other studies (Vo et al., 2022a; Han et al., 2021, 2023; Makhija et al., 2024; Guo et al., 2024) aim at transferring causal estimates from a source center to a target center.

The meta-analysis literature on combining estimates from multiple studies is extensive. One can mention the work of Morris et al. (2018) who discuss the differences and advantages of conducting meta-analysis with individual patient data on stratified (“two-stage”) versus pooled data (“one-stage”). However, they require sharing raw data and do not explore any federated strategy. Meta-analysis provides considerable flexibility in the choice of local models (Seo et al., 2021; Tan et al., 2022), but also comes with many subtle statistical challenges. These include the ecological fallacy bias Piantadosi et al. (1988), which occurs when incorrect conclusions about individuals are drawn from subgroup characteristics, as well as situations when ignoring center sizes can lead to biased ATE estimates (Kahan et al., 2023).

2 GENERAL FRAMEWORK

Notations. We consider a set of K centers, with H denoting the random variable with values in $\{1, \dots, K\}$ indicating membership to a center. Let $\mathcal{Z} = \{Z_i\}_{i=1}^n$ be a sample of n independent and identically distributed (i.i.d.) realizations of the quadruplet $Z = (X, W, Y, H)$, where X denotes a d -dimensional vector of covariates that belongs to a covariate space $\mathcal{X} \subset \mathbb{R}^d$, $W \in \{0, 1\}$ denote the binary treatment, and $Y \in \mathbb{R}$ is the observed outcome of interest.

We denote by \mathcal{Z}_k the local dataset of center k with $n_k = \sum_{i=1}^n \mathbb{1}_{\{H_i=k\}}$ observations, and by $\mathcal{Z}_k^{(w)}$ the $n_k^{(w)} = \sum_{i=1}^n \mathbb{1}_{\{H_i=k, W_i=w\}}$ observations in center k under treatment arm w . We further denote by $X_k^{(w)} = \{X_i \mid W_i = w, H_i = k\}_{i=1}^{n_k^{(w)}} \in \mathbb{R}^{n_k^{(w)} \times d}$ (resp. $Y_k(w) = \{Y_i \mid W_i = w, H_i = k\}_{i=1}^{n_k^{(w)}} \in \mathbb{R}^{n_k^{(w)}}$) the design matrix of the covariates (resp. the outcome vector) for treatment arm w in center k . Similarly, we denote by $\mathcal{Z}^{(w)}$ the $n^{(w)} = \sum_{k=1}^K n_k^{(w)}$ observations under treatment arm w in the pooled dataset \mathcal{Z} , and by $X^{(w)} \in \mathbb{R}^{n^{(w)} \times d}$ and $Y^{(w)} \in \mathbb{R}^{n^{(w)}}$ the corresponding design matrix and outcome vector.

Average treatment effect in K RCTs.

We consider the setting of K Bernoulli RCT trials, where each participant i in center k has a fixed probability $\mathbb{P}(W_i = 1 \mid H_i = k) = p_k$ of being assigned to the treatment group, which does not depend on X in this design. We denote $\rho_k = \mathbb{P}(H_i = k)$ the probability that an observation belongs to center k ($0 < \rho_k < 1$). Note that the probability p of being treated within the pooled dataset \mathcal{Z} is then given by $p = \sum_{k=1}^K \rho_k p_k$.

We consider the potential outcomes framework (Rubin, 1974) and we aim to estimate the ATE $\tau \in \mathbb{R}$ defined as the Risk Difference over the K centers as $\tau = \mathbb{E}(\mathbb{E}(Y_i(1) - Y_i(0) \mid H_i))$, where $Y(w)$ is the outcome had the subject received treatment w . We denote the local ATE in center k by $\tau_k = \mathbb{E}(Y_i(1) - Y_i(0) \mid H_i = k)$.

We assume that the classical identifiability assumptions for a RCT design hold locally at every center: (a) *SUTVA* (Stable Unit Treatment Value Assumption): $Y = WY(1) + (1 - W)Y(0)$, (b) *Positivity*: $\exists \eta_1 > 0$ such that, almost surely, $\eta_1 \leq \mathbb{P}(W = 1 \mid H) \leq 1 - \eta_1$ and (c) *Ignorability*: $W \perp\!\!\!\perp (Y(0), Y(1))$. Under these conditions, the ATE is identifiable and the simple ‘‘Difference-in-Means’’ (Splawa-Neyman et al., 1990) estimator defined as $\hat{\tau}_{\text{DM}} = \frac{1}{n^{(1)}} \sum_{i=1}^{n^{(1)}} Y_i W_i - \frac{1}{n^{(0)}} \sum_{i=1}^{n^{(0)}} Y_i (1 - W_i)$ is an unbiased estimator of the ATE. However, variance reduction can be obtained by adjusting from covariates and considering the ‘‘Plug-in G-formula’’ or ‘‘outcome-based regression’’ estimator.

Definition 1 (Robins, 1986). *The plug-in G-formula estimator is defined as $\hat{\tau} = \frac{1}{n} \sum_{i=1}^n (\hat{\mu}_1(X_i) - \hat{\mu}_0(X_i))$, where $\hat{\mu}_w(x)$ is an estimator of the surface response $\mu_w(x) = \mathbb{E}[Y \mid W = w, X = x]$.*

Throughout the paper, we will consider two different regimes for the sample sizes.

Condition 1 (Local Full Rank). *For all $(k, w) \in \llbracket 1, K \rrbracket \times \{0, 1\}$, we have $\text{rank}(X_k^{(w)\top} X_k^{(w)}) = d$, which implies that $\forall (k, w), n_k^{(w)} \geq d$.*

Condition 2 (Federated Full Rank). *For all $w \in \{0, 1\}$, we have $\text{rank}(X^{(w)\top} X^{(w)}) = d$, which implies that $\forall w, \sum_{k=1}^K n_k^{(w)} \geq d$.*




Figure 1: Graphical Model for Homogeneous Settings

3 HOMOGENEOUS POPULATION

In this section, we focus on estimating the ATE over K RCTs studying the same population. We assume that the joint distribution of $Z = (X, W, Y, H)$ decomposes as

$$\mathbb{P}(Z) = \mathbb{P}(H)\mathbb{P}(W|H)\mathbb{P}(X)\mathbb{P}(Y|X, W). \quad (1)$$

This corresponds to the graphical model shown in Figure 1. We refer to this setting as *homogeneous* because X and $Y|X, W$ are independent of H : in other words, there is no distributional shift for the covariates and the conditional outcomes across centers. *Heterogeneous* settings will be addressed in Section 4.

We consider a linear model for the potential outcomes

$$Y_{k,i}(w) = c^{(w)} + X_{k,i}\beta^{(w)} + \varepsilon_{k,i}(w), \quad (2)$$

with $c^{(w)} \in \mathbb{R}$ the intercept, $\beta^{(w)} \in \mathbb{R}^d$ the coefficients, $\mathbb{E}(\varepsilon_{k,i}(w) | X_{k,i}) = 0$ and $\mathbb{V}(\varepsilon_{k,i}(w) | X_{k,i}) = \sigma^2$.

Note that $\beta^{(1)}$ and $\beta^{(0)}$ can be different, so that the treatment effect can be heterogeneous (*i.e.* depends on the covariates). We denote by $\theta^{(w)} = (c^{(w)}, \beta^{(w)}) \in \mathbb{R}^{d+1}$ and $X' = (1, X) \in \mathbb{R}^{n,d+1}$ the covariate matrix augmented with a column of ones. The model parameters $\theta^{(w)}$ are the same in every center, in accordance with the homogeneous setting defined by the decomposition in Eq. 1. We assume that X has finite first two moments $\mathbb{E}(X) = \mu$ and $\mathbb{V}(X) = \Sigma = \mathbb{E}(X^\top X)$.

Under the above model, the ATE can be written as

$$\tau = \mathbb{E}(X'_i)(\theta^{(1)} - \theta^{(0)}) \quad (3)$$

and the ATEs per center $\tau_k = \mathbb{E}(X'_{k,i})(\theta^{(1)} - \theta^{(0)})$ are homogeneous across centers, *i.e.*, $\tau_1 = \dots = \tau_K = \tau$, as the covariate distribution is the same across centers.

The G-formula estimator on the pooled data \mathcal{Z} can be used to estimate the ATE over the K centers

$$\hat{\tau}_{\text{pool}} = \frac{1}{n} \sum_{i=1}^n (X'_i \hat{\theta}_{\text{pool}}^{(1)} - X'_i \hat{\theta}_{\text{pool}}^{(0)}), \quad (4)$$

where $\hat{\theta}_{\text{pool}}^{(1)}$ and $\hat{\theta}_{\text{pool}}^{(0)}$ are the regression coefficients estimated by fitting two OLS regressions over $\mathcal{Z}^{(1)}$ and $\mathcal{Z}^{(0)}$ respectively. This *pooled* estimator satisfies (see Appendix B.2.6 for a proof extending the standard result in Wager (2020) to non-centered covariates)

$$\sqrt{n}(\hat{\tau}_{\text{pool}} - \tau) \xrightarrow{d} \mathcal{N}(0, V_{\text{pool}}), \quad (5)$$

with $V_{\text{pool}} = \frac{\sigma^2}{p(1-p)} + \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2$. However, computing $\hat{\tau}_{\text{pool}}$ requires access to the *pooled* dataset \mathcal{Z} , which is not accessible in the decentralized setting we consider. Under Condition 1, each center in isolation can only estimate the ATE using its local dataset \mathcal{Z}_k

$$\hat{\tau}_k = \frac{1}{n_k} \sum_{i=1}^{n_k} (X'_{k,i} \hat{\theta}_k^{(1)} - X'_{k,i} \hat{\theta}_k^{(0)}), \quad (6)$$

where $\hat{\theta}_k^{(w)} = (X'_k{}^{(w)\top} X'_k{}^{(w)})^{-1} X'_k{}^{(w)\top} Y_k^{(w)}$ is the OLS estimator computed over $\mathcal{Z}_k^{(w)}$. To improve upon this baseline, we now introduce several estimation strategies of the ATE over the K centers, with the aim of obtaining estimates as if one had access to \mathcal{Z} .

3.1 Definition of the Estimators

3.1.1 Meta-Analysis Estimators

A first strategy under Condition 1 is to aggregate the local ATE estimates $\hat{\tau}_k$ in Eq. 6. The centers then send their local ATE estimates to the server, which aggregates them using non-negative weights ω_k that sum to 1 over the K centers to obtain a global ATE estimate. For selecting the weights ω_k , [Hunter and Schmidt \(2004\)](#) describes two common methods for absolute measures like the risk difference: sample size weighting (SW) and inverse variance weighting (IVW). These meta-analysis estimators involve a single round of communication: each center k sends $\hat{\tau}_k$ to the server.

Definition 2 (Meta-Analysis - SW Aggregation).

$$\hat{\tau}_{\text{Meta-SW}} = \sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k. \quad (7)$$

Definition 3 (Meta-Analysis - IVW Aggregation).

$$\hat{\tau}_{\text{Meta-IVW}} = \frac{\sum_{k=1}^K \mathbb{V}(\hat{\tau}_k)^{-1} \hat{\tau}_k}{\sum_{k=1}^K \mathbb{V}(\hat{\tau}_k)^{-1}}. \quad (8)$$

Proposition 1 (proof in Appendix B.1.2). $\hat{\tau}_{\text{Meta-IVW}}$ is the minimum-variance estimator of τ among the class of aggregation-based estimators.

Remark 1. In practice, $\mathbb{V}(\hat{\tau}_k)$ is often unknown and must be estimated, leading to an approximation of $\hat{\tau}_{\text{Meta-IVW}}$. In contrast, $\hat{\tau}_{\text{Meta-SW}}$ only requires knowledge of the local sample sizes.

3.1.2 One-Shot Federated Estimators

To go beyond the mere aggregation of local ATEs, we can follow [Xiong et al. \(2023\)](#) and aggregate the local outcome model parameters using a single round of communication (hence the term “one-shot” federated) to build better local ATE estimates, before aggregating them.

Step 1. Local estimation of outcome parameters: Under Condition 1, each center k estimates $\hat{\theta}_k^{(w)}$ locally with an OLS regression.

Step 2. One-shot federation of parameters: We perform a meta-analysis (local estimation then weighted aggregation) of the local outcome model parameters $\hat{\theta}_k^{(w)}$. Specifically, the centers send their local estimates to the server, which computes $\hat{\theta}_{1S}^{(w)} = \sum_{k=1}^K \omega_k^{(\theta)} \hat{\theta}_k^{(w)}$ (where “1S” stands for “one-shot”), with $\omega_k^{(\theta)}$ some federation weights (summing to 1 over the K centers) like SW or IVW. The server sends back the obtained $\hat{\theta}_{1S}^{(w)}$ to all the centers.

Definition 4 (SW Federation of $\hat{\theta}_k^{(w)}$).

$$\hat{\theta}_{1S-SW}^{(w)} = \sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \hat{\theta}_k^{(w)}. \quad (9)$$

Definition 5 (IVW Federation of $\hat{\theta}_k^{(w)}$).

$$\hat{\theta}_{1S-IVW}^{(w)} = \frac{\sum_{k=1}^K (\mathbb{V}(\hat{\theta}_k^{(w)})^{-1} \hat{\theta}_k^{(w)})}{\sum_{k=1}^K \mathbb{V}(\hat{\theta}_k^{(w)})^{-1}}. \quad (10)$$

with $\mathbb{V}(\hat{\theta}_k^{(w)})^{-1} = \frac{1}{\sigma^2} X_k'^{(w)\top} X_k'^{(w)}$.

Theorem 1 (proof in Appendix B.2.1). *Under Condition 1, $\hat{\theta}_{1S-IVW}^{(w)} = \hat{\theta}_{\text{pool}}^{(w)}$.*

Remarkably, Theorem 1 shows that one can obtain similar estimates as $\hat{\theta}_{\text{pool}}$ (which has the lowest variance among the class of linear unbiased estimators (Giraud et al., 2012)) by federating the local estimates with a one-shot IVW procedure, even in finite sample sizes, whenever the local datasets are of full rank. This gives a very strong argument in favor of this approach in comparison to the One-Shot SW which thus necessarily has higher variance than $\hat{\theta}_{1S-IVW}$ and $\hat{\theta}_{\text{pool}}$. However, note that the communication cost of computing $\hat{\theta}_{1S-IVW}^{(w)}$ is $O(d)$ times larger than for $\hat{\theta}_{1S-SW}^{(w)}$, as each center must send to the server its $(d+1) \times (d+1)$ local variance matrix $X_k'^{(w)\top} X_k'^{(w)}$.

Step 3. Aggregation of the ATEs: Each center estimates its local ATE using the federated outcome model parameters:

$$\hat{\tau}_k^{1S-\text{agg}} = \frac{1}{n_k} \sum_{i=1}^{n_k} (X_{k,i}' \hat{\theta}_{1S-\text{agg}}^{(1)} - X_{k,i}' \hat{\theta}_{1S-\text{agg}}^{(0)}) \quad (11)$$

with $\text{agg} \in \{\text{SW}, \text{IVW}\}$. Finally, a second communication round is used where centers each send their $\hat{\tau}_k^{1S-\text{agg}}$ to the server for aggregation with weights $\omega^{(\tau)}$:

$$\hat{\tau}_{1S-\text{agg}} = \sum_{k=1}^K \omega_k^{(\tau)} \hat{\tau}_k^{1S-\text{agg}}.$$

It turns out that using SW or IVW for $\omega^{(\tau)}$ is asymptotically equivalent (see Appendix B.2.6), so in the following we focus on sample size aggregation weights.

Definition 6 (1S SW Federation - SW Aggregation).

$$\hat{\tau}_{1S-SW} = \sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k^{1S-SW} \quad (12)$$

Definition 7 (1S IVW Federation - SW Aggregation).

$$\hat{\tau}_{1S-IVW} = \sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k^{1S-IVW} \quad (13)$$

3.1.3 Gradient-based Federated Estimators

Neither the meta nor the one-shot estimators can be used when Condition 1 does not hold (e.g., as soon as one center has its sample size $n_k^{(w)} < d$), since $\hat{\theta}_k^{(w)}$ is not defined. In such cases, we propose to leverage gradient-based federated estimators.

Step 1. Multi-shot federation of parameters: Centers jointly estimate $\hat{\theta}_{\text{pool}}^{(w)}$ by solving the underlying OLS problem in a federated fashion. To the best of our knowledge, we are the first to propose this approach to estimate the ATE in Federated Causal Inference. Finding $\hat{\theta}_{\text{pool}}^{(w)}$ amounts to minimizing the Mean Squared Error loss function, defined as $\ell(\theta^{(w)}, \bigcup_{k=1}^K \mathcal{Z}_k^{(w)}) = \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} (Y_i^{(w)} - X_i^{\prime(w)} \theta^{(w)})^2$. This optimization problem can be solved by a federated gradient descent-based algorithm. We propose to use the FedAvg algorithm (McMahan et al., 2017) for its intuitive simplicity and good performance on homogeneous (Stich, 2019; Khaled et al., 2020) and even heterogeneous data (Wang et al., 2024). FedAvg alternates for T rounds between performing E local gradient steps in each center and aggregating the parameters at the server, see Appendix D for the detailed algorithm. The output of this procedure is an estimate $\hat{\theta}_{\text{GD}}^{(w)}$ (“GD” stands for Gradient Descent).

Note that under an appropriate choice of learning rate and setting $T = 1$, FedAvg converges to the one-shot estimator $\hat{\theta}_{\text{SW}}^{(w)}$ as $E \rightarrow \infty$. On the other hand, under an appropriate choice of learning rate and setting $E = 1$, FedAvg converges to the pooled model $\hat{\theta}_{\text{pool}}^{(w)}$ as $T \rightarrow \infty$ (Kairouz et al., 2021; Wang et al., 2024). FedAvg thus allows to learn a better estimate of the outcome model parameters than one-shot approaches (in particular when Condition 1 does not hold) at the cost of more communication. In our homogeneous case, an extremely accurate estimate of $\hat{\theta}_{\text{pool}}^{(w)}$ can be obtained with a small number of communication rounds T (see Appendix D). In the following, we consider that the parameters of FedAvg are chosen such that $\hat{\theta}_{\text{GD}}^{(w)} = \hat{\theta}_{\text{pool}}^{(w)}$.

Step 2. Aggregation of the ATEs: Each center k then computes a local estimate of the ATE using $\hat{\theta}_{\text{GD}}^{(w)}$

$$\hat{\tau}_k^{\text{GD}} = \frac{1}{n_k} \sum_{i=1}^{n_k} (X'_{k,i} \hat{\theta}_{\text{GD}}^{(1)} - X'_{k,i} \hat{\theta}_{\text{GD}}^{(0)}) \quad (14)$$

Finally, these estimates are aggregated in a last round of communication with sample weighting (IVW is asymptotically equivalent, see Section 3.1.2), which yields the following GD estimator of the global ATE.

Definition 8 (GD Federation - SW Aggregation).

$$\hat{\tau}_{\text{GD}} = \sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k^{\text{GD}} \quad (15)$$

3.2 Comparison of the Federated Estimators

General case. Under the graphical model in Figure 1 and Condition 1, all ATE estimators presented so far are unbiased (as proved in Appendix B.2.5).

For each estimator, we report in Table 1 its asymptotic variance \mathbb{V}^∞ (with proofs in Appendix B.2.6), the sample size required (as per Condition 1 or the weaker Condition 2),

Table 1: Properties of the (unbiased) estimators of the ATE in the homogeneous setting: asymptotic variance, number of communication rounds and total communication cost (in number of floats per center).

Estimator	Notation	Condition \mathbb{V}^∞	Com. rounds	Com. cost
Local	$\hat{\tau}_k$ (Eq. 6)	Cond. 1 $\frac{\sigma^2}{n_k} \frac{1}{p_k(1-p_k)} + \frac{1}{n_k} \ \beta^{(1)} - \beta^{(0)}\ _\Sigma^2$	0	0
Meta-SW	$\hat{\tau}_{\text{Meta-SW}}$ (Eq. 7)	Cond. 1 $\frac{\sigma^2}{n} \sum_{k=1}^K \frac{\rho_k}{p_k(1-p_k)} + \frac{1}{n} \ \beta^{(1)} - \beta^{(0)}\ _\Sigma^2$	1	$O(1)$
Meta-IVW	$\hat{\tau}_{\text{Meta-IVW}}$ (Eq. 8)	Cond. 1 $\left(\sum_{k=1}^K \left(\sigma^2 \frac{n\rho_k}{p_k(1-p_k)} + \frac{1}{n_k} \ \beta^{(1)} - \beta^{(0)}\ _\Sigma^2 \right) \right)^{-1}$	1	$O(1)$
1S-SW	$\hat{\tau}_{\text{1S-SW}}$ (Eq. 12)	Cond. 1 $\frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \ \beta^{(1)} - \beta^{(0)}\ _\Sigma^2$	2	$O(d)$
1S-IVW	$\hat{\tau}_{\text{1S-IVW}}$ (Eq. 13)	Cond. 1 $\frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \ \beta^{(1)} - \beta^{(0)}\ _\Sigma^2$	2	$O(d^2)$
GD	$\hat{\tau}_{\text{GD}}$ (Eq. 15)	Cond. 2 $\frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \ \beta^{(1)} - \beta^{(0)}\ _\Sigma^2$	$T + 1$	$O(Td)$
Pool	$\hat{\tau}_{\text{pool}}$ (Eq. 4)	Cond. 2 $\frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \ \beta^{(1)} - \beta^{(0)}\ _\Sigma^2$	—	—

the number of communication rounds needed between the centers and the server, and the total communication cost per center (in number of floats). We observe that the one-shot and gradient-based federated estimators achieve the same variance of the pooled-data estimator. The differences lie in the sample size conditions and communication costs. While one-shot estimators require two communication rounds (and generally lower total communication costs), they require that the *local* sample size $n_k^{(w)}$ at each center k and arm w be larger than the dimension d . In contrast, the GD estimator requires this only for the *pooled* sample size $n^{(w)} = \sum_{k=1}^K n_k^{(w)}$.

Theorem 2 (Proof in Appendix B.2.7). *Under graphical model in Fig. 1, the estimators compare as:*

$$\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{fed}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}})$$

where $\text{fed} \in \{\text{GD}, \text{1S-IVW}, \text{1S-SW}\}$. Theorem 2 shows that meta-analysis estimators typically exhibit larger variance. This happens as soon as treatment probabilities are not equal across centers, and the variance difference increases as the treatment probabilities $\{p_k\}_k$ become more distinct. Moreover, $\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) < \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}})$ whenever $\{p_k(1-p_k)\}_k$ differ across centers, and the difference increase as the difference between these quantities increase. We provide examples in Appendix B.2.7.

Special case: one RCT conducted across K centers. Consider the special case where $W \perp\!\!\!\perp H$, i.e., there is no edge from H to W in the graphical model of Figure 1. This corresponds to a single Bernoulli RCT with treatment probability p implemented across multiple centers. Then, under Condition 1, all estimators are asymptotically equivalent (see Appendix B.2.8) and thus meta-analysis estimators should be used as they require a single round of communication.




Figure 2: Graphical Model in the Heterogeneous Distributions Setting.

4 HETEROGENEOUS SCENARIOS

4.1 Distributional Shifts in Covariates

In model (2), we do not assume $H \perp\!\!\!\perp X$ and consider the graphical model in Figure 2, depicting *Heterogeneous* Bernoulli Trials in the distribution of $X|H$.

Here, $\mathbb{P}(Z) = \mathbb{P}(Y|X, W)\mathbb{P}(X|H)\mathbb{P}(W|H)\mathbb{P}(H)$, hence the observations $(X_{k,i}, W_{k,i}, Y_{k,i})_i$ are i.i.d. within center k but not necessarily across centers. We denote $\mathbb{E}(X_{k,i}) = \mu_k$ and $\mathbb{V}(X_{k,i}) = \Sigma_k$ and we have $\Sigma = \mathbb{V}(X) = \sum_{k=1}^K \rho_k \Sigma_k$ and $\mu = \mathbb{E}(X) = \sum_{k=1}^K \rho_k \mu_k$. Furthermore, the local ATEs $\tau_k = \mathbb{E}(Y_k(1) - Y_k(0)) = \mathbb{E}(X'_k)(\theta^{(1)} - \theta^{(0)})$ generally differ from each other and from the population ATE $\tau = \mathbb{E}(Y(1) - Y(0)) = \mathbb{E}(X')(\theta^{(1)} - \theta^{(0)})$.

Meta estimators. The ATE can be written as $\tau = \sum_{k=1}^K \rho_k \tau_k$, so the Meta-SW estimator remains unbiased (as n_k/n is an unbiased estimate of ρ_k) and has the same variance as in Table 1 with $\Sigma = \sum_{k=1}^K \rho_k \Sigma_k$ (proof in Appendix B.3.1). In contrast, the Meta-IVW estimator becomes unsuitable under distributional shifts: IVW weights give biased estimates of the ρ_k 's, leading to a biased estimate of τ .

Pool and GD estimators. The pooled and GD estimators are robust to covariate shifts, leading them to be also unbiased with same variance as in Table 1 (as proved in Appendix B.3.1).

One-shot estimators. Although the One-Shot IVW outcome parameters estimator still enjoys Theorem 1, the variance of the One-Shot SW one is impacted by the difference in population means at each center.

Proposition 2 (Larger Variance of 1S-SW, proof in B.3.1). *Under Condition 1, the one-shot federated estimators are unbiased and*

$$\mathbb{V}(\hat{\theta}_{\text{pool}}) = \mathbb{V}(\hat{\theta}_{\text{GD}}) = \mathbb{V}(\hat{\theta}_{\text{1S-IVW}}) \leq \mathbb{V}(\hat{\theta}_{\text{1S-SW}})$$

which yields

$$\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{GD}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{1S-IVW}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{1S-SW}})$$

Theorem 3 (Comparison of asymptotic variances under distributional shift).

$$\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{GD}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{1S-IVW}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}})$$

Note that in this setting the DM estimators will no longer be unbiased as there is a path between W and Y in Figure 2. Here, X is a sufficient adjustment set and adding H could be harmful, particularly when the association between X and Y is weak and the association between H and W is strong (Rotnitzky and Smucler, 2020, Lemma 2 therein).




Figure 3: Graphical model for the heterogeneous center effects setting.

4.2 Center Effects

We now consider the graphical model shown in Figure 3, exhibiting an effect of the center H onto the outcome Y . The distribution of Z decomposes as $\mathbb{P}(Z) = \mathbb{P}(Y|X, W, H)\mathbb{P}(X|H)\mathbb{P}(W|H)\mathbb{P}(H)$. We modify model (2) to account for a constant center effect on individual outcomes by adding a term $h_k \in \mathbb{R}$:

$$Y_{k,i}(w) = c^{(w)} + h_k + X_{k,i}\beta^{(w)} + \varepsilon_i(w) \quad (16)$$

Model (16) accounts for the possibility that centers may have different baselines in individual outcomes resulting from varying practices or organizational contexts. Here, τ and τ_k are still defined as $\mathbb{E}(Y(1) - Y(0)) = c^{(1)} - c^{(0)} + \mathbb{E}(X)(\beta^{(1)} - \beta^{(0)})$ and $\mathbb{E}(Y_k(1) - Y_k(0)) = c^{(1)} - c^{(0)} + \mathbb{E}(X_k)(\beta^{(1)} - \beta^{(0)})$ respectively since the h_k terms cancel out in the differences. In this setting, aggregating multiple RCTs into the pooled data is not itself an RCT because H is now a confounder, affecting both the outcome variable $Y_{k,i}(w)$ through h_k and the treatment variable $W_{k,i}$ through the treatment probability p_k , thereby violating the ignorability assumption (c). Therefore, the (unadjusted) pooled OLS estimator (4) is biased (see proof in Appendix B.4.1). An unbiased estimator can be obtained by adjusting the model to incorporate the center effect.

Meta estimators. While most estimators need to be adjusted and thus require prior knowledge on the underlying model, Meta-analysis estimators can be applied directly without such modifications. We prove in Appendix B.4 that under Graphical Model 3 and model (16), the Meta-IVW (which is relevant here as $H \perp\!\!\!\perp X$) and Meta-SW estimators remain unbiased, with asymptotic variances as in Table 1.

Adjusted gradient-based estimator. We augment the design matrix X with $K - 1$ dummy variables $H = \{H_2, \dots, H_K\}$ and note $\tilde{X}'_{k,i} = (1, X_{k,i}, H_{2,i}, \dots, H_{K,i})$. Then, the method is the same as in Section 3.1.3: $\hat{\theta}_{\text{GD}\circ}^{(w)} = (\hat{c}^{(w)}, \hat{\beta}^{(w)}, \hat{h}_2, \dots, \hat{h}_K) \in \mathbb{R}^{d+K}$ is obtained with FedAvg and used to compute the local ATEs before final aggregation.

Definition 9 (Adjusted GD estimator).

$$\hat{\tau}_{\text{GD}\circ} = \frac{1}{n} \sum_{k=1}^K \sum_{i=1}^{n_k} (\tilde{X}'_{k,i} (\hat{\theta}_{\text{GD}\circ}^{(1)} - \hat{\theta}_{\text{GD}\circ}^{(0)}))$$

This estimator is unbiased, and we do not pay a price in terms of asymptotic variance in adjusting the variables H . Indeed, its asymptotic variance is equal to the unadjusted $\hat{\tau}_{\text{GD}}$'s

one in Table 1 (proof in Appendix B.4.2) since h_k is equal in both treatment arms and cancel out in the difference of the true parameters.

Adjusted one-shot federated estimators. Like the vanilla pooled and GD estimators, the one-shot estimators are biased in the presence of center effects. However, their adjustment procedure is different because the one-shot procedure does not allow the inclusion of membership variables, as the variance matrices of the local dummy-augmented datasets are not full rank, violating Condition 1. Instead, we compute the OLS $\hat{\theta}_k$ at each center, then share and aggregate only the coefficients $\hat{\beta}_k^{(w)}$, without federating the locally estimated intercepts $\hat{a}_k^{(w)} = \hat{c}^{(w)} + \hat{h}_k$.

Definition 10 (Adjusted one-shot ATE estimators).

$$\begin{aligned}\hat{\tau}_{1S-SW\circ} &= \frac{1}{n} \sum_{k=1}^K \sum_{i=1}^{n_k} (\hat{a}_k^{(1)} - \hat{a}_k^{(0)} + X_{k,i}(\hat{\beta}_{SW\circ}^{(1)} - \hat{\beta}_{SW\circ}^{(0)})) \\ \hat{\tau}_{1S-IVW\circ} &= \frac{1}{n} \sum_{k=1}^K \sum_{i=1}^{n_k} (\hat{a}_k^{(1)} - \hat{a}_k^{(0)} + X_{k,i}(\hat{\beta}_{IVW\circ}^{(1)} - \hat{\beta}_{IVW\circ}^{(0)}))\end{aligned}$$

with $\hat{\beta}_{IVW\circ}^{(w)} = \frac{\sum_{k=1}^K \mathbb{V}(\hat{\beta}_k^{(w)})^{-1} \hat{\beta}_k^{(w)}}{\sum_{l=1}^K \mathbb{V}(\hat{\beta}_l^{(w)})^{-1}}$, $\hat{\beta}_{SW\circ}^{(w)} = \sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \hat{\beta}_k^{(w)}$.


Note that, in general, $\hat{\beta}_{IVW\circ}^{(w)} \neq \hat{\beta}_{IVW}^{(w)}$ as the aggregation weights $\mathbb{V}(\hat{\beta}_k^{(w)})^{-1}$ and $\mathbb{V}(\hat{\theta}_k^{(w)})^{-1}$ are different.

The variances of the adjusted one-shot estimators are affected by the lack of federation of the local intercepts $\{\hat{a}_k^{(w)}\}_k$, which converge at a rate of $1/n_k^{(w)}$ rather than $1/n^{(w)}$. Additionally, they estimate the center effects twice in each center on independent data, whereas GD \circ and pool \circ estimate them on $n^{(w)}$ observations. As a result, adjusted one-shot estimators generally underperform compared to other methods (see simulations in Appendix C.2).

5 SIMULATIONS

We compare the following estimators: Meta-SW (Definition 2), Meta-IVW (Definition 3), 1S-SW (Definition 6), 1S-IVW (Definition 7) and GD (Definition 8), and the adjusted estimators GD \circ (Definition 9), 1S-SW \circ and 1S-IVW \circ (Definition 10). We also include the pooled estimator (Pool, Eq. 4) as a baseline. For the Meta-IVW estimator, we use empirical estimates of $\mathbb{V}^\infty(\hat{\tau}_k)^{-1}$ for the aggregation weights.

We generate data according to the graphical models in Figures (1) and (2), with $K = 5$ centers and $d = 10$ covariates. We consider two magnitudes of sample sizes, referred to as *Large* ($\forall k, n_k = 20*d$) and *Small* ($\forall k, n_k = 6*d$) local sample sizes. We consider the following treatment assignment probabilities for the *Large* setting: $p_1 = p_2 = p_3 = 0.9, p_4 = p_5 = 0.1$. In the *Small* regime we choose less extreme probabilities in order to guarantee Condition 1: $p_1 = p_2 = p_3 = 0.65, p_4 = p_5 = 0.35$. For each scenario considered, we perform 2000



(a) Homogeneous Case (Figure 1): Large (left) and Small (right) regimes (b) Fully Heterogeneous Case (Figure 5): Large regime, unadjusted (left); Large regime, adjusted (middle); Small regime, adjusted (right)

Figure 4: Multicentric ATE Estimation under Homogeneous and Heterogeneity Scenarios

simulations and display the distribution of the global estimates of the ATE given by each estimator. More details about the simulations can be found in Appendix C.1.

Homogeneous case. The results displayed in Figure 4a (left) are in agreement with Theorem 2: the variance of the meta estimators are larger than the pooled, one-shot and gradient-based estimators when the assignment probabilities p_k differ from one center to another, with the variance of the Meta-IVW being smaller than that of Meta-SW (Proposition 1). In the *Small* regime (Figure 4a, right), the variance of One-Shot SW is larger than compared to One-Shot IVW and GD, but we still have $\mathbb{V}(\hat{\tau}_{\text{pool}}) = \mathbb{V}(\hat{\tau}_{\text{GD}}) = \mathbb{V}(\hat{\tau}_{\text{1S-IVW}})$, in line with Theorem 1.

Heterogeneous case. We now consider a “fully heterogeneous” setting combining three sources of heterogeneity (see graphical model in Figure 5): different covariate distributions across centers, center effects, and different p_k (see Appendix C.3 for the chosen values of $\{(\mu_k, \Sigma_k, h_k, p_k)\}_k$). Figure 4b (left) shows that the pooled, one-shot and gradient-based estimators are biased when we do not adjust for the center effects. Here, only the Meta SW estimator is unbiased.




Figure 5: Graphical model in fully heterogeneous case.

Figure 4b (middle, right) shows the results of adjusted estimators, as described in Section 4.2. Adjusting removes their bias but comes at a cost for the one-shot estimators, whose variances are impacted by center effects, unlike the meta and gradient-based estimators. In the *Small* regime, the one-shot estimators further suffer from the combination of differences in covariate means and unshared local intercepts, which inflate $\mathbb{V}^\infty(\hat{a}_k^{(w)})$. On the other hand, the variance of the Meta-SW estimator is nearly equal to that of the adjusted pooled and

adjusted GD estimators.

6 CONCLUSION

After clearly defining the population and estimand of interest, our findings can be turned into clear guidelines for practitioners that we summarize as a decision diagram (Figure 6 in Appendix A). We recommend the one-shot IVW estimator when each center can perform a local OLS regression (Condition 1) and there are no center effects. However, this estimator requires centers to share sample covariance matrices, which can be impractical in high-dimensional settings or when privacy is a concern. On the other hand, meta estimators are preferable when center effects are present or when there is limited prior knowledge about the underlying model, provided Condition 1 holds. Caution is required with Meta-IVW, which has lower variance than Meta-SW but is biased when centers analyze different populations. Finally, our (adjusted) GD estimator allows us to estimate the ATE under the weaker Condition 2 with the same precision as if the data were pooled, regardless of the setting.

Several challenges remain for the deployment of these methods, particularly in medical contexts. Future work includes addressing covariate mismatch, where some features are missing in specific centers, estimating non-collapsible causal measures (e.g., odds ratio), and extensions to observational studies. Finally, a key practical challenge in multi-centric studies is to ensure consistent data encoding, especially for outcomes.

Acknowledgments

This work has been done in the frame of the PEPR SN SMATCH project and has benefited from a governmental grant managed by the Agence Nationale de la Recherche under the France 2030 programme, reference ANR-22-PESN-0003. Rémi Khellaf was supported by an AXIAUM fellowship.

References

- David Benkeser, Iván Díaz, Alex Luedtke, Jodi Segal, Daniel Scharfstein, and Michael Rosenblum. Improving precision and power in randomized trials for covid-19 treatments using covariate adjustment, for binary, ordinal, and time-to-event outcomes. *Biometrics*, 77(4): 1467–1481, 2021.
- Jesse A. Berlin and Robert M. Golub. Meta-analysis as evidence: Building a better pyramid. *JAMA*, 312(6):603–606, 2014. doi: 10.1001/jama.2014.8167.
- Christopher Blunt. *Hierarchies of evidence in evidence-based medicine*. PhD thesis, London School of Economics and Political Science, 2015.
- Michael Borenstein, Larry V Hedges, Julian PT Higgins, and Hannah R Rothstein. *Introduction to meta-analysis*. John Wiley & Sons, 2021.
- Carly Lupton Brantner, Ting-Hsuan Chang, Trang Quynh Nguyen, Hwanhee Hong, Leon Di Stefano, and Elizabeth A Stuart. Methods for integrating trials and non-experimental data to examine treatment effect heterogeneity. *Statistical Science*, 38(4):640–654, 2023.
- Sicong Che, Zhaoming Kong, Hao Peng, Lichao Sun, Alex Leow, Yong Chen, and Lifang He. Federated multi-view learning for private medical data integration and analysis. *ACM Transactions on Intelligent Systems and Technology (TIST)*, 13(4):1–23, 2022.
- Mackenzie J Edmondson, Chongliang Luo, and Yong Chen. Statistical analysis—meta-analysis/reproducibility. In *Clinical Applications of Artificial Intelligence in Real-World Data*, pages 125–139. Springer, 2023.
- European Medicines Agency. ICH E9 Statistical Principles for Clinical Trials: Scientific Guideline, 2024.
- French Health Authority. Pricing & reimbursement of drugs and hta policies in france, 2024. URL https://www.has-sante.fr/upload/docs/application/pdf/2014-03/pricing_reimbursement_of_drugs_and_hta_policies_in_france.pdf.
- Christophe Giraud, Sylvie Huet, and Nicolas Verzelen. High-dimensional regression with unknown variance. 2012.
- Tianyu Guo, Sai Praneeth Karimireddy, and Michael I. Jordan. Collaborative heterogeneous causal inference beyond meta-analysis. *arXiv preprint arXiv:2404.15746*, 2024.
- Gordon Guyatt, Drummond Rennie, Maureen O. Meade, and Deborah J. Cook. *Users’ Guides to the Medical Literature : A Manual for Evidence-Based Clinical Practice*. McGraw-Hill Education, New York, 2015.
- Larry Han, Jue Hou, Kelly Cho, Rui Duan, and Tianxi Cai. Federated adaptive causal estimation (face) of target treatment effects. *arXiv preprint arXiv:2112.09313*, 2021.

- Larry Han, Zhu Shen, and José R. Zubizarreta. Multiply robust federated estimation of targeted average treatment effects. In Alice Oh, Tristan Naumann, Amir Globerson, Kate Saenko, Moritz Hardt, and Sergey Levine, editors, *Advances in Neural Information Processing Systems 36: Annual Conference on Neural Information Processing Systems 2023, NeurIPS 2023, New Orleans, LA, USA, December 10 - 16, 2023*, 2023.
- JM Hernan, MA Robins. *Causal Inference: What If*. Boca Raton: Chapman & Hall/CRC., 2020.
- Julian PT Higgins, James Thomas, Jacqueline Chandler, Miranda Cumpston, Tianjing Li, Matthew J Page, and Vivian A Welch, editors. *Cochrane Handbook for Systematic Reviews of Interventions*. John Wiley & Sons, Chichester, UK, 2nd edition, 2019.
- John E Hunter and Frank L Schmidt. *Methods of meta-analysis: Correcting error and bias in research findings*. Sage, 2004.
- Guido W Imbens and Donald B Rubin. *Causal Inference in Statistics, Social, and Biomedical Sciences*. Cambridge University Press, Cambridge UK, 2015.
- Brennan C Kahan, Fan Li, Andrew J Copas, and Michael O Harhay. Estimands in cluster-randomized trials: choosing analyses that answer the right question. *International Journal of Epidemiology*, 52(1):107–118, 2023.
- Peter Kairouz, H Brendan McMahan, Brendan Avent, Aurélien Bellet, Mehdi Bennis, Arjun Nitin Bhagoji, Kallista Bonawitz, Zachary Charles, Graham Cormode, Rachel Cummings, et al. Advances and open problems in federated learning. *Foundations and trends® in machine learning*, 14(1–2):1–210, 2021.
- Ahmed Khaled, Konstantin Mishchenko, and Peter Richtárik. Tighter theory for local SGD on identical and heterogeneous data. In *AISTATS*, 2020.
- Anastasia Koloskova, Nicolas Loizou, Sadra Boreiri, Martin Jaggi, and Sebastian Stich. A unified theory of decentralized SGD with changing topology and local updates. In Hal Daumé III and Aarti Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 5381–5393. PMLR, 13–18 Jul 2020. URL <https://proceedings.mlr.press/v119/koloskova20a.html>.
- Lihua Lei and Peng Ding. Regression adjustment in completely randomized experiments with a diverging number of covariates. *Biometrika*, 108(4):815–828, 2021.
- A Liberati, DG Altman, J Tetzlaff, and C Mulrow. G0tzsche pc, ioannidis jp, et al. the prisma statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol*, 62(10):e1–34, 2009.
- Winston Lin. Agnostic notes on regression adjustments to experimental data: Reexamining freedman’s critique. 2013.

- Disha Makhija, Joydeep Ghosh, and Yejin Kim. Federated learning for estimating heterogeneous treatment effects. *CoRR*, abs/2402.17705, 2024. doi: 10.48550/ARXIV.2402.17705. URL <https://doi.org/10.48550/arXiv.2402.17705>.
- Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Aguera y Arcas. Communication-efficient learning of deep networks from decentralized data. In *Artificial intelligence and statistics*, pages 1273–1282. PMLR, 2017.
- David Moher, Deborah J Cook, Susan Eastwood, Ingram Olkin, Drummond Rennie, and Donna F Stroup. Improving the quality of reports of meta-analyses of randomised controlled trials: the quorum statement. *The Lancet*, 354(9193):1896–1900, 1999.
- Tim P Morris, David J Fisher, Michael G Kenward, and James R Carpenter. Meta-analysis of gaussian individual patient data: Two-stage or not two-stage? *Statistics in medicine*, 37(9):1419–1438, 2018.
- Steven Piantadosi, David P Byar, and Sylvan B Green. The ecological fallacy. *American journal of epidemiology*, 127(5):893–904, 1988.
- Mattia Proserpi, Yi Guo, Matt Sperrin, James S Koopman, Jae S Min, Xing He, Shannan Rich, Mo Wang, Iain E Buchan, and Jiang Bian. Causal inference and counterfactual prediction in machine learning for actionable healthcare. *Nature Machine Intelligence*, 2(7):369–375, 2020.
- James Robins. A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect. *Mathematical modelling*, 7(9-12):1393–1512, 1986.
- Andrea Rotnitzky and Ezequiel Smucler. Efficient adjustment sets for population average causal treatment effect estimation in graphical models. *Journal of Machine Learning Research*, 21(188):1–86, 2020. URL <http://jmlr.org/papers/v21/19-1026.html>.
- Donald B Rubin. Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of educational psychology*, 66(5):688–701, 1974. ISSN 0022-0663.
- Michael Seo, Ian R White, Toshi A Furukawa, Hissei Imai, Marco Valgimigli, Matthias Egger, Marcel Zwahlen, and Orestis Efthimiou. Comparing methods for estimating patient-specific treatment effects in individual patient data meta-analysis. *Statistics in medicine*, 40(6):1553–1573, 2021.
- Micah J Sheller, Brandon Edwards, G Anthony Reina, Jason Martin, Sarthak Pati, Aikaterini Kotrotsou, Mikhail Milchenko, Weilin Xu, Daniel Marcus, Rivka R Colen, et al. Federated learning in medicine: facilitating multi-institutional collaborations without sharing patient data. *Scientific reports*, 10(1):12598, 2020.
- Jerzy Splawa-Neyman, D. M. Dabrowska, and T. P. Speed. On the Application of Probability Theory to Agricultural Experiments. Essay on Principles. Section 9. *Statistical Science*, 5(4):465 – 472, 1990. doi: 10.1214/ss/1177012031. URL <https://doi.org/10.1214/ss/1177012031>.

- Sebastian U. Stich. Local sgd converges fast and communicates little. In *ICLR*, 2019.
- Xiaoqing Tan, Chung-Chou H Chang, Ling Zhou, and Lu Tang. A tree-based model averaging approach for personalized treatment effect estimation from heterogeneous data sources. In *International Conference on Machine Learning*, pages 21013–21036. PMLR, 2022.
- Jean Ogier du Terrail, Quentin Klopfenstein, Honghao Li, Imke Mayer, Nicolas Loiseau, Mohammad Hallal, Félix Balazard, and Mathieu Andreux. Fedeca: A federated external control arm method for causal inference with time-to-event data in distributed settings. *arXiv preprint arXiv:2311.16984*, 2023.
- Anastasios A Tsiatis, Marie Davidian, Min Zhang, and Xiaomin Lu. Covariate adjustment for two-sample treatment comparisons in randomized clinical trials: a principled yet flexible approach. *Statistics in medicine*, 27(23):4658–4677, 2008.
- U.S. Food and Drug Administration. Adjusting for covariates in randomized clinical trials for drugs and biological products, 2023.
- Kelly Van Lancker, Frank Bretz, and Oliver Dukes. Covariate adjustment in randomized controlled trials: General concepts and practical considerations. *Clinical Trials*, page 17407745241251568, 2024.
- Thanh Vinh Vo, Arnab Bhattacharyya, Young Lee, and Tze-Yun Leong. An adaptive kernel approach to federated learning of heterogeneous causal effects. *Advances in Neural Information Processing Systems*, 35:24459–24473, 2022a.
- Thanh Vinh Vo, Young Lee, Trong Nghia Hoang, and Tze-Yun Leong. Bayesian federated estimation of causal effects from observational data. In *UAI*, 2022b.
- Stefan Wager. Stats 361: Causal inference. Technical report, 2020.
- Jiayi Wang, Shiqiang Wang, Rong-Rong Chen, and Mingyue Ji. A new theoretical perspective on data heterogeneity in federated optimization. In *International Conference on Machine Learning (ICML)*, 2024.
- Ruoxuan Xiong, Allison Koenecke, Michael Powell, Zhu Shen, Joshua T Vogelstein, and Susan Athey. Federated causal inference in heterogeneous observational data. *Statistics in Medicine*, 42(24):4418–4439, 2023.

Contents

1	INTRODUCTION	1
2	GENERAL FRAMEWORK	3
3	HOMOGENEOUS POPULATION	5
3.1	Definition of the Estimators	6
3.1.1	Meta-Analysis Estimators	6
3.1.2	One-Shot Federated Estimators	6
3.1.3	Gradient-based Federated Estimators	8
3.2	Comparison of the Federated Estimators	8
4	HETEROGENEOUS SCENARIOS	10
4.1	Distributional Shifts in Covariates	10
4.2	Center Effects	11
5	SIMULATIONS	12
6	CONCLUSION	14
A	DECISION DIAGRAM	21
B	PROOFS	22
B.1	Weighting methods	22
B.1.1	Probability of treatment in pooled dataset	22
B.1.2	Meta-IVW has minimum variance among unbiased aggregation-based estimators	22
B.2	Homogeneous setting	23
B.2.1	Properties of federated outcome estimators	23
B.2.2	Bias of the outcome model estimators	24
B.2.3	(Non-asymptotic) variance comparison of the outcome model parameters	25
B.2.4	Asymptotic variances of the outcome model parameters	26
B.2.5	Bias of the ATE estimators	27
B.2.6	Asymptotic variances of the ATE estimators	29
B.2.7	Comparison of asymptotic variances - General Case	34

B.2.8	Comparison of asymptotic variances - Special Case	35
B.3	Heterogeneous Settings	36
B.3.1	Distributional Shift in Covariates	36
B.4	Center Effects	39
B.4.1	Unadjusted Federated Estimators	40
B.4.2	With membership variable in the dataset	42
C	SIMULATIONS	43
C.1	Simulation parameters	43
C.2	Additional scenarios	44
C.2.1	Balanced datasets	44
C.2.2	Imbalanced datasets	45
C.3	Shift in covariates distributions	45
C.4	Center effects	46
C.4.1	No adjustment	47
C.4.2	Adjusted	48
C.5	Full heterogeneity: shifts in covariates distributions, center effects and in treatment allocation	49
C.6	G-Formula OLS covariate adjustment in non-linearity	49
D	Algorithms	50

APPENDIX

A DECISION DIAGRAM




Figure 6: Decision Diagram for Practitioners. The sign ★ denotes scenarios where the DM estimator is biased.

Our results yield clear guidelines for practitioners to select the most suitable estimator for different scenarios, which we present as a decision diagram in Figure 6. Note that each time we mention the use of a meta estimator in second position, it means that this meta estimator yields a valid unbiased estimator, but with a higher variance than the estimator in first position.

It is also worth mentioning that the Difference-in-Means estimator on the pooled individual data is biased whenever H acts as a confounder between W and Y , which happens when centers have distinct treatment probabilities and center effects. The DM estimator is also biased in the graphical model in Figure 2, although H is not technically a confounder in this setting.

B PROOFS

B.1 Weighting methods

B.1.1 Probability of treatment in pooled dataset

We consider that a center is included in the study if its sample size is strictly larger than zero, i.e. $\forall k, n_k > 0$. Since n_k is a binomial random variable of parameters n and ρ_k , we have $\mathbb{E}(n_k) = n\rho_k$ which yields $\rho_k = \mathbb{E}\left(\frac{n_k}{n}\right)$.

B.1.2 Meta-IVW has minimum variance among unbiased aggregation-based estimators

Proof of Proposition 1: Let $\hat{\tau}_k \sim \mathcal{N}(\tau, \mathbb{V}(\hat{\tau}_k))$ and K independent centers. We denote as $\hat{\tau} = \frac{\sum_{k=1}^K w_k \hat{\tau}_k}{\sum_{k=1}^K w_k}$ a w -weighted average of the local estimators of τ . We have:

$$\begin{aligned} \mathbb{V}(\hat{\tau}) &= \mathbb{V}\left(\frac{\sum_{k=1}^K w_k \hat{\tau}_k}{\sum_{k=1}^K w_k}\right) \\ &= \frac{1}{\left(\sum_{k=1}^K w_k\right)^2} \sum_{k=1}^K w_k^2 \mathbb{V}(\hat{\tau}_k) \\ &= \sum_{k=1}^K \left(\frac{w_k}{\sum_{k=1}^K w_k}\right)^2 \mathbb{V}(\hat{\tau}_k) \\ &= \sum_{k=1}^K u_k^2 \mathbb{V}(\hat{\tau}_k) \end{aligned}$$

with $u_k = \frac{w_k}{\sum_{k=1}^K w_k}$ and $\sum_{k=1}^K u_k = 1$

We want to minimize $\mathbb{V}(\hat{\tau})$ under the constraint $\sum_{k=1}^K u_k = 1$. We can use the Lagrange multiplier method to find the minimum of $\mathbb{V}(\hat{\tau})$ under this constraint. We define the Lagrangian function:

$$\mathcal{L}(u_1, \dots, u_K, \lambda) = \sum_{k=1}^K u_k^2 \mathbb{V}(\hat{\tau}_k) + \lambda \left(1 - \sum_{k=1}^K u_k\right)$$

Then we have:

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial u_k} &= 2u_k \mathbb{V}(\hat{\tau}_k) - \lambda \\ \frac{\partial \mathcal{L}}{\partial \lambda} &= 1 - \sum_{k=1}^K u_k \end{aligned}$$

To cancel out the two derivatives above, we get that $\forall k \in [1; K], u_k = \frac{\lambda}{2\mathbb{V}(\hat{\tau}_k)}$. Then starting back from the constraint we have:

$$\begin{cases} \sum_{k=1}^K u_k = 1 \\ u_k = \frac{\lambda}{2\mathbb{V}(\hat{\tau}_k)} \end{cases} \Rightarrow \begin{cases} u_k = \frac{\mathbb{V}(\hat{\tau}_k)^{-1}}{\sum_{k=1}^K \mathbb{V}(\hat{\tau}_k)^{-1}} \end{cases}$$

Injecting this result in $\mathbb{V}(\hat{\tau})$, we get:

$$\begin{aligned} \mathbb{V}(\hat{\tau}) &= \sum_{k=1}^K u_k^2 \mathbb{V}(\hat{\tau}_k) \\ &= \sum_{k=1}^K \left(\frac{\mathbb{V}(\hat{\tau}_k)^{-1}}{\sum_{k=1}^K \mathbb{V}(\hat{\tau}_k)^{-1}} \right)^2 \mathbb{V}(\hat{\tau}_k) \\ &= \frac{1}{\sum_{k=1}^K \mathbb{V}(\hat{\tau}_k)^{-1}} \end{aligned}$$

Finally, we get that $\forall k \in [1; K], u_k = \frac{\mathbb{V}(\hat{\tau}_k)^{-1}}{\sum_{k=1}^K \mathbb{V}(\hat{\tau}_k)^{-1}}$.

Therefore, $\hat{\tau}_{\text{Meta-IVW}}$ is the minimum-variance unbiased estimator of τ among the class of aggregation-based estimators.

B.2 Homogeneous setting

We prove in this section the results in Section 3, assuming Condition 1 and the graphical model in Figure 1.

B.2.1 Properties of federated outcome estimators

Proof of Theorem 1:

$$\begin{aligned} \hat{\theta}_{\text{IVW}}^{(w)} &= \frac{\sum_{k=1}^K \mathbb{V}(\hat{\theta}_k^{(w)})^{-1} \hat{\theta}_k^{(w)}}{\sum_{k=1}^K \mathbb{V}(\hat{\theta}_k^{(w)})^{-1}} \\ &= \frac{\sum_{k=1}^K \left(\frac{1}{\sigma^2} X_k^{(w)\top} X_k^{(w)} \right) \left(X_k^{(w)\top} X_k^{(w)} \right)^{-1} X_k^{(w)\top} y_k^{(w)}}{\sum_{k=1}^K \left(\frac{1}{\sigma^2} X_k^{(w)\top} X_k^{(w)} \right)} \\ &= \frac{\sum_{k=1}^K X_k^{(w)\top} y_k^{(w)}}{\sum_{k=1}^K X_k^{(w)\top} X_k^{(w)}} \\ &= \frac{X_{\text{pool}}^{(w)\top} y_{\text{pool}}^{(w)}}{X_{\text{pool}}^{(w)\top} X_{\text{pool}}^{(w)}} && \text{by sum of matrix product} \\ &= \hat{\theta}_{\text{pool}}^{(w)} \end{aligned}$$

B.2.2 Bias of the outcome model estimators

Unbiasedness of $\hat{\theta}_{\text{pool}}^{(w)}$:

$$\begin{aligned}
\mathbb{E} \left(\hat{\theta}_{\text{pool}}^{(w)} \right) &= \mathbb{E} \left(\left(X'^{(w)\top} X'^{(w)} \right)^{-1} X'^{(w)\top} y \right) \\
&= \mathbb{E} \left(\mathbb{E} \left(\left(X'^{(w)\top} X'^{(w)} \right)^{-1} X'^{(w)\top} y \mid X'^{(w)} \right) \right) \\
&= \mathbb{E} \left(\left(X'^{(w)\top} X'^{(w)} \right)^{-1} X'^{(w)\top} \mathbb{E} \left(X'^{(w)} \theta + \varepsilon \mid X'^{(w)} \right) \right) \\
&= \mathbb{E} \left(\left(X'^{(w)\top} X'^{(w)} \right)^{-1} X'^{(w)\top} X'^{(w)} \theta + 0 \right) \\
&= \theta
\end{aligned}$$

Unbiasedness of $\hat{\theta}_{\text{GD}}^{(w)}$: under convergence of Algorithm D, $\hat{\theta}_{\text{GD}}^{(w)} = \hat{\theta}_{\text{pool}}^{(w)}$ which implies $\mathbb{E} \left(\hat{\theta}_{\text{GD}}^{(w)} \right) = \mathbb{E} \left(\hat{\theta}_{\text{pool}}^{(w)} \right) = \theta$.

Unbiasedness of $\hat{\theta}_{\text{1S-IVW}}^{(w)}$:

$$\begin{aligned}
\mathbb{E} \left(\hat{\theta}_{\text{IVW}}^{(w)} \right) &= \mathbb{E} \left(\hat{\theta}_{\text{pool}}^{(w)} \right) && \text{using Theorem 1} \\
&= \theta
\end{aligned}$$

Unbiasedness of $\hat{\theta}_{\text{1S-SW}}^{(w)}$:

We condition on the realization of H to account for the variability in the $\{n_k^{(w)}\}_k$ terms.

$$\begin{aligned}
\mathbb{E} \left(\mathbb{E} \left(\hat{\theta}_{\text{1S-SW}}^{(w)} \mid H \right) \right) &= \mathbb{E} \left(\mathbb{E} \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \hat{\theta}_k^{(w)} \mid H \right) \right) \\
&= \mathbb{E} \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \mathbb{E} \left(\hat{\theta}_k^{(w)} \mid H \right) \right) \\
&= \sum_{k=1}^K \frac{\mathbb{E}(n_k^{(w)})}{n^{(w)}} \theta \\
&= \theta
\end{aligned}$$

B.2.3 (Non-asymptotic) variance comparison of the outcome model parameters

Variance of the local outcome model OLS estimators:

$$\begin{aligned}
\mathbb{V}\left(\hat{\theta}_k^{(w)}\right) &= \mathbb{E}\left(\mathbb{V}\left(\left(\hat{\theta}_k^{(w)} \mid X_k^{\prime(w)}\right)\right)\right) + \mathbb{V}\left(\mathbb{E}\left(\hat{\theta}_k^{(w)} \mid X_k^{\prime(w)}\right)\right) \\
&= \mathbb{E}\left(\mathbb{V}\left(\left(X_k^{\prime(w)\top} X_k^{\prime(w)}\right)^{-1} X_k^{\prime(w)\top} y_k^{(w)} \mid X_k^{\prime(w)}\right)\right) + 0 \\
&= \mathbb{E}\left(\left(X_k^{\prime(w)\top} X_k^{\prime(w)}\right)^{-1} X_k^{\prime(w)\top} \mathbb{V}\left(y_k^{(w)} \mid X_k^{\prime(w)}\right) X_k^{\prime(w)} \left(X_k^{\prime(w)\top} X_k^{\prime(w)}\right)^{-1}\right) \\
&= \mathbb{E}\left(\left(X_k^{\prime(w)\top} X_k^{\prime(w)}\right)^{-1} X_k^{\prime(w)\top} \sigma^2 I_p X_k^{\prime(w)} \left(X_k^{\prime(w)\top} X_k^{\prime(w)}\right)^{-1}\right) \\
&= \sigma^2 \mathbb{E}\left(\left(X_k^{\prime(w)\top} X_k^{\prime(w)}\right)^{-1}\right)
\end{aligned}$$

Similarly we get $\mathbb{V}\left(\hat{\theta}_{\text{IVW}}^{(w)}\right) = \mathbb{V}\left(\hat{\theta}_{\text{pool}}^{(w)}\right) = \mathbb{V}\left(\hat{\theta}_{\text{GD}}^{(w)}\right) = \sigma^2 \mathbb{E}\left(\left(X^{\prime(w)\top} X^{\prime(w)}\right)^{-1}\right)$.

The variance of the One-Shot SW outcome model parameters estimator is obtained using the law of total variance over H :

$$\begin{aligned}
\mathbb{V}\left(\hat{\theta}_{\text{SW}}^{(w)}\right) &= \mathbb{E}\left(\mathbb{V}\left(\hat{\theta}_{\text{SW}}^{(w)} \mid H\right)\right) + \mathbb{V}\left(\mathbb{E}\left(\hat{\theta}_{\text{SW}}^{(w)} \mid H\right)\right) \\
&= \sum_{k=1}^K \mathbb{E}\left(\frac{n_k^2}{n^2}\right) \mathbb{V}\left(\hat{\theta}_k^{(w)}\right) + 0 \\
&= \sigma^2 \sum_{k=1}^K \frac{\mathbb{E}(n_k^2)}{n^2} \mathbb{E}\left(X_k^{\prime(w)\top} X_k^{\prime(w)-1}\right)
\end{aligned}$$

by independence of the centers for the last equality.

We now compare the variances above. First notice that:

$$\mathbb{E}\left(\left(X_{\text{pool}}^{\prime(w)\top} X_{\text{pool}}^{(w)}\right)^{-1}\right) = \frac{1}{n} \mathbb{E}\left(\frac{1}{\frac{1}{n} X_{\text{pool}}^{\prime(w)\top} X_{\text{pool}}^{(w)}}\right) = \frac{1}{n} \mathbb{E}\left(\frac{1}{\sum_{k=1}^K \frac{\mathbb{E}(n_k)}{n} \left(\frac{1}{\mathbb{E}(n_k)} X_k^{\prime(w)\top} X_k^{(w)}\right)}\right)$$

By applying Jensen's inequality on the inverse function over the space of semi positive definite matrices and with weights summing to 1 $\{\mathbb{E}(n_k)/n\}_{k \in [1, K]}$:

$$\begin{aligned}
\mathbb{E}\left(\left(X_{\text{pool}}^{\prime(w)\top} X_{\text{pool}}^{(w)}\right)^{-1}\right) &\preceq \frac{1}{n} \sum_{k=1}^K \frac{\mathbb{E}(n_k)}{n} \mathbb{E}\left(\left(\frac{1}{\mathbb{E}(n_k)} X_k^{\prime(w)\top} X_k^{(w)}\right)^{-1}\right) \\
&\preceq \sum_{k=1}^K \frac{\mathbb{E}(n_k)^2}{n^2} \mathbb{E}\left(\left(X_k^{\prime(w)\top} X_k^{(w)}\right)^{-1}\right) \\
&\preceq \sum_{k=1}^K \frac{\mathbb{E}(n_k^2)}{n^2} \mathbb{E}\left(\left(X_k^{\prime(w)\top} X_k^{(w)}\right)^{-1}\right) \quad \text{as } \mathbb{E}(n_k)^2 \leq \mathbb{E}(n_k^2)
\end{aligned}$$

which leads to the conclusion of Proposition 2: $\mathbb{V}\left(\hat{\theta}_{\text{pool}}^{(w)}\right) = \mathbb{V}\left(\hat{\theta}_{\text{GD}}^{(w)}\right) = \mathbb{V}\left(\hat{\theta}_{\text{IVW}}^{(w)}\right) \preceq \mathbb{V}\left(\hat{\theta}_{\text{SW}}^{(w)}\right)$.

B.2.4 Asymptotic variances of the outcome model parameters

Asymptotically, we have

$$X_k'^{(w)\top} X_k'^{(w)} = \frac{1}{n_k^{(w)}} \sum_{i=1}^{n_k^{(w)}} X_{k,i}'^{(w)\top} X_{k,i}'^{(w)}$$

$$\xrightarrow{n_k^{(w)} \rightarrow \infty} A_k$$

$$\left(X_k'^{(w)\top} X_k'^{(w)} \right)^{-1} \rightarrow A_k^{-1} \quad \text{by continuous mapping}$$

where $A_k = \begin{pmatrix} 1 & \mu_{k,1} & \cdots & \mu_{k,p} \\ \mu_{k,1} & & & \\ \vdots & & \Sigma_k & \\ \mu_{k,p} & & & \end{pmatrix}$ and $\mu_{k,j}$ is the mean of the j -th covariate and $\Sigma_k =$

$\mathbb{E} \left(X_k^{(w)\top} X_k^{(w)} \right)$ is the covariate matrix in center k .

Therefore we get the asymptotic variance of the local outcome model parameters:

$$\mathbb{V}^\infty \left(\hat{\theta}_k^{(w)} \right) = \frac{\sigma^2}{n_k^{(w)}} A_k^{-1}$$

$$\mathbb{V}^\infty \left(\hat{\beta}_k^{(w)} \right) = \frac{\sigma^2}{n_k^{(w)}} \Sigma_k^{-1}$$

Under the graphical model in Figure 1, $\Sigma_k = \Sigma$ and $A_k = A$ which yields:

$$\text{Local OLS} \quad \mathbb{V}^\infty \left(\hat{\theta}_k^{(w)} \right) = \sigma^2 \left(X_k'^{(w)\top} X_k'^{(w)} \right)^{-1} = \frac{\sigma^2}{n_k^{(w)}} A^{-1}$$

$$\text{Pool OLS} \quad \mathbb{V}^\infty \left(\hat{\theta}_{\text{pool}}^{(w)} \right) = \sigma^2 \left(X'^{(w)\top} X'^{(w)} \right)^{-1} = \frac{\sigma^2}{n^{(w)}} A^{-1}$$

$$\text{GD} \quad \mathbb{V}^\infty \left(\hat{\theta}_{\text{GD}}^{(w)} \right) = \sigma^2 \left(X'^{(w)\top} X'^{(w)} \right)^{-1} = \frac{\sigma^2}{n^{(w)}} A^{-1}$$

$$\text{1S-SW} \quad \mathbb{V}^\infty \left(\hat{\theta}_{\text{SW}}^{(w)} \right) = \mathbb{E} \left(\mathbb{V}^\infty \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \hat{\theta}_k^{(w)} \mid H \right) \right) + 0$$

$$= \mathbb{E} \left(\sum_{k=1}^K \frac{n_k^{(w)^2}{n^{(w)^2}} \mathbb{V}^\infty \left(\hat{\theta}_k^{(w)} \mid H \right) \right)$$

$$= \mathbb{E} \left(\sum_{k=1}^K \left(\frac{n_k^{(w)}}{n^{(w)}} \right)^2 \frac{\sigma^2}{n_k^{(w)}} A^{-1} \right)$$

$$= \frac{\sigma^2}{n^{(w)^2}} A^{-1} \sum_{k=1}^K \mathbb{E}(n_k^{(w)})$$

$$= \frac{\sigma^2}{n^{(w)}} A^{-1}$$

$$\text{1S-IVW} \quad \mathbb{V}^\infty \left(\hat{\theta}_{\text{IVW}}^{(w)} \right) = \mathbb{V}^\infty \left(\frac{\sum_{k=1}^K \left(\mathbb{V}^\infty(\hat{\theta}_k^{(w)})^{-1} \hat{\theta}_k^{(w)} \right)}{\sum_{k=1}^K \mathbb{V}^\infty(\hat{\theta}_k^{(w)})^{-1}} \right)$$

$$= \mathbb{V} \left(\frac{\sum_{k=1}^K \frac{1}{\sigma^2} X_k'^{(w)\top} X_k'^{(w)} \times \left(X_k'^{(w)\top} X_k'^{(w)} \right)^{-1} X_k'^{(w)\top} y_k^{(w)}}{\sum_{k=1}^K \frac{1}{\sigma^2} X_k'^{(w)\top} X_k'^{(w)}} \right)$$

$$= \mathbb{V} \left(\frac{\sum_{k=1}^K X_k'^{(w)\top} y_k^{(w)}}{\sum_{k=1}^K X_k'^{(w)\top} X_k'^{(w)}} \right)$$

$$= \frac{\sum_{k=1}^K X_k'^{(w)\top} \mathbb{V}(y_k^{(w)}) X_k'^{(w)}}{\left(\sum_{k=1}^K X_k'^{(w)\top} X_k'^{(w)} \right)^2}$$

$$= \frac{\sigma^2}{\sum_{k=1}^K X_k'^{(w)\top} X_k'^{(w)}}$$

$$= \frac{\sigma^2}{n^{(w)}} A^{-1}$$

So under this model, for $w \in \{0, 1\}$, $\mathbb{V}^\infty \left(\hat{\theta}_{\text{GD}}^{(w)} \right) = \mathbb{V}^\infty \left(\hat{\theta}_{\text{SW}}^{(w)} \right) = \mathbb{V}^\infty \left(\hat{\theta}_{\text{IVW}}^{(w)} \right) = \frac{\sigma^2}{n^{(w)}} A^{-1}$. These results, along with the associated communication costs, are summarized in Table 2.

B.2.5 Bias of the ATE estimators

We now prove that under the graphical model in Figure 1 all the estimators are unbiased.

Estimator	Notation	Condition	\mathbb{V}^∞	Com. rounds	Com. cost
Local	$\hat{\theta}_k^{(w)}$	Condition 1	$\frac{\sigma^2}{n_k^{(w)}} A^{-1}$	0	0
One-Shot SW	$\hat{\theta}_{\text{IS-SW}}^{(w)}$ (Eq. 9)	Condition 1	$\frac{\sigma^2}{n^{(w)}} A^{-1}$	1	$O(d+1)$
One-Shot IVW	$\hat{\theta}_{\text{IS-IVW}}^{(w)}$ (Eq. 10)	Condition 1	$\frac{\sigma^2}{n^{(w)}} A^{-1}$	1	$O(d^2)$
GD-Federated	$\hat{\theta}_{\text{GD}}^{(w)}$ (Alg. D)	Condition 2	$\frac{\sigma^2}{n^{(w)}} A^{-1}$	T	$O(Td)$
Pool	$\hat{\theta}_{\text{pool}}^{(w)}$	Condition 2	$\frac{\sigma^2}{n^{(w)}} A^{-1}$	—	—

Table 2: Properties of the (unbiased) estimators of the outcome model parameters in the homogeneous setting: asymptotic variance, number of communication rounds and total communication cost (in number of floats per center).

First, notice that for any random variable $U \in \mathbb{R}^{n \times d+1}$, $w \in \{0, 1\}$,

$$\begin{aligned}
\mathbb{E}(U\hat{\theta}^{(w)}) &= \mathbb{E}\left(U(X'^{(w)\top} X'^{(w)})^{-1} X'^{(w)} Y^{(w)}\right) \\
&= \mathbb{E}\left(U(X'^{(w)\top} X'^{(w)})^{-1} X'^{(w)} (X'^{(w)} \theta^{(w)} + \varepsilon)\right) \\
&= \mathbb{E}(U) \theta^{(w)}
\end{aligned}$$

Then,

$$\begin{aligned}
\mathbb{E}(\hat{\tau}_k) &= \mathbb{E}\left(\frac{1}{n_k} \sum_{i=1}^{n_k} (X'_{k,i} \hat{\theta}_k^{(1)} - X'_{k,i} \hat{\theta}_k^{(0)})\right) && \text{Defined in (6)} \\
&= \frac{1}{n_k} \sum_{i=1}^{n_k} \left(\mathbb{E}(X'_{k,i} \hat{\theta}_k^{(1)}) - \mathbb{E}(X'_{k,i} \hat{\theta}_k^{(0)})\right) \\
&= \frac{1}{n_k} \sum_{i=1}^{n_k} (\mathbb{E}(X'_{k,i}) (\theta^{(1)} - \theta^{(0)})) \\
&= \mathbb{E}(X'_i) (\theta^{(1)} - \theta^{(0)}) = \tau && \text{Defined in (3)}
\end{aligned}$$

Similarly, conditioning on H to account for the variability in the n_k random (binomial) terms:

$$\begin{aligned}
\mathbb{E}(\hat{\tau}_{\text{pool}} | H) &= \mathbb{E}\left(\frac{1}{n} \sum_{i=1}^n (X'_i \hat{\theta}_{\text{pool}}^{(1)} - X'_i \hat{\theta}_{\text{pool}}^{(0)}) \mid H\right) && \text{Defined in (4)} \\
&= \frac{1}{n} \sum_{i=1}^n \left(\mathbb{E}(X'_i \hat{\theta}_{\text{pool}}^{(1)} \mid H) - \mathbb{E}(X'_i \hat{\theta}_{\text{pool}}^{(0)} \mid H)\right) \\
&= \frac{1}{n} \sum_{k=1}^K n_k \frac{1}{n_k} \sum_{i=1}^{n_k} (\mathbb{E}(X'_{k,i} \hat{\theta}_{\text{IS-IVW}}^{(1)} \mid H) - \mathbb{E}(X'_{k,i} \hat{\theta}_{\text{IS-IVW}}^{(0)} \mid H)) = \mathbb{E}(\hat{\tau}_{\text{IS-IVW}} \mid H) && \text{Definition 7 + Theorem 1} \\
&= \frac{1}{n} \sum_{k=1}^K n_k \frac{1}{n_k} \sum_{i=1}^{n_k} (\mathbb{E}(X'_{k,i} \hat{\theta}_{\text{GD}}^{(1)} \mid H) - \mathbb{E}(X'_{k,i} \hat{\theta}_{\text{GD}}^{(0)} \mid H)) = \mathbb{E}(\hat{\tau}_{\text{GD}} \mid H) && \text{Definition 8} \\
&= \mathbb{E}(X'_i \mid H) (\theta^{(1)} - \theta^{(0)}) = \tau
\end{aligned}$$

And,

$$\begin{aligned}
\mathbb{E}(\hat{\tau}_{1S-SW} | H) &= \frac{1}{n} \sum_{k=1}^K n_k \frac{1}{n_k} \sum_{i=1}^{n_k} (\mathbb{E}(X'_{k,i} \hat{\theta}_{1S-SW}^{(1)}) - \mathbb{E}(X'_{k,i} \hat{\theta}_{1S-SW}^{(0)})) && \text{Definition 6} \\
&= \frac{1}{n} \sum_{i=1}^n \left(\mathbb{E} \left(X'_{k,i} \sum_{l=1}^K \frac{n_l}{n} \hat{\theta}_l^{(1)} \right) - \mathbb{E} \left(X'_{k,i} \sum_{l=1}^K \frac{n_l}{n} \hat{\theta}_l^{(0)} \right) \right) \\
&= \frac{1}{n} \sum_{i=1}^n \left(\sum_{l=1}^K \frac{n_l}{n} \mathbb{E} \left(X'_{k,i} \hat{\theta}_l^{(1)} | H \right) - \sum_{l=1}^K \frac{n_l}{n} \mathbb{E} \left(X'_{k,i} \hat{\theta}_l^{(0)} | H \right) \right) \\
&= \frac{1}{n} \sum_{i=1}^n \left(\sum_{l=1}^K \frac{n_l}{n} \mathbb{E} \left(X'_{k,i} | H \right) (\theta^{(1)} - \theta^{(0)} | H) \right) \\
&= \mathbb{E} \left(X'_i | H \right) (\theta^{(1)} - \theta^{(0)}) = \tau
\end{aligned}$$

Noticing that none of the expectations above depend on the n_k and that $X \perp\!\!\!\perp H$, we can remove the conditioning over H , so $\mathbb{E}(\hat{\tau}_{\text{pool}}) = \mathbb{E}(\hat{\tau}_{\text{GD}}) = \mathbb{E}(\hat{\tau}_{1S-IVW}) = \mathbb{E}(\hat{\tau}_{1S-SW}) = \tau$.

B.2.6 Asymptotic variances of the ATE estimators

We recall that $\rho_k = \mathbb{P}(H_i = k) = \mathbb{E} \left(\frac{n_k}{n} \right)$ and that $p = \mathbb{P}(W_i = 1) = \sum_{k=1}^K \mathbb{P}(H_i = k) \mathbb{P}(W_i = 1 | H_i = k) = \sum_{k=1}^K \rho_k p_k$.

Proof of Section 3.1.2 : we prove that the SW ($\omega_k^{\text{SW}} = \frac{n_k}{\sum_{k=1}^K n_k}$) and IVW ($\omega_k^{\text{IVW}} = \frac{\mathbb{V}^\infty(\hat{\tau}_k^{\text{fed}})^{-1}}{\sum_{k=1}^K \mathbb{V}^\infty(\hat{\tau}_k^{\text{fed}})^{-1}}$) weights are asymptotically equivalent for federated local estimators ($\hat{\tau}_k^{1S-SW}$, $\hat{\tau}_k^{1S-IVW}$ both defined in Eq. 11, and $\hat{\tau}_k^{\text{GD}}$ defined in Eq. 14). We denote by $\hat{\tau}_k^{\text{fed}}$ any of these estimators.

Then:

$$\begin{aligned}
\omega_k^{\text{IVW}} &= \frac{\left(\frac{\sigma^2}{n_k} \left(\frac{n}{n^{(1)}} + \frac{n}{n^{(0)}} \right) + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2 \right)^{-1}}{\sum_{k=1}^K \left(\frac{\sigma^2}{n_k} \left(\frac{n}{n^{(1)}} + \frac{n}{n^{(0)}} \right) + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2 \right)^{-1}} \\
&= \frac{n_k \left(\sigma^2 \left(\frac{n}{n^{(1)}} + \frac{n}{n^{(0)}} \right) + \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2 \right)^{-1}}{\sum_{k=1}^K n_k \left(\sigma^2 \left(\frac{n}{n^{(1)}} + \frac{n}{n^{(0)}} \right) + \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2 \right)^{-1}} \\
&= \frac{n_k}{\sum_{k=1}^K n_k} \\
&= \omega_k^{\text{SW}}
\end{aligned}$$

Therefore, asymptotically, aggregating the local federated estimates of the ATE with SW or IVW is the same.

Asymptotic Variance of local ATE estimator.

First, recall from Appendix B.2.4 that if $\mathbb{E}(X^\top \varepsilon \mid H) = 0$ and Condition 1 holds, the OLS estimator $\hat{\theta}_k^{(w)}$ is consistent and asymptotically normal:

$$\begin{aligned} \hat{\theta}_k^{(w)} &\xrightarrow{P} \theta_k^{(w)} \\ \sqrt{n_k^{(w)}} (\hat{\theta}_k^{(w)} - \theta_k^{(w)}) &\xrightarrow{d} \mathcal{N}(0, \sigma^2 A^{-1}) \end{aligned} \quad (17)$$

which gives:

$$\sqrt{n_k^{(w)}} (\hat{c}_k^{(w)} - c_k^{(w)}) \xrightarrow{d} \mathcal{N}(0, \sigma^2) \quad (18)$$

$$\sqrt{n_k^{(w)}} (\hat{\beta}_k^{(w)} - \beta_k^{(w)}) \xrightarrow{d} \mathcal{N}(0, \sigma^2 \Sigma_k^{-1}) \quad (19)$$

In particular, we have that $\hat{c}_k^{(1)}, \hat{c}_k^{(0)}, \hat{\beta}_k^{(1)}, \hat{\beta}_k^{(0)}, \bar{X}_k = \frac{1}{n_k} \sum_{i=1}^{n_k} X_{k,i}$ are all asymptotically independent.

Then we have that:

$$\begin{aligned} \hat{\tau}_k - \tau &= \hat{c}_k^{(1)} - c^{(1)} - (\hat{c}_k^{(0)} - c^{(0)}) + \bar{X}_k (\hat{\beta}_k^{(1)} - \hat{\beta}_k^{(0)}) - \mathbb{E}(X_k) (\beta_k^{(1)} - \beta^{(0)}) \\ &= \hat{c}_k^{(1)} - c^{(1)} - (\hat{c}_k^{(0)} - c^{(0)}) + \bar{X}_k \left((\hat{\beta}_k^{(1)} - \beta^{(1)}) - (\hat{\beta}_k^{(0)} - \beta^{(0)}) \right) \\ &\quad - \mathbb{E}(X_k) (\beta_k^{(1)} - \beta^{(0)}) - \bar{X}_k (\beta_k^{(1)} - \beta^{(0)}) \\ &= \underbrace{\hat{c}_k^{(1)} - c^{(1)}}_{A_1} - \underbrace{(\hat{c}_k^{(0)} - c^{(0)})}_{A_0} + \underbrace{\bar{X}_k \left((\hat{\beta}_k^{(1)} - \beta^{(1)}) - (\hat{\beta}_k^{(0)} - \beta^{(0)}) \right)}_B \\ &\quad + \underbrace{(\bar{X}_k - \mathbb{E}(X_k)) (\beta^{(1)} - \beta^{(0)})}_C \end{aligned} \quad (20)$$

- For $w \in \{0, 1\}$, $A_w \xrightarrow{d} \mathcal{N}(0, \frac{\sigma^2}{n_k^{(w)}})$ from Equation (18)
- Let $M > 0$ be a real number. Then,

$$\mathbb{P}(|n_k B| > M) = \mathbb{P}\left(\left| \underbrace{\sqrt{n_k} (\bar{X}_k - \mathbb{E}(X_k))}_{\xrightarrow{d} \mathcal{N}(0, \Sigma_k)} \underbrace{(\hat{\beta}_k^{(1)} - \beta_k^{(1)})}_{\xrightarrow{d} \mathcal{N}(0, \frac{\sigma^2}{n_k^{(1)}})} - \underbrace{\sqrt{n_k} (\hat{\beta}_k^{(0)} - \beta_k^{(0)})}_{\xrightarrow{d} \mathcal{N}(0, \frac{\sigma^2}{n_k^{(0)}})} \right| > M \right) \xrightarrow{n_k \rightarrow \infty} 0$$

so that $\mathbb{P}\left(\left|\frac{B}{1/n_k} - 0\right| > M\right) < \varepsilon$, meaning that $B = \mathcal{O}_P(1/n_k)$ by definition.

- C : from the central limit theorem, we have that $\sqrt{n_k} (\bar{X}_k - \mathbb{E}(X_k)) \xrightarrow{d} \mathcal{N}(0, \Sigma_k)$ and $\beta_k^{(1)} - \beta_k^{(0)}$ is a constant vector. However, for any p-multivariate $\sqrt{n}Z \sim \mathcal{N}(0, \Sigma_k)$ random variable and D a constant vector of size $p \times 1$, we have $\sqrt{n}ZD \sim \mathcal{N}(0, D^\top \Sigma_k D)$. Therefore, $\sqrt{n_k}C \xrightarrow{d} \mathcal{N}(0, \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma_k}^2)$.

Finally, we have that:

$$\begin{aligned}
\mathbb{V}^\infty(\hat{\tau}_k) &= \frac{1}{n_k} \mathbb{V}^\infty(\sqrt{n_k}(\hat{\tau}_k - \tau)) \\
&= \frac{1}{n_k} (\mathbb{V}^\infty(\sqrt{n_k}A_1) + \mathbb{V}^\infty(\sqrt{n_k}A_0) + \mathbb{V}^\infty(\sqrt{n_k}B) + \mathbb{V}^\infty(\sqrt{n_k}C)) \\
&= \frac{1}{n_k} \left(\frac{n_k}{n_k^{(1)}} \sigma^2 + \frac{n_k}{n_k^{(0)}} \sigma^2 + 0 + \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma_k}^2 \right) \\
&= \sigma^2 \left(\frac{1}{n_k^{(1)}} + \frac{1}{n_k^{(0)}} \right) + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma_k}^2 \\
&= \frac{\sigma^2}{n_k} \left(\frac{1}{p_k} + \frac{1}{1-p_k} \right) + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma_k}^2
\end{aligned} \tag{21}$$

with $p_k = \mathbb{P}(W_i = 1 | H_i = k)$.

Using Central Limit Theorem:

$$\boxed{\sqrt{n_k}(\hat{\tau}_k - \tau) \xrightarrow{d} \mathcal{N} \left(0, \frac{\sigma^2}{p_k(1-p_k)} + \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma_k}^2 \right)} \tag{22}$$

Proof of Table 1:

From Equation (22) we have that in a Bernoulli trial (and denoting $[H = k] = \{H_i = k\}_{i=1}^n$):

$$\mathbf{Local\ ATE\ aVar} \quad \mathbb{V}^\infty(\hat{\tau}_k | H = k) = \frac{\sigma^2}{n_k p_k (1-p_k)} + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma_k}^2$$

Let's apply this result to the pooled dataset \mathcal{Z} and considering it a Bernoulli trial with treatment probability p , and denoting $H = \{H_i\}_{i=1}^n$:

$$\mathbb{V}^\infty(\hat{\tau}_{\text{pool}} | H) = \frac{\sigma^2}{np(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2$$

Finally, because H is not associated with the outcome, we have that $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{pool}} | H)$, which allows us to conclude:

$$\mathbf{Pooled\ ATE\ aVar} \quad \mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \frac{\sigma^2}{np(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2$$

which proves Eq. 5 (even in the non-centered covariates case).

To compute the asymptotic variance of the local federated outcome parameters ATE estimator, we first compute the asymptotic variance of the federated-outcome model parameters estimated

individual treatment effect $\hat{\tau}_{k,i}^{\text{fed}} = X'_{k,i} \hat{\theta}_{\text{fed}}^{(1)} - X'_{k,i} \hat{\theta}_{\text{fed}}^{(0)}$. To do this, remark that:

$$\begin{aligned} \mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) &= \mathbb{V}^\infty \left(\frac{1}{n} \sum_{k=1}^K \sum_{i=1}^{n_k} \hat{\tau}_{k,i}^{\text{pool}} \right) \\ &= \mathbb{V}^\infty \left(\frac{1}{n} \sum_{k=1}^K \sum_{i=1}^{n_k} \hat{\tau}_{k,i}^{\text{fed}} \right) && \text{since } \hat{\theta}_{\text{fed}}^{(w)} \rightarrow \hat{\theta}_{\text{pool}}^{(w)} \\ &= \frac{1}{n^2} \sum_{k=1}^K \sum_{i=1}^{n_k} \mathbb{V}^\infty(\hat{\tau}_{k,i}^{\text{fed}}) + \frac{1}{n^2} \sum_{k=1}^K \sum_{l=1}^K \sum_{i=1}^{n_k} \sum_{j \neq i}^{n_l} \text{Cov}^\infty(\hat{\tau}_{k,i}^{\text{fed}}, \hat{\tau}_{l,j}^{\text{fed}}) \end{aligned}$$

Let us show that the $\text{Cov}(\hat{\tau}_{k,i}^{\text{fed}}, \hat{\tau}_{l,j}^{\text{fed}})$ are asymptotically null for all $(k, i) \neq (l, j)$. Denote by $\text{Cov}^\infty(a, b)$ the asymptotic covariance of a and b random variables. We further write $\hat{Y}_i^{(w)} = X'_i \hat{\theta}_{\text{fed}}^{(w)}$, and remark that asymptotically, $\hat{Y}_i^{(w)} = X'_i \hat{\theta}_{\text{pool}}^{(w)}$, then:

$$\begin{aligned} \text{Cov}^\infty(\hat{\tau}_{k,i}^{\text{fed}}, \hat{\tau}_{l,j}^{\text{fed}}) &= \text{Cov}^\infty \left(X'_{k,i} \left(\hat{\theta}_{\text{fed}}^{(1)} - \hat{\theta}_{\text{fed}}^{(0)} \right), X'_{l,j} \left(\hat{\theta}_{\text{fed}}^{(1)} - \hat{\theta}_{\text{fed}}^{(0)} \right) \right) \\ &= \text{Cov}^\infty \left(\hat{Y}_{k,i}(1) - \hat{Y}_{k,i}(0), \hat{Y}_{l,j}(1) - \hat{Y}_{l,j}(0) \right) \\ &= \text{Cov}^\infty \left(\hat{Y}_{k,i}(1), \hat{Y}_{l,j}(1) \right) - \text{Cov}^\infty \left(\hat{Y}_{k,i}(1), \hat{Y}_{l,j}(0) \right) \\ &\quad - \text{Cov}^\infty \left(\hat{Y}_{k,i}(0), \hat{Y}_{l,j}(1) \right) + \text{Cov}^\infty \left(\hat{Y}_{k,i}(0), \hat{Y}_{l,j}(0) \right) \\ \text{Cov}^\infty \left(\hat{Y}_{k,i}^{(w_a)}, \hat{Y}_{l,j}^{(w_b)} \right) &= \mathbb{E} \left[\left(\hat{Y}_{k,i}^{(w_a)} - \mathbb{E}(\hat{Y}_{k,i}^{(w_a)}) \right) \left(\hat{Y}_{l,j}^{(w_b)} - \mathbb{E}(\hat{Y}_{l,j}^{(w_b)}) \right) \right] && \forall w_a, w_b \in \{0, 1\} \\ &= \mathbb{E} \left[\left(\hat{Y}_{k,i}^{(w_a)} - Y_{k,i}^{(w_a)} \right) \left(\hat{Y}_{l,j}^{(w_b)} - Y_{l,j}^{(w_b)} \right) \right] && \text{unbiased estimators} \\ &= \mathbb{E} \left[\varepsilon_{k,i}^{(w_a)} \varepsilon_{l,j}^{(w_b)} \right] && \text{residuals} \\ &= \mathbb{E} \left[\varepsilon_{k,i}^{(w_a)} \right] \mathbb{E} \left[\varepsilon_{l,j}^{(w_b)} \right] && \text{independent errors} \\ &= 0 && \text{centered noise} \end{aligned}$$

So that $\text{Cov}^\infty(\hat{\tau}_i^{\text{fed}}, \hat{\tau}_j^{\text{fed}} | H_i, H_j) = 0$, which yields that $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \frac{1}{n^2} \sum_{k=1}^K \sum_{i=1}^{n_k} \mathbb{V}^\infty(\hat{\tau}_{k,i}^{\text{fed}} | H_i = k)$. Finally, since the individuals follow the same distribution across centers within the graphical model in Figure 1, the $\hat{\tau}_{k,i}^{\text{fed}}$ are *i.i.d.* across centers, so that their asymptotic variances are equal:

$$\mathbb{V}^\infty(\hat{\tau}_{k,i}^{\text{fed}} | H_i = k) = n \mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \sigma^2 \left(\frac{1}{p(1-p)} \right) + \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2$$

Therefore,

$$\begin{aligned} \text{Federated Local ATE} \quad \mathbb{V}^\infty(\hat{\tau}_k^{\text{fed}} | H = k) &= \mathbb{V}^\infty \left(\frac{1}{n_k} \sum_{i=1}^{n_k} \hat{\tau}_{k,i}^{\text{fed}} | H = k \right) \\ &= \frac{1}{n_k^2} \sum_{i=1}^{n_k} \mathbb{V}^\infty(\hat{\tau}_{k,i}^{\text{fed}} | H = k) + \frac{1}{n_k^2} \sum_{i=1}^{n_k} \sum_{j \neq i}^{n_k} \underbrace{\text{Cov}^\infty(\hat{\tau}_{k,i}^{\text{fed}}, \hat{\tau}_{k,j}^{\text{fed}} | H = k)}_{=0} \\ &= \frac{n}{n_k} \mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) \\ &= \frac{\sigma^2}{n_k} \left(\frac{1}{p(1-p)} \right) + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \end{aligned}$$

For the Meta estimators we use the law of total variance:

$$\begin{aligned}
\text{Meta-SW ATE} \quad \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}}) &= \mathbb{V}^\infty\left(\sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k | H\right) \\
&= \mathbb{E} \left(\sum_{k=1}^K \left(\frac{n_k}{n}\right)^2 \mathbb{V}^\infty(\hat{\tau}_k | H) \right) + \mathbb{V}^\infty \left(\mathbb{E} \left(\sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k | H \right) \right) \\
&= \sum_{k=1}^K \mathbb{E} \left[\left(\frac{n_k}{n}\right)^2 \left(\frac{\sigma^2}{n_k p_k (1-p_k)} + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \right) \right] + 0 \\
&= \frac{\sigma^2}{n} \sum_{k=1}^K \mathbb{E} \left[\frac{n_k}{n} \frac{1}{p_k (1-p_k)} \right] + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \\
&= \frac{\sigma^2}{n} \sum_{k=1}^K \frac{\rho_k}{p_k (1-p_k)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2
\end{aligned}$$

$$\begin{aligned}
\text{Meta-IVW ATE} \quad \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) &= \mathbb{E} \left(\mathbb{V}^\infty \left(\frac{\sum_{k=1}^K (\mathbb{V}^\infty(\hat{\tau}_k | H = k)^{-1} \hat{\tau}_k)}{\sum_{k=1}^K \mathbb{V}^\infty(\hat{\tau}_k | H = k)^{-1}} \middle| H \right) \right) + 0 \\
&= \mathbb{E} \left(\frac{\sum_{k=1}^K \mathbb{V}^\infty(\hat{\tau}_k | H = k)^{-2} \mathbb{V}^\infty(\hat{\tau}_k | H = k)}{\left(\sum_{k=1}^K \mathbb{V}^\infty(\hat{\tau}_k | H = k)^{-1} \right)^2} \right) \quad \hat{\tau}_k \perp\!\!\!\perp \hat{\tau}_l \\
&= \mathbb{E} \left(\left(\sum_{k=1}^K \mathbb{V}^\infty(\hat{\tau}_k | H = k)^{-1} \right)^{-1} \right) \\
&= \mathbb{E} \left(\left(\sum_{k=1}^K \left(\frac{\sigma^2}{n_k p_k (1-p_k)} + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \right)^{-1} \right)^{-1} \right) \\
&= \mathbb{E} \left(\left(\sum_{k=1}^K \frac{n_k}{\frac{\sigma^2}{p_k (1-p_k)} + \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2} \right)^{-1} \right) \\
&= \frac{1}{n} \left(\sum_{k=1}^K \frac{\rho_k}{\frac{\sigma^2}{p_k (1-p_k)} + \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2} \right)^{-1} \quad (\text{LLN})
\end{aligned}$$

where (LLN) refers to the law of large numbers, stating that given $n_k \sim \text{Binomial}(n, \rho_k)$ and $\mathbb{E}[n_k] = n\rho_k$, for large n , we have $\mathbb{E} \left(\frac{1}{n_k} \right) \approx \frac{1}{\mathbb{E}[n_k]} = \frac{1}{n\rho_k}$.

We denote by $\hat{\tau}_{\text{SW}}^{\text{fed}} = \sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k^{\text{fed}}$:

$$\begin{aligned}
\text{SS Weighted Fed. ATE } \mathbb{V}^\infty(\hat{\tau}_{\text{SW}}^{\text{fed}}) &= \mathbb{E} \left(\mathbb{V}^\infty \left(\sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k^{\text{fed}} \mid H \right) \right) + 0 \\
&= \mathbb{E} \left(\mathbb{V}^\infty \left(\sum_{k=1}^K \frac{n_k}{n} \frac{1}{n_k} \sum_{i=1}^{n_k} \left(X'_{k,i} \hat{\theta}_{\text{fed}}^{(1)} - X'_{k,i} \hat{\theta}_{\text{fed}}^{(0)} \right) \mid H \right) \right) \\
&= \mathbb{E} \left(\mathbb{V}^\infty \left(\frac{1}{n} \sum_{i=1}^n \left(X'_i \hat{\theta}_{\text{fed}}^{(1)} - X'_i \hat{\theta}_{\text{fed}}^{(0)} \right) \mid H \right) \right) \\
&= \mathbb{E} \left(\mathbb{V}^\infty \left(\frac{1}{n} \sum_{i=1}^n \left(X'_i \hat{\theta}_{\text{pool}}^{(1)} - X'_i \hat{\theta}_{\text{pool}}^{(0)} \right) \mid H \right) \right) \\
&= \mathbb{E} (\mathbb{V}^\infty (\hat{\tau}_{\text{pool}} \mid H)) \\
&= \mathbb{V}^\infty (\hat{\tau}_{\text{pool}}) \\
&= \frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2
\end{aligned}$$

We denote by $\hat{\tau}_{\text{IVW}}^{\text{fed}} = \frac{\sum_{k=1}^K (\mathbb{V}^\infty(\hat{\tau}_k^{\text{fed}})^{-1} \hat{\tau}_k^{\text{fed}})}{\sum_{k=1}^K \mathbb{V}^\infty(\hat{\tau}_k^{\text{fed}})^{-1}}$:

$$\begin{aligned}
\text{IV Weighted Fed. ATE } \mathbb{V}^\infty(\hat{\tau}_{\text{IVW-agg}}) &= \mathbb{E} \left(\mathbb{V}^\infty \left(\frac{\sum_{k=1}^K (\mathbb{V}^\infty(\hat{\tau}_k^{\text{fed}})^{-1} \hat{\tau}_k^{\text{fed}})}{\sum_{k=1}^K \mathbb{V}^\infty(\hat{\tau}_k^{\text{fed}})^{-1}} \right) \right) \\
&= \mathbb{V}^\infty(\hat{\tau}_{\text{SW-agg}}) && \text{from Section 3.1.2} \\
&= \frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2
\end{aligned}$$

B.2.7 Comparison of asymptotic variances - General Case

Proof of Theorem 2: Under the graphical model in Figure 1:

- $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-SW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-IVW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{GD}})$ (Table 1)
- From Proposition 1 we have: $\mathbb{V}^\infty(\hat{\tau}_{\text{meta-IVW}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{SW}})$.
- Proving that $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}})$ is equivalent to proving the following inequality:

$$\begin{aligned}
\frac{1}{\sum_{k=1}^K \left(\frac{\sigma^2}{n_k p_k (1-p_k)} + \frac{1}{n_k} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \right)^{-1}} &\geq \frac{\sigma^2}{np(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \\
\iff \frac{1}{D} &\geq \frac{\sigma^2}{p(1-p)} + a
\end{aligned}$$

with $D = \sum_{k=1}^K \frac{n_k}{n} \frac{x_k}{\sigma^2 + x_k a}$, $x_k = p_k(1-p_k)$ and $a = \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2$.

$x \mapsto \frac{x}{\sigma^2 + xa}$ is concave. Therefore, by Jensen's inequality $D \leq \frac{\bar{x}}{\sigma^2 + \bar{x}a}$, with $\bar{x} = \sum_{k=1}^K \frac{n_k}{n} x_k$. We then have:

$$\frac{1}{D} \geq \frac{\sigma^2 + \bar{x}a}{\bar{x}} = \frac{\sigma^2}{\bar{x}} + a = \frac{\sigma^2}{\sum_{k=1}^K \frac{n_k}{n} p_k (1-p_k)} + a$$

Since $p \mapsto p(1-p)$ is also concave, by Jensen's inequality again we have:

$$\sum_{k=1}^K \frac{n_k}{n} p_k (1-p_k) \leq \left(\sum_{k=1}^K \frac{n_k}{n} p_k \right) \left(\sum_{k=1}^K \frac{n_k}{n} (1-p_k) \right) = p(1-p)$$

So that $\frac{1}{D} \geq \frac{\sigma^2}{p(1-p)} + a$, which ends the proof.

In conclusion,

$$\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}})$$

which concludes into $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{GD}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-SW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-IVW}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}})$.

Illustrating examples of this property:

1. $\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) - \mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) \geq 0$ increases as the treatment probabilities $\{p_k\}_k$ become more distinct. For example, with $K = 2$ centers with balanced datasets ($n_1 = n_2 = n/2$), having $p_1 = 0.99$ and $p_2 = 0.01$ yields $\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}})$ to be 8 times larger than $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}})$ (where we chose $\sigma^2 = 1$ and $\|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 = 10$).
2. Similarly, $\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}}) - \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) \geq 0$ grows with the $\{p_k(1-p_k)\}_k$ terms being more and more different from one another. For $K = 2$ and $n_1 = n_2 = n/2$, having $p_1 = 0.99$ and $p_2 = 0.5$ gives a 2.5 larger asymptotic variance of the Meta-SW than that of the Meta-IVW estimator.

B.2.8 Comparison of asymptotic variances - Special Case

Proof that the estimators of the ATE have equal asymptotic variances when one RCT is conducted over K centers: Under this setting, we modify the graphical model in Figure 1 and remove the edge between H and W :




Figure 7: Graphical Model of One RCT Conducted Over K Centers

Under this graphical model (Figure 7), using the variances in Table 1 and as $\forall k, p_k = p$, we have:

$$\text{First, } \mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-SW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-IVW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{GD}}).$$

Then,

$$\begin{aligned}
\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}}) &= \frac{\sigma^2}{n} \sum_{k=1}^K \frac{\mathbb{E}(n_k)}{n} \frac{1}{p_k(1-p_k)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \\
&= \frac{\sigma^2}{n} \frac{1}{p(1-p)} \sum_{k=1}^K \frac{\mathbb{E}(n_k)}{n} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \\
&= \frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \\
&= \mathbb{V}^\infty(\hat{\tau}_{\text{pool}})
\end{aligned}$$

And

$$\begin{aligned}
\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) &= \frac{1}{n} \left(\sum_{k=1}^K \frac{\rho_k}{\frac{\sigma^2}{p_k(1-p_k)} + \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2} \right)^{-1} \\
&= \left(\sum_{k=1}^K \left(\frac{\sigma^2}{\mathbb{E}(n_k) p_k(1-p_k)} + \frac{1}{\mathbb{E}(n_k)} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \right)^{-1} \right)^{-1} \\
&= \left(\sum_{k=1}^K \left(\frac{\sigma^2}{\mathbb{E}(n_k) p(1-p)} + \frac{1}{\mathbb{E}(n_k)} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \right)^{-1} \right)^{-1} \\
&= \left(\sum_{k=1}^K \left(\frac{1}{\mathbb{E}(n_k)} \left(\frac{\sigma^2}{p(1-p)} + \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \right) \right)^{-1} \right)^{-1} \\
&= \frac{\sigma^2}{n} \frac{1}{p(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2 \\
&= \mathbb{V}^\infty(\hat{\tau}_{\text{pool}})
\end{aligned}$$

which gives $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{GD}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-IVW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{IS-SW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-IVW}}) = \mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}})$.

B.3 Heterogeneous Settings

B.3.1 Distributional Shift in Covariates

In this part, we consider the graphical model in Figure 2, and model 2.

Unbiasedness and asymptotic variance of Meta-SW under covariate heterogeneity:

$$\begin{aligned}
\mathbb{E}(\hat{\tau}_{\text{Meta-SW}}) &= \mathbb{E}_H \left(\mathbb{E} \left(\sum_{k=1}^K \frac{n_k}{n} \hat{\tau}_k | H \right) \right) \\
&= \sum_k \mathbb{E}_H \left(\frac{n_k}{n} \right) \mathbb{E}(\hat{\tau}_k) \\
&= \sum_k \rho_k \tau_k && \text{under Condition 1 and model (2)} \\
&= \tau
\end{aligned}$$

$$\begin{aligned}
\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}}) &= \mathbb{E}(\mathbb{V}^\infty(\hat{\tau}_{\text{Meta-SW}})) + \mathbb{V}^\infty(\mathbb{E}(\hat{\tau}_{\text{Meta-SW}})) \\
&= \sum_k \mathbb{E}\left(\left(\frac{n_k}{n}\right)^2 \mathbb{V}^\infty(\hat{\tau}_k)\right) \\
&= \sum_{k=1}^K \frac{\mathbb{E}(n_k)}{n^2} \left(\frac{\sigma^2}{p_k(1-p_k)} + \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma_k}^2\right) \\
&= \frac{1}{n} \sum_{k=1}^K \rho_k \frac{\sigma^2}{p_k(1-p_k)} + \frac{1}{n} \sum_{k=1}^K \rho_k (\beta^{(1)} - \beta^{(0)})^\top \Sigma_k (\beta^{(1)} - \beta^{(0)}) \\
&= \frac{1}{n} \sum_{k=1}^K \rho_k \frac{\sigma^2}{p_k(1-p_k)} + \frac{1}{n} (\beta^{(1)} - \beta^{(0)})^\top \sum_{k=1}^K \rho_k \Sigma_k (\beta^{(1)} - \beta^{(0)}) \\
&= \frac{\sigma^2}{n} \sum_{k=1}^K \frac{\rho_k}{p_k(1-p_k)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_\Sigma^2
\end{aligned}$$

Unbiasedness and asymptotic variance of GD under covariate heterogeneity: The computation of the GD estimator's bias and asymptotic variance are direct using exactly the same proof as in Appendix B.2.6 since the assumptions over X_{pool} are still met in the distributional shift setting.

Proof of Proposition 2:

We have $\hat{c}_k^{(w)} = \bar{Y}_k^{(w)} - \bar{X}_k^{(w)} \hat{\beta}^{(w)}$ so that:

$$\begin{aligned}
\mathbb{V}(\hat{c}_k^{(w)} | X_k^{(w)}) &= \mathbb{V}(\bar{Y}_k^{(w)} - \bar{X}_k^{(w)} \hat{\beta}^{(w)} | X_k^{(w)}) \\
&= \mathbb{V}(\bar{Y}_k^{(w)}) + \mathbb{V}(\bar{X}_k^{(w)} \hat{\beta}^{(w)} | X_k^{(w)}) - 2\text{Cov}(\bar{Y}_k^{(w)} | X_k^{(w)}, \bar{X}_k^{(w)} \hat{\beta}^{(w)} | X_k^{(w)}) \\
&= \frac{\sigma^2}{n_k^{(w)}} + \bar{X}_k^{(w)\top} \mathbb{V}(\hat{\beta}^{(w)} | X_k^{(w)}) \bar{X}_k^{(w)} - 2\text{Cov}(\bar{Y}_k^{(w)} | X_k^{(w)}, \bar{X}_k^{(w)} \hat{\beta}^{(w)} | X_k^{(w)})
\end{aligned}$$

Given $X_k^{(w)}$ we have:

$$\begin{aligned}
\text{Cov}(\bar{Y}_k^{(w)}, \hat{\beta}_k) &= \text{Cov}\left(\frac{1}{n_k^{(w)}} \sum_{i=1}^{n_k^{(w)}} Y_{k,i}, \frac{\sum_{j=1}^{n_k^{(w)}} (X_{k,j}^{(w)} - \bar{X}_k^{(w)}) Y_{k,j}}{\sum_{i=1}^{n_k^{(w)}} (X_{k,i}^{(w)} - \bar{X}_k^{(w)})^2}\right) \\
&= \frac{1}{n_k^{(w)} \sum_{i=1}^{n_k^{(w)}} (X_{k,i}^{(w)} - \bar{X}_k^{(w)})^2} \text{Cov}\left(\sum_{i=1}^{n_k^{(w)}} Y_{k,i}, \sum_{j=1}^{n_k^{(w)}} (X_{k,j}^{(w)} - \bar{X}_k^{(w)}) Y_{k,j}\right) \\
&= \frac{1}{n_k^{(w)} \sum_{i=1}^{n_k^{(w)}} (X_{k,i}^{(w)} - \bar{X}_k^{(w)})^2} \sum_{i=1}^{n_k^{(w)}} \sum_{j=1}^{n_k^{(w)}} (X_{k,j}^{(w)} - \bar{X}_k^{(w)}) \text{Cov}(Y_{k,i}, Y_{k,j}) \\
&= \frac{1}{n_k^{(w)} \sum_{i=1}^{n_k^{(w)}} (X_{k,i}^{(w)} - \bar{X}_k^{(w)})^2} \sum_{j=1}^{n_k^{(w)}} (X_{k,j}^{(w)} - \bar{X}_k^{(w)}) \sigma^2 \\
&= 0
\end{aligned}$$

Therefore, $\mathbb{V}(\hat{c}_k^{(w)} \mid X_k^{(w)}) = \sigma^2 \left(\frac{1}{n_k^{(w)}} + (\bar{X}_k^{(w)})^\top (X_k^{(w)\top} X_k^{(w)})^{-1} \bar{X}_k^{(w)} \right)$, which yields in a random design over the $X_k^{(w)}$:

$$\mathbb{V}^\infty(\hat{c}_k^{(w)}) = \sigma^2 \left(\frac{1}{n_k^{(w)}} + \mathbb{E} \left((\bar{X}_k^{(w)})^\top (X_k^{(w)\top} X_k^{(w)})^{-1} \bar{X}_k^{(w)} \right) \right)$$

where

$$\begin{aligned} \mathbb{E} \left((\bar{X}_k^{(w)})^\top (X_k^{(w)\top} X_k^{(w)})^{-1} \bar{X}_k^{(w)} \right) &= \mu_k^\top \Sigma_k^{-1} \mu_k + \mathbb{E} \left(\frac{1}{n_k^{(w)}} (\bar{X}_k^{(w)} - \mu_k)^\top \Sigma_k^{-1} (\bar{X}_k^{(w)} - \mu_k) \right) \\ &= \mu_k^\top \Sigma_k^{-1} \mu_k + \frac{1}{n_k^{(w)}} \text{Tr} \left(\Sigma_k^{-1} \sum_{i=1}^{n_k^{(w)}} \mathbb{E} \left((X_{ki}^{(w)} - \mu_k)(X_{ki}^{(w)} - \mu_k)^\top \right) \right) \\ &= \mu_k^\top \Sigma_k^{-1} \mu_k + \frac{1}{n_k^{(w)}} \text{Tr} \left(\Sigma_k^{-1} \sum_{i=1}^{n_k^{(w)}} \Sigma_k \right) \\ &= \mu_k^\top \Sigma_k^{-1} \mu_k + \frac{1}{n_k^{(w)}} \text{Tr} (\Sigma_k^{-1}) \end{aligned}$$

so that $\mathbb{V}^\infty(\hat{c}_k^{(w)}) = \sigma^2 \left(\frac{1}{n_k^{(w)}} + \mu_k^\top \Sigma_k^{-1} \mu_k + \frac{1}{n_k^{(w)}} \text{Tr} (\Sigma_k^{-1}) \right)$.

Then, we get $\mathbb{V}^\infty(\hat{c}_{\text{pool}}^{(w)}) = \sigma^2 \left(\frac{1}{n^{(w)}} + \mu^\top \Sigma^{-1} \mu + \frac{1}{n^{(w)}} \text{Tr} (\Sigma^{-1}) \right)$ and

$$\begin{aligned} \mathbb{V}^\infty(\hat{c}_{\text{1S-SW}}^{(w)}) &= \sigma^2 \sum_{k=1}^K \left(\frac{n_k^{(w)}}{n^{(w)}} \right)^2 \left(\frac{1}{n_k^{(w)}} + \mu_k^\top \Sigma_k^{-1} \mu_k + \frac{1}{n_k^{(w)}} \text{Tr} (\Sigma_k^{-1}) \right) \\ &= \sigma^2 \left(\frac{1}{n^{(w)}} + \sum_{k=1}^K \left(\frac{n_k^{(w)}}{n^{(w)}} \right)^2 \left(\mu_k^\top \Sigma_k^{-1} \mu_k + \frac{1}{n_k^{(w)}} \text{Tr} (\Sigma_k^{-1}) \right) \right) \end{aligned}$$

Finally,

$$\begin{aligned} \frac{1}{\sigma^2} \left(\mathbb{V}^\infty(\hat{c}_{\text{pool}}^{(w)}) - \mathbb{V}^\infty(\hat{c}_{\text{1S-SW}}^{(w)}) \right) &= \mu^\top \Sigma^{-1} \mu - \sum_{k=1}^K \left(\frac{n_k^{(w)}}{n^{(w)}} \right)^2 \mu_k^\top \Sigma_k^{-1} \mu_k \\ &= \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \mu_k^\top \right) \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} (\Sigma_k + (\mu_k - \mu)(\mu_k - \mu)^\top) \right)^{-1} \\ &\quad \times \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \mu_k \right) - \sum_{k=1}^K \left(\frac{n_k^{(w)}}{n^{(w)}} \right)^2 \mu_k^\top \Sigma_k^{-1} \mu_k \end{aligned}$$

With Jensen's inequality applied to the convex function $f(x) = x^\top \Sigma_k^{-1} x$ for any positive definite matrix Σ_k with positive and summing to 1 weights, we have:

$$\left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \mu_k \right)^\top \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \Sigma_k \right)^{-1} \left(\sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \mu_k \right) \leq \sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \mu_k^\top \Sigma_k^{-1} \mu_k.$$

Therefore,

$$\mu^\top \Sigma^{-1} \mu - \sum_{k=1}^K \left(\frac{n_k^{(w)}}{n^{(w)}} \right)^2 \mu_k^\top \Sigma_k^{-1} \mu_k \leq \sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \left(1 - \frac{n_k^{(w)}}{n^{(w)}} \right) \mu_k^\top \Sigma_k^{-1} \mu_k \leq 0$$

After applying the law of total variance over H , we get $\mathbb{V}^\infty(\hat{c}_{\text{pool}}^{(w)}) \leq \mathbb{V}^\infty(\hat{c}_{\text{1S-SW}}^{(w)})$, implying $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}}) \leq \mathbb{V}^\infty(\hat{\tau}_{\text{1S-SW}})$.

In particular, notice that:

- If the centers have equal means ($\forall k, \mu_k = \mu$), then $\mathbb{V}^\infty(\hat{c}_{\text{pool}}^{(w)}) = \mathbb{V}^\infty(\hat{c}_{\text{1S-SW}}^{(w)})$, even if the $\{\Sigma_k\}_k$ are different.
- If a small number of centers have very distinct means from the rest of the centers, the difference in variances between the Pool and 1S-SW estimators will be large.

B.4 Center Effects

In this part, we consider the graphical model in Figure 3 and model (16).

Proof of meta estimators' unbiasedness and asymptotic variance (Section 4.2):

First, let us compute the intercepts for the treated and control groups in center k under model (16).

First, $a_k^{(w)} = c^{(w)} + h_k$ is the intercept in model (16), so that $a_k^{(w)} = \mathbb{E}(Y_k^{(w)} - X_k \beta^{(w)} - \varepsilon_k^{(w)}) = \mathbb{E}(Y_k^{(w)}) - \mathbb{E}(X_k) \beta^{(w)}$, which yields a **locally estimated** intercept $\hat{a}_k^{(w)} = \overline{Y}_k^{(w)} - \overline{X}_k^{(w)} \hat{\beta}_k^{(w)}$ in center k for treatment group w .

Then, we can compute the Meta estimator with weights ω_k :

$$\begin{aligned} \hat{\tau}_{\text{Meta}} &= \sum_{k=1}^K \omega_k \hat{\tau}_k \\ &= \sum_{k=1}^K \omega_k \left(\hat{a}_k^{(1)} - \hat{a}_k^{(0)} + \overline{X}_k (\hat{\beta}^{(1)} - \hat{\beta}^{(0)}) \right) \\ &= \sum_{k=1}^K \omega_k \left((a_k^{(1)} + \hat{\varepsilon}^{(1)}) - (a_k^{(0)} + \hat{\varepsilon}^{(0)}) + \overline{X}_k (\hat{\beta}^{(1)} - \hat{\beta}^{(0)}) \right) \\ &= \sum_{k=1}^K \omega_k \left((c^{(1)} + h_k + \hat{\varepsilon}^{(1)}) - (c^{(0)} + h_k + \hat{\varepsilon}^{(0)}) + \overline{X}_k (\hat{\beta}^{(1)} - \hat{\beta}^{(0)}) \right) \\ &= \sum_{k=1}^K \omega_k \left(\hat{c}^{(1)} - \hat{c}^{(0)} + \overline{X}_k (\hat{\beta}^{(1)} - \hat{\beta}^{(0)}) \right) \end{aligned}$$

where $\hat{c}^{(w)} = c^{(w)} + \hat{\varepsilon}^{(w)}$ and $\hat{\varepsilon}^{(w)}$ is the estimation bias of the OLS, with expectancy 0.

Therefore,

$$\begin{aligned}
\mathbb{E}(\hat{\tau}_{\text{Meta}}) &= \sum_{k=1}^K \omega_k \mathbb{E} \left(\hat{a}_k^{(1)} - \hat{a}_k^{(0)} + \overline{X}_k (\hat{\beta}^{(1)} - \hat{\beta}^{(0)}) \right) \\
&= \sum_{k=1}^K \omega_k \mathbb{E} \left(\hat{c}^{(1)} - \hat{c}^{(0)} + \overline{X}_k (\hat{\beta}^{(1)} - \hat{\beta}^{(0)}) \right) \\
&= \sum_{k=1}^K \omega_k \mathbb{E}(\hat{\tau}_k) \\
&= \sum_{k=1}^K \omega_k \tau_k
\end{aligned}$$

which leads the Meta estimators to be unbiased and have the same asymptotic variance as in Table 1.

B.4.1 Unadjusted Federated Estimators

We define the pooled outcome model parameters estimator as $\hat{\theta}_{\text{pool}}^{(w)} = \arg \min_{\theta} \sum_{i=1}^n (Y_i^{(w)} - X_i' \theta)^2$, and define $\hat{a}_{\text{pool}}^{(w)}$ the estimated intercept in the pooled data without the membership variable for group w , *i.e.* we have $\hat{\theta}^{(w)} = \{\hat{a}_{\text{pool}}^{(w)}, \hat{\beta}_{\text{pool}}^{(w)}\}$ and the estimated intercept in the pooled data is:

$$\begin{aligned}
\hat{a}_{\text{pool}}^{(w)} &= \overline{Y^{(w)}} - \overline{X'^{(w)}} \hat{\theta}_{\text{pool}}^{(w)} \\
&= \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} Y_i^{(w)} - \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} X_i'^{(w)} \hat{\theta}_{\text{pool}}^{(w)} \\
&= \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} (c^{(w)} + h_{k,i} + X_{k,i}' \theta^{(w)} + \varepsilon_i^{(w)}) - \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} X_i'^{(w)} \hat{\theta}_{\text{pool}}^{(w)} \quad \text{with } h_{k,i} = h_k \mathbf{1}_{\{H_i=k\}} \\
&= c^{(w)} + \overline{h_k^{(w)}} + \overline{\varepsilon_i^{(w)}} + \overline{X'^{(w)}} (\beta^{(w)} - \hat{\beta}_{\text{pool}}^{(w)})
\end{aligned}$$

with:

- $\overline{h_k^{(w)}} = \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} h_{k,i} = \frac{1}{n^{(w)}} \sum_{k=1}^{n^{(w)}} n_k^{(w)} h_k$ is the average effect of center k in group w .
- $\overline{\varepsilon_i^{(w)}} = \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} \varepsilon_i^{(w)}$ is the average error in group w .
- $\overline{X'^{(w)}} = \frac{1}{n^{(w)}} \sum_{i=1}^{n^{(w)}} X_i'^{(w)}$ is the average covariate in group w .

Therefore, the estimate of the intercept in the pooled dataset in presence of center effects has expectancy:

$$\begin{aligned}\mathbb{E}(\hat{a}_{\text{pool}}^{(w)}) &= c^{(w)} + \mathbb{E}\left(\overline{h_k^{(w)}}\right) \\ &= c^{(w)} + \mathbb{E}\left(\frac{1}{n^{(w)}} \sum_{k=1}^K n_k^{(w)} h_k\right) \\ &= c^{(w)} + \sum_{k=1}^K \mathbb{E}\left(\frac{n_k^{(w)}}{n^{(w)}}\right) h_k\end{aligned}$$

Then the unadjusted pooled estimator is:

$$\hat{\tau}_{\text{pool}} = \frac{1}{n} \sum_{i=1}^n \left(\hat{a}_{\text{pool}}^{(1)} - \hat{a}_{\text{pool}}^{(0)} + X_i (\hat{\beta}_{\text{pool}}^{(1)} - \hat{\beta}_{\text{pool}}^{(0)}) \right)$$

We now prove that this unadjusted estimator is biased when the p_k probabilities are not equal across centers. Then, we have:

$$\begin{aligned}\mathbb{E}(\hat{\tau}_{\text{pool}}) &= \mathbb{E}\left(\hat{a}_{\text{pool}}^{(1)} - \hat{a}_{\text{pool}}^{(0)} + \overline{X}(\hat{\beta}_{\text{pool}}^{(1)} - \hat{\beta}_{\text{pool}}^{(0)})\right) \\ &= c^{(1)} - c^{(0)} + \sum_{k=1}^K \mathbb{E}\left(\frac{n_k^{(1)}}{n^{(1)}}\right) h_k - \sum_{k=1}^K \mathbb{E}\left(\frac{n_k^{(0)}}{n^{(0)}}\right) h_k \\ &= \tau + \sum_{k=1}^K \mathbb{E}\left(\frac{n_k^{(1)}}{n^{(1)}} - \frac{n_k^{(0)}}{n^{(0)}}\right) h_k \\ &= \tau + \underbrace{\sum_{k=1}^K \left(p_k \mathbb{E}\left(\frac{n_k}{\sum_{k=1}^K n_k p_k}\right) - (1-p_k) \mathbb{E}\left(\frac{n_k}{\sum_{k=1}^K n_k (1-p_k)}\right) \right)}_{\text{bias}} h_k\end{aligned}$$

Then consider the two following cases:

- If equal treatment assignment ($p_k = p$ for all k):
Then, the bias is equal to $\sum_{k=1}^K p \mathbb{E}\left(\frac{n_k}{pn}\right) h_k - (1-p) \mathbb{E}\left(\frac{n_k}{(1-p)n}\right) h_k = 0$ so that $\mathbb{E}(\hat{\tau}_{\text{pool}}) = \tau$.
- If unequal treatment assignment ($p_k \neq p$ for all k):
Then, the bias is $\neq 0$ so that $\mathbb{E}(\hat{\tau}_{\text{pool}}) \neq \tau$.

Therefore, the Pool estimator is biased when the treatment probabilities are not equal among centers. This happens because the variable H acts as a confounder between the treatment and the outcome in cases of center effects. We need to account for it by adding a membership variable H in the dataset, which will allow the model to estimate the center effects.

B.4.2 With membership variable in the dataset

Proof of unbiasedness of the adjusted GD estimator: Denoting $\beta_H = (h_1, \dots, h_K)^\top$ the coefficients of the membership variables, model (16) can be written as:

$$\begin{aligned} Y_{k,i}(w) &= c^{(w)} + h_k + X_{k,i}\beta^{(w)} + \varepsilon_i(w) \\ &= c^{(w)} + H_{k,i}\beta_H + X_{k,i}\beta^{(w)} + \varepsilon_i(w) \end{aligned} \quad (23)$$

Under Equation (23), the (adjusted) Pooled estimator then estimates the coefficients of the variables H as a substitute for the center effects. This technique does not allow to estimate distinctly the intercepts $c^{(w)}$ and the effects of the centers $\{h_k\}_k$, as it relies on a relative rescaling of the intercepts of each center with respect to a chosen center of reference. In practice, choosing center 1 as the reference center by not including H_1 in the dataset offers the advantages of avoiding the underdeterminacy of the solutions of $\{(c^{(w)}, h_k)\}_k$ in Equation (23), and is easy to implement. In any case, this is without loss of generality.

Results from OLS estimation yield that the estimated intercept is the mean of the outcomes in the reference center (we arbitrarily choose center 1 to be the reference center), and the estimated coefficients of the membership variables are the differences between the means of the outcomes in the reference center and the other centers, which writes as:

$$\begin{aligned} \hat{c}_{\text{pool}}^{(w)} &= \frac{1}{n_1^{(w)}} \sum_{i=1}^{n_1^{(w)}} Y_{1,i}(w) \\ \hat{h}_k &= \frac{1}{n_k^{(w)}} \sum_{i=1}^{n_k^{(w)}} Y_{k,i}(w) - \hat{c}_{\text{pool}}^{(w)} \quad \forall k \in \llbracket 2, K \rrbracket \end{aligned}$$

where $\hat{h}_k^{(w)}$ is the estimated effect of center k on the outcomes in group w . In our model, we have that $\mathbb{E}(\hat{c}_{\text{pool}}^{(w)}) = Y_1(w)$ so that:

$$\begin{aligned} \mathbb{E}(\hat{h}_k^{(w)}) &= \mathbb{E}(Y_k(w)) - \mathbb{E}(Y_1(w)) \\ &= \mathbb{E}(c^{(w)} + X_{k,i}\beta^{(w)} + h_k + \varepsilon_i(w)) - \mathbb{E}(c^{(w)} + X_{1,i}\beta^{(w)} + h_1 + \varepsilon_i(w)) \\ &= h_k \end{aligned}$$

Finally, the adjusted pool estimator is:

$$\begin{aligned} \hat{\tau}_{\text{pool}_0} &= \frac{1}{n} \sum_{i=1}^n \left(\hat{c}_{\text{pool}}^{(1)} - \hat{c}_{\text{pool}}^{(0)} + X_i(\hat{\beta}_{\text{pool}}^{(1)} - \hat{\beta}_{\text{pool}}^{(0)}) + \sum_{k=1}^K H_{k,i}(\hat{\beta}_H^{(1)} - \hat{\beta}_H^{(0)}) \right) \\ &= \frac{1}{n} \sum_{k=1}^K \sum_{i=1}^{n_k} (\tilde{X}'_{k,i}(\hat{\theta}_{\text{pool}_0}^{(1)} - \hat{\theta}_{\text{pool}_0}^{(0)})) \end{aligned}$$

with the augmented design matrix $\tilde{X}'_{k,i} = (1, X_{k,i}, H_{2,i}, \dots, H_{K,i})$, $\hat{\beta}_H^{(w)} = (\hat{h}_1^{(w)}, \dots, \hat{h}_K^{(w)})^\top$ the estimated effects of the centers on the outcomes in group w and $\hat{\theta}_{\text{pool}_0}^{(w)} = (\hat{c}_{\text{pool}}^{(w)}, \hat{\beta}_{\text{pool}}^{(w)}, \hat{\beta}_H^{(w)})$. We then have $\mathbb{E}(\hat{\beta}_H^{(1)}) = \mathbb{E}(\hat{\beta}_H^{(0)}) = \beta_H$.

Finally, we get an unbiased estimator:

$$\begin{aligned}
\mathbb{E}(\hat{\tau}_{\text{pool}\circ}) &= \mathbb{E} \left(\hat{c}_{\text{pool}}^{(1)} - \hat{c}_{\text{pool}}^{(0)} + \bar{X}(\hat{\beta}_{\text{pool}}^{(1)} - \hat{\beta}_{\text{pool}}^{(0)}) + \sum_{k=1}^K H_k(\hat{\beta}_H^{(1)} - \hat{\beta}_H^{(0)}) \right) \\
&= \tau + \sum_{k=1}^K \mathbb{E} \left(H_k(\hat{\beta}_H^{(1)} - \hat{\beta}_H^{(0)}) \right) \\
&= \tau + \sum_{k=1}^K \mathbb{E} \left(H_k \mathbb{E}(\hat{\beta}_H^{(1)} - \hat{\beta}_H^{(0)} \mid H_k) \right) \\
&= \tau + \sum_{k=1}^K \mathbb{E} (H_k(\beta_H - \beta_H)) \\
&= \tau
\end{aligned}$$

The asymptotic variance of the adjusted Pool estimator is given by (using Appendix B.2.6): $\mathbb{V}^\infty(\hat{\tau}_{\text{pool}\circ}) = \frac{\sigma^2}{p(1-p)} + \frac{1}{n} \|\tilde{\beta}^{(1)} - \tilde{\beta}^{(0)}\|_{\tilde{\Sigma}}^2$ with $\tilde{\beta}^{(w)} = (\beta^{(w)}, \beta_H)$ and $\tilde{\Sigma} = \mathbb{V}(\tilde{X})$. Furthermore, remark that the block covariance-variance matrix is of the form $\tilde{\Sigma} = \begin{pmatrix} \Sigma & 0 \\ 0 & \text{diag}(p_1, \dots, p_K) \end{pmatrix}$ because the center effects are independent of the covariates X . Therefore, the asymptotic variance of the adjusted Pooled estimator is:

$$\mathbb{V}^\infty(\hat{\tau}_{\text{pool}\circ}) = \frac{\sigma^2}{p(1-p)} + \frac{1}{n} \|\beta^{(1)} - \beta^{(0)}\|_{\Sigma}^2$$

Note that the asymptotic variance of the adjusted Pool estimator in this center-effect model is insensitive to the center effects, which means that the heterogeneity among the centers does not affect this estimator.

Therefore, we can build the federated Gradient Descent with H variables (“ $\hat{\tau}_{\text{GD}\circ}$ ”) by allowing the centers to add $K - 1$ columns to their local datasets, with one of them containing only ones, unique to their dataset, and under convergence of Alg. D, we get $\hat{\tau}_{\text{GD}\circ} = \hat{\tau}_{\text{pool}\circ}$.

C SIMULATIONS

C.1 Simulation parameters

We generate data as follows:

1. Generate $X_{k,i} \sim \mathcal{N}(\mu_k, \Sigma_k)$ for individual $i \in \llbracket 1, n_k \rrbracket$, center $k \in \llbracket 1, K \rrbracket$ and d covariates. Add a constant covariate 1 to each $X_{k,i}$ to account for the intercept.
2. Generate $W_{k,i} \sim \mathcal{B}(p_k)$
3. Generate $\varepsilon_{k,i} \sim \mathcal{N}(0, \sigma^2)$ (homoscedasticity)
4. Build $Y_{k,i}(w) = c^{(w)} + h_k + X_{k,i} \beta_k^{(w)} + \varepsilon_{k,i}$, with $W_{k,i} = w$.

- Centers:
 - $K = 5$ centers
 - Sample sizes: asymptotic settings: $\{n_k = 20d\}_k$ so $n = 20Kd$; finite samples settings: $\{n_k = 4d\}_k$ so that $n = 4Kd$
- Model:
 - $\sigma^2 = 1$
 - $\{\theta_k^{(w)}\}_k = \{(c^{(w)} + h_k, \beta_1^{(w)}, \dots, \beta_d^{(w)})\}$ with $c^{(1)} = -1.85, c^{(0)} = -2,$
 $\beta^{(1)} = (-1.75, -1.5, -1.25, -1.0, -0.75, -0.5, -0.25, 0.0, 0.25, 0.5),$
 $\beta^{(0)} = (-1.8, -1.6, -1.4, -1.2, -1, -0.8, -0.6, -0.4, -0.2, 0),$
 $h_k = 0$ (no center effect by default for model (2))
- Covariates:
 - Dimension $d = 10$
 - $\left\{ (\mu_k = (\underbrace{1, \dots, 1}_{[d/2]}, \underbrace{-1, \dots, -1}_{[d/2]}), \Sigma_k = 0.5I_d + 0.5J_d) \right\}_k$, $J \in \mathbb{1}^{d,d}$ the matrix of ones

which yields $\tau = c^{(1)} - c^{(0)} + \mu^\top (\beta^{(1)} - \beta^{(0)}) = -1.1$
- Treatment assignment:
 - RCT: $W_{k,i} \sim \mathcal{B}(p_k)$ with $\{p_k = 0.5\}_k$;

For the Meta-IVW estimator, we estimate the asymptotic variance of the local ATE estimates,

i.e. $\hat{\tau}_{\text{Meta-IVW}} = \frac{\sum_{k=1}^K \widehat{\mathbb{V}}^\infty(\hat{\tau}_k)^{-1} \hat{\tau}_k}{\sum_{k=1}^K \widehat{\mathbb{V}}^\infty(\hat{\tau}_k)^{-1}}$ with:

- $\widehat{\mathbb{V}}^\infty(\hat{\tau}_k) = \frac{\hat{\sigma}_k^2}{n_k} \left(\frac{1}{\widehat{p}_k(1-\widehat{p}_k)} \right) + \frac{1}{n_k} \|\hat{\beta}_k^{(1)} - \hat{\beta}_k^{(0)}\|_{\hat{\Sigma}_k}^2$
- Sample variance of the residuals
 $\hat{\sigma}_k^2 = \frac{1}{n_k - d - 1} \sum_{i=1}^{n_k} (Y_{k,i} - X_{k,i}(\hat{\beta}_k^{(1)} \mathbb{1}_{[W_i=1]} + \hat{\beta}_k^{(0)} \mathbb{1}_{[W_i=0]}))^2$
- $\hat{p}_k = \frac{n_k^{(1)}}{n_k}$ and the sample covariance matrix $\hat{\Sigma}_k = \frac{1}{n_k - 1} \sum_{i=1}^{n_k} (X_{k,i} - \bar{X}_k)(X_{k,i} - \bar{X}_k)^\top$

C.2 Additional scenarios

C.2.1 Balanced datasets

In this scenario, we generate data in the *homogeneous* setting according to the graphical model in Figure 1 and outcome model 2 with $K = 5$ centers and $d = 10$ covariates dimension. We consider a balanced setting where all centers have equal sample size $n_k = 100d$ in the *Large* regime and $n_k = 5d$ in the *Small* sample size regime. The treatment assignment is the same for all centers in this first scenario, with $\forall k, W_{k,i} \sim \mathcal{B}(p)$ and $p = 0.5$ for all k .




Figure 8: Estimation of the ATE in the balanced and homogeneous setting for RCT studies.

Figure 8 represents the distribution over 2000 simulations of the estimated ATEs using the Pool estimator on the concatenated data as well as the Meta estimators with both weighting strategies (sample weighting and inverse variance weighting), the One-Shot estimators (IVW and SW) and Gradient Descent estimator (GD).

In large sample size regime, it highlights that in the balanced and homogeneous setting, (left panel *Large*), all estimators are unbiased and have the same variance, as expected in the special case of one RCT conducted over K centers (Section 3.2).

In the small sample size regime (right panel *Small*), the Metas and One-Shot SW have larger variances than the One-Shot IVW and GD estimators which are both equal to the pooled data one as expected given Theorem 1.

C.2.2 Imbalanced datasets

We consider a case of imbalance in the sample sizes of the centers, where one center has more observations than the others. For the *Large* setting, $n_1 = 400d$ and $n_2 = \dots = n_5 = 25d$, leading to the same total number of observations of $n = 5000$ as in Figure 8, whereas in the *Small* case $n_1 = 13d$ and $n_2 = \dots = n_5 = 3d$ resulting in $n = 250$, similarly to the balanced scenario.

Figure 9 shows that in the *Large* case, the partition of the pooled data has no impact on the estimation of the ATE for all estimators as long as Condition 1 holds. The boxplots are similar to the ones obtained in the balanced case (Figure 8), as expected again with the “One RCT” scenario.

However, in the *Small* case, the variances of the Meta estimators and 1S-SW are greater than in the balanced setting due to the local estimates $\hat{\tau}_2, \dots, \hat{\tau}_5$ being obtained after performing two local OLS regressions on each of their treatment arms on very small datasets.

C.3 Shift in covariates distributions

We now consider an *heterogeneous* setting where the data are generated according to the graphical model in Figure 2 under model (2), where individuals follow different distributions according to the




Figure 9: Estimation of the ATE in the imbalanced and homogeneous setting for RCT studies.

center they belong to. We consider the case where the means and covariance matrices $\{(\mu_k, \Sigma_k)\}_k$ are different from one center to another, with values:

Means	Covariances
$\mu_1 = \underbrace{(1, \dots, 1)}_{\lfloor d/2 \rfloor}, \underbrace{(-1, \dots, -1)}_{\lfloor d/2 \rfloor}^\top$	$\Sigma_1 = I_d + 0.5 - 0.5I_d$
$\mu_2 = \underbrace{(-1, \dots, -1)}_{\lfloor d/2 \rfloor}, \underbrace{(1, \dots, 1)}_{\lfloor d/2 \rfloor}^\top$	$\Sigma_2 = 20 \cdot \Sigma_1 + 0.5I_d - 0.5$
$\mu_3 = (0, \dots, 0)^\top$	$\Sigma_3 = 0.02 \cdot \Sigma_1 + 0.7I_d$
$\mu_4 = \underbrace{(0.5, \dots, 0.5)}_{\lfloor d/2 \rfloor}, \underbrace{(-1, \dots, -1)}_{\lfloor d/2 \rfloor}^\top$	$\Sigma_4 = 1 \cdot \Sigma_1 + 0.5I_d - 0.15$
$\mu_5 = \underbrace{(1.2, \dots, 1.2)}_{\lfloor d/2 \rfloor}, \underbrace{(0.8, \dots, 0.8)}_{\lfloor d/2 \rfloor}^\top$	$\Sigma_5 = 1.5 \cdot \Sigma_1 + 0.5I_d - 0.15$

Note that in this scenario, the Meta-IVW is not relevant as explained in Section 4. In this setting, the targeted ATE is $\tau = \sum_{k=1}^K \rho_k (c^{(1)} - c^{(0)} + \mu_k^\top (\beta^{(1)} - \beta^{(0)})) \approx 0.45$.

Figure 10 illustrates the fact that when centers have different covariate means, the variance of the One-Shot SW estimator is enlarged compared to homogeneous settings (Proposition 2), both in *Small* and *Large* sample sizes regimes. Notice that the Meta-SW estimator achieves better performance than in the homogeneous setting since its weights n_k/n are good estimates of ρ_k under Condition 1 of the true weights of the local ATEs, as $\tau = \sum_{k=1}^K \rho_k \tau_k$.

C.4 Center effects

We now generate data according to model (16) under the graphical model in Figure 3, with center effects equal to $(h_1, h_2, h_3, h_4, h_5) = (1, .2, -1, 30, 2)$, and with the other parameters set to their default values (Appendix C.1).




Figure 10: Estimation of the ATE in the heterogeneous setting for RCT studies.

C.4.1 No adjustment




Figure 11: Estimation of the ATE in a homogeneous balanced setting with center effects for RCTs.




Figure 12: Estimation of the ATE in a homogeneous setting with center effects and different treatment allocation schemes for RCTs.

Figures 11 and 12 present the distribution of all estimators without any modification, respectively in the “One RCT” scenario, and in the different treatment probabilities p_k setting. The estimators are not adjusted to take into account the presence of center effects, leading to the presence of a confounder between the treatment and the outcome variables (Assumption c), violating the identifiability assumption.

This case illustrates the advantage of Meta estimators over other estimators, which require in this setting less a priori knowledge on the underlying model in order to be used.

C.4.2 Adjusted

We now consider the same center effects scenario when adjusting the estimators with the procedures described in Section 4.2: the Pool and GD estimators are computed with access to the membership variables, and the One-Shot estimators do not federate local intercepts. The computation of the Meta estimators remains unchanged.




Figure 13: Estimation of the ATE in a homogeneous setting with center effects for RCTs.

Figure 13 shows that all the estimators are now unbiased, and their variance are equal to the pooled data one in the *Large* sample size regime.

However, in the *Small* regime, the adjusted One-Shot IVW and SW are less efficient than others, as they suffer from the unshared intercepts' variances. The adjusted One-Shot IVW does not benefit from Theorem 1. Their variances are highly dependent on the partition of the data into K splits, unlike the Pool and GD estimators whose variances solely depend on the total amount of data. The meta estimators naturally handle the center effects and their variances do not suffer much from their magnitude.

This scenario still highlights the advantage of the Meta estimators, which do not require a specific modelling of the center effects. However, in this setting they have a higher variance than the (adjusted) GD and (adjusted) pool estimators, as in the simple homogeneous balanced setting (C.2.1).

C.5 Full heterogeneity: shifts in covariates distributions, center effects and in treatment allocation

We now combine the different scenarios of Appendix C.3, Appendix C.4 with center effects (specific intercept per center), distribution of the covariates that are different from one center to the other (different means and covariance matrices) and different probabilities of being treated by center, corresponding to the graphical model in Figure 5. The simulation parameters used in Figures 4b are displayed in Appendix C.6:

Means	Covariances	Center Effects	Treatment probabilities
$\mu_1 = \underbrace{(1, \dots, 1)}_{[d/2]}, \underbrace{(-1, \dots, -1)}_{[d/2]}^\top$	$\Sigma_1 = I_d + 0.5 - \frac{1}{2}I_d$	$h_1 = 1$	$p_1 = 0.75$
$\mu_2 = \underbrace{(-1, \dots, -1)}_{[d/2]}, \underbrace{(1, \dots, 1)}_{[d/2]}^\top$	$\Sigma_2 = 20 \cdot \Sigma_1 + 0.5I_d - 0.5$	$h_2 = 0.2$	$p_2 = 0.75$
$\mu_3 = (0, \dots, 0)^\top$	$\Sigma_3 = 0.02 \cdot \Sigma_1 + 0.7I_d$	$h_3 = -1$	$p_3 = 0.75$
$\mu_4 = \underbrace{(0.5, \dots, 0.5)}_{[d/2]}, \underbrace{(-1, \dots, -1)}_{[d/2]}^\top$	$\Sigma_4 = 1 \cdot \Sigma_1 + 0.5I_d - 0.15$	$h_4 = 30$	$p_4 = 0.25$
$\mu_5 = \underbrace{(1.2, \dots, 1.2)}_{[d/2]}, \underbrace{(0.8, \dots, 0.8)}_{[d/2]}^\top$	$\Sigma_5 = 1.5 \cdot \Sigma_1 + 0.5I_d - 0.15$	$h_5 = 2$	$p_4 = 0.25$

C.6 G-Formula OLS covariate adjustment in non-linearity

In this setting, we simulate one dataset where $Y_i(1) = 0x_1^2 + \frac{-1}{2}x_2^2 + \frac{1}{2}x_3^2 + \frac{3}{2}x_4^2 + x_3 * x_4$ and $Y_i(0) = -0.35x_1^2 + 0x_2^2 + \frac{1}{2}x_3^2 + \frac{3}{2}x_4^2 + x_1 * x_2$.

The pooled-data OLS G-Formula has reduced variance compared to the simple Difference-in-Means, even if the outcome model is non-linear, illustrating (U.S. Food and Drug Administration, 2023; European Medicines Agency, 2024; Tsiatis et al., 2008; Benkeser et al., 2021; Lin, 2013; Wager, 2020; Lei and Ding, 2021; Van Lancker et al., 2024).




Table 3: ATE Estimation: DM vs. G-Formula with OLS adjustment on covariates under non-linear outcome modeling

D Algorithms

We describe in Algorithm 1 the procedure to obtain the Gradient Descent estimator of outcome model parameters $\hat{\theta}_{\text{GD}}$:

For appropriate learning rates, choosing:

$$T = \tilde{\Omega} \left(\frac{LE}{m} \log(1/\varepsilon) + \frac{\sqrt{L}\zeta E}{m\sqrt{\varepsilon}} \right),$$

gives $\|\theta_T - \hat{\theta}_{\text{pool}}^{(w)}\|^2 \leq \varepsilon$, where $\theta_T^{(w)}$ is the output of FedAvg after T communication rounds, L, m are respectively the maximum and minimum eigenvalues of $X^{(w)\top} X^{(w)}$ ($m > 0$ under Condition 2), and $\zeta^2 = \frac{1}{K} \sum_{k=1}^K \left\| \frac{1}{n_k} \sum_{i=1}^{n_k} X_{k,i} (X_{k,i}^{(w)\top} \hat{\theta}_{\text{pool}}^{(w)} - Y_{k,i}^{(w)}) \right\|^2$, and the $\tilde{\Omega}$ notation hides constants and polylogarithmic factors (Koloskova et al., 2020).

Algorithm 1 Federated Averaging (FedAvg) algorithm to learn $\hat{\theta}_{\text{GD}}^{(w)}$

```

1: Input:  $K$  centers,  $E$  local steps,  $B$  batch size,  $\eta$  learning rate,  $T$  rounds of communication
2: Server executes:
3:   Initialize  $\theta_0^{(w)}$ 
4: for each round  $t = 0, 1, \dots, T$  do
5:   for each center  $k \in \llbracket 1, K \rrbracket$  in parallel do
6:      $\theta_{t+1}^{k(w)} \leftarrow \text{LocalUpdate}(k, \theta_t^{(w)})$ 
7:   end for
8:    $\theta_{t+1}^{(w)} \leftarrow \sum_{k=1}^K \frac{n_k^{(w)}}{n^{(w)}} \theta_{t+1}^{k(w)}$  // Federated Averaging
9: end for
10: LocalUpdate( $k, \theta_k^{(w)}$ ):
11: for each local step  $e = 0, 1, \dots, E - 1$  do
12:    $\mathcal{B}_k \leftarrow$  a random batch of  $B$  samples from  $\mathcal{Z}_k$ 
13:    $\nabla \ell(\theta_k^{(w)}, \mathcal{B}_k) \leftarrow -\frac{2}{B} X_{\mathcal{B}_k}^\top (Y_{\mathcal{B}_k} - X_{\mathcal{B}_k} \theta_k^{(w)})$  // Compute gradient on the mini-batch
14:    $\theta_k^{(w)} \leftarrow \theta_k^{(w)} - \eta \nabla \ell(\theta_k^{(w)}, \mathcal{B}_k)$  // Gradient descent step
15: end for
16: return  $\hat{\theta}_{\text{GD}}^{(w)} \leftarrow \theta_k^{(w)}$ 

```
