



HAL
open science

Risk assessments underestimate threat of pesticides to wild bees

René S Shahmohamadloo, Mathilde L Tissier, Laura Melissa Guzman

► **To cite this version:**

René S Shahmohamadloo, Mathilde L Tissier, Laura Melissa Guzman. Risk assessments underestimate threat of pesticides to wild bees. *Conservation Letters*, 2024, 17 (4), pp.e13022. 10.1111/conl.13022 . hal-04748365

HAL Id: hal-04748365

<https://hal.science/hal-04748365v1>

Submitted on 22 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LETTER

Risk assessments underestimate threat of pesticides to wild bees

René S. Shahmohamadloo¹  | Mathilde L. Tissier^{2,3}  | Laura Melissa Guzman⁴ 

¹School of Biological Sciences, Washington State University, Vancouver, Washington, USA

²Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

³Centre de recherche et d'innovation sur les végétaux, Université Laval, Quebec City, Québec, Canada

⁴Department of Biological Sciences, University of Southern California, Los Angeles, California, USA

Correspondence

Laura Melissa Guzman, Department of Biological Sciences, University of Southern California, 3616 Trousdale Parkway, Los Angeles, CA 90089-0371, USA. Email: guzmanur@usc.edu

René S. Shahmohamadloo, School of Biological Sciences, Washington State University, 14204 NE Salmon Creek Ave, Vancouver, WA 98686, USA. Email: rene.shahmohamadloo@wsu.edu

Ecological risk assessments underestimate pesticide threats to pollinators by overextending acute lethality data on surrogate species to natural populations.

Funding information

Liber Ero Foundation; USC Wrigley Faculty Innovation Award; Natural Sciences and Engineering Research Council of Canada

Abstract

Ecological risk assessments (ERAs) are crucial when developing national strategies to manage adverse effects from pesticide exposure to natural populations. Yet, estimating risk with surrogate species in controlled laboratory studies jeopardizes the ERA process because natural populations exhibit intraspecific variation within and across species. Here, we investigate the extent to which the ERA process underestimates the risk from pesticides on different species by conducting a meta-analysis of all records in the ECOTOX Knowledgebase for honey bees and wild bees exposed to neonicotinoids. We found the knowledgebase is largely populated by acute lethality data on the Western honey bee and exhibits within and across species variation in LD50 up to 6 orders of magnitude from neonicotinoid exposure. We challenge the reliability of surrogate species as predictors when extrapolating pesticide toxicity data to wild pollinators and recommend solutions to address the (a)biotic interactions occurring in nature that make such extrapolations unreliable in the ERA process.

KEYWORDS

Apis, neonicotinoids, pesticides, policy, risk assessment

1 | INTRODUCTION

The planetary scale of human impacts on nature has increased sharply since the 1970s, driven by the demands of a growing population that is challenged with mitigating and adapting to climate change (Díaz et al., 2019). Ecosys-

tem deterioration and biodiversity loss are exacerbated by anthropogenic pollutants dispersing both on the land and in the water (Bongaarts, 2019). One mechanism adopted by governments and regulatory agencies to mitigate biodiversity loss is the ecological risk assessment (ERA) process, which evaluates the likelihood and magnitude of risk that an environment might be impacted from exposure to one or more stressors (Levin et al., 1989). Reliably predicting how natural populations will respond to environmental

René S. Shahmohamadloo, Mathilde L. Tissier, and Laura Melissa Guzman contributed equally to this work.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. Conservation Letters published by Wiley Periodicals LLC.

stressors remains a defining feature of the ERA process (Levin et al., 1989).

Historically, the most influential metric of the ERA process has been the lethal dose at which a chemical will kill 50% of a test population (LD50) (Relyea & Hoverman, 2006). This metric is derived from controlled laboratory studies quantifying toxicological responses on individual organisms (Barata et al., 1998; Stark et al., 2015). For over half a century, LD50s have guided regulatory agencies on how best to protect biodiversity while maintaining ecosystem services and economic output (Shahmohammadloo et al., 2022; Stark et al., 2015; USEPA, 2014). Yet, environmental protection goals are designed to protect populations and communities, making it difficult to translate laboratory studies on individual organisms to interpretable field outcomes (Clements & Rohr, 2009; Hommen et al., 2010; Newman, 2009). This is especially true when assessing the effects of pesticides on target and nontarget species in agroecosystems, which must concomitantly suppress pests while having negligible and nontarget effects on beneficial species and humans (Stark et al., 2015).

Unsurprisingly, the ERA process for pesticides has come under scrutiny. Mounting criticisms (Chapman et al., 1998; Sánchez-Bayo, 2014; Stark et al., 2004, 2015; Witwicka et al., 2022) challenge the preferential treatment of acute lethality data derived from stringent laboratory experiments on select surrogate species (EFSA, 2013; Franklin & Raine, 2019). Under the directives of the United States Environmental Protection Agency (USEPA), European Food Safety Agency (EFSA), and similar regulatory agencies worldwide, toxicological responses of these surrogate species are used to extrapolate the ecological risks of pesticides using probabilistic approaches (NRC, 2013; PMRA, 2021; Siviter et al., 2023a; USEPA, 2004). However, this process remains highly debated because surrogate species may not be reliable predictors of what may happen to thousands of beneficial species exposed to pesticides in nature (Banks et al., 2010, 2011; Franklin & Raine, 2019; Raine & Rundlöf, 2024; Siviter et al., 2021; Witwicka et al., 2022). Additional considerations, including associated effects on heterogeneous and genetically diverse populations (Forfert et al., 2017), interactions between pesticides and climate change (Delcour et al., 2015), and the metabolic effects of pesticides on species life-history traits (Cook, 2019), to name but a few, add further complexities unaccounted for in the ERA process ostensibly designed to assess risks to natural populations. Paradoxically, the pursuit to assess comparative toxicity between species and understand the mechanisms of toxicity require maximal control of laboratory conditions. This approach ignores the complexity of interactions between pesticides and other (a)biotic stressors in nature, an issue that reduces confidence in extrapolating between lab and field studies and yields con-

siderable uncertainty among regulators in pesticide risk management (Monchanin et al., 2019).

Here, we investigate the extent to which LD50 assays on surrogate species represent the distribution of LD50s for related species. We profile the most widely used insecticides in the world (Sánchez-Bayo, 2014) and their documented toxicity on a pollinator driving policy reform: neonicotinoids and the honey bee (*Apis*). Applied to seeds, crop foliage, and soils (Sánchez-Bayo, 2014), neonicotinoids have been implicated in the decline of honey bee health (Gross, 2013; Tsvetkov et al., 2017), and more worryingly connected to the decline of wild bees (Rundlöf et al., 2015; Siviter et al., 2021; Woodcock et al., 2016). Demands across all sectors to protect bees—and more broadly pollinators as providers of vital ecosystem services—led to concerted efforts by governments around the world to utilize the Western honey bee (*Apis mellifera*) as a surrogate for both *Apis* and non-*Apis* bees (Thompson & Pamminger, 2019) via a multitiered approach embedded in the ERA process (Fisher et al., 2023; Siviter et al., 2023a; USEPA, 2004) (*SI Appendix*, SI 1). We conducted a meta-analysis of all acute and chronic lethality records in the ECOTOX Knowledgebase (hereafter, called ECOTOX) for honey bees exposed to neonicotinoids, the world's largest curated database of ecologically relevant toxicity tests to support environmental research and risk assessment (Olker et al., 2022). This knowledgebase is used by the USEPA and regulatory agencies worldwide in pollinator risk assessment, risk management, and research. We collated all LD50 data along with any accompanying environmental parameters measured in ECOTOX (e.g., origin of bee strain, min and max temperature, duration of studies, and route of neonicotinoid exposure, to name a few). We hypothesized that lethality data of the Western honey bee do not sufficiently represent the breadth of intraspecific variation experienced by other *Apis* and non-*Apis* species under more realistic conditions (Banks et al., 2010, 2011; Franklin & Raine, 2019; Raine & Rundlöf, 2024; Siviter et al., 2021; Witwicka et al., 2022) and underestimate risk characterization of pesticides by regulatory agencies.

2 | METHODS

All records from ECOTOX (<https://cfpub.epa.gov/ecotox/>; database accessed October 16, 2022) were downloaded with filters applied for neonicotinoids and bees. We evaluated the variation in LD50 both within *Apis* and between *Apis* and non-*Apis* species, and subsetted assays that reported an LD50. The LD50s reported across all assays were standardized to ng/organism, which were used to generate species and genus sensitivity curves. This resulted in a total of 252 assays that evaluated LD50 on any bee species.

We returned to each original study and recorded the sample size for each assay. We obtained sample sizes for 226 assays. Furthermore, ECOTOX contains information on 95% confidence intervals. For some studies where the 95% confidence intervals were missing from ECOTOX, we were able to obtain it from the original studies. In total, we had 162 assays with confidence intervals. For the remaining assays, we were able to obtain the standard error of the mean response for one assay, standard deviation (which we then converted into standard error by standardizing the units and combining with the sample size) for 18 assays, and finally 29 assays had a chi-squared and p -value. In total, we obtained 210 assays with a mean response, sample size, and some measure of variation. For the studies with p -values, we constrained these p -values to be between 0.0001 and 0.9999 to allow for meta-analysis convergence.

We evaluated the intergenus variation in LD50 via a mixed-effects meta-analysis regression using the *meta* package in R (Balduzzi et al., 2019), with genus and the duration of the study (in days) as predictors. All LD50s were log-transformed to meet normality assumptions. Each pesticide and route of exposure was evaluated separately.

We then selected *A. mellifera* and imidacloprid assays—the most commonly tested across all assays—for a case study to determine whether records from ECOTOX accounted for genetic and environmental interactions with the toxicant. Relevant parameters investigated were: the origin of bee strain; route of neonicotinoid exposure to bees; and min and max temperature. For all analysis, the p -level significance cutoff was 0.05. See *SI Appendix (SI 2–5)* for further details.

3 | RESULTS

3.1 | The ECOTOX Knowledgebase

We identified 252 assays encompassing 49 studies from ECOTOX to generate the LD50 species and genus sensitivity curves (Dataset S1). The bee genera included were *Apis* (200/252 assays; 38/49 studies), *Bombus* (24/252 assays; 8/49 studies), *Megachile* (7/252 assays; 1/49 studies), *Melipona* (3/252 assays; 2/49 studies), *Nannotrigona* (2/252 assays; 1/49 studies), *Osmia* (7/252 assays; 3/49 studies), *Partamona* (1/252 assays; 1/49 studies), *Plebeia* (1/252 assays; 1/49 studies), *Scaptotrigona* (4/252 assays; 2/49 studies), and *Tetragonisca* (3/252 assays; 2/49 studies). The route of neonicotinoid exposure also differed among studies in ECOTOX, which we broadly categorized via “dietary” (diet = 8/252 assays; food = 162/252 assays) and under “topical” (topical = 28/252 assays;

dermal = 41/252 assays; environmental = 3/252 assays; spray = 9/252 assays).

For the case study on imidacloprid and *A. mellifera*, no studies investigated the role of genetic diversity on phenotypic outcomes. In addition, although reporting the origin of a colony is a requirement for studies in Europe on the risk assessment of plant protection products on bees (EFSA, 2013), only 77/129 studies reported a location where their Western honey bee colony originated; yet, upon closer inspection, none of these studies reported strain-specific information. Regarding biotic and abiotic variables that could influence bee response to neonicotinoid exposure (Cook, 2019), they were considered in a very small portion of the studies from ECOTOX. Less than 6% of the case studies considered diet as potentially affecting honey bee responses to imidacloprid, despite substantial evidence of an interaction between nutrition and exposure to neonicotinoids (Costa et al., 2022; Klaus et al., 2021; Leza et al., 2018; Stuligross & Williams, 2020; Tosi et al., 2017). Nutritional and metabolic properties of neonicotinoids were not investigated in the studies, while min and max temperatures were reported in only 13/129 studies. The majority of studies were conducted on adult workers, with only two studies looking at imidacloprid impacts on larvae. Finally, only one study considered the impact of imidacloprid on the reproductive output of the queen bee. Because so few studies evaluated the impact on reproduction or the role of genetic diversity, life stage, nutritional properties, metabolism, and temperature, we did not examine these data further.

3.2 | LD50 variation within and across bees

To test whether LD50 varies due to genetic diversity within and across species, we ran a meta-analytical regression, and compared genus-level estimates for all bee genera to those of *Apis*. Because the predictor is categorical (bee genera), the first group was treated as the intercept, which in this case was *Apis*. We did not conduct post-hoc comparisons because we were mainly interested in the comparison between all genera to *Apis* given it is a surrogate species in toxicity testing. The model also provides an estimate for differences between each genus and the intercept; a negative estimate, for instance, shows the LD50 of that genus is lower compared to *Apis*, while a positive estimate shows the LD50 of that genus is higher than *Apis*. A lower estimate, therefore, suggests that toxicity increases compared to *Apis*. We also evaluated the effect of the study duration using the number of days of the study. A negative estimate for duration indicates the LD50 across all




FIGURE 1 Lethal dose (LD50) measures from dietary exposure to neonicotinoids within and across bees. The surrogate, *Apis*, is compared to wild bees of seven genera (*Bombus*, *Megachile*, *Melipona*, *Osmia*, *Partamona*, *Scaptotrigona*, and *Tetragonisca*). The duration of studies is color-coated and the sample size is indicated by the diameter of the circle. Raw data are presented on a y-axis that is log-transformed (see Methods). In instances where several points are clustered near each other, spacing was applied to show the breadth of data.

groups decreases as the duration of the study increases (i.e., toxicity significantly increases).

For dietary exposure, the model shows that bee genera exhibited significant variation in LD50 response to neonicotinoids, both within and across genera (Figure 1 and Dataset S1). *Bombus* was significantly more sensitive to imidacloprid and thiamethoxam than *Apis* (estimate = -6.03 , se = 1.73, p -value = 0.00050; estimate = -2.02 , se = 0.80, p -value = 0.011). *Megachile* was significantly more sensitive to clothianidin, imidacloprid, and thiamethoxam than *Apis* (estimate = -7.37 , se = 1.42, p -value < 0.0001; estimate = -4.20 , se = 2.12, p -value = 0.048; estimate = -4.99 , se = 1.19, p -value < 0.0001). *Partamona* was significantly more sensitive to imidacloprid than *Apis* (estimate = -6.49 , se = 2.71, p -value = 0.017). All estimates from dietary exposure to neonicotinoids were also negative, demonstrating toxicity significantly increases as the duration of studies increases. As for variation within species from dietary exposure, we found *Apis* exhibits variation in LD50 up to 3 orders of magnitude in acetamiprid (21.5–72,600 ng/organism), 6 orders of magnitude in clothianidin (0.00001–26.9 ng/organism), 5 orders of magnitude in imidacloprid (0.03–4,170 ng/organism), and 7 orders of magnitude in thiamethoxam (0.00002–229 ng/organism) (Figure 1).

For topical exposure, the model similarly shows that bee genera exhibited significant variation in LD50 response to neonicotinoids, both within and across genera (Figure 2 and Dataset S1). Several genera were significantly more sensitive to imidacloprid than *Apis*, including *Melipona* (estimate = -3.09 , se = 0.85, p -value = 0.00028) and *Nannotrigona* (estimate = -3.43 , se = 1.15, p -value = 0.0029), while *Osmia* was significantly less sensitive to imidacloprid (estimate = 4.46, se = 1.25, p -value = 0.00035). Estimates from topical exposure to neonicotinoids were also negative, demonstrating toxicity significantly increases as the duration of studies increases. As for variation within species from topical exposure, we found *Apis* exhibits variation in LD50 up to 3 orders of magnitude in acetamiprid (220–395,500 ng/organism), 2 orders of magnitude in clothianidin (3.4–110 ng/organism), 3 orders of magnitude in imidacloprid (3.6–1,400 ng/organism), and 2 orders of magnitude in thiamethoxam (2.4–460 ng/organism) (Figure 2).

3.3 | Duration of studies

Aware that the ECOTOX Knowledgebase is largely populated by acute lethality data between 1 and 5 days—particularly topical studies (Figure 2)—we further tested




FIGURE 2 Lethal dose (LD50) measures from topical exposure to neonicotinoids within and across bees. The surrogate, *Apis*, is compared to wild bees of eight genera (*Bombus*, *Megachile*, *Melipona*, *Nannotrigona*, *Osmia*, *Plebeia*, *Scaptotrigona*, and *Tetragonisca*). The duration of studies is color-coated and the sample size is indicated by the diameter of the circle. Raw data are presented on a y-axis that is log-transformed (see Methods). In instances where several points are clustered near each other, spacing was applied to show the breadth of data.

whether LD50 is influenced by the duration of studies. Imidacloprid LD50s from topical exposure were significantly influenced by the duration of studies (estimate = 0.26, $se = 0.11$, p -value = 0.019). The estimate from imidacloprid topically exposed was also positive. This means that toxicity decreases with the duration of studies, and becomes less lethal more rapidly. Although for imidacloprid only—owing to the fact that most studies in ECOTOX conducted assays with this neonicotinoid—this result nonetheless illuminates the importance of chronic versus acute studies on pesticide toxicity in bees.

4 | DISCUSSION

4.1 | The impact of interspecific variation on pollinator risk assessment

Pollinator risk assessments are predicated on studies that provide key lethality data for regulators to make decisions invested in protecting biodiversity while maintaining ecosystem services and economic output (Bongaarts, 2019; Levin et al., 1989; Relyea & Hoverman, 2006; Shahmohamadloo et al., 2022; Stark et al., 2015; USEPA, 2004).

The multitiered approach adopted by regulatory agencies worldwide provides a framework by which regulators translate study results to inform their decisions on pesticide risk management (Olker et al., 2022). For neonicotinoids and bees, the majority of studies found in ECOTOX used *Apis* and most were conducted in laboratories on an acute timescale. Although *Apis* is a model organism for pollinator risk assessment, growing concerns advocate it is giving false confidence on pesticide safety thresholds for other bee species as life histories and route of exposure across genera may vary widely (Boyle et al., 2019; Franklin & Raine, 2019; Raine & Rundlöf, 2024; Schmolke et al., 2021; Witwicka et al., 2022).

We validate these concerns by showing *Apis* does not accurately estimate lethality risks from dietary (Figure 1) or topical (Figure 2) exposure to neonicotinoids on native bees from the genera *Bombus*, *Megachile*, *Melipona*, *Nannotrigona*, *Partamona*, and *Osmia*. *Apis* makes up the majority of assays (79.4%) in ECOTOX, while native bees lag behind (20.6%). Trait-based vulnerability analyses suggest that non-*Apis* bees may be more vulnerable to pesticides than the Western honey bee due to traits impacting exposure and population recovery potential (Schmolke et al., 2021). Recent findings showing differ-

ences in expression of the neonicotinoid target receptor nicotinic acetylcholine receptor among tissues and life stages within species (*A. mellifera*) and between species (*A. mellifera* and *Bombus terrestris*) bolster our concerns about generalizing toxicity estimates across bees and highlight the need for a greater understanding of the molecular targets under selection and variation in those targets (Witwicka et al., 2022). As one example, Europe is moving toward a more holistic approach to risk assessment that acknowledges non-*Apis* species as integral to pollination services and wild bee diversity (Knapp et al., 2023; More et al., 2021)—though only through guidance and recommendations, thus lacking mandatory steps to mandate such a holistic approach. For North America and other regions, diligent work is needed to develop toxicity assays for native and wild bees who, as we have found, differ in their life histories and sensitivities to neonicotinoids by several orders of magnitude compared to the Western honey bee.

Demands for risk assessments to include more chronic toxicity studies are ongoing and widely debated for pesticides (Sanchez-Bayo & Goka, 2014; Sánchez-Bayo & Tennekes, 2017; Stuligross & Williams, 2021; Tsvetkov et al., 2017). Arguments in favor of maintaining the status quo suggest that “Tier I” studies in honey bee acute risk assessment guidelines are protective of other bee species and tests should only be conducted in cases of concern (Thompson & Pamminger, 2019)—a conclusion drawn from studies not found in ECOTOX. As is the case from our meta-analysis, parsing the route of toxicity uncovered a large bias of acute assays for topical studies. The chronic impacts on bees exposed topically remain largely unknown, and more chronic studies are needed for dietary exposure. Our findings should compel risk assessors to question whether the LD50 responses measured in *Apis* on an acute timescale meet the broader objectives of protecting wild bees and biodiversity chronically exposed to pesticides.

4.2 | Sources of intraspecific variation within *Apis*

The wide variation in LD50 by several orders of magnitude within *Apis* (Figures 1 and 2) brings into question whether lethal doses are truly knowable. Even when stringent evaluation criteria are applied on *A. mellifera* in the adult life stage, variation in LD50 still exists up to 3 orders of magnitude from acute, dietary exposure to imidacloprid (0.1–100 ng/organism; Figure S1) and up to 2 orders of magnitude from acute, topical exposure to imidacloprid (10–1,000 ng/organism; Figure S2). Several factors likely contributed to this variation up to 6 orders of magnitude

within *Apis*. Below, we highlight three factors—namely, genetic variation, temperature, and nutrition—which we believe are synergistic and deserve special consideration in the ERA process, and more broadly biodiversity conservation efforts.

Failure to incorporate intraspecific genetic variation in toxicity studies hinders the estimation and understanding of genetic diversity, as well as the representativeness of the tested strain within a species or genus (Shahmohammadloo et al., 2023). This limitation introduces significant uncertainty when applying laboratory results to real-world scenarios, which undermines the accuracy and effectiveness of risk management strategies (Coutellec & Barata, 2011; Forbes, 1998; Thompson et al., 2023). Oversight of intraspecific variation becomes particularly relevant in instances of pesticide resistance evolution, as the presence of genetic variation can significantly reduce the effectiveness of agrochemicals due to natural selection (Hawkins et al., 2019). Thus, evolutionary processes such as adaptation to pesticides can plausibly affect the outcome of toxicity tests (Coutellec & Barata, 2011) and need better representation in risk assessments. If the ERA process cannot quantify interactions between toxicants and evolutionary forces on a population scale (e.g., genetic drift, inbreeding, selection), an accompanying decrease in genetic diversity may have negative effects at the population level, including: a reduced ability of populations to evolve genetically based changes in traits in response to environmental fluctuations (Thompson et al., 2023); increased mutations (Coutellec & Barata, 2011); risks of extinction (Exposito-Alonso et al., 2022); and pesticide resistance (Gould et al., 2018). Pesticides act as a strong agent of natural selection, often capable of driving rapid adaptation (Gould et al., 2018; Rudman et al., 2018). The standing stock of intraspecific genetic variation as it sits today can be important in predicting pesticide responses because there are differences in genotype. Yet, it is reasonably difficult to predict evolutionary responses in nature where the selective landscape has multiple drivers of natural selection causing phenotypic variation, an outward expression of the genotype (Endler, 1986). The lack of strain-specific, genetic information from the studies we reviewed in ECOTOX represents a major gap and further demonstrates the need for toxicity testing to assess the additive genetic variance in response to pesticides. Additive genetic variance is measurable at a population level and, when combined with information about natural selection, can be used to predict the pace and direction of adaptation (Rudman et al., 2022). Collecting these data represents a crucial responsibility for scientists and more broadly supporting the ERA process.

Although the ERA process in its current state does not explicitly incorporate the interactive effects of climate and pesticides, there is growing recognition that




FIGURE 3 Recommendations to scientists, regulatory agencies, and policymakers overseeing ecological risk assessments of neonicotinoids on bees.

temperature can affect the risks of pesticides to pollinators (Albacete et al., 2023). For instance, honey bees were more sensitive to imidacloprid and thiamethoxam at 24°C than 35°C (Saleem et al., 2020). This reduction in toxicity at 35°C was not due to degradation of the pesticide itself; nonlinearities may arise when heat stress is high enough to induce physiological responses that allow bees to better tolerate neonicotinoids (Saleem et al., 2020). Increases in temperature can also affect the absorption and metabolism of pesticides. For instance, studies on the effect of organophosphorus insecticides on midge (*Chironomus tentans*) lethality showed that acute toxicity increased with temperature, likely due to the metabolic activation of organophosphorus insecticides to more toxic forms (Lydy et al., 1999). However, the trend may not always be straightforward. Studies evaluating the combined effect of thiamethoxam and temperature on the homing performance of honey bees found that lower temperatures (16–20°C) exacerbated homing failure induced by the pesticide (Monchanin et al., 2019). Further, the interaction between temperature and neonicotinoids may depend on the type of behavior being observed. Neonicotinoids had a greater effect on movement and food consumption at a low temperature (21°C) yet a greater effect on flight distance at a high temperature (30°C) (Kenna et al., 2023). Considering only 13/129 studies reported temperature from our analysis of ECOTOX, monitoring temperature can be critical to interpreting data used for risk assessments.

In their natural environment, bees are primarily exposed to neonicotinoids through their diet (e.g., nectar, pollen, water consumption), although ground-nesting bees can also be exposed to neonicotinoids and the associated byproducts that may leach into soils (Main et al., 2020; Willis Chan & Raine, 2021; Siviter et al., 2023b). Dietary

exposure to neonicotinoids comprised 70% of studies in ECOTOX for honey bees. Although most studies assessed honey bee lethality or behavioral responses following dietary exposure to imidacloprid specifically, they overlooked the interactions of these molecules with the diet, its nutrient content, or the presence of other biomolecules. This could help explain the observed differences between topical and dietary exposure (Figures 1 and 2). This observation is of special concern given nutritional stress can substantially increase bee sensitivity to pesticides, including neonicotinoids (Leza et al., 2018; Stuligross & Williams, 2020; Tosi et al., 2017), while an appropriate diet can alleviate or even buffer the detrimental effects of insecticides on bees (Costa et al., 2022; Klaus et al., 2021). Interestingly, primary and secondary compounds of plants (e.g., alkaloids or flavonoids) can help bee detoxification processes (Balieira et al., 2018; Johnson et al., 2012), ultimately offsetting the negative impacts of insecticides on bee physiology, health, and performance (Costa et al., 2022; Riveros & Gronenberg, 2022). Thus, controlled studies looking at bee responses to a pesticide in a nutritionally rich setting may not capture the complexity of pesticide exposure in the natural environment.

5 | RECOMMENDATIONS

Our meta-analysis of ECOTOX uncovered substantial areas of development in interpreting pesticide risks to pollinators. This analysis relied heavily on data from the honey bee as the surrogate species, which showed variation in LD50 by several orders of magnitude within *Apis* (Figure 1). However, the data on *Apis* underestimate threats to wild bees (Figures 1 and 2), and the median

LD50 for acute exposure in *Apis*, used as a level of concern by the USEPA, remains above the LD50 values for non-*Apis* bees (Figures S3 and S4). We present three recommendations, both in writing and illustration (Figure 3), aiming to assist scientists, regulatory agencies, and policy-makers in protecting biodiversity and ecosystem services while sustaining economic outputs.

- **Develop assays for non-*Apis* bees.** The ERA process relies heavily on *Apis* to predict risks from pesticides to other bees. Yet, our meta-analysis shows that North American native bees differ in their life histories and sensitivities to neonicotinoids by several orders of magnitude compared to *Apis*. Developing toxicity assays for wild native bees and integrating these into the ERA process is a promising step forward that acknowledges non-*Apis* species as integral to pollination services and wild bee diversity.
- **Prioritize more chronic studies.** The ERA process for pollinators exposed to neonicotinoids relies heavily on acute toxicity data. Our meta-analysis brings into question whether LD50 responses measured on an acute timescale meet the broader objectives of protecting native bees and biodiversity chronically exposed to pesticides. Chronic studies should be prioritized.
- **Collect genetic and environmental data.** Studies need to better account for and report genetic and environmental data (e.g., climate and nutrition), otherwise, it becomes difficult to interpret LD50s. Acquiring these data can assist with developing better models to characterize ecological risks and developing better mitigation strategies.

AUTHOR CONTRIBUTIONS

R.S.S., M.L.T., and L.M.G. designed the research, conducted the literature search, analyzed the data, and wrote the paper.

ACKNOWLEDGMENTS

This work was supported by Liber Ero Postdoctoral Fellowships, an NSERC Postdoctoral Fellowship, and the USC Wrigley Faculty Innovation Award. We thank S. Otto, W. Palen, P. Sibley, S. Rudman, K. Cox, and two anonymous reviewers for their constructive comments and discussions.

CONFLICT OF INTEREST STATEMENT

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available and can be found in the Supporting Information. All analyses were completed in R version 4.2.2. and

can be retrieved here: https://github.com/lmguzman/bee_pesticide_epa.

ORCID

René S. Shahmohamadloo  <https://orcid.org/0000-0002-5373-9808>

Mathilde L. Tissier  <https://orcid.org/0000-0002-2107-8064>

Laura Melissa Guzman  <https://orcid.org/0000-0003-2198-9738>

REFERENCES

- Albacete, S., Sancho, G., Azpiazu, C., Rodrigo, A., Molowny-Horas, R., Sgolastra, F., & Bosch, J. (2023). Bees exposed to climate change are more sensitive to pesticides. *Global Change Biology*, 29, 6248–6260. <https://doi.org/10.1111/gcb.16928>
- Balduzzi, S., Rücker, G., & Schwarzer, G. (2019). How to perform a meta-analysis with R: A practical tutorial. *Evidence-Based Mental Health*, 22, 153–160. <https://doi.org/10.1136/ebmental-2019-300117>
- Balieira, K. V. B., Mazzo, M., Bizerra, P. F. V., de Jesus Santos Guimarães, A. R., Nicodemo, D., & Mingatto, F. E. (2018). Imidacloprid-induced oxidative stress in honey bees and the antioxidant action of caffeine. *Apidologie*, 49, 562–572. <https://doi.org/10.1007/s13592-018-0583-1>
- Banks, J. E., Ackleh, A. S., & Stark, J. D. (2010). The use of surrogate species in risk assessment: Using life history data to safeguard against false negatives. *Risk Analysis*, 30, 175–182. <https://doi.org/10.1111/j.1539-6924.2009.01349.x>
- Banks, J. E., Stark, J. D., Vargas, R. I., & Ackleh, A. S. (2011). Parasitoids and ecological risk assessment: Can toxicity data developed for one species be used to protect an entire guild? *Biological Control*, 59, 336–339. <https://doi.org/10.1016/j.biocontrol.2011.09.005>
- Barata, C., Baird, D. J., & Markich, S. J. (1998). Influence of genetic and environmental factors on the tolerance of *Daphnia magna* Straus to essential and non-essential metals. *Aquatic Toxicology*, 42, 115–137. [https://doi.org/10.1016/S0166-445X\(98\)00039-3](https://doi.org/10.1016/S0166-445X(98)00039-3)
- Bongaarts, J. (2019). IPBES, 2019. Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. *Population and Development Review*, 45, 680–681. <https://doi.org/10.1111/padr.12283>
- Boyle, N. K., Pitts-Singer, T. L., Abbott, J., Alix, A., Cox-Foster, D. L., Hinarejos, S., Lehmann, D. M., Morandin, L., O'Neill, B., Raine, N. E., Singh, R., Thompson, H. M., Williams, N. M., & Steeger, T. (2019). Workshop on Pesticide Exposure Assessment Paradigm for Non-*Apis* Bees: Foundation and Summaries. *Environmental Entomology*, 48, 4–11. <https://doi.org/10.1093/ee/nvy103>
- Chapman, P. M., Fairbrother, A., & Brown, D. (1998). A critical evaluation of safety (uncertainty) factors for ecological risk assessment. *Environmental Toxicology and Chemistry*, 17, 99–108. <https://doi.org/10.1002/etc.5620170112>
- Clements, W. H., & Rohr, J. R. (2009). Community responses to contaminants: Using basic ecological principles to predict ecotoxicological effects. *Environmental Toxicology and Chemistry*, 28, 1789–1800. <https://doi.org/10.1897/09-140.1>
- Cook, S. C. (2019). Compound and dose-dependent effects of two neonicotinoid pesticides on honey bee (*Apis mellifera*)

- metabolic physiology. *Insects*, 10(1), 18. <https://doi.org/10.3390/insects10010018>
- Costa, C. P., Leza, M., Duennes, M. A., Fisher, K., Vollaro, A., Hur, M., Kirkwood, J. S., & Woodard, S. H. (2022). Pollen diet mediates how pesticide exposure impacts brain gene expression in nest-founding bumble bee queens. *Science of the Total Environment*, 833, 155216. <https://doi.org/10.1016/j.scitotenv.2022.155216>
- Coutellec, M.-A., & Barata, C. (2011). An introduction to evolutionary processes in ecotoxicology. *Ecotoxicology (London, England)*, 20, 493–496. <https://doi.org/10.1007/s10646-011-0637-x>
- Delcour, I., Spanoghe, P., & Uyttendaele, M. (2015). Literature review: Impact of climate change on pesticide use. *Food Research International*, 68, 7–15. <https://doi.org/10.1016/j.foodres.2014.09.030>
- Diaz, S., Settele, J., Brondizio, E. S., Ngo, H. T., Agard, J., Arneth, A., Balvanera, P., Brauman, K. A., Butchart, S. H. M., Chan, K. M. A., Garibaldi, L. A., Ichii, K., Liu, J., Subramanian, S. M., Midgley, G. F., Miloslavich, P., Molnár, Z., Obura, D., Pfaff, A., ... Zayas, C. N. (2019). Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science*, 366(6471), eaax3100. <https://doi.org/10.1126/science.aax3100>
- Endler, J. A. (1986). *Natural selection in the wild*. Princeton University Press.
- European Food Safety Authority. (2013). Guidance on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus* spp. and solitary bees). *EFSA Journal*, 11, 3295.
- Exposito-Alonso, M., Booker, T. R., Czech, L., Gillespie, L., Hateley, S., Kyriazis, C. C., Lang, P. L. M., Leventhal, L., Nogues-Bravo, D., Pagowski, V., Ruffley, M., Spence, J. P., Toro Arana, S. E., Weiß, C. L., & Zess, E. (2022). Genetic diversity loss in the Anthropocene. *Science*, 377, 1431–1435. <https://doi.org/10.1126/science.abn5642>
- Fisher, A., 2nd, Tadei, R., Berenbaum, M., Nieh, J., Siviter, H., Crall, J., Glass, J. R., Muth, F., Liao, L.-H., Traynor, K., DesJardins, N., Nocelli, R., Simon-Delso, N., & Harrison, J. F. (2023). Breaking the cycle: Reforming pesticide regulation to protect pollinators. *Bioscience*, 73, 808–813. <https://doi.org/10.1093/biosci/biad088>
- Forbes, V. E. (1998). *Genetics and Ecotoxicology*. Taylor & Francis, Philadelphia, PA, USA.
- Forfert, N., Troxler, A., Retschnig, G., Gauthier, L., Straub, L., Moritz, R. F. A., Neumann, P., & Williams, G. R. (2017). Neonicotinoid pesticides can reduce honeybee colony genetic diversity. *PLoS ONE*, 12, e0186109. <https://doi.org/10.1371/journal.pone.0186109>
- Franklin, E. L., & Raine, N. E. (2019). Moving beyond honeybee-centric pesticide risk assessments to protect all pollinators. *Nature Ecology & Evolution*, 3, 1373–1375. <https://doi.org/10.1038/s41559-019-0987-y>
- Gould, F., Brown, Z. S., & Kuzma, J. (2018). Wicked evolution: Can we address the sociobiological dilemma of pesticide resistance? *Science*, 360, 728–732. <https://doi.org/10.1126/science.aar3780>
- Gross, M. (2013). EU ban puts spotlight on complex effects of neonicotinoids. *Current Biology*, 23, R462–R464. <https://doi.org/10.1016/j.cub.2013.05.030>
- Hawkins, N. J., Bass, C., Dixon, A., & Neve, P. (2019). The evolutionary origins of pesticide resistance. *Biological Reviews of the Cambridge Philosophical Society*, 94, 135–155. <https://doi.org/10.1111/brv.12440>
- Hommen, U., Baveco, J. M. H., Galic, N., & van den Brink, P. J. (2010). Potential application of ecological models in the European environmental risk assessment of chemicals. I. Review of protection goals in EU directives and regulations. *Integrated Environmental Assessment and Management*, 6, 325–337. <https://doi.org/10.1002/ieam.69>
- Johnson, R. M., Mao, W., Pollock, H. S., Niu, G., Schuler, M. A., & Berenbaum, M. R. (2012). Ecologically appropriate xenobiotics induce cytochrome P450s in *Apis mellifera*. *PLoS ONE*, 7, e31051. <https://doi.org/10.1371/journal.pone.0031051>
- Kenna, D., Graystock, P., & Gill, R. J. (2023). Toxic temperatures: Bee behaviours exhibit divergent pesticide toxicity relationships with warming. *Global Change Biology*, 29(11), 2981–2998. <https://doi.org/10.1111/gcb.16671>
- Klaus, F., Tschardtke, T., Bischoff, G., & Grass, I. (2021). Floral resource diversification promotes solitary bee reproduction and may offset insecticide effects—Evidence from a semi-field experiment. *Ecology Letters*, 24, 668–675. <https://doi.org/10.1111/ele.13683>
- Knapp, J. L., Nicholson, C. C., Jonsson, O., de Miranda, J. R., & Rundlöf, M. (2023). Ecological traits interact with landscape context to determine bees' pesticide risk. *Nature Ecology & Evolution*, 7(4), 547–556. <https://doi.org/10.1038/s41559-023-01990-5>
- Levin, S. A., Harwell, M. A., Kelly, J. R., & Kimball, K. D. (1989). *Ecotoxicology: Problems and approaches*. Springer.
- Leza, M., Watrous, K. M., Bratu, J., & Woodard, S. H. (2018). Effects of neonicotinoid insecticide exposure and monofloral diet on nest-founding bumblebee queens. *Proceedings of the Royal Society B Biological Sciences*, 285(1880), 20180761.
- Lydy, M. J., Belden, J. B., & Ternes, M. A. (1999). Effects of temperature on the toxicity of m-parathion, chlorpyrifos, and pentachlorobenzene to *Chironomus tentans*. *Archives of Environmental Contamination and Toxicology*, 37(4), 542–547. <https://doi.org/10.1007/s002449900550>
- Main, A. R., Webb, E. B., Goynes, K. W., & Mengel, D. (2020). Reduced species richness of native bees in field margins associated with neonicotinoid concentrations in non-target soils. *Agriculture Ecosystems and Environment*, 287, 106693. <https://doi.org/10.1016/j.agee.2019.106693>
- Monchanin, C., Henry, M., Decourtye, A., Dalmon, A., Fortini, D., Bœuf, E., Dubuisson, L., Aupinel, P., Chevallereau, C., Petit, J., & Fourier, J. (2019). Hazard of a neonicotinoid insecticide on the homing flight of the honeybee depends on climatic conditions and Varroa infestation. *Chemosphere*, 224, 360–368. <https://doi.org/10.1016/j.chemosphere.2019.02.129>
- More, S. J., Auteri, D., Rortais, A., & Pagani, S. (2021). EFSA is working to protect bees and shape the future of environmental risk assessment. *EFSA Journal*, 19, e190101. <https://doi.org/10.2903/j.efsa.2021.e190101>
- Newman, M. C. (2009). *Fundamentals of ecotoxicology*. CRC Press.
- NRC. (2013). *Assessing risks to endangered and threatened species from pesticides*. National Academies Press.
- Olker, J. H., Elonen, C. M., Pilli, A., Anderson, A., Kinziger, B., Erickson, S., Skopinski, M., Pomplun, A., LaLone, C. A., Russom, C. L., & Hoff, D. (2022). The ECOTOXicology knowledgebase: A curated database of ecologically relevant toxicity tests to support environmental research and risk assessment. *Environmental Toxicology and Chemistry*, 41, 1520–1539. <https://doi.org/10.1002/etc.5324>
- PMRA. (2021). A framework for risk assessment and risk management of pest control products. Health Canada.

- Raine, N. E., & Rundlöf, M. (2024). Pesticide exposure and effects on non-*Apis* bees. *Annual Review of Entomology*, *69*, 551–576.
- Relyea, R., & Hoverman, J. (2006). Assessing the ecology in ecotoxicology: A review and synthesis in freshwater systems. *Ecology Letters*, *9*, 1157–1171. <https://doi.org/10.1111/j.1461-0248.2006.00966.x>
- Riveros, A. J., & Gronenberg, W. (2022). The flavonoid rutin protects the bumble bee *Bombus impatiens* against cognitive impairment by imidacloprid and fipronil. *Journal of Experimental Biology*, *225*(17), jeb.244526. <https://doi.org/10.1242/jeb.244526>
- Rudman, S. M., Barbour, M. A., Csilléry, K., Gienapp, P., Guillaume, F., Hairston, N. G., Jr, Hendry, A. P., Lasky, J. R., Rafajlović, M., Räsänen, K., Schmidt, P. S., Seehausen, O., Therkildsen, N. O., Turcotte, M. M., & Levine, J. M. (2018). What genomic data can reveal about eco-evolutionary dynamics. *Nature Ecology & Evolution*, *2*, 9–15. <https://doi.org/10.1038/s41559-017-0385-2>
- Rudman, S. M., Greenblum, S. I., Rajpurohit, S., Betancourt, N. J., Hanna, J., Tilk, S., Yokoyama, T., Petrov, D. A., & Schmidt, P. (2022). Direct observation of adaptive tracking on ecological time scales in *Drosophila*. *Science*, *375*, eabj7484. <https://doi.org/10.1126/science.abj7484>
- Rundlöf, M., Andersson, G. K. S., Bommarco, R., Fries, I., Hederström, V., Herbertsson, L., Jonsson, O., Klatt, B. K., Pedersen, T. R., Yourstone, J., & Smith, H. G. (2015). Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature*, *521*, 77–80. <https://doi.org/10.1038/nature14420>
- Saleem, M. S., Huang, Z. Y., & Milbrath, M. O. (2020). Neonicotinoid pesticides are more toxic to honey bees at lower temperatures: Implications for overwintering bees. *Frontiers in Ecology and Evolution*, *8*, 556856. <https://doi.org/10.3389/fevo.2020.556856>
- Sánchez-Bayo, F. (2014). Environmental science. The trouble with neonicotinoids. *Science*, *346*, 806–807. <https://doi.org/10.1126/science.1259159>
- Sanchez-Bayo, F., & Goka, K. (2014). Pesticide residues and bees—A risk assessment. *PLoS ONE*, *9*, e94482. <https://doi.org/10.1371/journal.pone.0094482>
- Sánchez-Bayo, F., & Tennekes, H. A. (2017). Assessment of ecological risks of agrochemicals requires a new framework. *Environmental Risk Assessment and Remediation*, *1*(3), 20–28. <https://doi.org/10.4066/2529-8046.100025>
- Schmolke, A., Galic, N., Feken, M., Thompson, H., Sgolastra, F., Pitts-Singer, T., Elston, C., Pamminger, T., & Hinarejos, S. (2021). Assessment of the vulnerability to pesticide exposures across bee species. *Environmental Toxicology and Chemistry*, *40*, 2640–2651. <https://doi.org/10.1002/etc.5150>
- Shahmohamadloo, R. S., Febria, C. M., Fraser, E. D. G., & Sibley, P. K. (2022). The sustainable agriculture imperative: A perspective on the need for an agrosystem approach to meet the United Nations Sustainable Development Goals by 2030. *Integrated Environmental Assessment and Management*, *18*, 1199–1205. <https://doi.org/10.1002/ieam.4558>
- Shahmohamadloo, R. S., Rudman, S. M., Clare, C. I., Westrick, J. A., De Meester, L., & Fryxell, J. M. (2023). Intraspecific genetic variation is critical to robust toxicological predictions in *Daphnia*. *bioRxiv*. <https://doi.org/10.1101/2023.06.06.543817>
- Siviter, H., Linguadoca, A., Ippolito, A., & Muth, F. (2023a). Pesticide licensing in the EU and protecting pollinators. *Current Biology*, *33*, R44–R48. <https://doi.org/10.1016/j.cub.2022.12.002>
- Siviter, H., Pardee, G. L., Baert, N., McArt, S., Jha, S., & Muth, F. (2023b). Wild bees are exposed to low levels of pesticides in urban grasslands and community gardens. *Science of the Total Environment*, *858*, 159839. <https://doi.org/10.1016/j.scitotenv.2022.159839>
- Siviter, H., Richman, S. K., & Muth, F. (2021). Field-realistic neonicotinoid exposure has sub-lethal effects on non-*Apis* bees: A meta-analysis. *Ecology Letters*, *24*, 2586–2597. <https://doi.org/10.1111/ele.13873>
- Stark, J. D., Banks, J. E., & Vargas, R. (2004). How risky is risk assessment: The role that life history strategies play in susceptibility of species to stress. *Proceedings of the National Academy of Sciences of the United States of America*, *101*, 732–736. <https://doi.org/10.1073/pnas.0304903101>
- Stark, J. D., Vargas, R. I., & Banks, J. E. (2015). Incorporating variability in point estimates in risk assessment: Bridging the gap between LC50 and population endpoints. *Environmental Toxicology and Chemistry*, *34*, 1683–1688. <https://doi.org/10.1002/etc.2978>
- Stuligross, C., & Williams, N. M. (2020). Pesticide and resource stressors additively impair wild bee reproduction. *Proceedings of the Royal Society B Biological Sciences*, *287*(1935), 20201390.
- Stuligross, C., & Williams, N. M. (2021). Past insecticide exposure reduces bee reproduction and population growth rate. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(48), e2109909118.
- Thompson, H. M., & Pamminger, T. (2019). Are honeybees suitable surrogates for use in pesticide risk assessment for non-*Apis* bees? *Pest Management Science*, *75*, 2549–2557. <https://doi.org/10.1002/ps.5494>
- Thompson, L. M., Thurman, L. L., Cook, C. N., Beever, E. A., Sgrò, C. M., Battles, A., Botero, C. A., Gross, J. E., Hall, K. R., Hendry, A. P., Hoffmann, A. A., Hoving, C., LeDee, O. E., Mengelt, C., Nicotra, A. B., Niver, R. A., Pérez-Jvostov, F., Quiñones, R. M., Schuurman, G. W., ... Whiteley, A. (2023). Connecting research and practice to enhance the evolutionary potential of species under climate change. *Conservation Science and Practice*, *5*(2), e12855. <https://doi.org/10.1111/csp2.12855>
- Tosi, S., Nieh, J. C., Sgolastra, F., Cabbri, R., & Medrzycki, P. (2017). Neonicotinoid pesticides and nutritional stress synergistically reduce survival in honey bees. *Proceedings of the Royal Society B Biological Sciences*, *284*(1869), 20171711.
- Tsvetkov, N., Samson-Robert, O., Sood, K., Patel, H. S., Malena, D. A., Gajiwala, P. H., Maciukiewicz, P., Fournier, V., & Zayed, A. (2017). Chronic exposure to neonicotinoids reduces honey bee health near corn crops. *Science*, *356*, 1395–1397. <https://doi.org/10.1126/science.aam7470>
- USEPA. (2004). Overview of the Ecological Risk Assessment Process in the Office of Pesticide Programs United States Environmental Protection Agency: Endangered and Threatened Species Effects Determinations. DIANE Publishing.
- USEPA. (2014). Guidance for assessing pesticide risks to bees. Office of Chemical Safety and Pollution Prevention.

- Willis Chan, D. S., & Raine, N. E. (2021). Population decline in a ground-nesting solitary squash bee (*Eucera pruinosa*) following exposure to a neonicotinoid insecticide treated crop (*Cucurbita pepo*). *Scientific Reports*, *11*, 4241. <https://doi.org/10.1038/s41598-021-83341-7>
- Witwicka, A., López-Osorio, F., Patterson, V., & Wurm, Y. (2022). Expression of subunits of an insecticide target receptor varies across tissues, life stages, castes, and species of social bees. *Molecular Ecology*, *32*(5), 1034–1044.
- Woodcock, B. A., Isaac, N. J. B., Bullock, J. M., Roy, D. B., Garthwaite, D. G., Crowe, A., & Pywell, R. F. (2016). Impacts of neonicotinoid use on long-term population changes in wild bees in England. *Nature Communications*, *7*, 12459. <https://doi.org/10.1038/ncomms12459>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Shahmohamadloo, R. S., Tissier, M. L., & Guzman, L. M. (2024). Risk assessments underestimate threat of pesticides to wild bees. *Conservation Letters*, e13022. <https://doi.org/10.1111/conl.13022>