



X-ray natural optical activity beyond chirality

Elisabeth A. Hillard, Elen Duverger-Nédellec, Patrick Rosa, Ph. Sainctavit,
Marie-Anne Arrio, Andrei Rogalev, Fabrice Wilhelm, James Ablett

► To cite this version:

Elisabeth A. Hillard, Elen Duverger-Nédellec, Patrick Rosa, Ph. Sainctavit, Marie-Anne Arrio, et al..
X-ray natural optical activity beyond chirality. Congrès de l'Association Française de Cristallographie
- AFC 2024, AFC, Jul 2024, Montpellier, France. hal-04752203

HAL Id: hal-04752203

<https://hal.science/hal-04752203v1>

Submitted on 24 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

X-ray Natural Optical Activity beyond Chirality

Elizabeth HILLARD (ICMCB-CNRS, Pessac)
 Elen DUVERGER-NéDELLEC (ICMCB-CNRS, Pessac)
 Patrick ROSA (ICMCB-CNRS, Pessac)
 Philippe SAINCTAVIT (IMPMC-Sorbonne Universités, Paris)
 Marie-Anne ARRIO (IMPMC-Sorbonne Universités, Paris)
 Andrei ROGALEV (ESRF, Grenoble)
 Fabrice WILHELM (ESRF, Grenoble)
 Ablett JAMES (SOLEIL Synchrotron, St. Aubin)

Abstract

While optical activity in achiral space groups was predicted by Gibbs as long ago as 1882 [1], linear and circular birefringence effects hamper the measurement of circular dichroism in the UV-vis energy ranges [2]. We have used tender X-rays at the ESRF beamline ID12 to examine Natural Circular Dichroism (XNCD) [3] at the K-edges of copper and iron coordination salts respectively belonging to the achiral, but optically active, point groups D_{2d} and S_4 . The angular dependence of the XNCD signals was revealed with respect to the polar axis, theta, as well as the azimuthal axis, phi, by careful orientation of millimetric-sized crystals in the X-ray beam. In the case of D_{2d} , the XNCD followed the expected angular dependence in theta and phi [3]. However, for S_4 , an offset of about 27° in phi was observed and the more complicated angular dependance was ostensibly not respected. This result was subsequently confirmed using Resonant Inelastic X-ray Scattering-NCD at the SOLEIL synchrotron, in the first example of this RIXS-NCD technique, and can be rationalized using symmetry arguments.

[1] J. W. Gibbs, Am. J. Sci., 1882, 23, 460–476.

[2] a) R. Gautier, J. M. Klingsporn, R. P. Van Duyne and K. R. Poeppelmeier, Nature Mater., 2016, 15, 591–592; b) B. Kahr, A. T. Martin and K.-H. Ernst, Chirality, 2018, 30, 378–382; c) L.-L. Xu, H.-F. Zhang, M. Li, S. W. Ng, J.-H. Feng, J.-G. Mao and D. Li, J. Am. Chem. Soc., 2018, 140, 11569–11572; d) Y. Li, X. Ma, X. Xu, Y. Ye and B. Wang, Chem. Eur. J., 2023, 29, e202203534.

[3] J. Goulon, A. Rogalev, F. Wilhelm, C. Goulon-Ginet, P. Carra, I. Marri and Ch. Brouder, J. Exp. Theor. Phys., 2003, 97, 402–431.

Acknowledgments: We thank the French Agence National de la Recherche for funding via the XIMTEX (PCR ANR-21-CE29-0032) project.