



HAL
open science

Evolution towards a two-way cooperation in a single symbiotic organ: the developmental chronicle of a multi-partner insect endosymbiosis

François Renoz, Karen Gaget, Hubert Charles, Nicolas Parisot, Patrice Baa-Puyoulet, Mélanie Ribeiro Lopes, Sylvain Benhamou, Marie-Gabrielle Duport, Hugo Terrasson, Patrick Callaerts, et al.

► To cite this version:

François Renoz, Karen Gaget, Hubert Charles, Nicolas Parisot, Patrice Baa-Puyoulet, et al.. Evolution towards a two-way cooperation in a single symbiotic organ: the developmental chronicle of a multi-partner insect endosymbiosis. International Congress In Ecology and Evolution, Société Française d'Ecologie et d'Evolution (SFE2), Oct 2024, Lyon, France. hal-04753005

HAL Id: hal-04753005

<https://hal.science/hal-04753005v1>

Submitted on 25 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evolution towards a two-way cooperation in a single symbiotic organ: the developmental chronicle of a multi-partner insect endosymbiosis

François Renoz^{*1}, Karen Gaget², Hubert Charles³, Nicolas Parisot⁴, Patrice Baa-Puyoulet⁵, Mélanie Ribeiro-Lopes⁶, Sylvain Benhamou⁷, Gabrielle Duport², Hugo Terrasson⁶, Patrick Callaerts⁸, Thierry Hance¹, and Federica Calevro³

¹Earth and life Institute, Biodiversity Research Centre, Université de Louvain – Belgique

²Biologie Fonctionnelle, Insectes et Interactions (BF2I) – Institut national de la recherche agronomique (INRA) : UMR0203, Institut National des Sciences Appliquées [INSA] - Lyon – F-69621 Villeurbanne, France

³Biologie Fonctionnelle, Insectes et Interactions (BF2I) – Institut National de la Recherche Agronomique : UR0203, Institut National des Sciences Appliquées Lyon – INSA Bat. L. Pasteur 20 avenue Albert Einstein 69621 Villeurbanne cedex, France

⁴INRAE, INSA Lyon, BF2I, UMR203 – Univ Lyon, INRAE, INSA Lyon, BF2I, UMR 203, F-69621, Villeurbanne, France – 69621 Villeurbanne, France

⁵Univ Lyon, INSA-Lyon, INRA, BF2i, UMR0203, F-69621, Villeurbanne, France – Institut national de la recherche agronomique (INRA), Institut National des Sciences Appliquées (INSA) - Lyon – France

⁶Biologie Fonctionnelle, Insectes et Interactions – Univ Lyon, INSA Lyon, INRAE, BF2I, UMR 203, F-69621, Villeurbanne, France – France

⁷Biologie Fonctionnelle, Insectes et Interactions – Institut National des Sciences Appliquées (INSA) - Lyon – France

⁸Laboratory of Behavioral and Developmental Genetics, Department of Human Genetics, KU Leuven, Leuven, Belgium – Belgique

Résumé

Dependence on multiple nutritional endosymbionts forming a metabolic unit has evolved many times in insect taxa that feed on nutritionally unbalanced diets. It has been shown that the distribution of nutrient biosynthesis between nutritional symbionts coexisting within the same host goes hand in hand with their assignment to distinct specialized host cells (bacteriocytes) within the same symbiotic organ (bacteriome), according to an architectural layout that prevails in many insect species. This compartmentalization enables symbiont populations to be finely tuned to the host's nutritional needs during development. However, the mechanisms underlying the coordination of partners in these multi-symbiotic systems are unknown. To unveil these mechanisms, we used a combination of cell imaging, qPCR and dual-RNA seq approaches on the cereal aphid *Sipha maydis*, which has evolved a nutritional dependence on the symbiotic unit *Buchnera aphidicola-Serratia symbiotica*. The two symbionts are rapidly internalized in distinct types of bacteriocytes during embryogenesis,

*Intervenant

forming a horseshoe-shaped bacteriome. This consists of an assembly of syncytial bacteriocytes containing *Serratia*, embedded between uninucleate bacteriocytes containing *Buchnera*, which tend to dissociate from each other as the aphid ages. Surprisingly, while the infection dynamics of the two nutritional symbionts are strictly parallel during the nymphal stages, they tend to diverge during the reproductive phase, with a decrease in *Buchnera* populations and an increase in *Serratia* populations until the death of the host. This suggests that the metabolic complementation of the two symbionts tends to slacken with host development, and that their populations are not controlled in the same way. Our observations also show that the nutritional symbionts are not regulated in the same way as in the model aphid *Acyrtosiphon pisum*. They also highlight the existence of a new cell type systematically associated with bacteriocyte clusters, whose function is unknown. Dual RNA-seq analyses of bacteriocyte clusters collected at different developmental stages show that control of both symbionts involves the regulation of a set of genes related to metabolism, development and virulence. In conclusion, our study provides in-depth insight into the developmental mechanisms that govern cooperative intracellular life between anatomically integrated nutritional symbionts, and provides a comprehensive base map for examining the evolutionary developmental biology of insect multi-symbiotic systems.

Mots-Clés: Aphids, *Buchnera*, *Serratia*, Bacteriocyte, Multipartner endosymbiosis, Dual RNAseq