



**HAL**  
open science

## Opportunities to synchronise and date archaeological and climate records in Northwest Africa using volcanic ash (tephra) layers

Danielle Mclean, Emma Horn, Simone Aguiar, Nick Barton, Richard Brown, Stacy Carolin, Christopher Day, Holger Kuhlman, Steffen Kutterolf, Bryce Mitsunaga, et al.

### ► To cite this version:

Danielle Mclean, Emma Horn, Simone Aguiar, Nick Barton, Richard Brown, et al.. Opportunities to synchronise and date archaeological and climate records in Northwest Africa using volcanic ash (tephra) layers. *Lybian Studies*, 2024, 2024, pp.1-21. doi:10.1017/lis.2024.18 . hal-04753423

**HAL Id: hal-04753423**

**<https://hal.science/hal-04753423v1>**

Submitted on 28 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Opportunities to synchronise and date archaeological and climate records in Northwest Africa using volcanic ash (tephra) layers**

Danielle McLean<sup>1\*</sup>, Emma L. Horn<sup>1</sup>, Simone Aguiar<sup>2</sup>, Nick Barton<sup>1</sup>, Richard Brown<sup>3</sup>, Stacy Carolin<sup>1</sup>, Christopher Day<sup>4</sup>, Holger Kuhlman<sup>5</sup>, Steffen Kutterolf<sup>6</sup>, Bryce Mitsunaga<sup>7</sup>, Nicholas A. O'Mara<sup>8,9</sup>, José M. Pacheco<sup>2</sup>, Adriano Pimentel<sup>2</sup>, Ricardo S. Ramalho<sup>10</sup>, Julie Christin Schindlbeck-Belo<sup>5</sup>, Emmanuelle Stoetzel<sup>11</sup>, Amy Styring<sup>1</sup>, Kevin T. Uno<sup>6</sup>, Vaneshree Vidyarthi<sup>1</sup>, Yunbei Xu<sup>1</sup>, Victoria C. Smith<sup>1</sup>

<sup>1</sup>School of Archaeology, University of Oxford, Oxford, OX1 3TG, UK

<sup>2</sup>Instituto de Investigação em Vulcanologia e Avaliação de Riscos (IVAR), University of the Azores, Ponta Delgada, 9500-321, Portugal

<sup>3</sup>Department of Earth Sciences, Durham University, Durham, DH1 3LE, UK

<sup>4</sup>Department of Earth Sciences, University of Oxford, Oxford OX1 3AN, UK

<sup>5</sup>University of Bremen, MARUM, Bremen, 28359, Germany

<sup>6</sup>GEOMAR Helmholtz Centre for Ocean Research, Kiel, 24148, Germany

<sup>7</sup>Department of Human Evolutionary Biology and Earth and Planetary Sciences, Harvard University, Cambridge, MA, 02138, USA

<sup>8</sup>School of the Environment, Yale University, New Haven, CT 06511, USA

<sup>9</sup>Yale Institute for Biospheric Studies, Yale University, New Haven, CT 06511, USA

<sup>10</sup>School of Earth and Environmental Sciences, Cardiff University, Cardiff, CF10 3AT, UK10

<sup>11</sup>Histoire Naturelle de l'Homme Préhistorique - UMR 7194, CNRS/MNHN/UPVD, Musée de l'Homme, 75016, Paris, France

**\*Corresponding author:** Danielle McLean, email: [danielle.mclean@arch.ox.ac.uk](mailto:danielle.mclean@arch.ox.ac.uk)

**Competing interests:** The authors declare none

**Journal:** LIS (Libyan Studies)

**Cite this article:** McLean D et al (2024). Opportunities to synchronise and date archaeological and climate records in Northwest Africa using volcanic ash (tephra) layers. *Libyan Studies* 1–20.

### **Abstract**

Archaeological sites in Northwest Africa are rich in human fossils and artefacts providing proxies for behavioural and evolutionary studies. However, these records are difficult to underpin on a precise chronology, which can prevent robust assessments of the drivers of cultural/behavioural transitions. Past investigations have revealed that numerous volcanic ash (tephra) layers are interbedded within the Palaeolithic sequences and likely originate from large volcanic eruptions in the North Atlantic (e.g., the Azores, Canary Islands, Cape Verde). Critically, these tephra layers offer a unique opportunity to provide new relative and absolute dating constraints (via tephrochronology) to synchronise key archaeological and palaeoenvironmental records in this region. Here, we provide an overview of the known eruptive histories of the potential source volcanoes capable of widespread ashfall in the region during the last ~300,000 years, and discuss the diagnostic glass compositions essential for robust tephra correlations. To investigate the eruption source parameters required for ash dispersal towards NW Africa, we simulate plausible ashfall distributions using Ashfall3D. This work constitutes the first step in developing a more robust tephrostratigraphic framework for distal ash layers in NW Africa and highlights how tephrochronology may be used to robustly synchronise and date key climatic and cultural transitions during the Palaeolithic.

**Keywords:** *Tephrochronology, NW Africa, Palaeolithic, Glass Geochemistry, Volcanic Ashfall Simulation*

## 1. Introduction

Despite our desire to learn more about the forces driving the biological and behavioural evolution of our species during the Middle and Late Pleistocene (~300,000 to 10,000 years ago), many fundamental questions remain unresolved. These concern both the timing and synchronicity of major cultural transitions that occurred across NW Africa, and the role that climatic variability and other natural forces played in the origin, migrations, and behaviour of *Homo sapiens*. For example, it is possible that periods of severe climatic conditions (e.g., aridity) may have led to the emergence of significant cultural developments (e.g., innovations in tool technology), as *H. sapiens* adapted to a life in the changing Late Pleistocene landscape (Potts, 2013; Chase et al., 2018; Kuhn, 2023). Unfortunately, it has not been possible to interrogate the archaeological record in this way, since fossil sequences are notoriously difficult to underpin by a robust chronology, and many critical developments took place during times that lie beyond the limits of radiocarbon ( $^{14}\text{C}$ ) dating (>50,000 years). Moreover, climatic variability is known to have occurred on extremely rapid timescales (even within a human lifespan) and can respond asynchronously in different regions (Lane et al., 2013; Shanahan et al., 2015; Abrook et al., 2020; Nakagawa et al., 2021; O'Mara et al., 2022). Exceptional chronological precision is therefore required to robustly compare archaeological (e.g., fossil assemblage) and highly detailed climate (e.g., marine isotope or sea surface temperature signal) records, and for robust causal-effect relationships to be established.

Archaeological sites in NW Africa are rich in Palaeolithic organic remains and lithic technology, and have recently emerged at the forefront of evolutionary studies of *H. sapiens* (Hublin et al., 2017; Scerri et al., 2018; Barton et al., 2021). Those in modern-day Morocco (e.g., Taforalt, Bizmoune, Dar es Soltan, Harhoura II; Jebel Irhoud; Figure 1) in particular have been providing a critical insight into early human occupation and behaviour, recording detailed continuous Middle Stone Age (MSA) or Middle Palaeolithic (MP) assemblages, as well as the subsequent and widespread transition into the Later Stone Age (LSA) (~250,000 to 40,000 years ago) (Figure 1). The region hosts some of the earliest evidence of behavioural innovations in modern humans, including the introduction of novel tool forms and hafting methods, as well as the onset of symbolism and artistic behaviour (Bouzouggar et al., 2007; Bouzouggar and Barton, 2012; Sehassseh et al., 2021). For example, several MSA sites dating back to as early as the last interglacial period, Marine Isotope Stage (MIS) 5e, (~130 ka), contain modified *Nassariidae gibbosulus* perforated shells that exhibit wear-patterns and red ochre colouration, consistent with their use as personal ornamentation (Vanhaeren et al., 2006; Dibble et al., 2012; Steele et al., 2019; Sehassseh et al., 2021). Despite the age uncertainties often associated with these findings, the modified shells provide some of the earliest known evidence of explicitly symbolic objects in the archaeological record, and a fundamental stage in the emergence of modern social behaviour in *H. sapiens* (Barton and d'Errico, 2012).




Figure 1: (a) Location of key MSA-LSA archaeological and climate records (e.g., MD03-2705; Skonieczny et al., 2019) in North West Africa and the key volcanic source regions of greatest relevance for this region. (b) Schematic showing the potential of identifying co-located volcanic ash (tephra) layers. These time-parallel markers permit possibilities to: (i) share chronological information and (ii) directly compare climatic and cultural changes.

The MSA in NW Africa is often sub-divided into the Maghrebian Mousterian and Aterian industries (Dibble et al., 2013), with the latter widely traced from the Atlantic coast to the fringes of the Nile Valley (see Bouzouggar and Barton, 2012). The origin, chronology and significance of the Aterian have been a source of long-standing debate, and its spread has been linked to the dispersal of behaviourally modern humans (McBrearty and Brooks, 2000). Aterian assemblages include lithics such as bi-pointed bifacial foliates and a wide range of tanged implements, as recorded at sites like Dar es-Soltan I, situated on the Atlantic coast of Morocco (Figure 1), dated to the later part of MIS 5b (80 – 90 ka; Barton et al., 2009). Although novelties including small bifacial tools are recorded in East and South Africa, it is not yet clear if these were used at different times and under different assorted conditions and selective pressures (Marean, 2015; Blome et al., 2012; Powell et al., 2009). The role of climatic factors in the distribution of the Aterian, and whether this might have been related to the opening of green corridors through the Sahara during more humid episodes of MIS 5 (132 – 74 ka) remains unresolved (Garcea and Giraudi, 2006; Osborne et al., 2008).

Chronological uncertainties are also pertinent for younger archaeological assemblages, in particular for the transformational MSA to LSA transition observed across NW Africa. This widespread cultural shift is most clearly marked by a change from MSA flake and blade technologies to a more standardised microlithic bladelet production (an industry also referred to as Iberomaurusian). This shift is well documented at the site of Grotte des Pigeons (Taforalt, eastern Morocco; Figure 1), where a clear break in MSA deposits precedes a rich and thick sequence of bladelet and composite tool technology. This has been constrained by several accelerator mass spectrometry (AMS) radiocarbon dates between 25 – 23 ka cal. years BP (Barton et al., 2013). Improved chronological frameworks for other such sites, however, are required to disentangle the intricacies of these changes and determine whether this significant shift also originated via independent behavioural pathways, and what role climate may have played.

Chronometric dating methods have developed considerably over recent years, allowing new opportunities to reinterpret archaeological data and hypotheses, particularly in the realm of evolutionary studies (Wood, 2015; Becerra-Valdivia and Higham, 2023, Grün and Stringer, 2023). Notably for radiocarbon dating, more reliable preparation methods, that allow the extraction of diagenetically unaffected organics have been established, permitting more robust chronologies for archaeological sites in Europe (Higham et al., 2014). Furthermore, more accurate calibration methods (e.g., Ramsey et al., 2010; Reimer et al., 2020) alongside their integration within detailed Bayesian age modelling techniques (e.g., OxCal Bayesian program; Ramsey, 1995) have allowed a more robust anchoring of such developments in time. Notwithstanding these latest advancements, robust age models for archaeological sites are still difficult to construct, with problems typically arising from the availability of directly dateable material through the sequence (or at least with the necessary quality for high-resolution dating), and the potential for discontinuous or disturbed sedimentation accumulation (see Hunt et al., 2015). Beyond the radiocarbon timeframe (>50,000 years) reconstructions are even more blurred by the greater age uncertainties that accompany other techniques suitable for archaeological sequences, such as multi- and single-grain optically stimulated luminescence (OSL), uranium-series, thermoluminescence (TL), and cosmogenic nuclide dating techniques, meaning it is difficult to isolate taphonomic issues. There can also be limited opportunities to generate detailed and continuous paleoclimatic reconstructions using archaeological sequences, dependent on suitable accumulation and preservation of proxy material, such as organic plant/microfossil remains (e.g., Scott and Neumann, 2018), or bone/enamel/shell (e.g., Stoetzel et al., 2011; 2019; Jeffrey, 2016; Barton et al., 2020; Terray et al., 2023).

To establish the long-term climatic framework of the archaeological and faunal assemblages, the record must have an adequate chronology allowing the sequence to be pinned against other high-resolution climate archives. These can typically include speleothems/stalagmites from cave sites (e.g., Ait Brahim et al., 2023; Day et al., 2023), or detailed sedimentary records, like offshore marine (e.g., MD03-2705; Skonieczny et al., 2019; O'Mara et al., 2022; Figure 1) or lacustrine sediments (e.g., Atlas Mountains; Rhoujjati et al., 2010). In particular, records from deep sea sedimentary cores can provide global to regional climate reconstructions that reflect both the marine and terrestrial realms. Further, these records can be tied to an orbitally-tuned chronology with millennial scale resolution using benthic oxygen isotope ( $\delta^{18}\text{O}$ ) data (Lisiecki and Raymo, 2005). Marine climate records include (global) atmospheric  $\text{CO}_2$  reconstructions from foraminiferal boron isotope or haptophyte algae alkenone  $\delta^{13}\text{C}$  values (e.g., Hönisch et al., 2023); regional marine climate variables include sea surface temperatures, which provide information about latitudinal temperature gradients, regional temperatures, and relative monsoon strength from proxies such as alkenones ( $U_k^{37}$ ), foraminiferal Mg/Ca, or glycerol dialkyl glycerol tetraether (GDGT) analyses (Brassell, et al, 1986; Elderfield and Ganssen, 2000; Schouten et al., 2002). Terrestrial vegetation is recorded in marine cores in the form of microbotanical remains (e.g., cuticle, pollen, or phytoliths) and carbon isotope ratios of *n*-alkyl lipids (Morley and Richards, 1993; Bonnefille, 2010; Uno et al., 2016). Continental hydroclimate can be discerned from hydrogen isotope ratios of *n*-alkyl lipids and dust fluxes off the Sahara (Sachse et al., 2012; Tierney et al., 2016; O'Mara et al., 2022). Finally, emerging methods for reconstructing past fires from molecular products of biomass burning add new dimensions to terrestrial paleoecological reconstructions (e.g., Karp et al., 2020; 2021). Critically, tephtras that occur in both a marine core and archaeological record provide a means to establish a direct temporal link between regional climate records and human behavioural and technological transitions, discussed above, with exceptional chronological precision.

Pilot investigations at several key archaeological sites in NW Africa have revealed that numerous microscopic volcanic ash layers (known as 'cryptotephra') are preserved in the sediments with long MSA and LSA cultural sequences (Lane et al., 2014; Barton et al., 2015; 2021; Figure 2). At Grotte de Pigeons (Taforalt; Figure 1), microscopic volcanic glass shards were found interbedded between the levels containing MSA and LSA technology (Barton et al., 2015; 2016). The dominantly alkalic chemical composition of the glass, which can be used to determine the volcanic source, suggests that the ash was erupted from ocean island volcanoes in the North Atlantic. Due to the strong prevailing westerly winds in this region, ash layers have the potential to be widely dispersed, and become deposited in a range of sedimentary environments, including the surrounding ocean basins and onshore in subaerial, peat, and lacustrine records. Indeed, ash erupted from the Azores has been identified ~5000 km from its volcanic source in lake sediments in Svalbard, Norway (van der Bilt and Lane, 2019; Figure 2), highlighting the opportunity to temporally link records over exceptionally large distances. However, prior to the utilisation of these volcanic ash layers as synchronous markers, it is critical to conduct a detailed assessment of the source regions and geochemical uniqueness of the layers, so the fingerprint of individual and well dated volcanic events is robustly identified. This assessment constitutes one of the cornerstones of tephrochronology, without which it is not possible to ensure that unequivocal correlations are established across archives.




Figure 2: (a) Location of the key volcanic regions located within the North Atlantic, including the islands of the (b) Azores (AZ), (c) Canaries (CA), and (d) Cape Verde (CV), of greatest relevance for ash dispersal across NW Africa. The distal sedimentary palaeoenvironmental records reported to contain volcanic ash erupted from the Azores are marked with a blue box (Chambers et al., 2004; Barton et al., 2015; Watson et al., 2017; van der Bilt and Lane, 2019; Wastegård et al., 2020; Kinder et al., 2020; Walsh et al., 2021).

This article explores the key volcanic sources that have the potential to disperse widespread ash to NW Africa and provide tephra layers suitable for linking key MSA-LSA archaeological and climate records. Here, we first outline the main prerequisites for utilising tephra layers as time-stratigraphic markers, particularly within an archaeological setting. Secondly, we explore the volcanic regions (the Azores, Canary Islands, and Cape Verde) known to have produced large ash-rich eruptions (sub-Plinian to Plinian in style) and the known chronology of several key and widespread units. We collate the geochemical datasets available for the key regions and eruptions to distinguish specific glass chemical ‘fingerprints’ (unique to different eruptions; Lowe, 2011), which are essential for correlating deposits in the distal zone with their source. Finally, we investigate the eruption source parameters (e.g., tephra volume, column height, eruption duration) and weather patterns that are required to generate ash fall dispersal towards NW Africa and mainland Europe, which support the locations of distal evidence of tephra.

## 2. Volcanic ash layers as time-stratigraphic markers

### 2.1. Prerequisites for utilising tephra layers

Volcanic ash (tephra) layers can provide ideal time-stratigraphic markers and a powerful way of overcoming problems in comparing disparate sedimentary records (e.g., archaeological and paleoenvironmental) on independent time scales (Davies et al., 2002; Turney et al., 2006; Lane et al.,

2014; McLean et al., 2016). In general, three basic prerequisites are required before they can be widely used as chronological markers (Davies et al., 2012; Lane et al., 2014; Lowe et al., 2015). First, it must be possible to robustly identify the primary ashfall event within the sequence (i.e., discriminate undisturbed volcanic deposits from reworked or secondary deposits). Second, for a tephra layer to be utilised, it must have a known and distinct geochemical ‘fingerprint’ that can be singled out and used for unequivocal correlations. Finally, the approximate (relative or absolute) eruption age of the event must be established, permitting the layer to be integrated within a detailed chronological framework. These key principles are explored further here.

The term ‘tephra’ encompasses all pyroclastic material (quenched melt and crystals) ejected during a volcanic eruption (Thórarinnsson, 1944), however in medial and distal regions (e.g., >100 km from source), tephra is typically comprised of ash-size particles (<2 mm in size), and to a lesser extent lapilli-size particles (2 – 64 mm), and is predominantly composed of volcanic glass. In order to produce a significant amount of ash, eruptions need to be highly explosive, typically of sub-Plinian to Plinian styles, i.e. they need to attain a Volcanic Explosivity Index (VEI; Newhall and Self, 1982) or an eruption magnitude (M; Pyle, 1989) greater than or equal to 4. This implies the extrusion of tephra volumes of >0.1 km<sup>3</sup> and eruptive columns >15 km in height. The usefulness of the ash as a time-stratigraphic marker is reliant on the tephra being quickly deposited following the eruption and remaining as a discrete, *in situ* horizon (i.e., recording an instantaneous event). Reworking processes (e.g., erosion and redeposition, bioturbation, site reoccupation, high-energy floods) can sometimes obscure the primary ashfall event, either by moving/skewing the stratigraphic positioning, or by the re-deposition of older units (e.g., Wastegård et al., 2006; McLean et al., 2018). It is therefore essential that all possible taphonomic processes are considered, and that secondary reworking events can be identified and avoided. This criterion is usually achieved on a site-specific basis and best accomplished by considering several lines of reworking evidence (e.g., visual features of the unit such as grading, microscopic components of minerals and glass, geochemical consistency, morphoscopy of the grains, etc; Gudmundsdóttir et al., 2011; Abbott et al., 2018; McLean et al., 2018).

Tephra markers can be preserved as non-visible (cryptotephra) layers, where the concentration of glass shards is diluted and does not form a clear macroscopic unit (e.g., Davies, 2015). Cryptotephra layers are almost exclusively composed of volcanic glass shards and are typically <125 µm thick in distal settings, and can contain very low concentrations (i.e., below 500 shards per gram of dried sediment). As such, they can provide evidence of lower magnitude and/or very distant events, allowing records over wider geographic footprints to be synchronised. In order to identify the primary stratigraphic position of cryptotephra layers, sequences are continuously scanned to identify a peak in glass shard concentrations. This can be achieved by non-destructive techniques, such as X-ray fluorescence (XRF) continuous scanning (e.g., Kylander et al., 2012; McCanta et al., 2015; McLean et al., 2022) or computed tomography (CT) (e.g., Griggs et al., 2015; van der Bilt et al., 2021) but findings can be inconsistent especially for low concentrations of glass shards or those with geochemical compositions similar to the host sediment (e.g., McLean et al., 2022). Arguably the most reliable but labour-intensive technique is achieved using density separation methods, which isolate the volcanic glass from the host sediment. Heavy liquid flotation methods effectively extract the volcanic shards from lighter (typically organic) and denser (minerogenic) components, allowing the glass to be microscopically counted to calculate concentrations (see Eden et al., 1996; Turney, 1998; Blockley et al., 2005; Iverson et al., 2017).

To correlate distal ash to its volcanic source and ensure robust correlations, individual glass shards must be geochemically analysed to determine the eruption’s diagnostic ‘fingerprint’. Glass shard



compositions, obtained through electron microprobe (EPMA) analyses, approximate the composition of the magma at the time of the eruption, meaning major (> 1 wt. %) and minor (0.1-1 wt. %) element concentrations can be used to discriminate different eruption events and the different tectonic settings where the magmas were formed. Since some volcanic centres erupt geochemically similar compositions through time (e.g., Óladóttir et al., 2011; Lane et al., 2012; Bourne et al., 2015; Albert et al., 2019; McLean et al., 2020), trace element (< 0.1 wt. %) element compositions are often analysed to further discriminate the deposits of different eruptions (e.g., Albert et al., 2018). Trace element compositions of individual glass shards can be determined using laser ablation-inductively coupled plasma-mass spectrometry (LA-ICP-MS) (e.g., Tomlinson et al., 2010; Pearce et al., 2011), with recent improvements in the spatial resolution (e.g., spot size) and machine sensitivity (e.g., precision and accuracy) allowing the reliable analysis of increasingly small shard sizes.

In addition to providing a valuable relative dating technique, once the eruption age is determined, it can be imported into other archives that contain distal tephra, to improve or independently test the existing chronology. Eruptions can be dated directly using radiometric-dating methods (e.g.,  $^{40}\text{Ar}/^{39}\text{Ar}$  dating or fission track; Brauer et al., 2014), but these techniques often rely on knowing the source volcano to obtain large quantities of minerals from the deposits which are suitable for dating. For example,  $^{40}\text{Ar}/^{39}\text{Ar}$  dating requires measurable quantities of radiogenic argon, formed from the decay of potassium, and therefore K-rich minerals are required for precise ages. In fact, some of the most precise eruption ages are obtained indirectly and are constrained in the medial or distal settings by sedimentary archives such as those provided by lacustrine or marine cores, for which is often possible to obtain detailed age models based on estimated sedimentation rates (e.g., OxCal Bayesian age-models; Ramsey, 2008; Staff et al., 2013), and/or are underpinned by incremental chronologies (e.g., annually laminated varves; Wulf et al., 2004; Smith et al., 2013) or other reference isotope chronostratigraphic curves (e.g. marine isotope geochronology).

The increasing number of distal archives found to contain cryptotephra isochrons, as well as a better understanding of the values and complexities of tephrochronology, has led to the development of regional tephrostratigraphic frameworks (also referred to as lattices) (e.g., Blockley et al., 2014; Davies et al., 2014; Lowe et al., 2015; Fontijn et al., 2016; McLean et al., 2018; Jensen et al., 2021). These frameworks can be constructed using the sequence of ash layers preserved in the geological record to create a network of sites that are bound by co-located markers. These are necessary to grasp a comprehensive understanding of the number and frequency of eruption events, generate constrained eruption chronologies, and build geochemical fingerprints for correlation purposes. Regional frameworks are usually underpinned by key reference sites (often termed tephrostratotypes), which offer both detailed tephrostratigraphic sequences and a precise chronology of eruptive events, usually from a range of volcanic sources (e.g., the Greenland ice cores; Abbott and Davies, 2012; Bourne et al., 2015; Cook et al., 2022).

## *2.2. Tephra dispersal from the North Atlantic region*

There are three key volcanic archipelagos within the North Atlantic with the potential to produce widespread ash dispersal towards NW Africa over the last 300 ka. These include the islands of the Azores, Canary Islands, and Cape Verde (Figure 2). Proximal outcrops of pyroclastic deposits indicate that these ocean island volcanoes have had a wide variety of explosive styles, ranging from low explosivity or largely effusive eruptions, to large caldera-forming eruptions. As discussed, in order to have generated a substantial amounts of ash, explosive eruptions would need to have attained a VEI or M greater or equal to 4 (see Section 4).

Distal cryptotephra layers compositionally attributed to ocean island volcanoes in the North Atlantic region have been identified in sedimentary records in NW Africa and Europe (as marked in Figure 2), showing the remarkable opportunity to link records over continental scales. There has been extensive work on the volcanic stratigraphies of the Azores, Canary Islands, and Cape Verde, but an integrated tephrostratigraphic framework, including detailed major and trace element glass chemistry, for widespread events has not yet been established. Currently, there are few distal ash layers from these volcanic archipelagos that can be confidently linked to specific individual eruptions or used as time-stratigraphic markers.

To date, evidence of Holocene-derived ash from the Azorean volcanoes has been reported in palaeoclimate records in the UK (Chambers et al., 2004; Watson et al., 2017; Walsh et al., 2021), Svalbard (Wastegård et al., 2019; van der Bilt and Lane, 2019), and eastern Europe (Kinder et al., 2020) (Figure 2). Moreover, as part of the 'Response of Humans to Abrupt Environmental Transitions' (RESET) Project (Lowe et al., 2015), pilot investigations at key Palaeolithic archaeological sites in NW Africa identified Atlantic and Mediterranean-derived glass shards in sites including Taforalt (Barton et al., 2015). The RESET Project was also key in verifying that widespread tephra layers have the potential to answer longstanding questions in archaeology (Lowe et al., 2015). For example, volcanic ash of the Campanian Ignimbrite eruption from Campi Flegrei in Italy (dated to ~40 ka; Giaccio et al., 2017) was identified in several palaeoenvironmental sites and archaeological cave sequences, synchronising these eastern Mediterranean records to show spatial and temporal variation in the start of the Upper Palaeolithic lithic industries associated with Anatomically Modern Humans (Lowe et al., 2012). Further work in sites in NW Africa may also locate widespread layers from Italian or Icelandic sources depending on the specific meteorological conditions and eruption source parameters.

### **3. Source regions for widespread ash dispersal in NW Africa**

Here, we explore the documented eruptive histories, eruption source parameters, and published glass geochemical data available for the three volcanic archipelagos, with a specific focus on large eruptions capable of generating widespread tephra fall (over the last ~300 ka). We highlight key references that offer additional information and primary datasets.

#### **3.1. The Azores**

##### *3.1.1. Geological Setting and Eruptive History*

The Azores Archipelago, located in the central North Atlantic ~1700 km from the coast of NW Africa (Figure 2), has an extensive record of explosive eruptions. Due to the prevailing strong south-westerly winds in this region, tephra of Azorean volcanoes has dispersed over wide areas reaching Europe and Africa, including the adjacent continental coastlines.

The Azores are formed of nine islands straddling the triple junction between the Eurasian, African (Nubian), and North American plates, and extending 600 km from WSW to ENE (between latitudes 37° – 40° N and longitudes 25° – 31° W) (see Figure 2b). The volcanic islands are arranged into the eastern (São Miguel and Santa Maria), central (Graciosa, Terceira, São Jorge, Faial, and Pico), and western (Flores and Corvo) groups (see Figure 2b). Volcanism in this region is thought to result from the interaction between a deep melting anomaly (often referred to as the Azores mantle plume) and

volcano-tectonic structures (e.g., Cannat et al., 1999; Trippanera et al., 2014; Storch et al., 2020). Eruptions occur along regional fault zones (volcanic fissure systems) or at the intersection of fault systems (central volcanoes) (Madeira and Brum da Silveira, 2003; Madeira et al., 2015). Almost all the islands consist of one or more central volcanoes intersected by fissure zones with WNW-ESE direction. Seven of the islands have active volcanic systems, most of which have erupted in historical times, i.e., since settlement in the 15th century. Since then, 28 volcanic eruptions (subaerial and submarine) have been recorded (Gaspar et al., 2015a), showcasing the highly active nature of these volcanoes.

Eruptions on the Azores islands have ranged from Hawaiian (effusive) to Plinian (explosive) in style, including Surtseyan-style events. At least four of the nine islands (São Miguel, Terceira, Faial and Graciosa) are known to have produced very recent large magnitude events, with eruption columns extending high into the atmosphere and generating ground-hugging pyroclastic density currents (PDCs), which formed massive pumiceous PDC deposits termed ignimbrites. The Plinian and sub-Plinian events originate from the active central volcanoes with calderas, where intermediate to small volume events are also recurrent in their eruptive histories (e.g., Self, 1976; Booth et al., 1978; Gaspar, 1996; Pacheco, 2001; Gertisser et al., 2010; Guest et al., 2015; Pimentel, 2015; Queiroz et al., 2015; Wallenstein et al., 2015). Given the short distances between the calderas and coastlines, large volumes of tephra are commonly deposited offshore. Thus, eruption volumes are poorly constrained and likely substantially underestimated. However, due to the small size of the Azorean calderas, it is estimated that the larger events probably did not involve more than 1 km<sup>3</sup> DRE of magma (Gertisser et al., 2010; Pimentel et al., 2015). The established island stratigraphies are typically grouped by volcano and delineated using an Upper/Superior group (younger) and Lower/Inferior group (older) scheme, as shown in Figure 3 and Table 1.

### *São Miguel (Eastern Group)*

São Miguel (part of the eastern group along with the inactive Santa Maria) is the largest and most populated of the Azorean islands with >137,000 inhabitants. It is comprised of three active trachytic central volcanoes with calderas, which dominate the island (listed west to east) – Sete Cidades, Fogo (also known as Água de Pau) and Furnas, which are linked by Picos and Congro fissure systems (Figure 3a). These trachytic central volcanoes are the sources of widespread ash layers relevant for linking distal records in this region. The eastern part of the island is formed by the older volcanic systems of Povoação and Nordeste (>878 ka; Johnson et al., 1998) (Figure 3a), which are considered extinct. Although their stratigraphy is poorly understood and radiometric dating has yet to resolve their chronology, it is thought that they have not erupted in at least several hundred thousand years (Johnson et al., 1998; Duncan et al., 2015). The active central volcanoes of São Miguel are characterised by explosive trachytic volcanism of Plinian and sub-Plinian style, while basaltic Hawaiian/Strombolian eruptions dominate in the fissure systems. There are several key studies which have established the volcanostratigraphy of São Miguel (Booth et al., 1978; Queiroz, 1997; Guest et al., 1999; Wallenstein, 1999). The eruptive histories of these centres include several caldera-forming events, and the more recent intra-caldera sub-Plinian and hydromagmatic eruptions (Table 1; Figure 3b).




Figure 3: (a) Map of São Miguel and the location of the three trachytic central volcanoes Sete Cidades, Fogo and Furnas (Basemap: NASA JPL; 2021). Simplified volcanostratigraphic scheme for (b) São Miguel, and (c) Terceira, Faial and Graciosa's volcanoes. These are based on those presented by Queiroz (1997), Queiroz et al. (2008; 2015), Wallenstein (1999), Wallenstein et al. (2015); Guest et al. (1999; 2015); Self (1976); Gertisser et al. (2010), Pimentel et al. (2021), Maderia (1998), Pacheco (2001); Maund (1985), Gaspar (1996), Larrea et al. (2014), respectively.

*Table 1: Simplified stratigraphy and key pyroclastic formations for the Azores Islands (São Miguel, Terceira, Faial, Graciosa islands). Key widespread units are highlighted grey and are used to distinguish groups and link across the islands. Key published geochemical datasets (whole rock and glass) available for the formations are listed.*

<b>Azores Island</b>	<b>Volcanic System</b>	<b>Key Groups/Formations</b>	<b>Key Formations and/or Members</b>	<b>Approximate Age Range</b>	<b>Compositions reference (whole rock and glass)</b>
São Miguel	Sete Cidades	Lagoas	Pepom (P1 to P17) and Cascalho Negro	c. 5 - 500 ka	Ellis et al. (2022)
		Superior Group (Santa Bárbara)	Santa Bárbara	~16 ka	Kueppers et al. (2019); Laeger et al. (2019)
		Superior Group (pre-Santa Bárbara)	Risco, Ajuda, Bretanha, Lombas,	36 - 16 ka	Queiroz et al. (2015)
		Inferior Group	<i>Numerous</i>	210 - 36 ka	Queiroz et al. (2015)
	Fogo (Água de Pau)	Upper Group (post-5 ka)	Fogo A to Fogo 1563	4.6 ka - AD 1563	Wallenstein et al. (2015)
		Upper Group (~40 ka)	Roída de Praia, Ribeira Chã, Pisão	40 - 4.6 ka	Wallenstein et al. (2015)
		Lower Group (pre-40 ka)	<i>Numerous</i>	181 - 40 ka	Wallenstein et al. (2015)
	Furnas	Upper Furnas Group	<i>Numerous including AD 1445 to Furnas A to Furnas AD 1630</i>	5 ka - AD 1630	Jeffery et al. (2016); Guest et al. (2015); Guest et al. (1999)
		Middle Furnas Group	<i>Numerous</i>	27 - 5 ka	Guest et al. (2015); Guest et al. (1999)
		Povoação Ignimbrite Formation	Povoação Ignimbrite	30 ka	Jeffery (2016) unpublished thesis
		Lower Furnas Group (pre-Povoação Ignimbrite)	<i>Numerous</i>	95 - 30 ka	Guest et al. (2015); Guest et al. (1999)
	Terceira	Pico Alto	Upper Terceira Group (including the Lajes-Angra Formation)	<i>Numerous</i>	< 25 - 1 ka
Lower Terceira Group (Old ignimbrite sequence)			<i>Numerous</i>	100 - 25 ka	Pimentel et al. (2021); Jeffery et al. (2017); D'Oriano et al. (2017); Gertisser et al. (2010)
Santa Bárbara		Upper Terceira Group	<i>Numerous</i>	<25 - AD 1761	Pimentel et al. (2016)
Faial	Caldeira	Cedros Volcanic Complex - Upper Group (14 eruptions)	<i>Numerous (C1 to C14)</i>	16 ka - AD 1958	Pimentel et al. (2015); Pacheco (2001); Zanon et al. (2013)
Graciosa	Central	Vitoria-Vulcão Central Volcanic Complex	<i>Numerous</i>	100 ka?	Larrea et al. (2014a; 2014b)

On São Miguel Island, Sete Cidades has been the most active in the last 5000 years (Booth et al., 1978, Queiroz, 1997; Queiroz et al., 2008), with at least seventeen trachytic explosive intracaldera eruptions, predominantly hydromagmatic in nature (part of the Lagoas Formation; eruptions are named P1 to P17; Figure 3). Today, Sete Cidades has a broadly circular caldera 5 km wide, occupied by lakes and several pumice cones, tuff rings and maars. The last paroxysmal eruption dated at ~16 ka (Table 1; Figure 3b) was related to the final phase of caldera-enlarging and is recorded by the Santa Bárbara Formation (Queiroz, 1997; Queiroz et al., 2015; Porreca et al., 2018). Prior to this event, two paroxysmal eruptions related to main phases of caldera collapse are identified, including the Risco Formation (~36 ka) and the Bretanha Formation (~29 ka) (Figure 3b; Queiroz, 1997). All three formations are dominated by ignimbrite members, but also include other members with fallout pumice and minor PDC deposits (e.g. pyroclastic surges and block-and-ash flow deposits). These major pyroclastic formations are intercalated with subordinate trachytic and basaltic products of the Ajuda and Lombas Formations. There are other thick ignimbrites interpreted to be older than these major formations (e.g., located at Rocha da Relva and Ponta da Ferrara) (Queiroz, 1997) and are ascribed to the Inferior Group (Figure 3b; Table 1).

Due to the short distance between the caldera rim and the coast (<2 – 5 km) and the thick cover of younger products, field data for estimation of source parameters of these major eruptions are somewhat limited. However, the existence of a distinctive deposit of the Santa Bárbara formation ~25 km east of Sete Cidades caldera (at Ponta do Cintrão) (Kueppers et al., 2019; Figure 3a) allowed the estimation of eruption source parameters and wind conditions. Numerical simulations suggest the last phase of the eruption was sub-Plinian with an eruption column that extended up to 17 km, dispersed towards E, and had an erupted volume of at least 0.27 km<sup>3</sup> (Kueppers et al., 2019; discussed further in Section 4). Moreover, simulated eruption scenarios of similar events would affect air traffic in the North Atlantic and ash could reach NW Africa, Europe and Central/South America depending on wind direction (Kueppers et al., 2019). The estimated eruption source parameters are in agreement with those by Cole et al. (2008), who modelled sub-Plinian events at Sete Cidades, using a P17 type eruption with a column height of 10 km, and a larger P11 type eruption with a column height of 18 km.

Fogo volcano, also known as Água de Pau, is situated in the central part of São Miguel (Figure 3a) and is responsible for several major explosive eruptions (Wallenstein, 1999; Wallenstein et al., 2015; Figure 3a). The stratigraphic sequences of the northern and southern flanks are described separately since most of the units cannot be correlated/integrated across the volcano (Figure 3b). The stratigraphy of the north flank is composed of several thick pyroclastic sequences, some of which contain ignimbrites and PDC deposits, including the Porto Formoso (~21 ka), the Barrosa, and the Fenais da Luz. Older (>40 ka) unnamed sequences of volcanoclastic deposits including tephra fall and thick ignimbrites have also been identified along the north coastal cliffs of Fogo (Wallenstein, 1999). The south flank sequence includes two thick pyroclastic formations also containing ignimbrites: the Róida da Praia (~34 to 15 ka) and the Ribeira Chã (constrained between 12 and 8 ka). One of the largest and most widespread eruption, named Fogo A (dated to ~4.5 ka), outcrops both north and south of Fogo caldera (Wallenstein, 1999; Pensa et al., 2015) and is commonly used as a time-stratigraphic marker to link the upper stratigraphies and eruptive histories of Sete Cidades and Furnas volcanoes (Figure 3b). This eruption corresponds to a paroxysmal Plinian event, associated with the formation of the caldera, that produced ignimbrites with distinct characteristics (Pensa et al., 2015). The Fogo A eruption parameters estimated by Pensa et al. (2015) suggest this was a VEI 5/M5.6 event with an erupted tephra volume of 4.4 km<sup>3</sup> (see Section 4). Post-Fogo A volcanism includes four trachytic sub-Plinian eruptions, named Fogo B to Fogo 1563 (Wallenstein, 1999; Wallenstein et al., 2015).

Furnas is a nested caldera-complex situated on the eastern side of São Miguel and its stratigraphy includes several major pyroclastic formations (Guest et al., 1999; 2015; Figure 3a). The lower sequence is dominated by trachytic pyroclastic deposits including ignimbrites and other PDC deposits. The largest known eruption of Furnas is represented by the Povoação Ignimbrite Formation (~30 ka) and is interpreted to record the first caldera-forming event (Duncan et al., 1999; Guest et al., 1999; Figure 3b). Other pyroclastic formations located stratigraphically above the Povoação Ignimbrite (i.e., within the Middle Furnas Group) include other key units (oldest to youngest): the Ribeira do Tufo Formation (~27 ka), the Ponta Garça Ignimbrite Formation (~17 ka), the Cancelinha Formation, and an unnamed younger ignimbrite (~12 ka) that outcrops below Pico do Ferro domes. The latter is believed to be associated with the formation of the inner caldera (Guest et al., 1999). Within the Upper Furnas Group (<5 ka), ten intracaldera sub-Plinian eruptions (Furnas A to J), with alternating magmatic and hydromagmatic activity, are recognized; a few of them generated dilute PDCs including the AD 1630 (also known as Furnas J), Furnas I and Furnas C eruptions (Cole et al., 1999).

#### *Terceira, Faial, Graciosa (Central Group)*

Terceira Island is formed by four overlapping central volcanoes (from east to west): Cinco Picos, Guilherme Moniz, Pico Alto, and Santa Bárbara, as well as a Fissure Zone that crosses the island from NW to SE (Self, 1976; Madeira, 2005). Cinco Picos forms the eastern third of the island and is dominated by a large eroded caldera (~7 km in diameter, the largest of the Azores). This extinct volcano is considered to be the oldest eruptive centre (401 ka; Hildenbrand et al., 2014). Guilherme Moniz, situated in the central part of the island, is characterised by a partially destroyed elliptical caldera. Together with Pico Alto (to the north), they form a twin caldera complex. Pico Alto has erupted highly evolved lavas and pyroclastic deposits (pantellerites and comendites) including several major ignimbrites (Gertisser et al., 2010). Santa Bárbara is located in the western third of the island and is the youngest eruptive centre (65 ka; Hildenbrand et al., 2014). The conical-shaped edifice has been truncated by two small overlapping calderas. As shown in Figure 3c, the volcanostratigraphy on Terceira is separated into two main groups following Self (1974) and is delineated by the Lajes-Angra Ignimbrite Formation, which represents the last ignimbrite-forming phase (~25 cal. Ka). These ignimbrites (Lajes and Angra) exhibit a relatively rare peralkaline composition (Pimentel et al., 2021; Figure 4). The Lajes-Angra Ignimbrite Formation is the most widespread formation on Terceira and is used as a key time-stratigraphic marker to link outcrops on the island (Pimentel et al., 2021). The Lower Terceira Group (>25 ka), includes the oldest directly-dated ignimbrite on the Azores with an approximate age of 86 ka, named Ignimbrite i (Gertisser et al., 2010). At least seven pyroclastic formations dominated by ignimbrites are recognized in the last 86 ka and are likely to have originated from the central part of the island (e.g., Pico Alto or Guilherme Moniz volcanoes). The Upper Terceira Group records numerous eruptive episodes from Santa Barbara, Pico Alto, and the Fissure Zone, including nine trachytic sub-Plinian eruptions named A to I (Self, 1974; 1976).

Faial, like other islands of the Azores, is built by central and fissure volcanism. It is composed of four volcanic systems, which include the extinct Ribeirinha shield volcano (850 ka; Hildenbrand et al., 2012), the Caldeira central volcano, and the fissure systems of Horta Platform and Capelo Peninsula (Madeira, 1998; Pacheco, 2001). Explosive volcanism on Faial Island is fairly recent (<16 ka) and restricted to Caldeira Volcano (Figure 3c). The stratigraphy reveals that at least 14 explosive events occurred in this timeframe, two of which have generated PDCs (Pacheco, 2001). The products of Caldeira constitute the Cedros Volcanic Complex (following Pacheco; 2001), which is divided into the

Lower (>16 ka) and the Upper (<16 ka) Groups (Figure 3c). The Upper Group, mainly of Holocene age, is of most relevance for widespread tephra dispersal as it is dominated by trachytic pyroclastic deposits (eruptions C1 to C12; Pacheco, 2001), including the pumice fall and ignimbrite of the major C11 eruption interpreted to represent the first stage of caldera formation (Pimentel et al., 2015).

Graciosa is the northernmost island of the Central Group and consists of a succession of volcanic edifices built one over the other, which have been partially dismantled by faulting and erosion (Maund, 1985; Gaspar, 1996; Larrea et al., 2014a). Three major volcanic complexes are recognized on the island, including (from the oldest to youngest), the Serra das Fontes Complex (>620 ka), the Serra Branca Complex, and the Vitória-Vulcão Central Complex (Gaspar, 1996; Figure 3c). The latter is divided into the Vulcão Central Unit, which comprises a range of rocks from basaltic to trachytic composition (subunits A to V), recording effusive and explosive volcanism, and the Vitória Unit constituted of basaltic products. Most of the products of Serra Branca Complex have been eroded and covered by the younger Vitória-Vulcão deposits. Only one major PDC-forming eruption has been identified on the island, as part of the Upper Hydromagmatic Sequence (subunit S:~12 ka) of the Vulcão Central and is thought to have resulted from a caldera-forming event (Gaspar, 1996).

### 3.1.2. *Glass Geochemistry*

Published geochemical glass compositions of key pyroclastic sequences from the Azorean islands are listed in Table 1 and plotted in Figure 4. Although datasets are missing for many individual eruptive units (particularly those for the lower/inferior groups from the central islands), those available can be used to generate general compositional fields (i.e., fingerprints) of the different volcanic centres and individual eruptions. The largest eruptions of the Azores are generally trachytic in composition, although some are peralkaline (i.e., those from Terceira; Pimentel et al., 2021) and can be easily discriminated from the more heterogeneous eruptive products from the Canary Islands and Cape Verde (Figure 4a). Those geochemically characterised from São Miguel Island (Sete Cidades, Furnas, and Fogo) compositionally overlap on all major elements. The known glass compositions of products erupted from Terceira, can best be discriminated using a  $\text{FeO}_T$  vs  $\text{SiO}_2$  biplot (Figure 4c).

### 3.1.3. *Distal Identifications*

As discussed in Section 2.2, several distal cryptotephra layers compositionally attributed to ocean island volcanoes in the North Atlantic region have been identified in sedimentary records in NW Africa and northern and central Europe (as marked in Figure 2). The glass compositions of these tephra are in agreement with the major element compositional field of the Azores (Figure 4d). However, there have been no visible medial ash deposits (i.e., within a couple of hundred kilometers) of Azores eruptions (i.e., those preserved in offshore marine cores) that can be used to further investigate the eruption source parameters.






Figure 4: (a-c) Published major element glass compositions of key eruptive units from the Azores (plotted with blue symbols and compositional fields), the Canary Islands (orange) and Cape Verde (purple). (d) Glass compositions from distal sedimentary records (1 – 6 on Figure 2) plotted with fields defined in a-c. Glass chemistry data for the Azores (Tomlinson et al., 2015; Johansson et al., 2017; Laeger et al., 2019; Wastegård et al., 2020; Pimentel et al., 2021; Ellis et al., 2022), Canary Islands (Brey and Schmincke, 1980; Bryan et al., 1998; Klügel et al., 2000; Gottsmann and Dingwell, 2001; Olin and Wolff, 2007; Klügel et al., 2005; Galipp et al., 2006; Stroncik et al., 2009; Clay et al., 2011; Del Moro et al., 2015; Di Roberto et al., 2020; Wolff et al., 2020; Romero et al., 2022; Diego González-García, 2022; Jagerup et al., 2023), Cape Verde (Eisele et al., 2015a; Eisele et al., 2015b), and distal records (Chambers et al., 2004; Barton et al., 2015; Watson et al., 2017; van der Bilt and Lane, 2019; Wåstegård et al., 2019; Kinder et al., 2020; Walsh et al., 2021).

## 3.2. The Canary Islands

### 3.2.1. Geological Setting and Eruptive History

The Canary Islands, situated ~300 km from NW Africa, are the second largest intraplate ocean island volcanic system after the Hawaiian chain. The seven islands lie in a complex non-linear age progressive E-W chain and include Lanzarote, Fuerteventura, Gran Canaria, Tenerife, La Gomera, La Palma and El Hierro (Figure 2b). As such, these volcanic islands are in different phases of evolution, ranging from a shield-building stage (El Hierro and La Palma), a rejuvenation phase (Tenerife), and, for the easternmost islands, an erosive phase (Lanzarote and Fuerteventura) (Schmincke, 1979; Carracedo et al., 1998, 1999, 2001). The region is characterised by long-lived volcanic activity, and each of the islands has been active since its formation. Canary Island volcanism is proposed to have stemmed from the upwelling of melt 'blobs' in the Canary plume (Hoernle and Schmincke, 1993). The majority of the Canary Islands' eruptions corresponded to effusive alkali basaltic events; however, some felsic eruptions have occurred at the largest islands of Tenerife and Gran Canaria owing to continued plume activity and magmatic evolution over >10 Myr.

Tenerife is both the largest (2058 km<sup>2</sup>) and tallest (3718 m) of the Canary Islands (Ancochea et al., 1990) and is at the peak of its development (Guillou et al., 2004). It is the only island known to have produced significant explosive events in the last 300 ka (Schmincke and Sumita, 2010; Troll and Carracedo, 2016), several of which were caldera-forming eruptions. The island has been constructed by numerous phases of volcanism, which has spanned more than 12 Myr (Marti and Wolf, 2000). The most recent phase of activity generated a new central stratovolcano complex (Pico Viejo-Pico Teide) within the older Las Cañadas caldera; however, the eruption chronology is currently poorly constrained. Over the island's eruptive history, Tenerife has displayed a large variation in eruption styles, ranging from basaltic lavas from monogenetic cones and fissures to Plinian eruptions producing fallouts and PDCs (Cas et al., 2022).

The stratigraphy, eruption chronology and whole-rock geochemistry of Tenerife are well-studied (e.g., Martí et al., 1994; Bryan et al., 1998; Ancochea et al., 1999; Huertas et al., 2002; Edgar, 2003; Figure 5a; Table 2). The most recent cycle (relevant for the last ~250 ka) is the Diego Hernandez Formation (DHF) (Marti et al., 1994; Wolff et al., 2000; Edgar et al., 2007; Cas et al., 2022), also referred to as the Bandas Del Sur (Brown et al., 2003; Davila-Harris, 2009) (Figure 5a). A number of caldera-forming events, with an estimated deposit volume of >190 km<sup>3</sup> over 11 distinct Members, excluding intracaldera volumes, are grouped into this formation (Cas et al., 2022; Figure 5; Table 2). Two of the largest events in this sequence were the Fasnía (312 ka; Edgar et al., 2007) and Abrigo (~170-190 ka; Brown et al., 2003; Edgar et al., 2007; Table 2), which erupted 30 km<sup>3</sup> and >11 km<sup>3</sup> of tephra respectively, with the column height for the Fasnía estimated at 25 km (Edgar et al., 2007). Similar to the Azores, many large fall deposits have been mostly dispersed offshore or are now significantly eroded on the Island, making the eruption stratigraphy and volume calculations more difficult. In general the formations include a complex sequence of eruptions, with Edgar et al. (2007) noting at least 6 (based on the maximum number of units observed at one location), and as many as 15 (based on the chemical stratigraphy) Plinian eruptions between the Caleta and Abrigo Members alone. There are also numerous minor eruptive units, which cannot be recognised at more than a few exposures, or are of uncertain stratigraphic position (Edgar et al., 2007). Several sub-Plinian events have also been recorded since the Abrigo (<170 ka) that are known to have produced pumice fall deposits and PDCs (Ablay et al., 1995; Gracia et al., 2011; Martí et al., 2012; García et al., 2014).



Table 2: Key pyroclastic formations/members from the Diego Hernandez (Edgar et al., 2007) and Bandas Del Sur (Brown et al., 2003) their ( $^{40}\text{Ar}/^{39}\text{Ar}$ ) geochronological and compositional datasets.

Island	Eruptions	Eruption Age (Brown et al., 2003)	Eruption Age (Edgar et al., 2002/2007/2017)	Compositions ref. (whole rock and glass)
Tenerife	Abrigo	169 ± 1 ka	196 ± 6 ka	Wolff et al. (2020), González-García (2022), Olin (2007)
	Battista	-	234 ± 7 ka	
	La Caleta	221 ± 5 ka	-	Olin (2007)
	Sabinita Formation	-	-	Olin (2007)
	Poris (member 9)	271 ± 6 ka	268 ± 8 ka	Wolff (2020), Edgar et al. (2002)
	Poris (member 2)	276 ± 9 ka	-	Wolff et al., (2020), Edgar et al. (2002)
	Fasnía	289 ± 6 ka	312 ± 6 ka	Olin (2007)
	Aldea Blanca	-	322 ± 5 ka	Olin (2007)

### 3.2.2. Glass Geochemistry

Published geochemical glass compositions relating to key eruption formations/members from the Canary Islands are collated in Table 2 and plotted in Figure 4. The compositional range is notably distinct from those erupted from the Azores and Cape Verde, leaning towards more alkalic compositions. Moreover, the individual eruptions exhibit greater heterogeneity, often spanning the entire compositional range.

## 3.3. Cape Verde

### 3.3.1. Geological Setting and Eruptive History

The Cape Verde (Cabo Verde) volcanic islands are situated ~1000 km SW of NW Africa and may also have dispersed widespread ash across the North Atlantic (Figure 2; Section 4). The archipelago consists of ten major islands, as well as several islets and a number of peripheral seamounts (Kwasnitschka et al., 2024). Volcanism is considered to have formed as a result of mantle plume activity on the Cape Verde Rise. The distribution of the islands forms a horseshoe shape with two island chains (Figure 2d). The eastern to southern chain include the islands of Sal, Boa Vista, Maio, Santiago, Fogo, and Brava. The northern chain include the islands of Santo Antão, São Vicente, Santa Luzia, and São Nicolau (Figure 2d). The eastern to southern chain shows an age progression from NE to SW (Ramalho et al., 2010a; 2010b), with the oldest volcanic activity known from Sal

(Torres et al., 2002). Late Pleistocene to Holocene volcanic activity within the southern chain is limited to the islands of Fogo and Brava and the adjacent Cadamosto seamount (Holm et al., 2008; Ramalho et al., 2010a; 2010b; Grevemeyer et al., 2010; Eisele et al., 2015a; Kwasnitschka et al., 2024), with Fogo and Brava being the most likely candidates for dispersal of ash towards NW Africa. Fogo is the most active and it has at least 28 reported eruptions since its discovery in the 15<sup>th</sup> century (Mata et al., 2017). The eruptive history of Fogo has been divided into four main phases (Day et al., 1999; Foeken et al., 2009), of which the Monte Amarelo Group is the main subaerial phase. This consists of highly alkaline mafic to intermediate lava (Foeken et al., 2009). This phase is known to have terminated with a giant lateral collapse of the Monte Amarelo volcano, which is estimated to have occurred at ~68 ka (Cornu et al., 2021).

Santo Antão, the westernmost island of the northern chain, was also the source of several large explosive eruptions in the last 250 ka. Noteworthy are the sub-Plinian to Plinian Cão Grande eruptions I and II (CG I and CG II), which ejected over 10.3 km<sup>3</sup> (VEI 6) and 3 km<sup>3</sup> (VEI 5) of tephra, respectively (Eisele et al., 2015a; 2016). These two eruptions, which have distinct geochemical fingerprints, happened in close succession at 106 ± 3 ka and 107 ± 15 ka, respectively.

Marine sediment sequences around the ocean islands have been successfully used to determine the stratigraphic order of the eruptive units and identify widespread events. Eisele et al. (2015a) utilised thirteen sediment gravity cores obtained from offshore the southern islands of Fogo and Brava and the Cadamosto seamount (Figure 5b). The tephrostratigraphy of these cores includes 43 mafic and five phonolitic tephra layers spanning the Late Pleistocene to the Holocene. Of these, ten layers could be stratigraphically identified across a region of at least 6200 km<sup>2</sup> to 17,650 km<sup>2</sup>. This revealed that tephra volumes were in order of 1 km<sup>3</sup>, equating to the VEI 5/M5 event and sub-Plinian to Plinian in style. Moreover, the tempo of these events could be elucidated, suggesting that a relatively large magnitude eruption occurred about every 300 years on Fogo during the last 150 kyr. One widespread event identified in these cores (named C12; Figure 5a) could be correlated to Brava, verifying that a widely dispersed eruption event occurred at 145 ka (Eisele et al., 2015a). Additional sedimentary cores north of the islands are required to determine the eruptions that were dispersed in a NW direction.

### 3.3.2. *Glass Geochemistry*

The glass shards in the ash layers preserved in the marine cores and proximal deposits from the southern islands of Fogo and Brava have been characterised using their major and trace elements by Eisele et al. (2015a) producing a detailed compositional field for these volcanic centres and the widespread events. As shown in Figure 4a, glasses are typically less evolved than those of the Canary Islands and the Azores, with higher alkali contents. The glass geochemistry for most eruptive events are homogenous in composition, covering a small section of the overall trend and have phonolitic, intermediate or mafic compositions (Eisele et al., 2015a; Figure 5a). For example, the widespread eruption from Brava (C12; Figure 5a) dated to 145 ka has a SiO<sub>2</sub> content of 53.5 – 57.5 wt. %, whereas the significant C4 eruption at 25 ka from Fogo has a SiO<sub>2</sub> content of 41.2 – 47.3 wt. %. Groups with overlapping major element compositions can be separated using their trace elements. It is not yet clear how the composition of the eruptive products of these islands (Brava and Fogo) compares to others in the Cape Verde Archipelago.

### 3.3.3. *Distal Identifications*

There are no known distal tephra associated with eruptions from the Cape Verde volcanoes in the NW Africa archaeological record. However, considering the eruption size of some of the events (e.g., ~VEI 5/M5) and the proximity of active centres like Fogo and Santo Antão to the western coast of Africa, the tephra must have made it onshore and should be preserved in records (see Section 4).

### **3.4. Italy: potential volcanic regions**

Various volcanic sources in southern Italy, ~1500 km NE of NW Africa, have been active in the last 250 ka, and include Campi Flegrei caldera (e.g., Costa et al., 2022), Roccamarina (De Rita and Giordano, 1996), Aeolian Islands (Lucchi et al., 2013), and Pantelleria (Jordan et al., 2018). Many of the large eruptions from these sources are preserved in distal records in Italy and towards the east (e.g., Lake Ohrid, North Macedonia; Leicher et al., 2019). The 40 ka Campanian Ignimbrite from Campi Flegrei, the largest eruption in Europe during the last 200 kyr, is found in Libya at Haua Fteah (Douka et al., 2014). However, given the prevailing westerly winds, it is unlikely that many of the Italian eruptions, if any, dispersed ash over NW Africa. Hence, the sources and their eruptive histories and compositions are not discussed in detail here.

## **4. Eruption source parameters for ash dispersal to NW Africa**

### *4.1. Ash3D Overview and Input Parameters*

Geological evidence in the distal realm reveals that volcanic ash from sources in the North Atlantic can be dispersed over distances of >5000 km, extending into northern Europe (Figure 2). Yet due to the limited exposure on the islands, eruption source parameters such as tephra volume, column height and mass eruption rate, particle grain size distribution, density, and shape, as well as dispersal direction are poorly constrained. Atmospheric tephra dispersal models, that use databases of known meteorological conditions (e.g., wind speed and direction), offer an excellent means of investigating and forecasting the likely dispersal and limits, for a wide range of eruptive conditions (e.g., tephra volume, column height, eruption duration). Here, we use Ash3D, a three-dimensional Eulerian atmospheric model for tephra transport, dispersal, and deposition, which is frequently used to study and forecast hazards of volcanic ash clouds and ash fallout (Mastin et al., 2014; Schwaiger et al., 2012). This model can be used to predict airborne volcanic ash concentration and tephra deposition during volcanic eruptions. Ash3D models ash transport by dividing the atmosphere into 3D grid cells and calculating the flow of mass through the cells. The model simulates downwind advection, turbulent diffusion, and settling of ash injected into the atmosphere by a volcanic eruption column. The model uses a wind field taken from the global NCEP/NCAR Reanalysis 1 model with a 2.5-degree resolution (Kalnay et al., 1996). Several studies have utilised Ash3D and the modern wind profiles to effectively simulate the tephra dispersal of a range of prehistoric explosive events (e.g., Chang and Yun, 2017; Barker et al., 2019; Buckland et al., 2022).

Numerical simulations using Ash3D allow us to investigate the effects of variable eruption size and meteorological conditions on ash dispersal and critically, ashfall likelihoods across NW Africa. The eruption scenarios are determined using the following criteria:

- 1) Characteristics of explosive eruptions suitable to produce large amounts of ash (e.g., at least  $0.23 \text{ km}^3$  tephra, M4.4), with other source parameters partly constrained by analogous historical eruptions.
- 2) Geological parameters calculated for past eruptions from the source volcanoes (e.g., evidence of previous activity and eruptive volumes known for the last 200 ka).

Here, we simulate ash dispersal using three plausible eruption scenarios, similar to the approach by Barker et al. (2019). These are run for three exemplary volcanic centres in the investigated source regions, including Sete Cidades (São Miguel, Azores), Teide (Tenerife, Canary Islands) and Fogo (Cape Verde). For simplification purposes, eruption events are considered as single phase of activity. Eruption Scenario 1 (S1) is the smallest magnitude scenario, and uses a tephra volume of  $0.23 \text{ km}^3$  ( $0.1 \text{ km}^3$  dense rock equivalent (DRE) magma), with a column height or umbrella cloud top height of 15 km above sea level, and an eruption duration of 6 hours (Figure 6). These parameters are closest to Scenario 1 of Barker et al. (2019). Scenario 2 (S2) uses a tephra volume of  $2.3 \text{ km}^3$  ( $1 \text{ km}^3$  DRE), with the same column height or umbrella cloud top height (15 km above sea level), but with an increased eruption duration of 12 hours. These parameters are similar to Scenario 2 of Barker et al. (2019). Scenario 2b (S2b) has the same eruption source parameters as S2, but the full ash dispersal simulation is run for 72 hours. A longer simulation run is of relevance to understand the entire dispersal footprint potentially depositing enough ash for cryptotephra preservation. To crudely investigate changes in relation to historical wind patterns, each of the Scenarios for the three volcanoes was run on the 1<sup>st</sup> of each month in 2022, as well as several other variations (e.g., runs on consecutive days, varying start times). As shown in Figure 6, we use the volcanic ash concentration results, which simulate possible extents for cryptotephra deposition in distal sedimentary archives (e.g., a tephra load of  $\sim 100 \text{ g/m}^2$  equates to a tephra thickness of  $\sim 1 \text{ mm}$ ).




Figure 6: Selected examples of Ash3D airborne ash (end results) simulated for moderate-sized eruptions from (a) Sete Cidades (Azores; AZ), (b) Teide (Canary Islands; CA) and (c) Fogo (Cape Verde; CV). The input parameters were changed for the three scenarios, as listed on the upper row. The examples show the wind conditions suitable for ash dispersal towards NW Africa and mainland Europe and therefore of relevance for the deposition of time-stratigraphic markers in archaeological and climate records. The vent location is shown by a blue marker. A tephra load of ~100 g/m<sup>2</sup> equates to a tephra thickness of ~1 mm. The rose diagrams show the typical wind directions and speeds in m/s across a vertical section extending to 5000 m (based on measurements from 1990 through to 2009; see header) The diagrams are adapted from those in Mastin (2017).



The three Scenarios (S1, S2 and S2b) follow eruption source parameters estimates for well-studied eruptions in the source regions, as outlined in Section 3. For example, those calculated for a moderate size Azorean eruption (e.g., the sub-Plinian phase of the ~16 ka Santa Bárbara eruption from Sete Cidades) by Kueppers et al. (2019) suggested a tephra volume of 0.27 km<sup>3</sup> and column height of 17 km, falling closest to S1. These parameters also reflected in ash fall models investigated by Cole et al. (2008), who modelled sub-Plinian events at Sete Cidades, with the P17 type eruption (column height of 10 km) and larger P11 type eruption (column height of 18 km). Estimated eruption source parameters for the Fogo A Plinian eruption (~4.6 ka) include a higher tephra volume of 4.4 km<sup>3</sup> (Pensa et al., 2015), thus closer to the larger magnitude scenario S2.

#### *4.2. Model Simulation Observations*

Representative simulations of eruptions with ash dispersal towards NW Africa (e.g., January and November 2022 for the Azores) are shown in Figure 6. As expected, the factors that most influence the ash dispersal relate to the run date and time of day, and are therefore related to the specific meteorological conditions (e.g., wind direction and/or speed). Simulations run on the 1<sup>st</sup> of each month through 2022 show differing ash cloud directions and extents for all regions. The climate across the Atlantic Ocean is highly changeable, with complex weather patterns that can change over relatively short timescales. For the Azores, the dominant ash cloud direction often led towards the E or NE, reflecting the dynamic wind patterns of this region, dominated by the prevailing westerlies (Figure 6a). Although ash dispersal frequently extended towards Morocco (e.g., November), or towards Europe (January) under both S1 and S2 conditions, coverage across both regions required an increase in simulation duration (i.e., the number of hours the ash cloud is tracked) to >24 hours (i.e., as shown in Scenario 2b; Figure 6a). Indeed, the ash cloud direction and extent are highly variable within the same season (although they would have different likelihoods of possibility; Pimentel et al., 2006; Gaspar et al., 2015b), as well as on subsequent days. For example, between the period of 1 – 3<sup>rd</sup> November 2022, ash erupted under S1 conditions was able to reach Morocco on the 1<sup>st</sup>, Portugal on the 2<sup>nd</sup> and remained clustered within the Atlantic Ocean on the 3<sup>rd</sup>. Such stark ash cloud directions are also evident even between simulations run on the same day (e.g., 12 am vs. 12 pm).

In general, if the wind conditions are favourable, the model simulations showed that an ash cloud from all three regions, under both S1 and S2, can reach NW Africa. Specifically, the November 2022 S1 simulation estimates that ash from Sete Cidades would reach NW Africa in <24 hours (Agadir in 16 hours, and Rabat in 21 hours). For S2, the ash cloud arrival times are almost identical (Agadir in 17 hours, and Rabat in 20 hours). For S2b (with an extended run time), ash would arrive in mainland Europe within 23 hours (Gibraltar in ~22 hours, Lisbon in ~33 hours, Cornwall UK in ~38 hours). Such dominant directions also align with geological evidence (e.g., isopach and isopleth maps) for the large eruption mapped on São Miguel (e.g., Santa Bárbara Fm, Fogo A, and others; Booth et al., 1978; Kueppers et al., 2019) and other islands like Flores (Funda Volcanic System; Andrade et al., 2022), which suggest dispersal orientated to the E and NE.

Although the potential ash concentrations are marginally lower in S1 than those in S2, the concentrations correspond to tephra thicknesses around 1 mm, and eruptions should be recorded as cryptotephra over a similar dispersal footprint. It is likely this modelled footprint provides an underestimation of the true extent. The specific conditions for fine ash dispersal and deposition are still poorly understood and modelled (Stevenson et al., 2015; Cashman and Rust, 2020; Krüger and van de Bogaard, 2021). Buckland et al. (2022) discuss many of the contributing reasons why numerical simulations may differ from the field data and cryptotephra extent. Large eruptions can

have multiple phases of activity and complex dynamics. Especially those with significant co-PDC phases can be associated with large volumes of fine ash, and the behaviour of far-travelled ash has been difficult to reconcile with the geological record. The grain size in distal settings can be <100 mm, which has a low particle settling velocity that rarely exceeds the vertical component of air velocity (atmospheric turbulence). Therefore, sedimentation is suppressed, and other mechanisms such as particle aggregation is required, which can be difficult to model numerically. Cryptotephra data are measured by shard counts that lie close to the mass loading limit, and there are also discrepancies between tephrochronology and satellite infrared measurements of volcanic ash (Stevenson et al., 2015). Examples show: (1) 10 – 20 % of the eruptive mass is typically deposited outside the mapped limits, (2) estimates of the ash mass transported in volcanic clouds cannot account for all this unmapped ash, (3) ash fall observed at distances beyond mapped deposits can have measurable impacts, and can form cryptotephra deposits with high shard counts (see Cashman and Rust, 2020).

This work demonstrates that comprehensive ash dispersal models offer a powerful method of investigating the likely source parameters of the prehistoric eruptions from North Atlantic ocean island volcanoes, and assessment of the likelihood of ash reaching NW Africa. It is clear that even moderate-sized eruptions may generate substantial amounts of ash with a widespread dispersal, which is greatly facilitated by the strong and variable winds prevalent in this region. Furthermore, these simulations underscore the necessity for detailed, integrated records of past eruptions from this region. Such records are crucial to prevent the misinterpretation or assumption that distal tephra relate to the largest magnitude events preserved in the proximal realm.

## **5. Conclusions: Building a Tephrostratigraphic Framework for NW Africa**

In this paper, we demonstrate the significant opportunity for utilising tephrochronology in NW Africa to advance the chronology of environmental and behavioural changes in humans over the last 300,000 years. This provides much needed chronological control beyond the radiocarbon limit of ~50,000 years. The likely source regions within the North Atlantic include volcanoes of the Azores, Canary Islands and Cape Verde. As explored, these ocean island volcanoes have undergone a diverse range of eruption styles, many of which could produce widespread ash dispersal over NW Africa if the wind conditions were simultaneously favourable. Collating the available published geochemical glass data reveals that the major element compositions are conducive to fingerprint and discriminate the different source regions. However, the current dataset is still limited and therefore it is still unclear how the compositional diversity has varied through time for the specific volcanoes. Examination of the geological evidence and its integration within atmospheric tephra dispersal models indicates that markers from each of the three regions could be remarkably widespread, serving as critical time-stratigraphic layers to link a broad array of Palaeolithic archaeological and climate sequences (e.g., marine core records). Synchronising these archives would allow the Palaeolithic sequences to be contextualised within their climatic backdrop and shed new light on the role it may have played in shaping the observed behavioural and technological pathways.

The chronological evidence for the largest eruption events indicates tephra would ideally frame pivotal advancements within the archaeological sequences, providing numerous chronological constraints and anchors for synchronising. Eruptions within the Upper/Superior Groups (<50 ka years) from São Miguel (Azores), such as the Povoação Ignimbrite (from Furnas volcano dated to ~30 ka), would be interbedded within key MSA layers (e.g., characterised by core and flake technology). Similarly, the Santa Bárbara eruption (from Sete Cidades volcano dated to ~16 ka), would align with the notable

and widespread transition to the LSA, and a shift towards a more standardised microlithic bladelet production. For older sequences outside the radiocarbon timeframe, the caldera-forming succession from Tenerife (e.g., the Abrigo Formation, dated to ~190-170 ka) could also offer a valuable chronological anchor, particularly for the onset of symbolism and artistic behaviour in the MSA, as observed at key sites like Bizmoune in Morocco (Bouzouggar et al., 2007; 2012; Sehasseh et al., 2021). Importantly, the North Atlantic tephrostratigraphic framework can facilitate future cryptotephra identifications in two key ways: (1) by serving as a predictive tool to determine the likely positioning/age-range for locating key isochrons within archaeological and climate sequences, and (2) by assisting in addressing common taphonomic questions, such as the likelihood of primary tephra deposits and thus, proxy remobilisation (i.e., the possibility of upward or younger reworking within the sequence).

Collating this data underscores the need for further detailed investigations in both proximal and distal zones before tephra layers can be solidly utilised as discrete time-stratigraphic markers. Crucially, additional pyroclastic samples from at least the major eruptions are necessary to generate detailed major and trace element glass fingerprints for individual volcanic centres and eruptions. Presently, the identified distal ash layers in NW Africa can only be tentatively linked to their volcanic source, lacking firm association with dated events. As demonstrated by the dispersal simulations, favourable wind conditions could mean that moderately explosive eruptions may produce widespread ash, further highlighting the importance of compositional fingerprints through the complete stratigraphy. Moreover, it is imperative to establish an integrated tephrostratigraphic framework, or lattices, for the region to accurately determine the relative and absolute timing of widespread events. This is crucial for identifying some of the eruptions that may have limited/no exposure on the relatively small volcanic islands. Developing such a framework entails detailed tephrostratigraphic studies in the medial and distal regions, utilising offshore marine cores spanning the coast of NW Africa. These cores hold immense potential for constructing a clear and integrated record and importantly identify the isochrons suitable to precisely synchronise marine core climate records and the Palaeolithic sequences in NW Africa for the first time.

## Acknowledgments

DM is supported by an Early Career Fellowship funded by the Leverhulme Trust (grant: ECF-2020-151) and the John Fell Fund (#9438), held at the University of Oxford, and a British Institute for Libyan & Northern African Studies (BILNAS) Postdoctoral Writing Fellowship Award (2024). Many researchers on this paper are funded by a Leverhulme Trust Award entitled “Agencies of behavioural change in early modern humans in NW Africa (CAVES Project)” (grant: RPG-2022-0490; NB is the Principal Investigator; <https://caves.web.ox.ac.uk>). AP acknowledges the CEEC Institutional contract Ref. CEECINST/00024/2021/CP2780/CT0003 funded by Fundação para a Ciência e a Tecnologia (<https://doi.org/10.54499/CEECINST/00024/2021/CP2780/CT0003>). The authors would also like to acknowledge additional financial support from BILNAS, awarded to DM (in 2022) and SC, ELH (2023), and the Boise Trust (Research Award, University of Oxford) that were awarded to VV (2022) and DM (2023). The authors would like to thank Prof. Abdeljalil Bouzouggar (Director of the Institut National des Sciences de l'Archéologie et du Patrimoine, Morocco) for his valuable assistance throughout the project. We also acknowledge fieldwork assistance and expertise provided by Margarida Ramalho and Dr. Mariana Andrade. We thank Prof. Stefan Wastegård and an anonymous reviewer for their constructive comments which helped to improve the manuscript, and Dr. Victoria Leitch for editorial handling.

## References

- Abbott, P.M. and Davies, S.M., 2012. Volcanism and the Greenland ice-cores: the tephra record. *Earth-Science Reviews*, 115(3), pp.173-191.
- Abbott, P. M., Griggs, A. J., Bourne, A. J. and Davies, S. M. 2018. Tracing marine cryptotephra in the North Atlantic during the last glacial period: Protocols for identification, characterisation and evaluating depositional controls. *Marine Geology*, 401: 81-97
- Ablay, G. J., Ernst, G. G. J., Marti, J., & Sparks, R. S. J. (1995). The ~ 2 ka subplinian eruption of Montaña Blanca, Tenerife. *Bulletin of Volcanology*, 57, 337-355.
- Abrook, A.M., Matthews, I.P., Candy, I., Palmer, A.P., Francis, C.P., Turner, L., Brooks, S.J., Self, A.E. and Milner, A.M., 2020. Complexity and asynchrony of climatic drivers and environmental responses during the Last Glacial-Interglacial Transition (LGIT) in north-west Europe. *Quaternary Science Reviews*, 250, p.106634.
- Albert, P.G., Smith, V.C., Suzuki, T., Tomlinson, E.L., Nakagawa, T., McLean, D., Yamada, M., Staff, R.A., Scholout, G., Takemura, K. and Nagahashi, Y., 2018. Constraints on the frequency and dispersal of explosive eruptions at Sambe and Daisen volcanoes (South-West Japan Arc) from the distal Lake Suigetsu record (SG06 core). *Earth-science reviews*, 185, pp.1004-1028.
- Albert, P.G., Smith, V.C., Suzuki, T., McLean, D., Tomlinson, E.L., Miyabuchi, Y., Kitaba, I., Mark, D.F., Moriwaki, H., Members, S.P. and Nakagawa, T., 2019. Geochemical characterisation of the Late Quaternary widespread Japanese tephrostratigraphic markers and correlations to the Lake Suigetsu sedimentary archive (SG06 core). *Quaternary Geochronology*, 52, pp.103-131.
- Ancochea, E., Fuster, J., Ibarrola, E., Cendrero, A., Coello, J., Hernan, F., Cantagrel, J.M. and Jamond, C., 1990. Volcanic evolution of the island of Tenerife (Canary Islands) in the light of new K-Ar data. *Journal of Volcanology and Geothermal Research*, 44(3-4), pp.231-249.
- Andrade, M., Pimentel, A., Ramalho, R., Kutterolf, S. and Hernández, A., 2022. The recent volcanism of Flores Island (Azores): Stratigraphy and eruptive history of Funda Volcanic System. *Journal of Volcanology and Geothermal Research*, 432, p.107706.
- Barker, S.J., Van Eaton, A.R., Mastin, L.G., Wilson, C.J., Thompson, M.A., Wilson, T.M., Davis, C. and Renwick, J.A., 2019. Modeling ash dispersal from future eruptions of Taupo supervolcano. *Geochemistry, Geophysics, Geosystems*, 20(7), pp.3375-3401.
- Barton, R.N.E., Bouzouggar, A., Hogue, J.T., Lee, S., Collcutt, S.N. and Ditchfield, P., 2013. Origins of the Iberomaurusian in NW Africa: new AMS radiocarbon dating of the Middle and Later Stone Age deposits at Tafalet Cave, Morocco. *Journal of Human Evolution*, 65(3), pp.266-281.
- Barton, R.N.E., Lane, C.S., Albert, P.G., White, D., Collcutt, S.N., Bouzouggar, A., Ditchfield, P., Farr, L., Oh, A., Ottolini, L. and Smith, V.C., 2015. The role of cryptotephra in refining the chronology of Late Pleistocene human evolution and cultural change in North Africa. *Quaternary Science Reviews*, 118, pp.151-169.

Barton, R.N.E., Bouzouggar, A., Colcutt, S.N., Marco, Y.C., Clark-Balzan, L., Debenham, N.C. and Morales, J., 2016. Reconsidering the MSA to LSA transition at Taforalt Cave (Morocco) in the light of new multi-proxy dating evidence. *Quaternary International*, 413, pp.36-49.

Barton, R.N.E., Bouzouggar, A., Colcutt, S. N., Humphrey, L. T. (Hrsg.): *Cemeteries and Sedentism in the Later Stone Age of NW Africa: Excavations at Grotte des Pigeons, Taforalt, Morocco*, Heidelberg: Propylaeum, 2020 (Monographien des RGZM, Band 147).

Barton, R.N.E., Belhouchet, L., Colcutt, S.N., Aouadi, N., Albert, P.G., Douka, K., Drake, N., Linderholm, L., Macphail, R.I., McLean, D. and Mekki, H., 2021. New insights into the late Middle Stone Age occupation of Oued el Akarit, southern Tunisia. *Libyan Studies*, 52, pp.12-35.

Barton, N. and d'Errico, F., 2012. North African origins of symbolically mediated behaviour and the Aterian. In *Developments in Quaternary Sciences* (Vol. 16, pp. 23-34). Elsevier.

Becerra-Valdivia, L., & Higham, T. (2023). New Developments in Radiocarbon Dating. *Handbook of Archaeological Sciences*, 1, 25-35.

Blockley, S.P., Pyne-O'Donnell, S.D., Lowe, J.J., Matthews, I.P., Stone, A., Pollard, A.M., Turney, C.S. and Molyneux, E.G., 2005. A new and less destructive laboratory procedure for the physical separation of distal glass tephra shards from sediments. *Quaternary Science Reviews*, 24(16-17), pp.1952-1960.

Blockley, S.P., Bourne, A.J., Brauer, A., Davies, S.M., Hardiman, M., Harding, P.R., Lane, C.S., MacLeod, A., Matthews, I.P., Pyne-O'Donnell, S.D. and Rasmussen, S.O., 2014. Tephrochronology and the extended intimate (integration of ice-core, marine and terrestrial records) event stratigraphy 8–128 ka b2k. *Quaternary Science Reviews*, 106, pp.88-100.

Blome, M.W., Cohen, A.S., Tryon, C.A., Brooks, A.S. and Russell, J., 2012. The environmental context for the origins of modern human diversity: a synthesis of regional variability in African climate 150,000–30,000 years ago. *Journal of human evolution*, 62(5), pp.563-592.

Bonnefille, R., 2010. Cenozoic vegetation, climate changes and hominid evolution in tropical Africa. *Glob. Planet. Change* 72(4), 390-411.

Booth, B., Croasdale, R., and Walker, G. P. L., 1978. A quantitative study of five thousand years of volcanism on S. Miguel, Azores, *Philos. T. R. Soc. A.*, 228, 271319,

Bourne, A.J., Cook, E., Abbott, P.M., Seierstad, I.K., Steffensen, J.P., Svensson, A., Fischer, H., Schüpbach, S. and Davies, S.M., 2015. A tephra lattice for Greenland and a reconstruction of volcanic events spanning 25–45 ka b2k. *Quaternary Science Reviews*, 118, pp.122-141.

Bouzouggar, A., Barton, N., Vanhaeren, M., d'Errico, F., Colcutt, S., Higham, T., Hodge, E., Parfitt, S., Rhodes, E., Schwenninger, J.L. and Stringer, C., 2007. 82,000-year-old shell beads from North Africa and implications for the origins of modern human behavior. *Proceedings of the National Academy of Sciences*, 104(24), pp.9964-9969.

Bouzouggar, A. and Barton, R.N.E., 2012. The identity and timing of the Aterian in Morocco. Modern origins: a North African perspective, pp.93-105.

Brahim, Y.A., Sha, L., Wassenburg, J.A., Azennoud, K., Cheng, H., Cruz, F.W. and Bouchaou, L., 2023. The spatiotemporal extent of the Green Sahara during the last glacial period. *Iscience*, 26(7).

Brassell, S.C., Eglinton, G., Marlowe, I.T., Pflaumann, U., Sarnthein, M., 1986. Molecular stratigraphy: a new tool for climatic assessment. *Nature* 320(6058), 129-133. doi.org/10.1038/320129a0.

Brauer, A., Hajdas, I., Blockley, S. P., Ramsey, C. B., Christl, M., Ivy-Ochs, S., Moseley, G. E., Nowaczyk, N. N., Rasmussen, S. O., Roberts, H. M. and Spötl, C. 2014. The importance of independent chronology in integrating records of past climate. 228 change for the 60–8 ka INTIMATE time interval. *Quaternary Science Reviews*, 106: 47-66

Brey, G. and Schmincke, H.U., 1980. Origin and diagenesis of the Roque Nublo breccia, Gran Canaria (Canary Islands)—Petrology of roque nublo volcanics, II. *Bulletin Volcanologique*, 43, pp.15-33.

Brown, R.J., Barry, T.L., Branney, M.J., Pringle, M.S. and Bryan, S.E., 2003. The Quaternary pyroclastic succession of southeast Tenerife, Canary Islands: explosive eruptions, related caldera subsidence, and sector collapse. *Geological Magazine*, 140(3), pp.265-288.

Bryan, S.E., Cas, R.A. and Martí, J., 1998. Lithic breccias in intermediate volume phonolitic ignimbrites, Tenerife (Canary Islands): constraints on pyroclastic flow depositional processes. *Journal of Volcanology and Geothermal Research*, 81(3-4), pp.269-296.

Buckland, H.M., Mastin, L.G., Engwell, S.L. and Cashman, K.V., 2022. Modelling the transport and deposition of ash following a magnitude 7 eruption: the distal Mazama tephra. *Bulletin of Volcanology*, 84(9), p.87.

Cannat, M., Briais, A., Deplus, C., Escartín, J., Georgen, J., Lin, J., Mercouriev, S., Meyzen, C., Muller, M., Pouliquen, G. and Rabain, A., 1999. Mid-Atlantic Ridge–Azores hotspot interactions: along-axis migration of a hotspot-derived event of enhanced magmatism 10 to 4 Ma ago. *Earth and Planetary Science Letters*, 173(3), pp.257-269.

Carracedo, J.C., Day, S., Guillou, H., Badiola, E.R., Cañas, J.A. and Torrado, F.P., 1998. Hotspot volcanism close to a passive continental margin: the Canary Islands. *Geological magazine*, 135(5), pp.591-604.

Carracedo, J.C., Day, S.J., Guillou, H. and Gravestock, P., 1999. Later stages of volcanic evolution of La Palma, Canary Islands: Rift evolution, giant landslides, and the genesis of the Caldera de Taburiente. *Geological Society of America Bulletin*, 111(5), pp.755-768.

Carracedo, J.C., Badiola, E.R., Guillou, H., de La Nuez, J. and Torrado, F.P., 2001. Geology and volcanology of la Palma and el Hierro, western Canaries. *Estudios Geologicos-Madrid*, 57, pp.175-273.

Cas, R.A., Wolff, J.A., Martí, J., Olin, P.H., Edgar, C.J., Pittari, A. and Simmons, J.M., 2022. Tenerife, a complex end member of basaltic oceanic island volcanoes, with explosive polygenetic phonolitic calderas, and phonolitic-basaltic stratovolcanoes. *Earth-Science Reviews*, 230, p.103990.

Cashman, K.V. and Rust, A.C., 2020. Far-travelled ash in past and future eruptions: combining tephrochronology with volcanic studies. *Journal of Quaternary Science*, 35(1-2), pp.11-22.

Chambers, F.M., Daniell, J.R., Hunt, J.B., Molloy, K. and O'Connell, M., 2004. Tephrostratigraphy of An Loch Mor, Inis Oirr, western Ireland: implications for Holocene tephrochronology in the northeastern Atlantic region. *The Holocene*, 14(5), pp.703-720.

Chang, C. and Yun, S.H., 2017. The numerical simulation of volcanic ash dispersion at Aso Caldera volcano using Ash3D model. *Journal of the Korean earth science society*, 38(2), pp.115-128.

Chase, B.M., Faith, J.T., Mackay, A., Chevalier, M., Carr, A.S., Boom, A., Lim, S. and Reimer, P.J., 2018. Climatic controls on Later Stone Age human adaptation in Africa's southern Cape. *Journal of Human Evolution*, 114, pp.35-44.

Clay, P.L., Kelley, S.P., Sherlock, S.C. and Barry, T.L., 2011. Partitioning of excess argon between alkali feldspars and glass in a young volcanic system. *Chemical Geology*, 289(1-2), pp.12-30.

Cole, P.D., Pacheco, J.M., Gunasekera, R., Queiroz, G., Gonçalves, P. and Gaspar, J.L., 2008. Contrasting styles of explosive eruption at Sete Cidades, São Miguel, Azores, in the last 5000 years: hazard implications from modelling. *Journal of Volcanology and Geothermal Research*, 178(3), pp.574-591

Cook, E., Abbott, P.M., Pearce, N.J., Mojtabavi, S., Svensson, A., Bourne, A.J., Rasmussen, S.O., Seierstad, I.K., Vinther, B.M., Harrison, J. and Street, E., 2022. Volcanism and the Greenland ice cores: A new tephrochronological framework for the last glacial-interglacial transition (LGIT) based on cryptotephra deposits in three ice cores. *Quaternary Science Reviews*, 292, p.107596.

Cornu, M.N., Paris, R., Doucelance, R., Bachélery, P., Bosq, C., Auclair, D., Benbakkar, M., Gannoun, A.M. and Guillou, H., 2021. Exploring the links between volcano flank collapse and the magmatic evolution of an ocean island volcano: Fogo, Cape Verde. *Scientific Reports*, 11(1), p.17478.

Costa, A., Di Vito, M.A., Ricciardi, G.P., Smith, V.C. and Talamo, P., 2022. The long and intertwined record of humans and the Campi Flegrei volcano (Italy). *Bulletin of Volcanology*, 84, pp.1-27.

Davies, S.M., 2015. Cryptotephra: the revolution in correlation and precision dating. *Journal of Quaternary Science*, 30(2), pp.114-130.

Davies, S.M., Branch, N.P., Lowe, J.J. and Turney, C.S., 2002. Towards a European tephrochronological framework for Termination 1 and the Early Holocene. *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 360(1793), pp.767-802.

Davies, S.M., Abbott, P.M., Pearce, N.J., Wastegård, S. and Blockley, S.P., 2012. Integrating the INTIMATE records using tephrochronology: rising to the challenge. *Quaternary Science Reviews*, 36, pp.11-27.

Davies, S.M., Abbott, P.M., Meara, R.H., Pearce, N.J., Austin, W.E., Chapman, M.R., Svensson, A., Bigler, M., Rasmussen, T.L., Rasmussen, S.O. and Farmer, E.J., 2014. A North Atlantic tephrostratigraphical framework for 130–60 ka b2k: new tephra discoveries, marine-based correlations, and future challenges. *Quaternary Science Reviews*, 106, pp.101-121.

Dávila Harris, P., 2009. Explosive ocean-island volcanism: the 1.8–0.7 Ma explosive eruption history of Cañadas volcano recorded by the pyroclastic successions around Adeje and Abona, southern Tenerife, Canary Islands (Doctoral dissertation, University of Leicester).

Day, C.C., Couper, H.O., Barrott, J.J., Carolin, S.A., Bouzouggar, A., Barton, R.N.E. and Henderson, G.M. Timing and cause of increased water availability in arid South-of-Atlas region - reconstruction from cave stalagmites. *Goldschmidt 2023 Conference*, 2023.

Day, S.J., Da Silva, S.H. and Fonseca, J.F.B.D., 1999. A past giant lateral collapse and present-day flank instability of Fogo, Cape Verde Islands. *Journal of Volcanology and Geothermal Research*, 94(1-4), pp.191-218.

De Rita, D. and Giordano, G., 1996. Volcanological and structural evolution of Roccamonfina volcano (Italy): origin of the summit caldera. *Geological Society, London, Special Publications*, 110(1), pp.209-224.

Del Moro, S., Di Roberto, A., Meletlidis, S., Pompilio, M., Bertagnini, A., Agostini, S., Ridolfi, F. and Renzulli, A., 2015. Xenopumice erupted on 15 October 2011 offshore of El Hierro (Canary Islands): a subvolcanic snapshot of magmatic, hydrothermal and pyrometamorphic processes. *Bulletin of Volcanology*, 77, pp.1-19.

Di Roberto, A., Risica, G., Del Carlo, P., Pompilio, M., Speranza, F. and Meletlidis, S., 2020. The forgotten eruption: The basaltic scoria cone of Montaña Grande, Tenerife. *Journal of Volcanology and Geothermal Research*, 401, p.106918.

Dibble, H.L., Reed, D., Rezek, Z., Richter, D., Roberts, R.G., Sandgathe, D., Schurmans, U., Skinner, A.R., Steele, T.E., Reed, K. and Olszewski, D.I., 2012. New Excavations at the Site of Contrebandiers Cave, Morocco. *PaleoAnthropology*, 2012, pp.145-201.

D'Oriano, C., Landi, P., Pimentel, A. and Zanon, V., 2017. Magmatic processes revealed by anorthoclase textures and trace element modeling: The case of the Lajes Ignimbrite eruption (Terceira Island, Azores). *Journal of Volcanology and Geothermal Research*, 347, pp.44-63.

Douka, K., Jacobs, Z., Lane, C., Grün, R., Farr, L., Hunt, C., Inglis, R.H., Reynolds, T., Albert, P., Aubert, M. and Cullen, V., 2014. The chronostratigraphy of the Haua Fteah cave (Cyrenaica, northeast Libya). *Journal of Human Evolution*, 66, pp.39-63.



Duncan, A.M., Queiroz, G., Guest, J.E., Cole, P.D., Wallenstein, N. and Pacheco, J.M., 1999. The Povoação Ignimbrite, Furnas Volcano, São Miguel, Azores. *Journal of volcanology and geothermal research*, 92(1-2), pp.55-65.

Duncan, A.M., Guest, J.E., Wallenstein, N., Chester, D.K., 2015. The older volcanic complexes of São Miguel, Azores: Nordeste and Povoação. In: Gaspar, J.L., Guest, J.E., Duncan, A.M., Barriga, F.J.A.S., Chester, D.K. (Eds.), *Volcanic Geology of São Miguel Island (Azores Archipelago)*. Geological Society, London, *Memoirs*, vol. 44, pp. 147–153.

Eden, D.N., Froggatt, P.C., Zheng, H. and Machida, H., 1996. Volcanic glass found in Late Quaternary Chinese loess: A pointer for future studies?. *Quaternary International*, 34, pp.107-111.

Edgar, C.J., Wolff, J.A., Nichols, H.J., Cas, R.A. and Martí, J., 2002. A complex Quaternary ignimbrite-forming phonolitic eruption: the Poris member of the Diego Hernández Formation (Tenerife, Canary Islands). *Journal of Volcanology and Geothermal Research*, 118(1-2), pp.99-130.

Edgar, C.J., 2003. *The Stratigraphy & Eruption Dynamics of a Quaternary Phonolitic Plinian Eruption Sequence: The Diego Hernández Formation, Tenerife, Canary Islands (Spain)* (Doctoral dissertation, Monash University).

Edgar, C.J., Wolff, J.A., Olin, P.H., Nichols, H.J., Pittari, A., Cas, R.A.F., Reiners, P.W., Spell, T.L. and Martí, J., 2007. The late Quaternary Diego Hernandez Formation, Tenerife: Volcanology of a complex cycle of voluminous explosive phonolitic eruptions. *Journal of Volcanology and Geothermal Research*, 160(1-2), pp.59-85.

Eisele, S., Freundt, A., Kutterolf, S., Ramalho, R.S., Kwasnitschka, T., Wang, K.L. and Hemming, S.R., 2015. Stratigraphy of the Pleistocene, phonolitic Cão Grande Formation on Santo Antão, Cape Verde. *Journal of Volcanology and Geothermal Research*, 301, pp.204-220.

Eisele, S., Reißig, S., Freundt, A., Kutterolf, S., Nürnberg, D., Wang, K.L. and Kwasnitschka, T., 2015. Pleistocene to Holocene offshore tephrostratigraphy of highly explosive eruptions from the southwestern Cape Verde Archipelago. *Marine Geology*, 369, pp.233-250.

Eisele, S., Freundt, A., Kutterolf, S., Hansteen, T.H., Klügel, A. and Irion, I.M., 2016. Evolution of magma chambers generating the phonolitic Cão Grande Formation on Santo Antão, Cape Verde Archipelago. *Journal of Volcanology and Geothermal Research*, 327, pp.436-448.

Elderfield, H., Ganssen, G., 2000. Past temperature and  $\delta^{18}\text{O}$  of surface ocean waters inferred from foraminiferal Mg/Ca ratios. *Nature* 405(6785), 442-445. doi.org/10.1038/35013033.

Ellis, B.S., Pimentel, A., Wolff, J.A., Etter, A., Cortes-Calderon, E.A., Harris, C., Mark, D.F., Neukampf, J. and Bachmann, O., 2022. Geochemistry of the Pepom tephra deposits: The most recent intracaldera volcanism of Sete Cidades volcano, São Miguel, Azores. *Journal of Volcanology and Geothermal Research*, 432, p.107673.

Foeken, J.P., Day, S. and Stuart, F.M., 2009. Cosmogenic  $^3\text{He}$  exposure dating of the Quaternary basalts from Fogo, Cape Verdes: implications for rift zone and magmatic reorganisation. *Quaternary Geochronology*, 4(1), pp.37-49.

Fontijn, K., Rawson, H., Van Daele, M., Moernaut, J., Abarzúa, A.M., Heirman, K., Bertrand, S., Pyle, D.M., Mather, T.A., De Batist, M. and Naranjo, J.A., 2016. Synchronisation of sedimentary records using tephra: A postglacial tephrochronological model for the Chilean Lake District. *Quaternary Science Reviews*, 137, pp.234-254.

Galipp, K., Klügel, A. and Hansteen, T.H., 2006. Changing depths of magma fractionation and stagnation during the evolution of an oceanic island volcano: La Palma (Canary Islands). *Journal of Volcanology and Geothermal Research*, 155(3-4), pp.285-306.

Garcea, E.A. and Giraudi, C., 2006. Late Quaternary human settlement patterning in the Jebel Gharbi. *Journal of Human Evolution*, 51(4), pp.411-421.

García, O., Guzman, S.R. and Martí, J., 2014. Stratigraphic correlation of Holocene phonolitic explosive episodes of the Teide-Pico Viejo Complex, Tenerife.

García, O., Martí, J., Aguirre, G., Geyer, A., & Iribarren, I. (2011). Pyroclastic density currents from Teide–Pico Viejo (Tenerife, Canary Islands): implications for hazard assessment. *Terra Nova*, 23(3), 220-224.

Gaspar, J.L., 1996. Ilha Graciosa (Açores): História vulcanológica e avaliação do hazard. PhD thesis, Universidade dos Açores, Ponta Delgada, 261 pp.

Gaspar, J.L., Queiroz, G., Ferreira, T., Medeiros, A.R., Goulart, C., Medeiros, J., (2015a). Earthquakes and volcanic eruptions in the Azores region: Geodynamic implications from major historical events and instrumental seismicity. In: Gaspar, J.L., Guest, J.E., Duncan, A.M., Barriga, F.J.A.S., Chester, D.K. (Eds.), *Volcanic Geology of São Miguel Island (Azores Archipelago)*, Geological Society, London, *Memoirs*, vol. 44, pp. 33–49.

Gaspar, J. L., Guest, J. E., Queiroz, G., Pacheco, J., Pimentel, A., Gomes, A., Marques, R., Felpeito, A., Ferreira, T., and Wallenstein, N. (2015b) – Eruptive frequency and volcanic hazards zonation in São Miguel Island, Azores, in: *Volcanic Geology of São Miguel Island (Azores Archipelago)*, *Memoirs*, 44, edited by: Gaspar, J. L., Guest, J. E., Duncan, A. M., Barriga, F. J. A. S., and Chester, D. K., Geological Society, London,

Gertisser, R., Self, S., Gaspar, J.L., Kelley, S.P., Pimentel, A., Eikenberg, J., Barry, T.L., Pacheco, J.M., Queiroz, G. and Vespa, M., 2010. Ignimbrite stratigraphy and chronology on Terceira Island, Azores. *The Geological Society of America, Special Paper*, 464, pp.133-154.

Giaccio, B., Hajdas, I., Isaia, R., Deino, A. and Nomade, S., 2017. High-precision  $^{14}\text{C}$  and  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of the Campanian Ignimbrite (Y-5) reconciles the time-scales of climatic-cultural processes at 40 ka. *Scientific reports*, 7(1), p.45940.

González-García, D., Petrelli, M., Perugini, D., Giordano, D., Vasseur, J., Paredes-Mariño, J., Marti, J. and Dingwell, D.B., 2022. Pre-eruptive conditions and dynamics recorded in banded pumices from the El Abrigo caldera-forming eruption (Tenerife, Canary Islands). *Journal of Petrology*, 63(3), p.egac009.

Gottsmann, J. and Dingwell, D.B., 2001. Cooling dynamics of spatter-fed phonolite obsidian flows on Tenerife, Canary Islands. *Journal of Volcanology and Geothermal Research*, 105(4), pp.323-342.

Griggs, A.J., Davies, S.M., Abbott, P.M., Coleman, M., Palmer, A.P., Rasmussen, T.L. and Johnston, R., 2015. Visualizing tephra deposits and sedimentary processes in the marine environment: The potential of X-ray microtomography. *Geochemistry, Geophysics, Geosystems*, 16(12), pp.4329-4343.

Guest, J.E., Gaspar, J.L., Cole, P.D., Queiroz, G., Duncan, A.M., Wallenstein, N., Ferreira, T. and Pacheco, J.M., 1999. Volcanic geology of Furnas Volcano, São Miguel, Azores. *Journal of Volcanology and Geothermal Research*, 92(1-2), pp.1-29.

Guest, J.E., Pacheco, J.M., Cole, P.D., Duncan, A.M., Wallenstein, N., Queiroz, G., Gaspar, J.L. and Ferreira, T., 2015. Chapter 9 The volcanic history of Furnas Volcano, São Miguel, Azores. *Geological Society, London, Memoirs*, 44(1), pp.125-134.

Grevemeyer, I., Helffrich, G., Faria, B., Booth-Rea, G., Schnabel, M. and Weinrebe, W., 2010. Seismic activity at Cadamosto seamount near Fogo Island, Cape Verdes—formation of a new ocean island?. *Geophysical Journal International*, 180(2), pp.552-558.

Grün, R. and Stringer, C., 2023. Direct dating of human fossils and the ever-changing story of human evolution. *Quaternary Science Reviews*, 322, p.108379.

Gudmundsdóttir, E. R., Eiríksson, J. and Larsen, G. 2011. Identification and definition of primary and reworked tephra in Late Glacial and Holocene marine shelf sediments off North Iceland. *Journal of Quaternary Science*, 26: 589-602

Guillou, H., Carracedo, J.C., Paris, R. and Torrado, F.J.P., 2004. Implications for the early shield-stage evolution of Tenerife from K/Ar ages and magnetic stratigraphy. *Earth and Planetary Science Letters*, 222(2), pp.599-614.

Higham, T., Douka, K., Wood, R., Ramsey, C.B., Brock, F., Basell, L., Camps, M., Arrizabalaga, A., Baena, J., Barroso-Ruiz, C. and Bergman, C., 2014. The timing and spatiotemporal patterning of Neanderthal disappearance. *Nature*, 512(7514), pp.306-309.

Hildenbrand, A., Weis, D., Madureira, P., and Marques, F. O. (2014). Recent plate re-organization at the Azores Triple Junction: evidence from combined geochemical and geochronological data on Faial, S. Jorge and Terceira volcanic islands. *Lithos* 210-211, 27–39

Hildenbrand A, Marques FO, Costa ACG, Sibrant ALR, Silva PF, Henry B, Miranda JM, Madureira P (2012) Reconstructing the architectural evolution of volcanic islands from combined K/Ar, morphologic, tectonic, and magnetic data: the Faial Island example (Azores). *J Volcanol Geotherm Res* 241–242:39–48.

Hoernle, K.A.J. and Schmincke, H.U., 1993. The petrology of the tholeiites through melilite nephelinites on Gran Canaria, Canary Islands: crystal fractionation, accumulation, and depths of melting. *Journal of Petrology*, 34(3), pp.573-597.

Holm, P.M., Wilson, J.R., Christensen, B.P., Hansen, L., Hansen, S.L., Hein, K.M., Mortensen, A.K., Pedersen, R., Plesner, S. and Runge, M.K., 2003, December. Sampling the Cape Verde Mantle Plume: Evolution of Santo Antão, Cape Verde Islands. In *AGU Fall Meeting Abstracts* (Vol. 2003, pp. V32A-0998).

Holm, P.M., Grandvuinet, T., Friis, J., Wilson, J.R., Barker, A.K. and Plesner, S., 2008. An  $^{40}\text{Ar}$ - $^{39}\text{Ar}$  study of the Cape Verde hot spot: Temporal evolution in a semistationary plate environment. *Journal of Geophysical Research: Solid Earth*, 113(B8).

Consortium\*†, T.C.C.P.I.P., Hönisch, B., Royer, D.L., Breecker, D.O., Polissar, P.J., Bowen, G.J., Henehan, M.J., Cui, Y., Steinthorsdottir, M., McElwain, J.C., Kohn, M.J., Pearson, A., Phelps, S.R., Uno, K.T., Ridgwell, A., Anagnostou, E., Austermann, J., Badger, M.P.S., Barclay, R.S., Bijl, P.K., Chalk, T.B., Scotese, C.R., de la Vega, E., DeConto, R.M., Dyez, K.A., Ferrini, V., Franks, P.J., Giulivi, C.F., Gutjahr, M., Harper, D.T., Haynes, L.L., Huber, M., Snell, K.E., Keisling, B.A., Konrad, W., Lowenstein, T.K., Malinverno, A., Guillermic, M., Mejía, L.M., Milligan, J.N., Morton, J.J., Nordt, L., Whiteford, R., Roth-Nebelsick, A., Rugenstein, J.K.C., Schaller, M.F., Sheldon, N.D., Sosdian, S., Wilkes, E.B., Witkowski, C.R., Zhang, Y.G., Anderson, L., Beerling, D.J., Bolton, C., Cerling, T.E., Cotton, J.M., Da, J., Ekart, D.D., Foster, G.L., Greenwood, D.R., Hyland, E.G., Jagniecki, E.A., Jasper, J.P., Kowalczyk, J.B., Kunzmann, L., Kürschner, W.M., Lawrence, C.E., Lear, C.H., Martínez-Botí, M.A., Maxbauer, D.P., Montagna, P., Naafs, B.D.A., Rae, J.W.B., Raitzsch, M., Retallack, G.J., Ring, S.J., Seki, O., Sepúlveda, J., Sinha, A., Tesfamichael, T.F., Tripathi, A., van der Burgh, J., Yu, J., Zachos, J.C., Zhang, L., 2023. Toward a Cenozoic history of atmospheric  $\text{CO}_2$ . *Science* 382(6675), eadi5177. doi.org/doi:10.1126/science.adi5177.

Hublin, J.J., Ben-Ncer, A., Bailey, S.E., Freidline, S.E., Neubauer, S., Skinner, M.M., Bergmann, I., Le Cabec, A., Benazzi, S., Harvati, K. and Gunz, P., 2017. New fossils from Jebel Irhoud, Morocco and the pan-African origin of *Homo sapiens*. *Nature*, 546(7657), pp.289-292.

Huertas, M.J., Arnaud, N.O., Ancochea, E., Cantagrel, J.M. and Fúster, J.M., 2002.  $^{40}\text{Ar}/^{39}\text{Ar}$  stratigraphy of pyroclastic units from the Cañadas Volcanic Edifice (Tenerife, Canary Islands) and their bearing on the structural evolution. *Journal of Volcanology and Geothermal Research*, 115(3-4), pp.351-365.

Hunt, C.O., Gilbertson, D.D., Hill, E.A. and Simpson, D., 2015. Sedimentation, re-sedimentation and chronologies in archaeologically-important caves: problems and prospects. *Journal of Archaeological Science*, 56, pp.109-116.

Iverson, N.A., Kalteyer, D., Dunbar, N.W., Kurbatov, A. and Yates, M., 2017. Advancements and best practices for analysis and correlation of tephra and cryptotephra in ice. *Quaternary Geochronology*, 40, pp.45-55.

Jägerup, S.B., Troll, V.R., Geiger, H., Deegan, F.M., Harris, C., Carracedo, J.C., Meade, F.C., Omidian, S., Zaczek, K. and Van der Zwan, F.M., 2023. Silicic frothy xenoliths (xeno-pumice) in recent volcanics from Gran Canaria, Canary Islands. *Journal of Volcanology and Geothermal Research*, 440, p.107857.

Jeffery, A.J., Gertisser, R., Self, S., Pimentel, A., O'Driscoll, B. and Pacheco, J.M., 2017. Petrogenesis of the peralkaline ignimbrites of Terceira, Azores. *Journal of Petrology*, 58(12), pp.2365-2402.

Jeffery, A.J., Gertisser, R., Self, S., Pimentel, A., O'Driscoll, B. and Pacheco, J.M., 2017. Petrogenesis of the peralkaline ignimbrites of Terceira, Azores. *Journal of Petrology*, 58(12), pp.2365-2402.

Jeffrey, A.: Exploring palaeoaridity using stable oxygen and carbon isotopes in small mammal teeth: a case study from two Late Pleistocene archaeological cave sites in Morocco, North Africa, <https://ora.ox.ac.uk/objects/uuid:5443f540-1049-4f89-8240-970afd5e59f5> (last access: 2 May 2024), 2016.

Jensen, B.J., Davies, L.J., Nolan, C., Pyne-O'Donnell, S., Monteath, A.J., Ponomareva, V., Portnyagin, M., Booth, R., Bursik, M., Cook, E. and Plunkett, G., 2021. A latest Pleistocene and Holocene composite tephrostratigraphic framework for northeastern North America. *Quaternary Science Reviews*, 272, p.107242.

Johnson, C.L., Wijbrans, J.R., Constable, C.G., Gee, J., Staudigel, H., Tauxe, L., Forjaz, V.H. and Salueiro, M., 1998.  $^{40}\text{Ar}/^{39}\text{Ar}$  ages and paleomagnetism of Sao Miguel lavas, Azores. *Earth and Planetary Science Letters*, 160(3-4), pp.637-649.

Johansson, H., Lind, E.M. and Wastegård, S., 2017. Compositions of glass in proximal tephra from eruptions in the Azores archipelago and their links with distal sites in Ireland. *Quaternary Geochronology*, 40, pp.120-128.

Jordan, N.J., Rotolo, S.G., Williams, R., Speranza, F., McIntosh, W.C., Branney, M.J. and Scaillet, S., 2018. Explosive eruptive history of Pantelleria, Italy: Repeated caldera collapse and ignimbrite emplacement at a peralkaline volcano. *Journal of Volcanology and Geothermal Research*, 349, pp.47-73.

Kalnay, E., 1996. The NCEP/NCAR 40-year reanalysis project. *Bulletin of the American Meteorological Society*, 77, pp.437-470.

Karp, A.T., Holman, A.I., Hopper, P., Grice, K., Freeman, K.H., 2020. Fire distinguishers: Refined interpretations of polycyclic aromatic hydrocarbons for paleo-applications. *Geochimica et Cosmochimica Acta* 289, 93-113.

Karp, A.T., Uno, K.T., Polissar, P.J., Freeman, K.H., 2021. Late Miocene C4 Grassland Fire Feedbacks on the Indian Subcontinent. *Paleoceanography and Paleoclimatology* 36(4), e2020PA004106. [doi.org/](https://doi.org/)

Kinder, M., Wulf, S., Appelt, O., Hardiman, M., Żarczyński, M. and Tylmann, W., 2020. Late-Holocene ultra-distal cryptotephra discoveries in varved sediments of Lake Żabińskie, NE Poland. *Journal of Volcanology and Geothermal Research*, 402, p.106988.

Klügel, A., Hoernle, K.A., Schmincke, H.U. and White, J.D., 2000. The chemically zoned 1949 eruption on La Palma (Canary Islands): Petrologic evolution and magma supply dynamics of a rift zone eruption. *Journal of Geophysical Research: Solid Earth*, 105(B3), pp.5997-6016.

Klügel, A., Hansteen, T.H. and Galipp, K., 2005. Magma storage and underplating beneath Cumbre Vieja volcano, La Palma (Canary Islands). *Earth and Planetary Science Letters*, 236(1-2), pp.211-226.

Krüger, S. and van den Bogaard, C., 2021. Small shards and long distances—three cryptotephra layers from the Nahe palaeolake including the first discovery of Laacher See Tephra in Schleswig-Holstein (Germany). *Journal of Quaternary Science*, 36(1), pp.8-19.

Kuhn, S.L., 2023. Trajectories of Culture Change in Northwest Africa and the Levant: Parallels and Contrasts. In *Handbook of Pleistocene Archaeology of Africa: Hominin behavior, geography, and chronology* (pp. 2103-2119). Cham: Springer International Publishing.

Kueppers, U., Pimentel, A., Ellis, B., Forni, F., Neukampf, J., Pacheco, J., Perugini, D. and Queiroz, G., 2019. Biased volcanic hazard assessment due to incomplete eruption records on ocean islands: an example of Sete Cidades Volcano, Azores. *Frontiers in Earth Science*, 7, p.122.

Kwasnitschka, T., Hansteen, T.H., Ramalho, R.S., Devey, C.W., Klügel, A., Samrock, L.K. and Wartho, J.A., 2024. Geomorphology and age constraints of seamounts in the Cabo Verde Archipelago, and their relationship to island ages and geodynamic evolution. *Geochemistry, Geophysics, Geosystems*, 25(3), p.e2023GC011071.

Kylander, M.E., Lind, E.M., Wastegård, S. and Löwemark, L., 2012. Recommendations for using XRF core scanning as a tool in tephrochronology. *The Holocene*, 22(3), pp.371-375.

Laeger, K., Petrelli, M., Morgavi, D., Lustrino, M., Pimentel, A., Paredes-Mariño, J., Astbury, R.L., Kueppers, U., Porreca, M. and Perugini, D., 2019. Pre-eruptive conditions and triggering mechanism of the ~ 16 ka Santa Bárbara explosive eruption of Sete Cidades Volcano (São Miguel, Azores). *Contributions to Mineralogy and Petrology*, 174, pp.1-21.

Lane, C.S., Blockley, S.P., Mangerud, J.A.N., Smith, V.C., Lohne, Ø.S., Tomlinson, E.L., Matthews, I.P. and Lotter, A.F., 2012. Was the 12.1 ka Icelandic Vedde Ash one of a kind?. *Quaternary Science Reviews*, 33, pp.87-99.

Lane, C. S., Brauer, A., Blockley, S. P., & Dulski, P. 2013. Volcanic ash reveals time-transgressive abrupt climate change during the Younger Dryas. *Geology*, 41(12), 1251-1254.

Lane, C.S., Cullen, V.L., White, D., Bramham-Law, C.W.F. and Smith, V.C., 2014. Cryptotephra as a dating and correlation tool in archaeology. *Journal of Archaeological Science*, 42, pp.42-50.

Larrea, P., Galé, C., Ubide, T., Widom, E., Lago, M. and França, Z., 2014a. Magmatic evolution of graciosa (Azores, Portugal). *Journal of Petrology*, 55(11), pp.2125-2154.

Larrea, P., Wijbrans, J.R., Galé, C., Ubide, T., Lago, M., França, Z. and Widom, E., 2014b. 40 Ar/39 Ar constraints on the temporal evolution of Graciosa Island, Azores (Portugal). *Bulletin of Volcanology*, 76, pp.1-15.

Leicher, N., Giaccio, B., Zanchetta, G., Wagner, B., Francke, A., Palladino, D.M., Sulpizio, R., Albert, P.G. and Tomlinson, E.L., 2019. Central Mediterranean explosive volcanism and tephrochronology during the last 630 ka based on the sediment record from Lake Ohrid. *Quaternary Science Reviews*, 226, p.106021.

Lisiecki, L.E. and Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records. *Paleoceanography*, 20(1).

Lowe, D.J., 2011. Tephrochronology and its application: a review. *Quaternary Geochronology*, 6(2), pp.107-153.

Lowe, J., Barton, N., Blockley, S., Ramsey, C.B., Cullen, V.L., Davies, W., Gamble, C., Grant, K., Hardiman, M., Housley, R. and Lane, C.S., 2012. Volcanic ash layers illuminate the resilience of Neanderthals and early modern humans to natural hazards. *Proceedings of the National Academy of Sciences*, 109(34), pp.13532-13537.

Lowe, J.J., Ramsey, C.B., Housley, R.A., Lane, C.S., Tomlinson, E.L., Associates, R.E.S.E.T. and RESET Team, 2015. The RESET project: constructing a European tephra lattice for refined synchronisation of environmental and archaeological events during the last c. 100 ka. *Quaternary Science Reviews*, 118, pp.1-17.

Lucchi, F., Peccerillo, A., Keller, J., Tranne, C.A. and Rossi, P.L. eds., 2013, December. The Aeolian islands volcanoes. Geological Society of London.

Madeira, J., 1998. Estudos de neotectónica nas ilhas do Faial, Pico e S. Jorge: uma contribuição para o conhecimento geodinâmico da junção tripla dos Açores. *Unpublished PhD Thesis*.

Madeira, J., Brum da Silveira, A., 2003. Active tectonics and first paleoseismological results in Faial, Pico and S. Jorge Islands (Azores, Portugal). *Ann. Geophys.* 46, 733–761.

Madeira J, Brum da Silveira A, Hipólito A, Carmo R (2015) Active tectonics in the central and eastern Azores islands along the Eurasia–Nubia boundary: a review, In: Gaspar, J.L., Guest, J.E., Duncan, A.M., Barriga, F.J.A.S., Chester, D.K. (Eds.), *Volcanic Geology of São Miguel Island (Azores Archipelago)*. Geological Society, London, *Memoirs*, vol. 44, pp. 15–32.

Mata, J., Martins, S., Mattielli, N., Madeira, J., Faria, B., Ramalho, R.S., Silva, P., Moreira, M., Caldeira, R., Rodrigues, J. and Martins, L., 2017. The 2014–15 eruption and the short-term geochemical evolution of the Fogo volcano (Cape Verde): Evidence for small-scale mantle heterogeneity. *Lithos*, 288, pp.91-107.

Marean, C.W., 2015. An evolutionary anthropological perspective on modern human origins. *Annual Review of Anthropology*, 44, pp.533-556.

Martí, J., Mitjavila, J. and Araña, V., 1994. Stratigraphy, structure and geochronology of the Las Cañadas caldera (Tenerife, Canary Islands). *Geological Magazine*, 131(6), pp.715-727.

Martí, J. and Wolff, J.A., 2000. Introduction: the geology and geophysics of Tenerife. *Journal of Volcanology and Geothermal Research*, 103(1-4), pp.vii-x.

Martí, J., Sobradelo, R., Felpeto, A. and García, O., 2012. Eruptive scenarios of phonolitic volcanism at Teide–Pico Viejo volcanic complex (Tenerife, Canary Islands). *Bulletin of Volcanology*, 74, pp.767-782.

Mastin, L. G. (2017). Plots of Wind Patterns of the World's Volcanoes [Data set]. U.S. Geological Survey. <https://doi.org/10.5066/F7SQ8XKT>

Mastin, L.G., Van Eaton, A.R. and Lowenstern, J.B., 2014. Modeling ash fall distribution from a Yellowstone supereruption. *Geochemistry, Geophysics, Geosystems*, 15(8), pp.3459-3475.

Maund, J. (1985). The Volcanic Geology, Petrology and Geochemistry of Cal-P. Larrea et al. deira Volcano, Graciosa, Azores, and its bearing on contemporaneous felsic- mafic oceanic island volcanism. PhD Thesis, Univ. of Reading, 333 p.

McBrearty, S. and Brooks, A.S., 2000. The revolution that wasn't: a new interpretation of the origin of modern human behavior. *Journal of human evolution*, 39(5), pp.453-563.

McCanta, M.C., Hatfield, R.G., Thomson, B.J., Hook, S.J. and Fisher, E., 2015. Identifying cryptotephra units using correlated rapid, nondestructive methods: VSWIR spectroscopy, X-ray fluorescence, and magnetic susceptibility. *Geochemistry, Geophysics, Geosystems*, 16(12), pp.4029-4056.

McLean, D., Albert, P.G., Nakagawa, T., Staff, R.A., Suzuki, T. and Smith, V.C., 2016. Identification of the Changbaishan 'Millennium'(B-Tm) eruption deposit in the Lake Suigetsu (SG06) sedimentary archive, Japan: synchronisation of hemispheric-wide palaeoclimate archives. *Quaternary Science Reviews*, 150, pp.301-307.

McLean, D., Albert, P.G., Nakagawa, T., Suzuki, T., Staff, R.A., Yamada, K., Kitaba, I., Haraguchi, T., Kitagawa, J., Members, S.P. and Smith, V.C., 2018. Integrating the Holocene tephrostratigraphy for East Asia using a high-resolution cryptotephra study from Lake Suigetsu (SG14 core), central Japan. *Quaternary Science Reviews*, 183, pp.36-58.

McLean, D., Albert, P.G., Suzuki, T., Nakagawa, T., Kimura, J.I., Chang, Q., Miyabuchi, Y., Manning, C.J., MacLeod, A., Blockley, S.P.E. and Staff, R.A., 2020. Constraints on the timing of explosive volcanism at Aso and Aira calderas (Japan) between 50 and 30 ka: New insights from the Lake Suigetsu sedimentary record (SG14 core). *Geochemistry, Geophysics, Geosystems*, 21(8), p.e2019GC008874.

McLean, D., Albert, P.G., Schlolaut, G., Lamb, H.F., Marshall, M.H., Brauer, A., Wade, J., Nakagawa, T. and Smith, V.C., 2022. How reliable is  $\mu$ XRF core scanning at detecting tephra layers in sedimentary records? A case study using the Lake Suigetsu archive (central Japan). *Journal of Quaternary Science*, 37(7), pp.1189-1206.

Morley, R., Richards, K., 1993. Gramineae cuticle: a key indicator of Late Cenozoic climatic change in the Niger Delta. *Rev. Palaeobot. Palynology* 77(1), 119-127.

Nakagawa, T., Tarasov, P., Staff, R., Ramsey, C.B., Marshall, M., Schlolaut, G., Bryant, C., Brauer, A., Lamb, H., Haraguchi, T. and Gotanda, K., 2021. The spatio-temporal structure of the Lateglacial to early Holocene transition reconstructed from the pollen record of Lake Suigetsu and its precise correlation with other key global archives: Implications for palaeoclimatology and archaeology. *Global and Planetary Change*, 202, p.103493.



Newhall, C.G. and Self, S., 1982. The volcanic explosivity index (VEI) an estimate of explosive magnitude for historical volcanism. *Journal of Geophysical Research: Oceans*, 87(C2), pp.1231-1238.

Óladóttir, B.A., Larsen, G. and Sigmarsson, O., 2011. Holocene volcanic activity at Grímsvötn, Bárðarbunga and Kverkfjöll subglacial centres beneath Vatnajökull, Iceland. *Bulletin of Volcanology*, 73, pp.1187-1208.

Olin, P. H. and Wolff, J. A. Magma dynamics of the phonolitic Diego Hernández Formation, Tenerife, Canary Islands. PhD thesis, (published Ph.D dissertation) (2007).

O'Mara, N.A., Skonieczny, C., McGee, D., Winckler, G., Bory, A.J.M., Bradtmiller, L.I., Malaizé, B. and Polissar, P.J., 2022. Pleistocene drivers of Northwest African hydroclimate and vegetation. *Nature communications*, 13(1), p.3552.

Osborne, A.H., Vance, D., Rohling, E.J., Barton, N., Rogerson, M. and Fello, N., 2008. A humid corridor across the Sahara for the migration of early modern humans out of Africa 120,000 years ago. *Proceedings of the National Academy of Sciences*, 105(43), pp.16444-16447.

Pacheco, J.M.R., 2001. Processos associados ao desenvolvimento de erupções vulcânicas hidromagmáticas explosivas na ilha do Faial e sua interpretação numa perspectiva de avaliação do hazard e minimização do risco (Doctoral dissertation, Universidade dos Acores (Portugal)).

Pearce, N.J., Denton, J.S., Perkins, W.T., Westgate, J.A. and Alloway, B.V., 2007. Correlation and characterisation of individual glass shards from tephra deposits using trace element laser ablation ICP-MS analyses: current status and future potential. *Journal of Quaternary Science*, 22(7), pp.721-736.

Pensa, A., Cas, R., Giordano, G., Porreca, M. and Wallenstein, N., 2015. Transition from steady to unsteady Plinian eruption column: the VEI 5, 4.6 ka Fogo A Plinian eruption, São Miguel, Azores. *Journal of Volcanology and Geothermal Research*, 305, pp.1-18.

Pimentel, A., Pacheco, J.M. and Felpeto, A., 2006. Influence of wind patterns on the dispersal of volcanic plumes in the Azores region: test study of the 1630 eruption of Furnas Volcano (S. Miguel, Azores). In *Geophys. Res. Abstr* (Vol. 8, pp. 1029-7006).

Pimentel, A.H.G., 2016. Pyroclastic density current-forming eruptions on Faial and Terceira Islands, Azores.

Pimentel, A., Pacheco, J. and Self, S., 2015. The ~ 1000-years BP explosive eruption of Caldeira Volcano (Faial, Azores): the first stage of incremental caldera formation. *Bulletin of volcanology*, 77, pp.1-26.

Pimentel, A., Self, S., Pacheco, J.M., Jeffery, A.J. and Gertisser, R., 2021. Eruption style, emplacement dynamics and geometry of peralkaline ignimbrites: Insights from the Lajes-Angra Ignimbrite Formation, Terceira Island, Azores. *Frontiers in Earth Science*, 9, p.673686.

Plesner, S., Holm, P.M. and Wilson, J.R., 2003.  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  geochronology of Santo Antão, Cape Verde Islands. *Journal of Volcanology and Geothermal Research*, 120(1-2), pp.103-121.

Porreca, M., Pimentel, A., Kueppers, U., Izquierdo, T., Pacheco, J., and Queiroz, G. (2018). Event stratigraphy and emplacement mechanisms of the last major caldera eruption on Sete Cidades Volcano (São Miguel, Azores): the 16 ka Santa Bárbara Formation. *Bull. Volcanol.* 80:76.

Potts, R. (2013). Hominin evolution in settings of strong environmental variability. *Quaternary Science Reviews*, 73, 1-13.

Powell, A., Shennan, S. and Thomas, M.G., 2009. Late Pleistocene demography and the appearance of modern human behavior. *Science*, 324(5932), pp.1298-1301.

Pyle, D.M., 1989. The thickness, volume and grainsize of tephra fall deposits. *Bulletin of Volcanology*, 51, pp.1-15.

Queiroz, G. (1997). *Vulcão das Sete Cidades (S.Miguel, Açores): História Eruptiva e Avaliação do Hazard*. Ph.D. Thesis, Azores University, Portugal.

Queiroz, G., Pacheco, J.M., Gaspar, J.L., Aspinall, W.P., Guest, J.E., Ferreira, T. (2008). The last 5000 years of activity at Sete Cidades volcano (São Miguel Island, Azores): implications for hazard assessment. *J. Volcanol. Geotherm. Res.* 178, 562–573.

Queiroz, G., Gaspar, J.L., Guest, J.E., Gomes, A. and Almeida, M.H., 2015. Chapter 7 Eruptive history and evolution of Sete Cidades Volcano, São Miguel Island, Azores. *Geological Society, London, Memoirs*, 44(1), pp.87-104.

Ramalho, R.A. and Ramalho, R.A.D.S., 2011. Tracers of uplift and subsidence in the Cape Verde Archipelago. *Building the Cape Verde Islands*, pp.79-138.

Ramalho, R.S., Helffrich, G., Cosca, M., Vance, D., Hoffmann, D. and Schmidt, D.N., 2010a. Vertical movements of ocean island volcanoes: Insights from a stationary plate environment. *Marine Geology*, 275(1-4), pp.84-95

Ramalho, R., Helffrich, G., Cosca, M., Vance, D., Hoffmann, D. and Schmidt, D.N., 2010b. Episodic swell growth inferred from variable uplift of the Cape Verde hotspot islands. *Nature Geoscience*, 3(11), pp.774-777.

Ramsey, C.B., 1995. Radiocarbon calibration and analysis of stratigraphy: the OxCal program. *Radiocarbon*, 37(2), pp.425-430.

Ramsey, C.B., 2008. Deposition models for chronological records. *Quaternary Science Reviews*, 27(1-2), pp.42-60.

Ramsey, C.B., Dee, M., Lee, S., Nakagawa, T. and Staff, R.A., 2010. Developments in the calibration and modeling of radiocarbon dates. *Radiocarbon*, 52(3), pp.953-961.

Reimer, P.J., Austin, W.E., Bard, E., Bayliss, A., Blackwell, P.G., Ramsey, C.B., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M. and Grootes, P.M., 2020. The IntCal20 Northern Hemisphere radiocarbon age calibration curve (0–55 cal kBP). *Radiocarbon*, 62(4), pp.725-757.

Rhoujjati, A., Cheddadi, R., Taieb, M., Baali, A. and Ortu, E., 2010. Environmental changes over the past c. 29,000 years in the Middle Atlas (Morocco): a record from Lake Ifrah. *Journal of arid environments*, 74(7), pp.737-745.

Romero, J.E., Burton, M., Cáceres, F., Taddeucci, J., Civico, R., Ricci, T., Pankhurst, M.J., Hernández, P.A., Bonadonna, C., Llewellyn, E.W. and Pistolesi, M., 2022. The initial phase of the 2021 Cumbre Vieja ridge eruption (Canary Islands): Products and dynamics controlling edifice growth and collapse. *Journal of Volcanology and Geothermal Research*, 431, p.107642.

Sachse, D., Billault, I., Bowen, G.J., Chikaraishi, Y., Dawson, T.E., Feakins, S.J., Freeman, K.H., Magill, C.R., McInerney, F.A., Van der Meer, M.T., Polissar, P.J., Robins, R.J., Sachs, J.P., Schmidt, H.-L., Sessions, A.L., White, J.W.C., West, J.B., Kahmen, A., 2012. Molecular paleohydrology: interpreting the hydrogen-isotopic composition of lipid biomarkers from photosynthesizing organisms. *Annual Review of Earth and Planetary Sciences* 40, 221-249.

Scerri, E.M., Thomas, M.G., Manica, A., Gunz, P., Stock, J.T., Stringer, C., Grove, M., Groucutt, H.S., Timmermann, A., Rightmire, G.P. and d'Errico, F., 2018. Did our species evolve in subdivided populations across Africa, and why does it matter?. *Trends in ecology & evolution*, 33(8), pp.582-594.

Schouten, S., Hopmans, E.C., Schefuß, E., Damste, J.S.S., 2002. Distributional variations in marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water temperatures? *Earth Planet. Sci. Lett.* 204(1-2), 265-274.

Scott, L. and Neumann, F.H., 2018. Pollen-interpreted palaeoenvironments associated with the Middle and Late Pleistocene peopling of Southern Africa. *Quaternary International*, 495, pp.169-184.

Schmincke, H.U., 1979. Age and crustal structure of the Canary Islands. *Journal of Geophysics*, 46(1), pp.217-224.

Schmincke, H.U. and Sumita, M., 2010. Geological evolution of the Canary Islands: a young volcanic archipelago adjacent to the old African Continent. *Görres*.

Schwaiger, H.F., Denlinger, R.P. and Mastin, L.G., 2012. Ash3d: A finite-volume, conservative numerical model for ash transport and tephra deposition. *Journal of Geophysical Research: Solid Earth*, 117(B4).

Sehassseh, E.M., Fernandez, P., Kuhn, S., Stiner, M., Mentzer, S., Colarossi, D., Clark, A., Lanoe, F., Pailes, M., Hoffmann, D. and Benson, A., 2021. Early middle stone age personal ornaments from Bizmoune Cave, Essaouira, Morocco. *Science Advances*, 7(39), p.eabi8620.

Self S (1974) Recent volcanism on Terceira, Azores (unpublished). PhD thesis, Imperial College, London

Self, S. (1976). The Recent Volcanology of Terceira, Azores. *J. Geol. Soc.* 132, 645–666.

Shanahan, T.M., McKay, N.P., Hughen, K.A., Overpeck, J.T., Otto-Bliesner, B., Heil, C.W., King, J., Scholz, C.A. and Peck, J., 2015. The time-transgressive termination of the African Humid Period. *Nature Geoscience*, 8(2), pp.140-144.

Skonieczny, C., McGee, D., Winckler, G., Bory, A., Bradtmiller, L., Kinsley, C., Polissar, P., De Pol-Holz, R., Rossignol, L., Malaizé, B., 2019. Monsoon-driven Saharan dust variability over the past 240,000 years. *Science Advances* 5(1), eaav1887.

Smith, V.C., Staff, R.A., Blockley, S.P., Ramsey, C.B., Nakagawa, T., Mark, D.F., Takemura, K. and Danhara, T., 2013. Identification and correlation of visible tephra in the Lake Suigetsu SG06 sedimentary archive, Japan: chronostratigraphic markers for synchronising of east Asian/west Pacific palaeoclimatic records across the last 150 ka. *Quaternary Science Reviews*, 67, pp.121-137.

Staff, R.A., Scholaut, G., Ramsey, C.B., Brock, F., Bryant, C.L., Kitagawa, H., Van der Plicht, J., Marshall, M.H., Brauer, A., Lamb, H.F. and Payne, R.L., 2013. Integration of the old and new Lake Suigetsu (Japan) terrestrial radiocarbon calibration data sets. *Radiocarbon*, 55(4), pp.2049-2058.

Steele, T.E., Álvarez Fernández, E. and Hallett, E., 2019. Early personal ornaments: A review of shells as personal ornamentation during the African Middle Stone Age. *PaleoAnthropology*, (2019), pp.24-51.

Stevenson, J.A., Millington, S.C., Beckett, F.M., Swindles, G.T. and Thordarson, T., 2015. Big grains go far: reconciling tephrochronology with atmospheric measurements of volcanic ash. *Atmospheric Measurement Techniques Discussions*, 8(1).

Stoetzel, E., Marion, L., Nespoulet, R., El Hajraoui, M.A. and Denys, C., 2011. Taphonomy and palaeoecology of the late Pleistocene to middle Holocene small mammal succession of El Harhoura 2 cave (Rabat-Témara, Morocco). *Journal of Human Evolution*, 60(1), pp.1-33.

Stoetzel, E., Lalis, A., Nicolas, V., Aulagnier, S., Benazzou, T., Dauphin, Y., El Hajraoui, M.A., El Hassani, A., Fahd, S., Fekhaoui, M. and Geigl, E.M., 2019. Quaternary terrestrial microvertebrates from mediterranean northwestern Africa: State-of-the-art focused on recent multidisciplinary studies. *Quaternary Science Reviews*, 224, p.105966.

Storch, B., Haase, K.M., Romer, R.H.W., Beier, C., Koppers, A.A.P., 2020. Rifting of the oceanic Azores Plateau with episodic volcanic activity. *Sci. Rep.* 10, 19718.

Stroncik, N.A., Klügel, A. and Hansteen, T.H., 2009. The magmatic plumbing system beneath El Hierro (Canary Islands): constraints from phenocrysts and naturally quenched basaltic glasses in submarine rocks. *Contributions to Mineralogy and Petrology*, 157, pp.593-607.

Tomlinson, E.L., Smith, V.C., Albert, P.G., Aydar, E., Civetta, L., Cioni, R., Çubukçu, E., Gertisser, R., Isaia, R., Menzies, M.A. and Orsi, G., 2015. The major and trace element glass compositions of the productive Mediterranean volcanic sources: tools for correlating distal tephra layers in and around Europe. *Quaternary Science Reviews*, 118, pp.48-66.

Thórarinnsson, S. 1944. Tefrokronologiska studier på Island: fíjósárdalur och dess förödelse. *Geografiska annaler*, 26: 1-217

Terray, L., Stoetzel, E., Ben Arous, E., Kageyama, M., Cornette, R. and Braconnot, P., 2023. Refinement of the environmental and chronological context of the archeological site El Harhoura 2 (Rabat, Morocco) using paleoclimatic simulations. *Climate of the Past*, 19(6), pp.1245-1263.

Tierney, J.E., Pausata, F.S.R., deMenocal, P.B., 2017. Rainfall regimes of the Green Sahara. *Science Advances* 3(1). doi.org/10.1126/sciadv.1601503.

Tomlinson, E.L., Thordarson, T., Müller, W., Thirlwall, M. and Menzies, M.A., 2010. Microanalysis of tephra by LA-ICP-MS—Strategies, advantages and limitations assessed using the Thorsmörk ignimbrite (Southern Iceland). *Chemical Geology*, 279(3-4), pp.73-89.

Torres, P., Silva, L., Serralheiro, A., Tassinari, C., Munhá, L., 2002. Enquadramento geocronológico pelo método K/Ar das principais sequências vulcano-estratigráficas da Ilha do Sal — Cabo Verde. *Garcia de Orta, Serviços Geológicos* 18, 9–13.

Tripanera, D., Porreca, M., Ruch, J., Pimentel, A., Acocella, V., Pacheco, J., Salvatore, M., 2014. Relationships between tectonics and magmatism in a transtensive/transform setting: an example from Faial Island (Azores, Portugal). *Geol. Soc. Am. Bull.* 126, 164–181

Troll, V.R. and Carracedo, J.C., 2016. *The geology of the Canary Islands*. Elsevier.

Turney, C.S., 1998. Extraction of rhyolitic component of Vedde microtephra from minerogenic lake sediments. *Journal of Paleolimnology*, 19: 199-206

Turney, C.S.M., Den Burg, K.V., Wastegård, S., Davies, S.M., Whitehouse, N.J., Pilcher, J.R. and Callaghan, C., 2006. North European last glacial–interglacial transition (LGIT; 15–9 ka) tephrochronology: extended limits and new events. *Journal of Quaternary Science: Published for the Quaternary Research Association*, 21(4), pp.335-345.

Uno, K.T., Polissar, P.J., Jackson, K.E., deMenocal, P.B., 2016. Neogene biomarker record of vegetation change in eastern Africa. *Proc Natl Acad Sci U S A* 113(23), 6355-6363. doi.org/10.1073/pnas.1521267113.

van der Bilt, W.G. and Lane, C.S., 2019. Lake sediments with Azorean tephra reveal ice-free conditions on coastal northwest Spitsbergen during the Last Glacial Maximum. *Science Advances*, 5(10), p.eaaw5980.

van der Bilt, W.G., Cederstrøm, J.M., Støren, E.W., Berben, S.M. and Rutledal, S., 2021. Rapid tephra identification in geological archives with computed tomography: experimental results and natural applications. *Frontiers in Earth Science*, 8, p.622386.

Vanhaeren, M., d'Errico, F., Stringer, C., James, S.L., Todd, J.A. and Mienis, H.K., 2006. Middle Paleolithic shell beads in Israel and Algeria. *Science*, 312(5781), pp.1785-1788.

Wallenstein, N. (1999). Estudo da História Recente e do Comportamento Eruptivo do Vulcão do fogo (s. Miguel, açores). Avaliação Preliminar do Hazard. Ph.D. thesis, Azores University, Portugal.

Wallenstein, N., Chester, D., Coutinho, R., Duncan, A. and Dibben, C., 2015. Chapter 16 Volcanic hazard vulnerability on São Miguel Island, Azores. *Geological Society, London, Memoirs*, 44(1), pp.213-225.

Walsh, A.A., Blockley, S.P., Milner, A.M., Matthews, I.P. and Martin-Puertas, C., 2021. Complexities in European Holocene cryptotephra dispersal revealed in the annually laminated lake record of Diss Mere, East Anglia. *Quaternary Geochronology*, 66, p.101213.

Wastegård, S., Rasmussen, T. L., Kuijpers, A., Nielsen, T. and van Weering, T. C. 2006. Composition and origin of ash zones from Marine Isotope Stages 3 and 2 in the North Atlantic. *Quaternary Science Reviews*, 25: 2409-2419

Wastegård, S., Johansson, H. and Pacheco, J.M., 2020. New major element analyses of proximal tephra from the Azores and suggested correlations with cryptotephra in North-West Europe. *Journal of Quaternary Science*, 35(1-2), pp.114-121.

Watson, E.J., Swindles, G.T., Lawson, I.T., Savov, I.P. and Wastegård, S., 2017. The presence of Holocene cryptotephra in Wales and southern England. *Journal of Quaternary Science*, 32(4), pp.493-500.

Wolff, J. A., J. S. Grandy, and P. B. Larson. "Interaction of mantle-derived magma with island crust? Trace element and oxygen isotope data from the Diego Hernandez Formation, Las Cañadas, Tenerife." *Journal of Volcanology and Geothermal Research* 103.1-4 (2000): 343-366.

Wolff, J.A., Forni, F., Ellis, B.S. and Szymanowski, D., 2020. Europium and barium enrichments in compositionally zoned felsic tuffs: A smoking gun for the origin of chemical and physical gradients by cumulate melting. *Earth and Planetary Science Letters*, 540, p.116251.

Wood, R. (2015). From revolution to convention: the past, present and future of radiocarbon dating. *Journal of Archaeological Science*, 56, 61-72.

Wulf, S., Kraml, M., Brauer, A., Keller, J. and Negendank, J.F., 2004. Tephrochronology of the 100 ka lacustrine sediment record of Lago Grande di Monticchio (southern Italy). *Quaternary International*, 122(1), pp.7-30.

Zanon, V. and Frezzotti, M.L., 2013. Magma storage and ascent conditions beneath Pico and Faial islands (Azores archipelago): A study on fluid inclusions. *Geochemistry, Geophysics, Geosystems*, 14(9), pp.3494-3514.