

The concept of critical age group for density dependence: bridging the gap between demographers, evolutionary biologists and behavioral ecologists

Marlène Gamelon, Yimen G Araya-Ajoy, Bernt-Erik Sæther

To cite this version:

Marlène Gamelon, Yimen G Araya-Ajoy, Bernt-Erik Sæther. The concept of critical age group for density dependence: bridging the gap between demographers, evolutionary biologists and behavioral ecologists. Philosophical Transactions of the Royal Society B: Biological Sciences, 2024, 379 (1916), pp.20220457. 10.1098/rstb.2022.0457. hal-04755689

HAL Id: hal-04755689 <https://hal.science/hal-04755689v1>

Submitted on 28 Oct 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Abstract

 Density dependence plays an important role for population regulation in the wild. It involves a decrease in population growth rate when the population size increases. Fifty years ago, Charlesworth introduced the concept of "critical age group", denoting the age classes in which variation in the number of individuals most strongly contribute to density regulation. Since this pioneering work, the use of this concept has remained scarce. In light of Charlesworth's concept, we discuss the need to develop work between behavioral ecology, demography and evolutionary biology to better understand the mechanisms acting in density- regulated age-structured populations. We highlight demographic studies that explored age- specific contributions to density dependence and discuss the underlying evolutionary processes. Understanding competitive interactions among individuals is pivotal to identify the ages contributing most strongly to density regulation, highlighting the need to move towards behavioral ecology to decipher mechanisms acting in density-regulated age-structured populations. Because individual characteristics other than age can be linked to competitive abilities, expanding the concept of critical age to other structures (e.g. sex, dominance rank) offers interesting perspectives. Linking research fields based on the concept of the critical age group is key to move from a pattern-oriented view of density regulation to a process-oriented approach.

 Keywords: age-structured populations, density regulation, dominance rank, competition, personality, senescence, sociality, stage-structured populations, viability selection

1. Introduction

 Understanding fluctuations in population size over time is a major goal in both conservation and evolutionary ecology. Density-independent and density-dependent factors both shape fluctuations in population size over time [1]. Their relative importance has long been debated [2–5], but empirical evidence from a wide variety of taxa shows that both processes contribute to fluctuations in population size [6–8].

 Density dependence generates fluctuations in population size around a mean value, the carrying capacity [9], and thus plays an important role in population regulation. From a demographic viewpoint, negative density dependence (hereafter called density dependence) describes the negative relationship between a population's growth rate and the number of individuals currently present or at an earlier time step [4,10–13]. This decrease in population growth rate when the population increases in numbers may be explained by several factors. For instance, at high population sizes, contests for the access to critical resources or space, or higher risk of disease transmission [14] can negatively influence reproduction and survival [15] with cascading effects on the growth rate of the whole population.

 Assessing density dependence by exploring the strength of the negative relationship between population growth rate and population size relies on several assumptions, one of them being that all individuals are expected to have the same competitive effects on other individuals [16]. However, in age-structured populations, this assumption may be violated. Individuals may differently contribute to the density regulation according to their age. In the 1970's, Charlesworth introduced the concept of "critical age group" in a model with density- dependent selection. In his seminal works in 1972 and 1973 [17,18], he wrote: "*In any particular case, this density regulation must occur in response to the number of individuals in a specific group of ages, which I shall call the "critical age group". The critical age group might, for example, be composed of individuals in the first age class only, or of all individuals*

 of reproductive age, or of any other combination of age classes, depending on the exact mode of density regulation in the population under consideration. Mortality at all or some ages will be increased and/or fecundity decreased in relation to the number of individuals existing in the critical age group, either at the present time, or at a specific time in the past, or over a whole range of past times."

 This pioneering work on density-dependent selection was primarily rooted in evolutionary studies of natural selection in populations with overlapping generations. Beyond density-dependent selection, the definition of "critical age group" given in Charlesworth's studies ["*this density regulation must occur in response to the number of individuals in a specific group of ages"*] can be more broadly interpreted in the context of density-dependent regulation. Until recently, the use of this important concept in demographic studies has remained scarce [19,20], which is surprising as it can shed light on the underlying mechanisms acting on density-regulated age-structured populations. Similarly, this concept has to our knowledge not been explored by behavioral ecologists. This field has a long history of studying biotic interactions and can therefore provide key insights on the behavioral processes determining the critical age group. Linking evolution, demography and behavior based on the concept of the critical age group is particularly relevant to move from pattern- oriented view of density regulation to a process-oriented approach in age-structured populations (Figure 1).

 In the first part of this paper, we thus discuss some of the empirical demographic studies that explored age-specific contributions to density dependence in light of Charlesworth's concept of the critical age group. We discuss the implications for accurate predictions of fluctuations in population size over time. In the second part, we discuss how the critical age group has been included in evolutionary biology and describe the processes that may affect the evolutionary consequences of variation in the critical age class. In the third

 part, we discuss some of the behavioral mechanisms that could explain age-specific contributions to density dependence, including age-related changes in personality traits and dominance rank. Improved understanding of competitive interactions among individuals can be useful in identifying which age or group of ages will contribute the most strongly to the strength of density dependence. Finally, in the last part of the paper, we discuss the need to improve the interactions between behavioral ecology, population dynamics, demography and, ultimately evolutionary biology to address the concept of critical age group in a more comprehensive way across research fields. We conclude by discussing some new exciting research perspectives.

-
-

2. The concept of critical age group for density dependence in demographic studies

 In age-structured populations, individuals may have different contributions ("weights") to density regulation, depending on their age. In other words, adding one individual of age *i* in a population is not necessarily equivalent to adding one individual of age *j* in terms of negative effects on the vital rates of other individuals in the population. Surprisingly, age-specific contributions to density dependence have been rarely investigated in demographic studies [11,21,22]. Gamelon et al. [19] considered age-specific population 104 sizes N_i instead of the total population size N_{tot} to explore the influence of density dependence on the dynamics of a great tit (*Parus major*) population. Partitioning the contribution of *Ntot* 106 into the age-specific contributions N_i allows identifying the age(s) contributing the most to density regulation by estimating the "per capita" effects on density dependence. In other words, partitioning the contribution of *Ntot* allows answering the question: "what is the density-dependent effect of an individual of a certain age on the vital rates of other individuals?". With such an approach focusing on the per capita effects, the critical age group is not necessarily the largest age class. They found that the number of females in their first

 year of breeding had the strongest negative effects on survival and recruitment of all ages, and concluded that the youngest females were the critical age class (sensu Charlesworth).

 More recently, Lande et al. [20] expanded this approach and provided a general framework to quantify age-specific contributions to density dependence in (st)age-structured populations. In this study, density dependence in vital rates is exerted by a function called *g(N)*, of a weighted sum of stage abundances, *N*. In accordance with the previous study on great tits, they found that in this great tit population, the estimated relative weights of stage abundances decrease strongly for the older stages. Therefore, 1-year olds had the strongest impact on density regulation.

 Strikingly, some demographic studies have assessed the influence of the number of individuals in a specific group of age on density regulation acting on individuals' vital rates (survival, reproduction) or on the population growth rate, but did not explicitly use the term "critical age group". For instance, in their empirical study on Alpine plants, Cotto et al. [23] developed an eco-evolutionary model in which the survival probability of seedlings depends specifically on the number of adult individuals previously established. Similarly, Schmid et al. [24] simulated contrasting life history strategies, and assumed a decline of juvenile survival due to intraspecific competition, with increasing adult number for overlapping generations, and with juvenile number for non-overlapping generations. These examples highlight that despite not always been mentioned explicitly, the concept of critical age group for density dependence is growing in use in demographic studies. These density-dependence models imply that a group of age instead of all ages drives density regulation, and are thus more biologically realistic, depending strongly on the biology of the species.

 Identifying the critical age group is key to unravel the demographic mechanisms leading to density regulation at the population level, but it is also crucial for accurately predicting fluctuations in population size over time. Gamelon et al. [25] identified the age(s)

 contributing the most to density regulation in 12 great tit populations and in 12 blue tit (*Cyanistes caeruleus*) populations in Europe. They found that a density-dependence model with age-specific contributions to density dependence significantly improves predictions of fluctuations in population size, compared to the "classical" density-dependence model including only the total population size *Ntot*. Accordingly, including age in demographic analyses of density dependence seems particularly relevant in a management or in a conservation context to identify the age groups limiting the most the population growth rate and make accurate predictions of fluctuations in population size.

 While a few demographic studies have started to apply the concept of critical age group, its roots reside in evolutionary biology [17]. Therefore, unsurprisingly, this is in this research field that the term "critical age group" has been used more frequently (about fifty times) compared to demographic and behavioral studies. Putting more broadly the concept of critical age group back into the evolutionary theatre will improve our understanding of the evolutionary causes and consequences of the critical age class.

3. The concept of critical age group for density dependence in evolutionary biology

 Evolutionary studies that coin the term "critical age group" have primary focused on density-dependent selection, not on density-dependent population regulation [18,26,27]. However, different evolutionary processes may result in age classes differing in their competitive ability and therefore in their relative contribution to density regulation (Figure 2). Age-specific contributions to density regulation may be underpinned by phenotypes with different competitive abilities either appearing in or disappearing from the population in a particular sequence. For instance, viability selection can cause changes in the phenotypic composition of different ages. This has been extensively studied in the context of selective disappearance, a form of viability selection whereby specific phenotypes are selected for,

 creating a change in phenotypic composition across ages [28]. Selective disappearance of a given phenotype can result in older classes being composed of individuals with a larger or a lower impact on density regulation. For instance, figure 2A depicts a hypothetical example in which phenotype A has high competitive ability and high survival, whereas phenotype D has opposite characteristics (but possibly high reproductive output). Viability selection can therefore cause phenotypes D (frail) to be less common in older age classes, leading the average competitive ability of the different age classes to be higher in the older age classes, and the variance in competitive ability to be lower for the older age classes (Figure 2B). In this hypothetical scenario, the older age classes have a stronger per-capita effect on population 171 growth rate (Figure 2C). The same reasoning applies when a phenotype has high competitive ability, enabling it to better acquire resources for reproduction at the expense of low survival. In this scenario, viability selection will cause phenotypes with lower competitive abilities to be more common in older age classes, the youngest age classes thus having a stronger per-capita effect on the population growth rate (Figure 2D, E, F).

 Age-related changes in competitive ability and therefore age-specific variation in the contribution to density regulation can be also caused by senescence. This is a within- individual decline with age in residual reproductive value owing to deteriorating survival probability and reproductive performance, caused by a progressive loss of physiological and cellular function late in life [29,30]. Evidence for within-individual phenotypic deterioration is accumulating for various traits across a range of animal taxa [31]. In the hypothetical scenario where senescence causes a similar decline in competitive ability with age for all individuals (depicted in different colors on Figure 2G), i.e. same senescence rates, the average competitiveness decreases with age (Figure 2H), leading to a lower per-capita effect on the population growth rate for the older age classes (Figure 2I). Note that different phenotypes can have contrasting senescence patterns (e.g. different slopes, shapes, onsets) possibly

 resulting in complex relationships between competitive ability and age. The mounting evidence of a decline in trait values that may reflect competitive ability with age suggests that senescence can shape the contribution to density regulation of the different age classes.

 Both senescence and selective disappearance through viability selection can happen at the same time: competitive ability can decline within an individual's lifespan, and individuals with specific competitive abilities may disappear from the older age classes. This can result in complex non-linear relationships between the average competitive ability and age (Figures 2H, K), as well as with the per-capita effect on the growth rate and age (Figure 2I, L). Accordingly, population-level patterns do not necessarily provide insights into within- individual processes such as senescence [32], or age-specific changes in the phenotypic composition of a population, per se. For instance, selective disappearance of poor-quality individuals has been found to partly mask age-specific declines in reproductive performance measures in roe deer *Capreolus capreolus* [28] and mute swans *Cygnus olor* [33]. It is thus of key importance to separate these two processes in order to understand how evolution may affect the critical age class and its phenotypic characteristics.

 Concomitantly to Charlesworth's concept of critical age group for density dependence, the idea that several phenotypes with different competitive strategies co-exist and interact was introduced in the 70's. Indeed, in their seminal article [34], published at the early stages of the evolutionary game theory, Maynard Smith and Price introduced the concept of evolutionary stable strategy (ESS) that became an important tool for studying animal contests [35] in both evolutionary and behavioral studies. However, to the best of our knowledge, "conflicts" among individuals within the evolutionary game theory framework have not been studied in light of age-specific abilities to compete and Charlesworth's concept of critical ages.

 Moving from population to individual level may help understand why some ages are more "critical" than others for the density regulation in age-structured populations. Important

 knowledge can be gained from behavioral studies, that have long been interested in biotic interactions in natural populations.

4. Insights from behavioral studies on the critical age group

 To our knowledge, the term "critical age group" has not been used yet in behavioral studies to describe the age class contributing the most to density regulation. This is surprising as behavioral ecologists have extensively explored among-individual differences in competitive abilities in animals to get access to food resources, territory, mating partners, etc. [36,37] and the consequences on their vital rates [38] and the population growth rate. Competitive abilities can be related to personality traits (or temperament, [39]), like aggressiveness, exploration, problem-solving performance, etc. [40]. These traits can be under selection [41], and may have strong influence on density regulation and thus on population dynamics [42,43].

 Personality traits related to competitive abilities can be age-specific, changing over an individual lifespan due to senescence [44], as mentioned earlier (Figure 2G). In a study on blue tits, Class and Brommer [46] explored age-related changes in two traits, aggression and breath rate within individuals' lifespan, and found evidence for a decline in aggression with increasing ages. This observation of a decrease in performance of an individual with age may indicate senescence. This means that some traits associated with competitive capacity could decline with age, possibly weakening the contribution of older ages to density dependence. If traits associated with competitive abilities may decline over an individual lifespan, they can, in contrast, improve as individuals become older through social learning for instance $[40,47,48]$ + Hasenjager & Fefferman, this issue. In that case, expectations are opposite to Figures 2G, H, I, with a positive relationship between competitive ability and age resulting in an increase in per-capita effect on population growth rate with age. For instance, in beef cattle

 (*Bos taurus*), social dominance and thus the access to critical resources is mostly explained by age, that prevails over body mass in the structuring of the dominance network [49]. Similarly, in zebra (*Equus burchelli*), the social rank of mares increases with age, and mares with higher social rank appear to be the ones that have access to resources and have higher reproductive success [50]. Thus, older individuals may have the highest contribution to density regulation in this specific case. Note however that some personality traits related to competitive abilities can remain stable within life stages (see [51] for a review). This is the case for the yellow- bellied marmots (*Marmota flaviventris*), for which docility is established early in life and is a good predictor of docility at adulthood [52]. Similarly, in horses (*Equus caballus*), the social rank of an individual throughout its life is determined by the social rank of its mother [53]. Therefore, age-related changes in personality traits associated with competitive abilities can strongly differ among species. These species-specific patterns may provide a crucial explanation for the role played by different critical age classes in their population dynamics. Another important factor that could explain age-specific contributions to density dependence is social organization. Coulson and Godfray [15] contrasted two main types of competition . In the *contest* type, a few individuals are able to monopolize critical resources or space and exclude other individuals. In contrast, in the *scramble* competition type, resources are equally divided among the individuals in the population. In practice, most social systems are classified along a gradient between these two extremes. In their comparative study of altricial birds mostly regulated by territorial behavior, Sæther et al. [54] showed that in long- lived species in which adults may breed in the same territory for several years, the number of new recruits was negatively related to the return rate of adults from the previous breeding season. This indicates a contest type of social organization, with consequences for the definition of critical age group.

 The patterns of space use of species can also shed light on which ages are more likely to contribute to density dependence. Spatial assortment by age can lead to age-based social groups or hierarchies. Within these groups, individuals of similar ages may interact more frequently. For instance, assortative mating by age is commonly documented and one of the explanations is that individuals with similar ages are more likely to be in the same location [55]. Alternatively, older individuals may interact more with younger individuals, for instance individuals without a territory are generally younger and are constantly interacting with older territory owners to gain access to their territory [56]. In some vertebrate species, individuals use different habitat types sequentially throughout their life span to maximize fitness. This is known as ontogenetic niche shift in habitat use [57,58]. This could also result in non-random spatial distribution with age that can potentially affect the critical age class. More generally, patterns of non-random association of different age groups can result in both stronger or weaker density-dependent effects on population growth as compared to scenarios where interactions are at random.

5. Concluding remarks and perspectives

 Since the 1930's and Nicholson's pioneering work [59], density dependence has been intensively studied both theoretically and empirically in several fields of research, including demography, behavioral ecology and evolutionary biology. In demography, this is an important mechanism contributing to population size regulation. In behavioral ecology, this is a key aspect of biotic interaction, involving competition and cooperation (in case of positive density dependence)(see Woodman et al., this issue). In evolutionary biology, this is often studied in the context of density-dependent selection shaping phenotypic traits in natural populations [60–66] and affecting the rate of genetic drift [67]. As discussed earlier, there are many ecological situations that can lead to age-specific contributions to density dependence.

 Despite important advances and the seminal work of Charlesworth [17,18], it still remains an open question: which are the ages contributing the most to density regulation. Whether the critical age groups depend on the age-specific responses to density dependence (Box 1), the species life history strategy, climate conditions (Box 2), or other factors remain to be carefully explored. Addressing these questions is critical to identify general principles of how density regulation affects the eco-evolutionary dynamics of age-structured populations.

 Some methodological challenges can explain the lack of studies on this topic: quantifying the age-specific contributions to density dependence requires knowing the number of individuals in each age class [21], which is notoriously difficult as both determining individuals' age and counting individuals in the wild is a real challenge. Some new methodological tools such as integrated population models IPM [68,69] now allow accurately estimating age-specific population sizes from demographic data, while accounting for imperfect detection. Once age-specific population sizes are estimated, another methodological challenge is to quantify the age-specific contributions to density regulation. 300 Indeed, partitioning the contribution of N_{tot} into the age-specific contributions N_i increases the degrees of freedom and thus requires more data. Again, IPMs can be a suitable framework.

 After having identified "which are" the critical age classes, the question then becomes: why do some ages contribute more than others to density regulation? To interpret and understand the role played by the critical age group for density dependence, behavioral studies bring important insights. We argue that there is a clear need to fill the gap between demography, behavioral ecology and evolutionary biology. Such a transdisciplinary framework would deepen our understanding of the role played by age in biotic interactions (processes) and their consequences on population dynamics (patterns) (Figure 1).

 The need to bring together demography, behavioral ecology and evolution to explore the concept of age group for density dependence is even clearer when going back to

 Charlesworth's work [17]: "*the critical age group might, for example, be composed of all individuals of reproductive age, or of newborn individuals, depending on the biology of the population"*. Thus, the critical group is not necessarily an age class but more broadly a group of individuals, depending on the biology of the population. Age is obviously one key structuring factor in natural populations, possibly associated with contrasting abilities to compete. Furthermore, other individual characteristics than age can be closely linked to competitive abilities, as previously demonstrated by behavioral ecologists: body mass, sex, etc. For instance, in the common lizard (*Lacerta vivipara*), the number of males specifically reduces female's survival and reproduction, due to female's aggression [70]. Therefore, density dependence in that specific example seems primarily driven by the number of reproductive males. To what extent the pioneering work of Charlesworth [17,18] can be expanded to two-sex age-structured populations, or even generalized to other type of traits (behavioral, morphological, etc.) offers exciting research perspectives. Thus, instead of focusing on the female segment of the population as classically done in demographic studies, one should include age- and sex-structured models to assess density regulation acting on the population growth rate.

 While Charlesworth's work [17,18] preliminary focused on intraspecific density dependence, individuals do not live in isolation but interact with other species within a community. Since the 1920's and the demographic models of competition between two species of Lotka [71] and Volterra [72], it is well accepted that the density of species *A* may influence the population growth rate of species *B*, and vice versa. In behavioral ecology too, there is a large amount of literature including field and experimental studies, providing evidence for competition between species (see e.g. [73,74]). Expanding the concept of critical age group for density dependence to interspecific density dependence would shed light on density regulation acting in age-structured populations within a community. In that case, the

 number of individuals in a given age class for species *A* would influence not only the demographic performances of individuals of certain ages of species *A*, but also of species *B*, and vice versa. Gamelon et al. [25] developed such a demographic model and applied it on two competing species, the great tit and the blue tit living in sympatry. They considered that the number of great tits in four different age classes may influence the demographic performances of great tit and also the vital rates of blue tit, and vice versa. They found that blue tits (this species is smaller than great tits) were strongly influenced by the number of great tits, whereas great tit populations were mostly regulated by intraspecific density dependence. However, great and blue tits live in a community with other potential competitors, and expanding the approach to more than two competing species is a natural next step [75]. Here as well, knowledge gained by behavioral ecologists will help to understand biotic interactions among multiple species within a community, and help building realistic demographic models to characterize the critical age group for density dependence.

Box 1: What about age-specific response to density dependence?

 Until now, age-specific responses to density (i.e. who is more strongly affected by an increase in density in its vital rates?) have received more interests in both theoretical and empirical studies than age-specific contributions to density regulation (i.e. who is affecting the most the vital rates of others?). In the first case (i.e. who is more strongly affected by an increase in density in its vital rates?), it means for instance that increasing the total population 357 size N_{tot} affects the survival rates *S* of all ages *i* differently:

358
$$
logit(S_{i,t}) = \alpha_i + \beta_{Ntot\,i} \times N_{tot,t} \quad (1),
$$

 the β coefficients assessing the strength of density dependence acting on the survival of each age *i*. Importantly, density regulation can act on various vital rates (e.g. survival,

 reproduction), differently according to the age of the individuals, with potentially different (non-linear) forms. In the second case (i.e. who is affecting the most the vital rates of others?), it means for instance that increasing the number of individuals in age class *j=1* does not have the same effect on survival rate than increasing the number of individuals in age class *j>1*:

365
$$
logit(S_t) = \alpha + \sum_{j=1}^{j=number \text{ of } age \text{ groups}} \beta_j \times N_{j,t} (2)
$$

366 We define the age classes *j* with the largest negative β_i as the critical age groups. More complex density regulation patterns can exist. For instance, an increase in the number of individuals in a given age group *j* can differently affect the survival rates of individuals according to their age (mixture of Eq. 1 & 2):

370
$$
logit(S_{i,t}) = \alpha_{i,j} + \sum_{j=1}^{j=number \ of \ age \ groups} \beta_{i,j} \times N_{j,t} \ (3)
$$

 In age-structured populations, all individuals do not equally respond to an increase in density (Eq. 1). Since the 70's, it is well known that some ages are more strongly influenced in their vital rates (survival, reproduction) than others. The pioneering work of Eberhardt [76] for instance predicts sequential demographic responses to an increase in density, with first an

 increase in mortality rate for immatures, followed by a change in reproductive rate of adult females and finally an increase in adult mortality rate. This means that adults are affected by density only when the population reaches the carrying capacity, whereas the youngest individuals seem to be the first age group affected by density dependence [77]. Since then, this theoretical model has been supported by empirical studies (e.g. [77–81]). More generally, there is increasing empirical evidence that individuals, depending on their age, are not equally influenced by an increase in density (see e.g. [7,22,82]). Another striking example of age- specific response to increasing density has been reported in a German great tit population, in which low density favors fast explorer individuals that are the youngest ones, whereas high densities rather select for slower explorers generally the oldest ones [44,83]. This adds to the spate of studies highlighting that depending on their age, individuals may not equally respond to an increase in density. Furthermore, it has been shown that the evolutionary dynamics of species depend on the life stage on which density is acting [84]. How does this age-specific response to density interact with the critical age group? Let's consider that age class *j* is the critical age group. One can expect that an increase in the density of age group *j* could lead to different density-dependent patterns at the population level if i) individuals of different ages do not respond equally to such an increase in density; ii) all individuals in the population equally respond to increasing density; iii) only individuals of age class *j* are influenced by an increase of the density in their own age class. Exploring more deeply this question would offer interesting insights into the role played by age in density regulation, not only in terms of contribution but also in terms of response to density regulation. ---

-
-
-
-

Box 2: The concept of critical age group for density dependence in the context of climate change

 Density dependence in age-structured natural populations is a dynamical process, its strength is expected to vary across space and time in relation to climatic conditions. For instance, empirical evidence for stronger strength of density dependence under harsh environmental conditions is accumulating in the literature (see e.g. [85–87]). This means that climate conditions can mediate competitive abilities and, thus, density dependence [88], which is important in the current context of global changes. Critical age groups make no exception: within a species, the critical age group for density dependence varies among populations, depending on environmental conditions. For instance, in their study involving multiple great tit and blue tit populations across Europe, Gamelon et al. [25] found that even if young females consistently contribute to density regulation, older individuals also play an important role in one-third of the populations. They concluded that the fact that some ages appear to be important in driving density regulation at some sites but not others could be attributed to differences in local environmental conditions. Accordingly, the critical age class can be driven by local abiotic conditions and can change over habitat types, i.e. across space. Along the same line, one can expect that the critical age group during a period of time may change if environmental conditions are changing. Assessing to what extent age-specific contributions to density dependence depends on climate conditions offers exciting research perspectives, especially now, while climate is changing at an unprecedented rate.

 Further support for the key role of climate conditions in mediating biotic interactions is provided in behavioral studies. Some have explored the influence of climate conditions on personality traits potentially related to competitive abilities. For instance, environmental temperature during early life has been shown to shape the personality of mosquitofish

References

- 1. Lande R, Engen S, Sæther B-E. 2003 *Stochastic Population Dynamics in Ecology and Conservation*. Oxford ; New York: Oxford University Press.
- 2. Andrewartha HG, Birch LC. 1954 *The distribution and abundance of animals*. University of Chicago Press.
- 3. Nicholson AJ. 1957 The Self-Adjustment of Populations to Change. *Cold Spring Harb. Symp. Quant. Biol.* **22**, 153–173. (doi:10.1101/SQB.1957.022.01.017)
- 4. Turchin P. 1995 Population regulation: old arguments and a new synthesis. In *Population Dynamics*, pp. 19–40. New York: N. Cappucino & P.W. Price.
- 5. Coulson T, Rohani P, Pascual M. 2004 Skeletons, noise and population growth: the end of an old debate? *Trends Ecol. Evol.* **19**, 359–364. (doi:10.1016/j.tree.2004.05.008)
- 6. Leirs H, Stenseth NC, Nichols JD, Hines JE, Verhagen R, Verheyen W. 1997 Stochastic seasonality and nonlinear density-dependent factors regulate population size in an African rodent. *Nature* **389**, 176–180. (doi:10.1038/38271)
- 7. Coulson T, Catchpole EA, Albon SD, Morgan BJT, Pemberton JM, Clutton-Brock TH, Crawley MJ, Grenfell BT. 2001 Age, sex, density, winter weather, and population crashes in Soay sheep. *Science* **292**, 1528–1531. (doi:10.1126/science.292.5521.1528)
- 8. Boyce MS, Haridas CV, Lee CT, The Nceas Stochastic Demography Working Group. 2006 Demography in an increasingly variable world. *Trends Ecol. Evol.* **21**, 141–148. (doi:10.1016/j.tree.2005.11.018)
- 9. Sinclair ARE. 1989 Population regulation of animals. In *Ecological concepts*, pp. 197–241. J. M. Cherrett.
- 10. Royama T. 1992 *Analytical Population Dynamics*. Springer Science & Business Media.
- 11. Lande R, Engen S, Saether B-E, Filli F, Matthysen E, Weimerskirch H. 2002 Estimating density dependence from population time series using demographic theory and life-history data. *Am. Nat.* **159**, 321–337. (doi:10.1086/338988)
- 12. Berryman AA, Lima Arce M, Hawkins BA. 2002 Population regulation, emergent properties, and a requiem for density dependence. *Oikos* **99**, 600–606. (doi:10.1034/j.1600-0706.2002.12106.x)
- 13. Brook BW, Bradshaw CJA. 2006 Strength of evidence for density dependence in abundance time series of 1198 species. *Ecology* **87**, 1445–1451. (doi:10.1890/0012- 9658(2006)87[1445:SOEFDD]2.0.CO;2)
- 14. Sinclair ARE, Pech RP. 1996 Density Dependence, Stochasticity, Compensation and Predator Regulation. *Oikos* **75**, 164–173. (doi:10.2307/3546240)
- 15. Coulson T, Godfray HCJ. 2007 Single-species dynamics. *Theor. Ecol.* , 17.
- 16. Krebs CJ. 2002 Two complementary paradigms for analysing population dynamics. *Philos. Trans. R. Soc. B Biol. Sci.* **357**, 1211–1219. (doi:10.1098/rstb.2002.1122)
- 17. Charlesworth B. 1973 Selection in populations with overlapping generations. V. Natural selection and life histories. *Am. Nat.* **107**, 303–311. (doi:10.1086/282832)
- 18. Charlesworth B. 1972 Selection in populations with overlapping generations. III. Conditions for genetic equilibrium. *Theor. Popul. Biol.* **3**, 377–395. (doi:10.1016/0040-5809(72)90011-1)
- 19. Gamelon M, Grøtan V, Engen S, Bjørkvoll E, Visser ME, Sæther B. 2016 Density dependence in an age-structured population of great tits: identifying the critical age classes. *Ecology* **97**, 2479– 2490. (doi:10.1002/ecy.1442)
- 20. Lande R, Grøtan V, Engen S, Visser ME, Sæther B-E. 2023 Estimating Density Dependence, Environmental Variance, and Long-Term Selection on a Stage-Structured Life History. *Am. Nat.* **201**, 557–573. (doi:10.1086/723211)
- 21. Lande R, Engen S, Sæther B, Coulson T. 2006 Estimating Density Dependence from Time Series of Population Age Structure. *Am. Nat.* **168**, 76–87. (doi:10.1086/504851)
- 22. Sæther B-E *et al.* 2016 Demographic routes to variability and regulation in bird populations. *Nat. Commun.* **7**, 12001. (doi:10.1038/ncomms12001)
- 23. Cotto O, Wessely J, Georges D, Klonner G, Schmid M, Dullinger S, Thuiller W, Guillaume F. 2017 A dynamic eco-evolutionary model predicts slow response of alpine plants to climate warming. *Nat. Commun.* **8**, 15399. (doi:10.1038/ncomms15399)
- 24. Schmid M, Paniw M, Postuma M, Ozgul A, Guillaume F. 2022 A Trade-Off between Robustness to Environmental Fluctuations and Speed of Evolution. *Am. Nat.* **200**, E16–E35. (doi:10.1086/719654)
- 25. Gamelon M, Vriend SJG, Engen S, Adriaensen F, Dhondt AA, Evans SR, Matthysen E, Sheldon BC, Sæther B-E. 2019 Accounting for interspecific competition and age structure in demographic analyses of density dependence improves predictions of fluctuations in population size. *Ecol. Lett.* **22**, 797–806. (doi:10.1111/ele.13237)
- 26. Charlesworth B. 1971 Selection in Density-Regulated Populations. *Ecology* **52**, 469–474. (doi:10.2307/1937629)
- 27. Engen S, Sæther B-E. 2016 Optimal age of maturity in fluctuating environments under r- and K-selection. *Oikos* **125**, 1577–1585. (doi:10.1111/oik.03111)
- 28. Nussey DH, Coulson T, Festa-Bianchet M, Gaillard J-M. 2008 Measuring Senescence in Wild Animal Populations: Towards a Longitudinal Approach. *Funct. Ecol.* **22**, 393–406.
- 29. Williams GC. 1957 Pleiotropy, natural selection, and the evolution of senescence. *Evolution* **11**, 398–411. (doi:10.2307/2406060)
- 30. Hamilton WD. 1966 The moulding of senescence by natural selection. *J. Theor. Biol.* **12**, 12–45.
- 31. Nussey DH, Froy H, Lemaître J-F, Gaillard J-M, Austad SN. 2013 Senescence in natural populations of animals: widespread evidence and its implications for bio-gerontology. *Ageing Res. Rev.* **12**, 214–225. (doi:10.1016/j.arr.2012.07.004)
- 512 32. Cam E, Link WA, Cooch EG, Monnat J, Danchin E, 2002 Individual Covariation in Life-History Traits: Seeing the Trees Despite the Forest. *Am. Nat.* **159**, 96–105. (doi:10.1086/324126)
- 33. McCleery R h, Perrins C m, Sheldon B c, Charmantier A. 2008 Age-specific reproduction in a long- lived species: the combined effects of senescence and individual quality. *Proc. R. Soc. B Biol. Sci.* **275**, 963–970. (doi:10.1098/rspb.2007.1418)
- 34. Smith JM. 1973 The logic of Animal COnflict. **246**.
- 35. Richter X-YL, Lehtonen J. 2023 Half a century of evolutionary games: a synthesis of theory, application and future directions. *Philos. Trans. R. Soc. B Biol. Sci.* **378**, 20210492. (doi:10.1098/rstb.2021.0492)
- 36. Krebs JR, Davies NB, Parr J. 1993 *An introduction to behavioural ecology, 3rd ed*. Cambridge, MA, US: Blackwell Scientific Publications.
- 37. Westneat D, Fox C, Westneat D, Fox C, editors. 2010 *Evolutionary Behavioral Ecology*. Oxford, New York: Oxford University Press.
- 38. Moiron M, Laskowski KL, Niemelä PT. 2020 Individual differences in behaviour explain variation in survival: a meta-analysis. *Ecol. Lett.* **23**, 399–408. (doi:10.1111/ele.13438)
- 39. Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. 2007 Integrating animal temperament within ecology and evolution. *Biol. Rev.* **82**, 291–318. (doi:10.1111/j.1469-185X.2007.00010.x)
- 40. Cole EF, Quinn JL. 2011 Personality and problem-solving performance explain competitive ability in the wild. *Proc. R. Soc. B Biol. Sci.* **279**, 1168–1175. (doi:10.1098/rspb.2011.1539)
- 41. Linnenbrink M. 2022 Competitive ability is a fast-evolving trait between house mouse populations (Mus musculus domesticus). *Front. Zool.* **19**, 31. (doi:10.1186/s12983-022-00476-7)
- 42. López‐Sepulcre A, Kokko H. 2005 Territorial Defense, Territory Size, and Population Regulation. *Am. Nat.* **166**, 317–325. (doi:10.1086/432560)
- 43. Mougeot F, Redpath SM, Leckie F, Hudson PJ. 2003 The effect of aggressiveness on the population dynamics of a territorial bird. *Nature* **421**, 737–739. (doi:10.1038/nature01395)
- 44. Dingemanse NJ, Moiron M, Araya-Ajoy YG, Mouchet A, Abbey-Lee RN. 2020 Individual variation in age-dependent reproduction: Fast explorers live fast but senesce young? *J. Anim. Ecol.* **89**, 601–613. (doi:10.1111/1365-2656.13122)
- 45. Jones OR *et al.* 2013 Diversity of ageing across the tree of life. *Nature* **505**, 169–173. (doi:10.1038/nature12789)
- 46. Class B, Brommer JE. 2016 Senescence of personality in a wild bird. *Behav. Ecol. Sociobiol.* **70**, 733–744. (doi:10.1007/s00265-016-2096-0)
- 47. Carere C, Locurto C. 2011 Interaction between animal personality and animal cognition. *Curr. Zool.* **57**, 491–498. (doi:10.1093/czoolo/57.4.491)
- 48. Wright J, Haaland TR, Dingemanse NJ, Westneat DF. 2022 A reaction norm framework for the evolution of learning: how cumulative experience shapes phenotypic plasticity. *Biol. Rev.* **97**, 1999–2021. (doi:10.1111/brv.12879)
- 49. Šárová R, Špinka M, Stěhulová I, Ceacero F, Šimečková M, Kotrba R. 2013 Pay respect to the elders: age, more than body mass, determines dominance in female beef cattle. *Anim. Behav.* **86**, 1315–1323. (doi:10.1016/j.anbehav.2013.10.002)
- 50. Pluháček J, Bartoš L, Čulík L. 2006 High-ranking mares of captive plains zebra Equus burchelli have greater reproductive success than low-ranking mares. *Appl. Anim. Behav. Sci.* **99**, 315–329. (doi:10.1016/j.applanim.2005.11.003)
- 51. Cabrera D, Nilsson JR, Griffen BD. 2021 The development of animal personality across ontogeny: a cross-species review. *Anim. Behav.* **173**, 137–144. (doi:10.1016/j.anbehav.2021.01.003)
- 52. Petelle MB, McCoy DE, Alejandro V, Martin JGA, Blumstein DT. 2013 Development of boldness and docility in yellow-bellied marmots. *Anim. Behav.* **86**, 1147–1154. (doi:10.1016/j.anbehav.2013.09.016)
- 53. Araba BD, Crowell-Davis SL. 1994 Dominance relationships and aggression of foals (Equus caballus). *Appl. Anim. Behav. Sci.* **41**, 1–25. (doi:10.1016/0168-1591(94)90048-5)
- 54. Sæther B-E, Engen S, Matthysen E. 2002 Demographic Characteristics and Population Dynamical Patterns of Solitary Birds. *Science* **295**, 2070–2073. (doi:10.1126/science.1068766)
- 55. Jiang Y, Bolnick DI, Kirkpatrick M. 2013 Assortative Mating in Animals. *Am. Nat.* **181**, E125–E138. (doi:10.1086/670160)
- 56. Sergio F, Blas J, Hiraldo F. 2009 Predictors of floater status in a long-lived bird: A cross-sectional and longitudinal test of hypotheses. *J. Anim. Ecol.* **78**, 109–18. (doi:10.1111/j.1365- 2656.2008.01484.x)
- 57. Werner EE, Gilliam JF. 1984 The Ontogenetic Niche and Species Interactions in Size-Structured Populations. *Annu. Rev. Ecol. Syst.* **15**, 393–425. (doi:10.1146/annurev.es.15.110184.002141)
- 58. Wilbur HM. 1980 Complex Life Cycles. *Annu. Rev. Ecol. Syst.* **11**, 67–93. (doi:10.1146/annurev.es.11.110180.000435)
- 59. Nicholson AJ. 1933 Supplement: the Balance of Animal Populations. *J. Anim. Ecol.* **2**, 131–178. (doi:10.2307/954)
- 60. Sæther B-E, Visser ME, Grøtan V, Engen S. 2016 Evidence for *r* and *K* -selection in a wild bird population: a reciprocal link between ecology and evolution. *Proc. R. Soc. B Biol. Sci.* **283**, 20152411. (doi:10.1098/rspb.2015.2411)
- 61. Charlesworth B. 1994 *Evolution in Age-Structured Populations*. 2 edition. Cambridge England ; New York: Cambridge University Press.
- 62. Gamelon M, Tufto J, Nilsson ALK, Jerstad K, Røstad OW, Stenseth NC, Sæther B-E. 2018 Environmental drivers of varying selective optima in a small passerine: a multivariate, multiepisodic approach. *Evolution* **72**, 2325–2342. (doi:10.1111/evo.13610)
- 63. Hunter DC, Pemberton JM, Pilkington JG, Morrissey MB. 2018 Quantification and decomposition of environment-selection relationships. *Evol. Int. J. Org. Evol.* **72**, 851–866. (doi:10.1111/evo.13461)
- 64. Travis J, Bassar RD, Coulson T, Reznick D, Walsh M. 2023 Density-Dependent Selection. *Annu. Rev. Ecol. Evol. Syst.* **54**, 85–105. (doi:10.1146/annurev-ecolsys-110321-055345)
- 65. Travis J, Bassar RD, Coulson T, Lopez-Sepulcre A, Reznick D. 2023 Population Regulation and Density-Dependent Demography in the Trinidadian Guppy. *Am. Nat.* **202**, 413–432. (doi:10.1086/725796)
- 66. Sæther B-E. 2024 How population size shapes the evolution of guppy fish. *Nature* **626**, 725–726. (doi:10.1038/d41586-024-00276-5)
- 67. Myhre AM, Engen S, Sæther B-E. 2016 Effective size of density-dependent populations in fluctuating environments. *Evolution* **70**, 2431–2446.
- 68. Schaub M, Abadi F. 2011 Integrated population models: a novel analysis framework for deeper insights into population dynamics. *J. Ornithol.* **152**, 227–237. (doi:10.1007/s10336-010-0632-7)
- 69. Zipkin EF, Saunders SP. 2018 Synthesizing multiple data types for biological conservation using integrated population models. *Biol. Conserv.* **217**, 240–250. (doi:10.1016/j.biocon.2017.10.017)

 70. Le Galliard J-F, Fitze PS, Ferrière R, Clobert J. 2005 Sex ratio bias, male aggression, and population collapse in lizards. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 18231–18236. (doi:10.1073/pnas.0505172102)

- 71. Lotka AJ. 1925 *Elements of physical biology*. Baltimore: Williams and Wilkins Company.
- 72. Volterra V. 1926 Fluctuations in the abundance of a species considered mathematically. *Nature* **118**, 558.

- 74. Dhondt AA. 2011 *Interspecific competition in birds*. Oxford University Press.
- 75. Levine JM, Bascompte J, Adler PB, Allesina S. 2017 Beyond pairwise mechanisms of species coexistence in complex communities. *Nature* **546**, 56–64. (doi:10.1038/nature22898)
- 76. Eberhardt LL. 1977 Optimal policies for conservation of large mammals, with special reference to marine ecosystems. *Environ. Conserv.* **4**, 205–212. (doi:10.1017/S0376892900025819)
- 77. Bonenfant C *et al.* 2009 Empirical evidence of density‐dependence in populations of large herbivores. In *Advances in Ecological Research* (ed H Caswell), pp. 313–357. Academic Press.
- 78. Fowler CW. 1987 A review of density dependence in populations of large mammals. *Curr. Mammal.* **1**, 401–441.
- 79. Fowler CW. 1981 Density Dependence as Related to Life History Strategy. *Ecology* **62**, 602–610. (doi:10.2307/1937727)
- 80. Gaillard J-M, Festa-Bianchet M, Yoccoz NG. 1998 Population dynamics of large herbivores: variable recruitment with constant adult survival. *Trends Ecol. Evol.* **13**, 58–63. (doi:10.1016/S0169-5347(97)01237-8)

 73. Wauters LA, Gurnell J, Martinoli A, Tosi G. 2002 Interspecific competition between native Eurasian red squirrels and alien grey squirrels: does resource partitioning occur? *Behav. Ecol. Sociobiol.* **52**, 332–341. (doi:10.1007/s00265-002-0516-9)

- 81. Araya-Ajoy YG *et al.* 2021 Variation in generation time reveals density regulation as an important driver of pace of life in a bird metapopulation. *Ecol. Lett.* **24**, 2077–2087. (doi:10.1111/ele.13835)
- 82. Pardo D, Barbraud C, Authier M, Weimerskirch H. 2013 Evidence for an age-dependent influence of environmental variations on a long-lived seabird's life-history traits. *Ecology* **94**, 208–220.
- 83. Nicolaus M, Tinbergen JM, Ubels R, Both C, Dingemanse NJ. 2016 Density fluctuations represent a key process maintaining personality variation in a wild passerine bird. *Ecol. Lett.* **19**, 478–486. (doi:10.1111/ele.12584)
- 84. Mylius SD, Diekmann O. 1995 On Evolutionarily Stable Life Histories, Optimization and the Need to Be Specific about Density Dependence. *Oikos* **74**, 218–224. (doi:10.2307/3545651)
- 85. Peeters B *et al.* 2022 Harvesting can stabilise population fluctuations and buffer the impacts of extreme climatic events. *Ecol. Lett.* **25**, 863–875. (doi:10.1111/ele.13963)
- 86. Gamelon M *et al.* 2017 Interactions between demography and environmental effects are important determinants of population dynamics. *Sci. Adv.* **3**, e1602298. (doi:10.1126/sciadv.1602298)
- 87. Loison A, Sæther B-E, Jerstad K, Røstad OW. 2002 Disentangling the sources of variation in the survival of the European dipper. *J. Appl. Stat.* **29**, 289–304.
- 88. Stenseth NChr *et al.* 2015 Testing for effects of climate change on competitive relationships and coexistence between two bird species. *Proc. R. Soc. B Biol. Sci.* **282**, 20141958. (doi:10.1098/rspb.2014.1958)
- 89. Li H, Zhang X, Wu Y, Zhang F, Li C. 2021 Environmental temperature during early life affects the personality of mosquitofish in adulthood. *Curr. Zool.* **67**, 481–488. (doi:10.1093/cz/zoab003)
- 90. de Jong M, Phillips BL, Llewelyn J, Chapple DG, Wong BBM. 2022 Effects of developmental environment on animal personality in a tropical skink. *Behav. Ecol. Sociobiol.* **76**, 137. (doi:10.1007/s00265-022-03240-3)
- 91. Biro PA, Beckmann C, Stamps JA. 2009 Small within-day increases in temperature affects boldness and alters personality in coral reef fish. *Proc. R. Soc. B Biol. Sci.* **277**, 71–77. (doi:10.1098/rspb.2009.1346)
-

Figure legends

Figure 1

 Schematic summary of the unified approach integrating evolution, behavior and demography to study density regulation in age-structured populations. On the left, the "classical" view of density dependence is pattern-oriented. The total population size (*N*) negatively influences mean vital rates (e.g. survival, reproduction), that in turn influence the population growth rate. On the right, the suggested process-oriented framework linking demography, evolution and behavior. The total population size is divided into age-specific population sizes (*Ni*) (example with 4 age classes black, blue, orange and green), that negatively influence age-specific vital rates more or less strongly depending on their competitive abilities, driven by senescence, learning, sociality, etc. In turn, age-specific demographic rates influence the population growth rate. The critical age group corresponds to the age(s) contributing the most to the density regulation acting on the population growth rate.

Figure 2

 Simplified hypothetical scenarios leading to different critical age groups. Each color corresponds to one phenotype. The first row (A, B, C) corresponds to a scenario of selective disappearance of phenotypes with the lowest competitive abilities, resulting in higher competitive abilities and per-capita effect on growth rate for older age classes (age 4 is the critical age group). The second row (D, E, F) corresponds to a scenario of selective disappearance of phenotypes with the highest competitive abilities, resulting in lower competitive abilities and per-capita effect on growth rate for older age classes (age 1 is the critical age group). The third row (G, H, I) corresponds to a scenario of senescence with equal rate for all phenotypes, resulting in lower competitive abilities and per-capita effect on growth

- 676 rate for older age classes (age 1 is the critical age group). The fourth row (J, K, L)
- corresponds to a scenario in which both senescence and selective disappearance of
- phenotypes with the lowest competitive abilities happen, resulting in complex non-linear
- relationships between competitive ability and age (age 3 is the critical age group).

