

#### A single-step genomic evaluation of maturing rate index in French beef breeds

Armance Lepers, Pauline Martin, Sébastien Taussat

#### ► To cite this version:

Armance Lepers, Pauline Martin, Sébastien Taussat. A single-step genomic evaluation of maturing rate index in French beef breeds. 75. Annual Meeting of the European Federation of Animal Science (EAAP), Sep 2024, Florence (ITA), France. hal-04759255

#### HAL Id: hal-04759255 https://hal.science/hal-04759255v1

Submitted on 29 Oct 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# A single-step genomic evaluation of maturing rate index in French beef breeds



<sup>1</sup>Institut de l'Elevage, 149 Rue de Bercy, 75012 Paris, France, <sup>2</sup>INRAE GABI, UMT e-bis, 78352 Jouy-en-Josas, France, <sup>3</sup> Eliance, 149 Rue de Bercy, 75012 Paris, France

# INTRODUCTION

Particularity of the maturing rate index:



- This traits is interesting to :
- Maximise early beef production
- Reduce fattening time and added concentrates  $\mathbf{i}$
- Reduce the environmental impact of finished products >
- Facilitate 2-year-old calving through early development >



Moderate heritability of 0.24 and genetic variability of 9%



Estimated phenotype after slaughter (at least 4 years old)

# **MATERIAL AND METHODS**

Performance of maturing rate index : Estimated on adult females of 5 beef breeds using individuals weighings and the Brody equation.



L'ELEVAGE **idele** 

#### **Brody equation:**

 $Weight(time) = Weight_{adult} - (Weight_{adult} - Weight_{birth})e^{-maturing rate index*time}$ 

## **Genomic evaluation:**

- Univariate Single-Step model with HSSGBLUP software
- Fixed effects related to birth and slaughter + age at slaughter as a covariate.

Genomic selection on the maturing rate index is essential

## **RESULTS AND DISCUSSION**

## Number of animals used in genomic evaluation

Breed	Animals in pedigree	Animals phenotyped	Reference population
Aubrac	23,155	7,269	414
Blonde d'Aquitaine	171,204	93,718	3,856
Charolaise	811,536	460,627	7,360
Limousine	449,507	270,417	2,179
Parthenaise	61,851	38,929	1,060

**Reliability of GEBVs of genotyped candidates in each breeds:** 



## Impact of a +1 standard deviation selection on performance









an average gain of -1.3 months to reach 50% of the adult weight across all breeds (-1 to -1.7 month)



The reliability of candidates is fairly low, so it is important to take this into account during selection, particularly for the Aubrac.



Increasing the size of the reference population by genotyping females with performance data would be a good strategy for improving the accuracy of estimates



# OR

% of adult an average gain of 3.2% of the adult weight weight at 15m reached at the age of 15 months across all breeds  $\bigcirc$ (+3% to +3.5 %)

# **BUT ALSO**



an average loss of -40.6 kg of the adult weight across all breeds (-35 kg to -50.7kg)

# CONCLUSION

- Selecting on the maturing rate index is of interest for the productivity and profitability of the cattle beef farm
- A genomic evaluation is important to select as early as possible
- Evaluation requires regular genotyping of females to improve the reliability of GEBVs  $\mathbf{i}$

Many thanks to the 6 breed societies who participated.





















MINISTÈRE

la contribution

(co)funded by APIS-GENE

**Research program** 

**APIS-GENE** 

Invest Innovate Exploi

La responsabilité du ministère en charge de l'agriculture ne saurait être engagée