



**HAL**  
open science

## **The contribution of ultrasound to the synthesis of colloidal uranium oxide nanoparticles**

Matthieu Viro, Manon Cot-Auriol, Thomas Dumas, Philippe Moisy, Sergey I. Nikitenko

### ► **To cite this version:**

Matthieu Viro, Manon Cot-Auriol, Thomas Dumas, Philippe Moisy, Sergey I. Nikitenko. The contribution of ultrasound to the synthesis of colloidal uranium oxide nanoparticles. 18th Meeting of the European Society of Sonochemistry, May 2024, Louvain, Belgium. <hal-04764748>

**HAL Id: hal-04764748**

**<https://hal.science/hal-04764748v1>**

Submitted on 4 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## The contribution of ultrasound to the synthesis of colloidal uranium oxide nanoparticles

M. Viro<sup>1</sup>, M. Cot-Auriol<sup>1</sup>, T. Dumas<sup>2</sup>, P. Moisy<sup>2</sup>, S. I. Nikitenko<sup>1</sup>

<sup>1</sup>*ICSM, Univ Montpellier, CEA, CNRS, ENSCM, Marcoule, France*

<sup>2</sup>*CEA, DES, ISEC, DMRC, Univ Montpellier, Marcoule, France*

The tetravalent state is the main oxidation state of uranium in the nuclear fuel, and it is also the most prevalent reduced form under environmental conditions, particularly in the anoxic and reducing conditions of deep natural waters.[1-3] The speciation of uranium under these conditions is complex, and the potential formation of intrinsic colloids resulting from the hydrolysis of U(IV) is rarely reported in the literature. However, it has been shown that colloidal species can significantly contribute to the migration of radionuclides in the environment.[4-6] Therefore, the preparation of reproducible and size-controlled nanoparticles is of particular interest for environmental studies, but also for the development of materials tailoring specific properties, particularly when organized at the nanoscale.

This presentation will describe the synthesis and characterization of intrinsic colloids of U(IV), prepared by hydrolysis (65 °C, Ar) or ultrasonic irradiation (20 kHz, 65 °C, Ar/10%CO) from a HCl-stabilized U(IV) solution. The application of ultrasound in the presence of carbon monoxide, to inhibit the formation of hydrogen peroxide and thus the oxidation of U(IV), allowed the preparation of colloidal suspensions of uranium in aqueous solution. Characterization of these suspensions by SAXS, TEM-HR, and XRD revealed that they are composed of quasi-spherical nanoparticles with a crystalline structure similar to that of UO<sub>2+x</sub>, measuring 18.0 ± 0.1 nm and 7.8 ± 0.1 nm after sonolysis and hydrolysis, respectively. The application of power ultrasound appears as an original and relevant synthesis route for the preparation of actinide oxide nanoparticles with controlled size and crystalline properties.

### Keywords

Nanoparticle, Uranium, Colloid

### References

- [1] W. Cha, H. K. Kim, H. Cho, H. R. Cho, E. C. Jung, S. Y. Lee, *RSC Advances*, 2020, 10, 36723-36733.
- [2] R. C. Ewing, *Nature Materials*, 2015, 14, 252-257.
- [3] A. Y. Romanchuk, I. E. Vlasova, S. N. Kalmykov, *Frontiers in Chemistry*, 2020, 8, 630.
- [4] H. Zänker, C. Hennig, *Journal of Contaminant Hydrology*, 2014, 157, 87-105.
- [5] N. Priyadarshini, M. Sampath, S. Kumar, U. K. Mudali, R. Natarajan, *Journal of Nuclear Chemistry*, 2014, 2014, 1-10.
- [6] C. Walther, M. A. Denecke, *Chemical Reviews*, 2013, 113, 995-1015.