



HAL
open science

Methodological developments for monitoring the estimation of flammability indicators using airborne hyperspectral images in a Mediterranean forest context

Mathilda Porterie, Karine Adeline, Xavier Briottet

► To cite this version:

Mathilda Porterie, Karine Adeline, Xavier Briottet. Methodological developments for monitoring the estimation of flammability indicators using airborne hyperspectral images in a Mediterranean forest context. 8e colloque scientifique du groupe SFPT-GH, Jul 2023, Paris, France. 2023. hal-04765171

HAL Id: hal-04765171

<https://hal.science/hal-04765171v1>

Submitted on 4 Nov 2024


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONTEXT

Global Warming and Mediterranean Ecoregion


- Mediterranean ecoregion is a **biodiversity hotspot** with «drought high risk» as stated by IPCC.
- Savanna ecosystem** (about 20% of continental surfaces) is one of Mediterranean habitat that offers an important biodiversity because of its landscape mosaic.
- Even if **regular wildfires** take part of the regeneration of this Mediterranean ecosystem, the effects of Global Warming are affecting fire regime with more intensity and longer droughts.
- Coupling with anthropogenic pressures, fires are and will be **bigger and more frequent** in future years.



FAQ8.3 IPCC report 2022: prediction of most vulnerable regions against frequent and intense droughts

« Fire disturbance continuum »


From Veraverbeke, 2018



- Fire prevention (**pre-fire** phase) requires an accurate knowledge of fuel stand and **fire risk monitoring**.
- Modelisation** is mainly focused on fire behavior (active-fire phase) but is limited by the lack of spatial precision of the information on vegetation conditions.
- Fuel stand is generally assessed either by empirical approaches (spectral index) or by quantitative estimations through modelisation
- The challenge by pre-fire modelisation is to take into account the fuel stand behavior throughout the entire phenological cycle and to improve its spatial resolution

State of art

- Fuel stand estimations at the end of **spring** provide information on potential fire season tendency (Huesca et al., 2009)
- Spring (beginning of fire-season) studies are scarce and challenging in **open forests** due to the **mixed** spectral signals of under and overstory.



From Brown, 2016

The objective is to estimate biochemical and structural variables of woody vegetation in open Mediterranean forests based on a hybrid method during biomass peak phenophase when the two vegetation layers are active

Fuel Type

LAI

Leaf area index

Vegetation type

- leaf area relative to a surface unit.
- Variable without unit correlated to foliar density of a vegetative system at different scales: tree, population, stand, ecosystem, biome
- Essential climate variable** for observing and monitoring climate change effects (GCOS Global Climate Observing System, 2011).



Fuel spatial distribution

Maps of species or plant functional type classification and canopy cover (CC) to estimate fuel spatial continuity

VARIABLES

Fuel Stand

LFMC

Live Fuel Moisture Content

Hydric

- Leaf water concentration accessible with:
 - **EWT**: Equivalent Water Thickness
 - This concentration must be normalized with respect to the structure of the vegetation defined by:
 - **LMA**: Leaf Mass per Area

$$LFMC (\%) = \frac{EWT - LMA}{LMA} \times 100$$


Pigments

Physiologic

Hydric stress → changes of chlorophylls concentration [Cab] and carotenoids [Car] + reduction of intercellular spaces occupied by air in the mesophyll = changes in spectral behaviour



METHODOLOGY

Hybrid approach



1 Background spectral extraction and classification from Démoulin (internship 2022)



Background classification with 3 components for SJER



- Selection of the pixel with maximum proportion of background information (thresholding from MCARI2 image)
- Selection of the adequate number of components from PCA applied on an image stack composed of 3 dates per year for different phenological periods
- GMM classification and selection of number of classes


2 LUT generation with RTMs

- Simple Forest Representation with woody structure from LiDAR = DETAIL (from Miraglio PhD 2021)
- Leaf reflectance obtained with PROSPECT implemented in DART
- Background parameterized with AVIRIS CLASSIC spectra using classification from 1
- Design of Experiments generated for 700 combinations of [Cab], [Car], LAI, Average Leaf Angle, EWT, LMA, CC and background spectra



DART parameters

Trees geometric properties		Scene size (m)	
Crown shape	Ellipsoid composed	CC10	45,95 x 45,95
Crown diameter	8m	CC30	26,5 x 26,5
Trunk height below canopy	2m	CC50	20,5 x 20,5
Trunk Height with the crown	6m	CC70	17,4 x 17,4
		CC90	15,3 x 15,3





Study site

Measures of the functional and structural vegetation traits

These 2 sites are Mediterranean savannas (**Tree Grass Ecosystem = TGE**) where hyperspectral airborne data were performed since 2006 (AVIRIS CLASSIC). Coupled with these overflights, field data was collected by the CSTARS laboratory as part of a NASA-University of Davis – California project for the HySpIRI mission. These field bio-physio-chemical measurements will serve as calibration and validation data.

- 1 - Tonzi Ranch
- 2 - San Joaquin Experimental Range




Dataset description	
Biochemical	Chlorophylls and carotenoids pigment contents
Structural	LAI derived from hemispherical photo processing
	Forest Inventory

From Miraglio PhD 2021

Remote sensing

- Hyperspectral airborne images are very useful because they allow to estimate most vegetation traits
- Proof of local correlations between LFMC estimations from AVIRIS CLASSIC and LFMC from field (Roberts et al., 2006)

Dates		
Site	Image	Field data
SJER	20140407	LAI: 20140409-10 Bio: 20140411
TZ	20140410	LAI: 20140415 Bio: 20140410



AVIRIS – CLASSIC products characteristics:


- Airborne data acquired at 4-20 km from the level of the sea
- Hyperspectral data with 224 contiguous spectral channels from 400 → 2500 nm
- Reflectance images (L2) resampled at 18 m

Limits:

- Coarse spatial resolution to estimate variables at tree scale: majority of mixed pixels composed of trees + background
- No systematic revisiting: fuel monitoring not possible

3 Biophysical and structural vegetation traits estimation and validation: PHYTREE

- Generation of Species mask (*Quercus douglasii* and/or *Quercus wislizeni* >50% of pixel)
- Generation of Canopy Cover mask (CC >10%)
- Comparison of estimation performance with different machine learning algorithms (PLSR, RLR, RF, neural network ...)
- Validation criterion = comparison of variables estimated with field data values (R², RMSE)



The authors are grateful to the John Muir Institute of the Environment team (Susan Ustin) for collecting and processing the field data, and to NASA JPL from providing AVIRIS-C data (NASA grant No. NNX12AP08G). They also thank Jean-Philippe Gastellu-Etchegorry from CESBIO for his insight and help concerning the DART simulations and the HyperMED project (CNES)

Gale, Matthew G., Geoffrey J. Cary, Albert I. J. M. Van Dijk, et Marta Yebra. 2021. « Forest Fire Fuel through the Lens of Remote Sensing: Review of Approaches, Challenges and Future Directions in the Remote Sensing of Biotic Determinants of Fire Behaviour ». *Remote Sensing of Environment* 255 (mars): 112282.

Huesca et al (2009). Assessment of forest fire seasonality using MODIS fire potential: A time series approach. *Agricultural and Forest Meteorology*, 149(11), 1946-1955.

Miraglio, Thomas, Karine Adeline, Margarita Huesca, Susan Ustin, et Xavier Briottet. 2020. « Monitoring LAI, Chlorophylls, and Carotenoids Content of a Woodland Savanna Using Hyperspectral Imagery and 3D Radiative Transfer Modeling ». *Remote Sensing* 12 (1): 28.

Veraverbeke, Sander, Philip Dennison, Ioannis Gitas, Glynn Hulley, Olga Kalashnikova, Thomas Katagis, Le Kua, Ran Meng, Dar Roberts, et Natasha Stavros. 2018. « Hyperspectral Remote Sensing of Fire: State-of-the-Art and Future Perspectives ». *Remote Sensing of Environment* 216 (octobre): 105-21.

Yebra, Marta, Philip E. Dennison, Emilio Chuvieco, David Riaño, Philip Zylstra, E. Raymond Hunt, F. Mark Danson, Yi Qi, et Sara Jurdao. 2013. « A Global Review of Remote Sensing of Live Fuel Moisture Content for Fire Danger Assessment: Moving towards Operational Products ». *Remote Sensing of Environment* 136 (septembre): 455-68.