



HAL
open science

Sonoluminescence of Tb(III) at the extended solid-Liquid interface

Matthieu Viot

► **To cite this version:**

Matthieu Viot. Sonoluminescence of Tb(III) at the extended solid-Liquid interface. ATALANTE 2012 - International Conference on Nuclear Chemistry for Sustainable Fuel Cycles, Sep 2012, Montpellier, France. <hal-04765333>

HAL Id: hal-04765333

<https://hal.science/hal-04765333v1>

Submitted on 4 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Sonoluminescence of Tb(III) at the extended Solid-Liquid interface

Matthieu Viroot*, Rachel Pflieger, Johann Ravaux, Sergey I. Nikitenko

*Institute for Separation Chemistry of Marcoule (ICSM), UMR 5257 CEA/CNRS/UM2/ENSCM,
ICSM Site de Marcoule, BP 17171, 30207 Bagnols-sur-Cèze, France.*

*e-mail: matthieu.viroot@cea.fr

Abstract

The propagation of ultrasound in a liquid media may lead to acoustic cavitation, which is the nucleation, growth, and rapid collapse of vapor-filled micro bubbles. The collapse of these bubbles is known to generate extreme conditions in the medium (“hot spots”), going with high temperatures and pressures, shock waves, chemical reactions, and possibly the emission of light, known as sonoluminescence (SL). Recent investigations considered the SL as a source of excitation for photoactive species contained in the sonicated media. For instance, Ashokkumar and Grieser studied the 515 kHz excitation of pyranine in aqueous solution; and showed that the observed emission spectra were resulting from SL excitation and not from chemical reactions that could occur in the sonicated media. This phenomenon, referred to as “sonophotoluminescence”, was further observed during the sonication of fluorescent species in various aqueous and organic solutions. Later, these investigations led to the 20 kHz sonication of lanthanide salts in aqueous solutions. The Ln(III) excitation was suggested to occur *via* two mechanisms: (i) sonophotoluminescence, and (ii) collisional excitation of Ln(III) species at the bubble-solution interface with “hot” particles (radicals, excited molecules, etc.).

More recently, studies showed that the light emitted during the sonication of organic and inorganic slurries in appropriate solvent was resulting from triboluminescence (emission of light resulting from the mechanical action on a solid) rather than “true” SL. In view of the above, the excitation of luminescent species with ultrasound, or SL appears to be a quite new topic, scarcely reported in the literature. Most of the studies concentrated on homogeneous systems, and investigations dealing with heterogeneous media are even scarcer. To our knowledge, SL phenomenon has never been studied at the vicinity of an extended solid-liquid interface. In the present work, we investigated the 20 kHz excitation of Tb(III) contained in a lanthanide phosphate pellet: $(\text{Ce}_{0.9}, \text{Tb}_{0.1})\text{PO}_4$. Experiments were performed at 20 kHz in cooled aqueous solution sparged with argon, in a home-made reactor designed for the present purpose. Spectroscopic investigations revealed the possible observation of Tb(III) $f-f$ $^3\text{D}_4-^7\text{F}_5$ emission (Fig. 1). The emission origin was evidenced to result from a solid state excitation with the light emitted from the acoustically-generated cloud of cavitation bubbles, i.e. sonophotoluminescence.




Fig. 1 : SL spectra observed during the sonication of $(\text{Ce}_{0.9}\text{Tb}_{0.1})\text{PO}_4$ pellet