



## isomiRs as a feature for comparing microRNA regulatory networks in aggressive lymphomas

Romain Piucco, Ghislain Fievet, Julien Broséus, Sébastien Hergalant

### ► To cite this version:

Romain Piucco, Ghislain Fievet, Julien Broséus, Sébastien Hergalant. isomiRs as a feature for comparing microRNA regulatory networks in aggressive lymphomas. 23rd European Conference on Computational Biology - ECCB2024, Sep 2024, Turku (Finland), Finland. hal-04767947


HAL Id: hal-04767947

<https://hal.science/hal-04767947v1>

Submitted on 5 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




# IsomiRs as a feature for comparing microRNA regulatory networks in aggressive lymphomas

**ROMAIN PIUCCO<sup>1</sup>, GHISLAIN FIÉVET<sup>1</sup>, JULIEN BROSEUS<sup>1,2</sup> AND SÉBASTIEN HERGALANT<sup>1</sup>**

<sup>1</sup>NUTRITION, GENETICS AND ENVIRONMENTAL RISK EXPOSURE (NGERE) – INSERM U1256, UNIVERSITY OF LORRAINE, NANCY, FRANCE


<sup>2</sup>DEPARTMENT OF BIOLOGICAL HEMATOLOGY, LABORATORY CENTER, UNIVERSITY HOSPITAL OF NANCY, NANCY, FRANCE



## MIRNA & ISOMIRS

MicroRNAs (miRNAs) constitute a class of small non-coding RNAs that regulate protein translation by targeting messenger RNAs.



Mature miRNAs sequences are subject to variations, creating a panel of isomiRs with unique target repertoires and turnover rates. Recent studies increasingly underscore the involvement of isomiRs in cancer development.



## ISOMIRS IN AGGRESSIVE LYMPHOMAS

We applied our methodology to a publicly available small-RNA-seq dataset of 102 diffuse large B-cell lymphoma (DLBCL) and healthy samples<sup>1</sup>.

Comparative analysis between relapsed DLBCL and non-relapsed DLBCL cases revealed variations in the average count of unique isotypes per miRNA, as well as differential isomiR proportions and SNVs.



## CONCLUSION

These findings suggest a global loss in the isotype repertoire diversity and a disruption of regulatory mechanisms controlling expression of miRNAs in relapsed DLBCls.

This approach redefines isomiRs as intricate networks shaping the expression of miRNAs. It opens novel opportunities for investigating miRNA-seq data and for their use in multi-omics analyses.