



HAL
open science

Development of auditory cognition in 5- to 10-year-old children: Focus on speech-in-babble-noise perception

Jérémie Ginzburg, Lesly Fornoni, Pierre-Emmanuel Aguera, Caroline Pierre, Anne Caclin, Annie Moulin

► **To cite this version:**

Jérémie Ginzburg, Lesly Fornoni, Pierre-Emmanuel Aguera, Caroline Pierre, Anne Caclin, et al.. Development of auditory cognition in 5- to 10-year-old children: Focus on speech-in-babble-noise perception. *Child Development*, 2025, 96 (1), pp.407-425. 10.1111/cdev.14178 . hal-04769620

HAL Id: hal-04769620

<https://hal.science/hal-04769620v1>

Submitted on 18 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Development of auditory cognition in 5- to 10-year-old children: focus on speech-in-**
2 **babble-noise perception**

3 J. Ginzburg¹, L. Fornoni¹, P.E. Aguera¹, C. Pierre¹, A. Caclin¹, A. Moulin¹

4 ¹ Université Claude Bernard Lyon 1, CNRS, INSERM, Centre de Recherche en
5 Neurosciences de Lyon CRNL U1028 UMR5292, F-69500, Bron, France

6 Corresponding author: J. Ginzburg, jeremie.ginzburg@inserm.fr

7 Acknowledgments: We thank all the children who took part in the study, along with their
8 parents, and their welcoming teachers. The data necessary to reproduce the analyses presented
9 here are not publicly accessible due to privacy or ethical restrictions. The data are available on
10 request from the corresponding author. The analytic code necessary to reproduce the analyses
11 presented in this paper is publicly accessible at the following URL:
12 https://github.com/jeremieginzburg/cd_analysis_sample_code. The materials necessary to
13 attempt to replicate the findings presented here are not publicly accessible. The analyses
14 presented here were not preregistered.

15
16 Funding: This work was conducted within the framework of the LabEx CeLyA (“Centre
17 Lyonnais d’Acoustique”, ANR-10-LABX-0060) and of the LabEx Cortex (“Construction,
18 Cognitive Function and Rehabilitation of the Cortex”, ANR-11-LABX-0042) of Université de
19 Lyon, within the program “Investissements d’avenir” (ANR-16-IDEX-0005) operated by the
20 French National Research Agency (ANR). This work was funded by a Pack Ambition
21 Recherche (COGAUDYS project) from the Auvergne-Rhône-Alpes Region to AC and AM.

22

23

24

25

Abstract

26 Speech-in-noise perception is consistently reported to be impaired in learning disorders, which
27 stresses the importance of documenting its developmental course in young children. In this
28 cross-sectional study, ninety children (41 females, 5.5-11.6 years-old) and nineteen normal-
29 hearing adults (15 females, 20-30 years-old), were tested with a newly-developed closed-set
30 speech perception in babble-noise test, combining two levels of phonological difficulty and two
31 noise levels. Results showed that speech-in-babble-noise perception takes a definite maturation
32 step around 7 years-of-age ($d = 1.17$, grade effect) and is not mature at 10 years-of-age when
33 compared to young adults ($d = .94$, group effect). Developmental trajectories of both accuracy
34 and response times were evaluated, with influences of psycholinguistic factors, to foster the
35 development of adequate screening tests.

36 The ability to make perceptual sense of a complex sound environment (Auditory scene
37 analysis, Bregman, 1994) is a key component of auditory cognition and a major element for
38 harmonious cognitive development in children. Hearing screening has become widely
39 implemented in numerous countries, beginning at birth and continuing at regular intervals
40 throughout children's primary school years. Its primary objective is to identify peripheral
41 hearing loss, which is of paramount importance (Skarżyński & Piotrowska, 2012). However,
42 these screenings typically do not routinely include the assessment of speech perception in noisy
43 environments, despite an estimated 7% of children with normal audiograms experiencing
44 specific difficulties in listening in noise (Bamiou et al., 2001; Wilson & Arnott, 2013). Speech-
45 in-noise perception, along with other key auditory cognition processes (e.g., auditory short-term
46 memory, STM), are consistently reported to be impaired in learning disorders (Bradlow et al.,
47 2003; Majerus & Cowan, 2016; Plaza et al., 2002; Ziegler et al., 2005, 2009). This fact stresses
48 the need for systematically screening such listening difficulties specific to noisy environments
49 (Skarzynski et al., 2015), as they are often associated with learning disorders that will
50 sometimes be diagnosed years later. Dyslexia often eludes diagnosis until formal reading
51 instruction begins. For instance, Boets et al. (2011) found a link between speech-in-noise
52 perception deficits in kindergarten and dyslexia diagnosis in 3rd grade, with frequency
53 modulation sensitivity and speech-in-noise perception predicting reading ability growth. Guzek
54 & Iwanicka-Pronicka (2022) emphasized the importance of screening for auditory processing
55 disorders at age 6, with speech perception in babble noise being the most commonly impaired
56 among 1012 children tested, with and without auditory processing disorder.

57 Children display more difficulties than adults in recognizing speech when faced with
58 background sounds, and this trend is consistent across various stimuli and listening conditions
59 (Kidd & Colburn, 2017). Studies on children's speech recognition in noise reveal that these
60 differences between children and adults persist noticeably until adolescence, with the specific

61 age range varying depending on the nature of the noise. Various factors contribute collectively
62 to the progressive improvement of speech perception in noise as children develop. These
63 include the ability to leverage spatial separation between the target and masker (D. K. Brown
64 et al., 2010; Ellen Peng & Litovsky, 2021; Litovsky, 2005), frequency differences between the
65 target and masker (Flaherty et al., 2019; Leibold et al., 2018; Leibold & Buss, 2019), and the
66 use of glimpses in the noise masker (Buss et al., 2017). Beyond these low-level factors,
67 immature allocation of attention is suggested to play a pivotal role in children's pronounced
68 difficulties with speech in noise (Leibold & Buss, 2019). At the cortical level, it has been
69 proposed that children's challenges in taking advantage of acoustic cues to discriminate between
70 target and masker, may correspond to the maturation timeline of the thalamus, thalamo-cortical
71 radiations, and the primary auditory cortex, with structural maturation concluding around 10
72 years of age (Devous et al., 2006; Pujol et al., 2006; Werner et al., 2012). Since speech in noise
73 maturation has been observed to continue until adolescence, it is conceivable that this late
74 maturation aligns with the delayed development of the secondary auditory cortex and prefrontal
75 regions, known to underlie attentional processes involved in speech-in-noise perception and
76 maturing into adulthood (Kolk & Rakic, 2022).

77 Screening this process throughout development poses a significant challenge, given that
78 both the nature of the speech perception task and the balance of energetic/informational
79 masking within the noise masker exert substantial influence on the resulting scores (Buss et al.,
80 2016, 2022; Corbin et al., 2016). Classically, energetic masking refers to noise maskers with
81 the same spectro-temporal structure as the signal, hence reducing the availability of signal cues
82 and affecting its neural representation, mostly at a peripheral level (e.g., white noise, speech-
83 shaped noise). In contrast, informational masking refers typically to the situation of attending
84 to one talker in the midst of another or several talkers (termed speech-on-speech masking, or
85 babble noise or cocktail party noise), with masking occurring even when the target talker is

86 perfectly audible. Most studies in children have used energetic masking (mostly speech-shaped
87 noise) and showed that differences between children and adults remain noticeable until around
88 9–10 years of age (for a review on development of speech recognition in steady-state noise, see
89 Leibold & Buss, 2019). However, in realistic environments, environmental noises and speech
90 sources from other children within the class add up to form a noise masker closer to a multi-
91 talker babble noise than to speech-shaped broadband noise. In addition, listening to the
92 teacher's voice in a classroom poses challenges due to voice reflections on various surfaces,
93 creating multiple speech signals at the child's ear with different delays. While short
94 reverberation times (<0.5 s) are not problematic, longer delays can lead to temporal smearing
95 and partial masking, reducing word recognition (Wróblewski et al., 2012). Moreover,
96 classroom reverberation times often exceed the recommended 0.6 s, sometimes reaching up to
97 1.5 s (Anderson, 2004; Crandell & Smaldino, 2000), making the noise close to speech-on-
98 speech. Two-talker babble noise is known to be more challenging than speech-shaped noise in
99 young adults as well as in children (Corbin et al., 2016; Lewis et al., 1988) and various studies
100 indicate a more extended maturation of speech in multi-talker noise than speech in speech-
101 shaped noise (Corbin et al., 2016; Elliott, 1979; Hall et al., 2002). Hall et al. (2002) used a
102 forced-choice, picture-pointing task to measure the recognition of spondee words (words
103 comprising two equally stressed syllables, e.g., “popcorn”) within the presence of either speech-
104 shaped noise or speech from two talkers. Participants were children aged 5 to 10 years and
105 adults aged 19 to 48 years. On average, children needed an additional 3 dB to match the
106 performance level of adults in the noise masker. In contrast, there was an 8-dB difference
107 between adults and children in the two-talker masker to reach similar performance. Using a
108 similar task to assess recognition of consonant-vowel stimuli differing only by consonants,
109 Leibold & Buss (2013) showed that performance was adult-like at 11 years of age for the
110 speech-shaped noise. However, in a two-talker masker, performance was lower than adults until

111 the oldest age group (13 years old). Corbin et al. (2016) measured children's speech reception
112 thresholds (SRT, thresholds for 50% correct recognition) of monosyllabic words by varying the
113 masker's intensity with an open-set paradigm. Comparably to the previously described studies,
114 children reached adult-like performance by 10 years of age when the masker was a speech-
115 shaped noise, while children's performance increased up until 13 years of age in a two-talker
116 masker. These studies suggest that with a speech masker comprising two talkers, regardless of
117 the stimuli used for recognition and of the task (open/closed set), children's performance reach
118 adult's performance later than for speech-shaped noise.

119 Babble noise is considered to be the most challenging noise for speech perception and
120 the closest to ecological conditions. Assessing speech perception in babble noise might be more
121 sensitive to listening difficulties associated with learning disorders than using speech shaped
122 noise. The number of talkers (Freyman et al., 2004; Rosen et al., 2013; Simpson & Cooke,
123 2005), the distance in fundamental frequency and the difference in language type (native versus
124 non native, Calandruccio et al., 2016; Calandruccio & Zhou, 2014) between the target talker
125 and the multi-talker competitive babble, have all shown major influence on the masking
126 efficiency in adults. As the number of competitive talkers increase from 1 to 6, consonant
127 identification scores of vowel-consonant-vowel syllables in young normally hearing adults
128 decrease for up to 6 talkers, and then remain fairly stable (or slightly increase) from 8 to 128
129 talkers (Simpson & Cooke, 2005). Similarly, Rosen et al. (2013) obtained a sharp decrease in
130 natural sentence identification scores from 1 to 2 talkers maskers, and a small increase of scores
131 with 4 to 16 multi-talker babble as in Simpson & Cooke (2005)'s study. This observed increase
132 in scores with an increasing number of talkers is unexpected as one would anticipate a decrease
133 in scores due to reduced gaps in noise amplitude, which would decrease glimpsing opportunities
134 that would help speech perception. These results thus indicate that while energetic masking
135 increases with more talkers, informational masking decreases because words spoken in babble

136 noise become progressively less comprehensible as the number of talkers increases. Despite
137 this trend, it is important to note that multi-talker babble retains a significant element of
138 informational masking. Even with a high number of talkers, such as 16 (Rosen et al., 2013) or
139 128 (Simpson & Cooke, 2005), the babble noise remains more challenging for speech
140 perception than a speech-shaped noise masker. Research on the developmental trajectory of
141 speech perception in two-talker babble noise is less common compared to studies on speech-
142 shaped noise. To our knowledge, none have explored this aspect using babble noise involving
143 more than two talkers, despite the indication that this balance between informational and
144 energetic masking aligns more closely with children's real-world auditory environments.

145 Defining the type of masker to use is only part of the story to assess speech-in-noise
146 development in children. Indeed, most previous studies use open-set response paradigms, which
147 can limit their application in children. For instance, in the first “babble noise in children study”,
148 Elliott (1979) used sentences with different predictability of words, and the lower age limit
149 considered for such test was 9 years of age. McCreery et al. (2020) showed a significant
150 influence of vocabulary in sentence recognition in noise for children aged 5 to 6, compared to
151 children older than 9 years of age. Open-set speech tests, that require verbal responses, need
152 clear articulation by the child to allow precise scoring (Stiles et al. 2012), which can make them
153 problematic for hearing-impaired young children, children with disorders of speech sound
154 production (Cabbage & Hitchcock, 2022), or children whose verbal responses can be difficult
155 to obtain (for instance in some children with non-speaking autism). By using picture-pointing
156 response requiring to choose an answer amongst several foils, alternative forced choice tests
157 (nAFC) alleviate the need for a verbal response and are very valuable in paediatric assessments
158 (Leibold et al., 2024; Mendel, 2008; Vickers et al., 2018). Another advantage of alternative
159 forced choice tests for children, in addition to not relying on a verbal response, is the lower
160 sensitivity of this test format to top-down lexical influences, as shown in adults (Sommers et

161 al., 1997), and in young children (S. Jerger et al., 1983). Indeed, alternative forced-choice tests
162 show better reproducibility than open response tests (Gelfand, 1998, 2003), are not sensitive to
163 learning and are less sensitive to talker variability and lexical competition (Clopper et al., 2006;
164 Sommers et al., 1997). By providing the child with a set of possible answers, closed-set tests
165 restrict the set of potential answers, which can reduce the linguistic bias linked to language
166 development (J. Jerger et al., 1968). Jerger et al.'s Pediatric speech intelligibility test (S. Jerger
167 et al., 1981), which consists in a five-alternative forced choice test, with monosyllabic words
168 presented as pictures, has been shown to be independent of receptive language abilities, in
169 children aged 3 to 6 years. In adults, Clopper et al. (2006) showed that increasing the number
170 of alternatives (from 6 to 12) increased the test difficulty, and the sensitivity of closed-set tests
171 to talker variability and lexical competition.

172 In this study, our aim was to evaluate the developmental trajectory of speech perception
173 in multiple-talker ($n > 2$) noise conditions, employing a task designed to address previously
174 identified challenges. This approach enabled us to investigate the potential impact of crucial
175 factors such as phonological proximity, fatigability, and various psycholinguistic factors on
176 speech perception in noise. As this study focused on children, we adhered to the standard format
177 of a four-alternative forced-choice test commonly employed for children (e.g., Calandruccio et
178 al., 2014; Vickers et al., 2018) by using a French language adaptation for children of the English
179 versions of 4AFC tests (Buss et al., 2016; Vickers et al., 2018). We chose a word recognition
180 closed-set task over an open-response sentence-based test like QuickSINTM (Killion et al., 2004)
181 to mitigate contextual influences. We used a 16-talker babble noise in order to bring us closer
182 to real-world situations involving both energetic and informational masking, in contrast to the
183 conventional speech-shaped noises used in traditional clinical audiometry. In order to evaluate
184 the influence of the phonological proximity between target word and foils in nAFC tasks, we
185 manipulated the task demands by modifying the phonological proximity between target word

186 and foils: the same target-word was presented with either phonologically close foils (imposing
187 a more difficult phonological discrimination task), or phonologically distant foils (an easier
188 discrimination task). Indeed, while closed-set paradigms reduce top-down lexical influences
189 (Clopper et al., 2006; S. Jerger et al., 1983; Sommers et al., 1997), this type of task involves
190 directly comparing an auditorily perceived word with the words represented by the pictures in
191 the response set. The phonological proximity between the target words and foils has been shown
192 to influence speech perception in noise for both adults and children in forced-choice paradigms
193 (Clopper et al., 2006, Buss et al., 2016). The ability to discriminate efficiently between
194 phonologically close and distant words is a phonological skill related to the broader concept of
195 phonological awareness, defined as the ability to manipulate spoken language sound units
196 (Anthony & Lonigan, 2004). Phonological awareness undergoes significant and early
197 developmental changes with age (Snowling & Hulme, 1994; Stekić et al., 2023) and is related
198 to literacy acquisition, often being impaired in dyslexia (e.g., Gijbels et al., 2024; Stahl &
199 Murray, 1994; Stekić et al., 2023; Watson & Miller, 1993). Using an adaptive 4AFC in children
200 from 5 to 13 years old, Buss et al. (2016) showed that phonological similarity had a significant
201 effect for both speech-shaped noise and two-talker maskers. While phonological proximity
202 affected children regardless of age for speech-shaped noise, younger children were more
203 impacted by phonological proximity in the two-talker masker. In the present study, we used a
204 16-talker masker, a type of masker that has not been tested for such effects to our knowledge.
205 Therefore, it is important to assess, in our dataset, whether phonological proximity between
206 foils interacts with the developmental trajectory of speech-in-noise abilities, as it does with two-
207 talker maskers, but not speech-shaped noise (Buss et al., 2016). Additionally, we employed a
208 non-adaptive method with an equal number of trials for each phonological proximity condition,
209 allowing for a thorough comparison of its effects.

210 To ensure strict comparability between the two phonological conditions, identical
211 targets were randomly presented in both conditions, with fixed Signal-to-Noise Ratios (SNRs)
212 in separate blocks. This approach was preferred over an adaptive method, involving trial-by-
213 trial SNR modifications, which is considered more susceptible to fatigue and inattention
214 (Wightman & Allen, 1992) and would not allow us a balanced design between the different
215 levels of phonological proximity. Additionally, using this fixed SNR paradigm enabled the
216 examination of potential training effects with task repetition and the assessment of
217 psycholinguistic factors. Lastly, the use of a touch tablet allowed us to take advantage of the
218 greater motivational aspect of such computerized tests in children, and gave us easy access to
219 response time, in addition to accuracy data. This allowed us to investigate, in a systematic
220 manner, developmental aspects of speech perception in babble noise, in 5 to 10 years old
221 children, relative to a group of young normal-hearing adults.

222 Our expectations are as follows: Firstly, we anticipate a steep maturation trajectory in
223 children's speech in babble noise abilities, with younger children demonstrating lower
224 performance compared to older counterparts. Furthermore, we expect that children will not
225 reach adult-level performance even at 10 years of age, given our use of babble noise instead of
226 the typical speech-shaped noise. Secondly, we hypothesize a significant performance difference
227 between phonological proximity conditions, highlighting the importance of this factor in
228 alternative-forced choice tests. Lastly, as we used a forced-choice paradigm with carefully
229 selected pictures and words specifically designed for young children, we do not anticipate
230 observing any effects of learning, vocabulary or psycholinguistic factors throughout the task.
231 Due to the clear and well-founded hypotheses, as well as the straightforward expectations
232 regarding the direction of effects based on existing literature, this study qualifies as
233 confirmatory in nature.

234 **Methods**

235 *Participants*

236 One hundred children (mean age = 7.6 years; min = 4.5 years, max = 10.6 years, 6 left-
237 handed) attending a public primary school participated in this study in June 2019, . The children
238 were tested in June 2019 during school hours. Participation in the study was proposed to a total
239 of 114 children and they were included in the study only if both parents or legal guardians
240 provided a written informed consent. In addition, prior to each test, verbal assent from each
241 child was sought by the experimenters. The study was approved by the relevant services of the
242 French public education services (Inspection de l'Education Nationale [IEN] and Direction
243 Académique des Services de l'Education Nationale [DASEN] of the Isere department).
244 Children from Kindergarten (KG) grade to 5th grade performed the experiment. Before testing,
245 parents filled out a questionnaire about their child's level of education and their own level of
246 education, the child's laterality (e.g., dominant hand), possible vision or auditory impairments,
247 musical activities, bilingualism, learning disabilities, and 11 questions adapted from adults'
248 musical listening questionnaires (Lévêque et al., 2018; Tillmann et al., 2014). Among the 100
249 children, parents' responses to questionnaires revealed that 5 children had a diagnosed learning
250 disability and their data were excluded from the current analysis. Thirty children had seen at
251 least once a speech-therapist, six had already worn ear tubes, eight had a few years of musical
252 education, and three were bilingual. As we aimed here to explore the cognitive abilities of
253 children among a representative set of the population, we present the results including data from
254 all these children. Data from five children were excluded because of technical difficulties during
255 recordings. Overall, data from 10 children were thus excluded from the analysis, the sample
256 sizes per grade of the remaining 90 children are reported in Table 1. Data on participant
257 ethnicity were not collected because the collection of such data is illegal in France and would
258 require an exceptional waiver from government agencies that we did not seek. Proportion of
259 parental level of education did not significantly differ by grade (see Figure 1 in Ginzburg et al.,

2022). Due to the limited number of children with musical education, a history of ear tubes, and bilingualism, these variables could not be included in the main analysis. However, for each of these factors, we separately conducted the main developmental trajectory analysis by excluding data from children with musical education, a history of ear tubes, or bilingualism. We found that none of these exclusions altered the main result, aside from a small decrease in statistical power.

Nineteen adults (four men), with a mean age of 24.11 years (SD = 3 years) were also included in the study. None of them presented any visual or auditory impairment, nor neurological or psychiatric troubles, and none of them had any musical background.

Table 1: Number of participants and mean age (SD in parentheses) for each grade and matching English and French labels for educational level.

English label	Kindergarten (KG)	1 st Grade	2 nd Grade	3 rd Grade	4 th Grade	5 th Grade
French label	Grande Section de Maternelle (GS)	Cours Préparatoire (CP)	Cours Élémentaire 1 (CE1)	Cours Élémentaire 2 (CE2)	Cours Moyen 1 (CM1)	Cours Moyen 2 (CM2)
N (n females)	11 (5)	16 (9)	17 (8)	16 (6)	15 (5)	15 (8)
Mean age in years (SD)	5.01 (0.31)	6.03 (0.29)	6.93 (0.30)	8.01 (0.34)	8.92 (0.26)	9.94 (0.29)

270

271 ***Stimulus construction and task design***

272 Each trial of the speech-in-babble task consisted in matching an aurally-presented word
 273 with its corresponding image among four images. The material consisted of 20 monosyllabic
 274 and 4 disyllabic words, easily representable by a drawing, selected as a function of their
 275 concreteness, their frequency of occurrence in the French language, their age of acquisition,
 276 and their phonological similarity. The 24 selected words were recorded by a French native

277 female talker, in a sound proof booth, using a Rode NT1 microphone. Sound files had an
278 average duration of 540 ms (SD=171 ms) and their amplitudes were equated in root mean
279 square (RMS) level. Each word had a corresponding hand-drawn image. From this list of 24
280 words, six sets of four phonologically close words were created, e.g., /ku/, /kul/, /ru/, /ruz/ (*cou*,
281 *coule*, *roue*, *rouge* in French, corresponding to *neck*, *flow*, *wheel*, *red* in English). Within each
282 phonologically close set, only consonants could change and up to three phonemes were
283 modified. With the same 24 words, six sets of four phonologically distant words were created,
284 e.g., /bã/, /flœR/, /mẽ/, /trwa/ (*banc*, *fleur*, *main*, *trois* in French, corresponding to *bench*, *flower*,
285 *hand*, *three* in English), where several phonemes, including the vowels, were different. For the
286 sake of clarity throughout the remainder of the article, we will refer to the phonologically distant
287 condition as the "easy condition" (easy to discriminate) and to the phonologically close
288 condition as the "difficult condition" (difficult to discriminate). The average of pairwise
289 Levenshtein distance (i.e., the minimal possible number of insertions, deletions, and
290 substitutions needed to transform one chain of phonemes into another) was 3.03 phonemes (SD
291 = 0.4) for the easy condition and 1.39 phonemes (SD = 0.29) for the difficult condition. Using
292 the French database 'Lexique' (New et al., 2004), we calculated the number of phonological
293 neighbors (i.e., number of existing words by changing one phoneme without changing the
294 others) for each of the 24 words used, and then extracted the number of unique phonological
295 neighbors for each 4-word set. In the easy condition, the four words of each set were
296 phonologically distant (each word having phonological neighbors different from the other
297 three) thus the number of unique phonological neighbors within a set was relatively high (mean
298 = 83, sd = 6.7). Conversely, in the difficult condition, the four words were quasi-phonological
299 neighbors (each word in a set having several common phonological neighbors with the other
300 three) thus having a relatively small number of unique phonological neighbors within a set
301 (mean = 59.7, sd = 14.8). Lastly, we derived for each of the target words their frequency of

302 occurrence (per million occurrences) in the French language (mean = 85.9, SD=119) calculated
303 from a movie subtitles corpus (New et al., 2007) (for details about the stimuli, see supplemental
304 Tables S1 and S2).

305 *Procedure*

306 Children were tested by groups of five or six in the sport room of their school. Before
307 testing, children sat in front of their tables, listening to the experimenter's instructions. The
308 experimenter explained the speech-in-babble noise task with cardboard signs: children would
309 see 4 images appear on the touchpad and at the same time, they would hear a word through their
310 headphones. They had to find the spoken word in one of the four images and tap on it as quickly
311 as possible. They had to ignore the people talking in the background (babble noise). The task
312 was implemented on a touch-tablet and auditory stimuli were displayed binaurally through
313 Sennheiser HD-250-Pro circum-aural headphones. Children underwent a training block
314 comprising eight trials with a SNR of +3 dB. Then they underwent two 48-trial blocks with a
315 SNR of +3 dB for the first block and a SNR of -3 dB for the second block. In the subsequent
316 sections of the article, we will refer to the +3 dB SNR condition as the "favorable noise
317 condition" and to the -3 dB SNR condition as the "unfavorable noise condition". The target-
318 word was systematically presented at 66 dB SPL (A-weighted). During each trial, one of the
319 four-image sets was presented, and 800 ms later, the target word was aurally presented. The
320 subject had to tap with a finger on the matching image as quickly as possible but without any
321 time limit. The next trial began immediately after the response. In both blocks, twelve four-
322 image sets were used: six consisting of phonologically distant words (easy condition) and six
323 consisting of phonologically close words (difficult condition, see above). Within a block, each
324 set was presented four times, each time with a different target word (since the same words were
325 used to create the easy and the difficult condition, each word was used twice as the target: once
326 in each of the two phonological difficulty conditions). A block was thus divided in four series

327 of 12 trials with each of the four-image set presented once per series. The order of presentation
328 of the 12 sets was randomized for each subject and each block, and the same randomized order
329 was then used for the four series. This was done to avoid, as much as possible, the potential
330 advantage linked to the limitation of the number of remaining response alternatives once items
331 have been eliminated from the list of possibilities during the first series. The position of the
332 images presented on the screen was randomized. Lastly, no feedback was given on a trial per
333 trial basis. During the task, a babble noise made of 16 unintelligible French native male and
334 female voices (Moulin et al., 2013) was presented in a continuous manner. The use of 16 talkers
335 enabled us to achieve a balanced representation of both masculine and feminine voices,
336 reducing the risk of any single voice standing out or being disproportionately closer or further
337 from the signal voice. In contrast, with only a few talkers, the pitch (fundamental frequency)
338 differences between the maskers and the signal can become especially relevant. The subjects
339 could listen to several seconds of the babble noise before triggering the start of the block.

340 The duration of each block was around 2 minutes, so that the total hearing-in-noise test
341 was performed in about 10 minutes briefing and training included. The unfavorable noise
342 condition was always given last, as it was anticipated that the children listening behavior might
343 be modified by the challenge represented by this difficult SNR (i.e., inducing discouragement
344 before the next block).

345 Adults underwent the task individually in an experimental room in the lab with the same
346 equipment as children. They performed the same eight-trial training block as the children and
347 then eight 48-trial blocks with babble noise ranging from -12 dB SNR to +9 dB SNR by steps
348 of 3 dB SNR in order to calculate their individual detection thresholds. As for children, each
349 block was performed at a constant SNR with a constant-stimulus procedure. The order of the
350 blocks was counterbalanced.

351 *Data analyses*

352 All data analysis were performed on R (R Core Team, 2019). For each participant,
353 accuracy (percentage of correct responses) and response times (RT) from sound onset were
354 extracted for each phonological difficulty and for each noise condition. Trials with RTs above
355 8 seconds and when the participant answered before the onset of the sound were excluded (i.e.
356 below 800 ms), representing 1.4% of children's data and 0.1% of adult's data. For all analysis
357 on RTs, we used response times on correct trials only and on trials with common words between
358 the phonological easy and difficult condition (to avoid imbalance due to the fact that more trials
359 were successful in easy than in difficult condition), which represented 75.4% of the data for
360 children and 93.5% for adults.

361 Adult data: psychometric parameters

362 To evaluate if detection thresholds differed according to the phonological difficulty of
363 the task, psychometric curves were fitted to responses as a function of the signal-to-noise ratio
364 using a logistic function with a guess rate λ set to .25 (because of the 4AFC task's nature) for
365 each difficulty condition and for each participant by using the *quickpsy* package in R (Linares
366 & López-Moliner, 2016). It allowed the estimation of detection thresholds (i.e., SNR for which
367 each participant obtains $\lambda + (1 - \lambda) * 0.5 = 62.5\%$ of correct responses thus half-way
368 between chance level (25%) and perfect performance (100%)). Easy and difficult phonological
369 conditions were compared for each subject with the *thresholdcomparisons* function from the
370 *quickpsy* package in R which performs a bootstrap comparison between conditions to determine
371 if the difference between them falls outside a 95% confidence interval. Comparison between
372 conditions at the group level was tested by comparing the average detection thresholds across
373 all subjects for both phonological difficulties with a pairwise t-test after checking for normality
374 with a Shapiro-Wilk test.

375 Development of hearing in babble noise

376 Developmental aspects of children's performance were tested as a function of children's
 377 school grade, allowing for homogeneous groups composed of children being equally scholarly
 378 educated regarding reading and other abilities. In order to assess the effects of grade,
 379 phonological difficulty, and noise condition on children's performance, we used general linear
 380 models (GLM) for accuracy and general linear mixed models (GLMM) for response times
 381 (Bates et al., 2014). We thus analyzed the influence of three main effects on percent correct
 382 scores and RTs:

- 383 - Between-subject factor *Grade*: six levels (KG, 1st, 2nd, 3rd, 4th and 5th grade)
- 384 - Within-subject factor *Phonological Difficulty*: two levels (easy/difficult
 385 phonological condition)
- 386 - Within-subject factor *Noise condition*: two levels (unfavorable/favorable)

387 GLM(M) were fitted using the *lme4* package in R (Bates et al., 2014, p. 4). Statistical
 388 tests were performed using type II Wald chi-square analysis of variance using the *car* package
 389 in R (Fox & Weisberg, 2018), all effect-sizes were calculated for each fixed effect with the *esc*
 390 package in R (Lüdtke, 2019).

391 When a significant main effect or interaction was found, estimated marginal means post-
 392 hoc tests were performed using the *emmeans* package in R (Lenth, 2021) and corrected for
 393 multiple comparisons using the False Discovery Rate (FDR) method (Benjamini & Hochberg,
 394 1995).

395 Two measures were considered to investigate children's performance:

- 396 - *Accuracy*: percent correct scores were fitted with a binomial distribution as
 397 GLMs can handle non-normally distributed data and, in particular, count data (Bates et
 398 al., 2014). Data were thus fitted with a binomial distribution with a logit link function

399 using the *glm* function from the core *stats* package in R. We did not use mixed models
400 for the accuracy because one proportion of correct response was obtained for each
401 condition and each subject, thus rendering the intra-subject variability null.

402 - *Response times*: As recommended by Lo and Andrews (2015), RTs were not log-
403 transformed to fit a gaussian distribution which can produce additive effects. Since raw
404 RTs distribution can be fitted either with an inverse gaussian or a gamma distribution,
405 we compared four models, fitted with either a gamma or an inverse gaussian distribution
406 and with either an identity link function or a log-link function. The model with the
407 lowest AIC was then chosen. The random-effects structure was built according to Bates
408 et al. (2015)'s specifications: we used the subject variable as an intercept and we started
409 by including all within-subject fixed-effects as random slopes (i.e., all effects were
410 allowed to vary across subjects) and dropped slopes factors one by one. After rejecting
411 the non-converging models, we chose the model with the lowest AIC.

412 We also compared measures between the oldest children (5th grade) and adults. We
413 extracted adults' performance for the two noise conditions that children underwent (-3 and +3
414 dB SNR) and we performed similar GLM(M)s than the previously described ones (on accuracy
415 and RTs), replacing the *Grade* factor by a *Group* factor with two levels: 5th grade and adults.
416 We kept the *phonological difficulty* and *noise condition* factors.

417 Impact of parameters potentially influencing performance

418 - Repetition and learning: Influence of trials repetition within a block

419 The effect of trials repetition within a block was assessed by analyzing children's
420 accuracy (percentage of correct responses) as a function of the series of 4-image sets presented
421 during a block (as each block consisted of four consecutive 12-trial series). Accuracies were
422 calculated per subject, per block, and per series and were fitted using the *glm* function from the

423 core *stats* package in R with a binomial distribution and a logit link function with grade (KG to
424 5th grade), phonological difficulty (easy/difficult), noise condition (unfavorable/favorable) and
425 series (1 to 4) as factors. In addition, the reproducibility within a block was assessed by a two-
426 way random effect interclass correlation.

427 - Word/picture familiarity

428 To assess potential variations in word/picture familiarity across different grades and to
429 rule out the influence of differences in picture familiarity on potential group variations in speech
430 perception, we ranked the 24 target words based on average accuracy (percentage of correct
431 responses) within each age group. The ranking ranged from the highest score (rank=1) to the
432 lowest score (rank=24) for both easy and difficult phonological conditions, using the same
433 method as Fallon et al. (2000). We then performed ranked correlations (Spearman method)
434 between the ranks for the different age groups, and between the rank of each age group and the
435 overall average ranking.

436 - Psycholinguistic parameters

437 The effect of two psycholinguistic factors (word frequency of occurrence and number
438 of phonological neighbors, see above) calculated from the Lexique database (New et al., 2004),
439 was assessed on children's accuracy. For frequency of occurrence, mean accuracy was
440 calculated for each target-word (24 in total), each phonological difficulty (easy/difficult) and
441 each noise condition (unfavorable/favorable). A GLMM was fitted on proportion data (number
442 of correct responses over number of trials) and fitted with a binomial distribution, a logit link
443 function and target-word as random factor. In each GLMM, the effect of the frequency of
444 occurrence factor was tested in addition with the grade factor (six levels: kindergarten to 5th
445 grade), the phonological difficulty factor (easy/difficult) and the noise condition factor
446 (unfavorable/favorable).

447 For phonological neighbors, mean accuracy was calculated for each four-image set (six
448 sets of the easy difficulty, six sets for the difficult difficulty). One GLMM was fitted for each
449 difficulty level, as by construction, the number of unique phonological neighbors for a set is
450 smaller for easy sets than difficult sets. These two GLMMs were performed on accuracy and
451 fitted with a binomial distribution, a logit link function, and four-image set as random factor.
452 In each GLMM, the effect of the phonological neighbors factor was tested in addition with the
453 grade factor (six levels: kindergarten to 5th grade) and the noise condition factor
454 (unfavorable/favorable). Statistical tests were performed using type II Wald chi-square analysis
455 of variance using the *car* package in R (Fox & Weisberg, 2018).


456 All effect sizes are reported as Cohen's *d* (Cohen, 1992).

457 **Results**

458 *Adults data: psychometric parameters*

459 Detection thresholds were computed for each participant and for each phonological
460 condition. At the individual level, 74% of the adult participants displayed significantly different
461 detection thresholds between the easy and the difficult conditions (fourteen out of the nineteen
462 participants, Figure 1a). At the group level, detection thresholds averaged across participants
463 were significantly different between the easy and the difficult condition (mean threshold for the
464 easy condition = -10.6 dB; mean threshold for the difficult condition = -6.3 dB; $t(18) = 10.57$,
465 $p < .001$, Figure 1b). Thus, in adults, to achieve similar performance in easy and difficult
466 phonological conditions, an increase of ~4.3 dB of SNR is necessary in the difficult condition
467 relative to the easy one. The average scores achieved by adults at the SNR levels used with
468 children were as follows: in the favorable noise condition (+3 dB SNR), 97% (SD = 16%) for
469 the easy condition and 95% (SD = 22%) for the difficult condition. In the unfavorable noise
470 condition (-3 dB SNR), the scores were 96% (SD = 19%) for the easy condition and 86% (SD
471 = 35%) for the difficult condition. As the adults' data were collected across multiple SNRs to

472 specifically obtain psychometric parameters using accuracy, no RT analysis was performed for
 473 all SNRs in the adults' data. However, adults' RTs were analyzed in comparison to children's
 474 RTs for the same fixed SNR levels (see the Comparison between children and adults'
 475 performance section).



476

477 *Fig. 1: subject-level and group-level effect of the phonological difficulty on psychometric*
 478 *thresholds in adults. a: fitted psychometric functions on the proportion of correct response for*
 479 *each subject (numbered panels) and each phonological difficulty (blue lines: easy/orange lines:*
 480 *difficult) as a function of the SNR during the task. Dotted lines represent the projection of the*
 481 *estimated SNR threshold needed to achieve 62.5% correct response for each condition.*
 482 *Asterisks above the participant number indicate significantly different thresholds for the two*
 483 *conditions at the individual level. b: average detection thresholds across participants as a*
 484 *function of the phonological difficulty. Individual data are represented by black lines. Asterisk*
 485 *indicate a significant difference between the two conditions.*

486 ***Development of hearing-in-babble noise***

487 GLM(M) were performed on accuracy data and response times with Grade
 488 (kindergarten to 5th grade), Phonological Difficulty (easy/difficult), and Noise Condition
 489 (unfavorable/favorable) as factors. Mean accuracy and response times are shown in Figure 2a
 490 and 2b. Complete statistical results are shown in Table 2 and we present significant effects,
 491 interactions, and their follow-up post-hoc tests below.

492

Table 2: Type II Wald chi-square analysis of deviance results on the GLM with binomial modeling for the accuracy measure and on the GLMM with inverse gaussian modeling for the correct response time measure for children data. All main effects were tested for each measure: Grade (six

levels: KG to 5th grade), Phonological Difficulty (two levels: easy/difficult), Noise condition (two levels: unfavorable/favorable) and their interactions (see text for details). Significant effects are depicted in bold font (p -values < .05). df : degrees of freedom. χ^2 : chi-square test. p : p -value. Cohen's d : standardized mean difference.


<i>Measure</i>	<i>Factor</i>	<i>Df</i>	χ^2	<i>p</i>	<i>Cohen's d</i>
Accuracy	Grade	5	22.90	3.5e - 04	1.17
	Phonological Difficulty	1	61.90	3.6e - 15	2.97
	Noise condition	1	32.02	1.5e - 08	1.49
	Grade:Difficulty	5	0.50	.99	0.15
	Grade:Noise	5	1.30	.94	0.24
	Difficulty:Noise	1	2.43	.12	0.33
	Grade:Difficulty:Noise	5	0.83	.98	0.20
Response times	Grade	5	12.90	.024	0.49
	Phonological Difficulty	1	9.49	.0021	0.69
	Noise Condition	1	0.0080	.93	0.02
	Grade:Difficulty	5	3.43	.63	0.10
	Grade:Noise	5	3.52	.62	0.10
	Difficulty:Noise	1	0.40	.53	0.13
	Grade:Difficulty:Noise	5	0.39	.99	0.0012

493 *Accuracy.* The following model was used for accuracy: *correct responses/number of*
494 *trials ~ grade* noise * difficulty, family = binomial(link=logit).* The main effect of Grade was
495 significant ($\chi^2(5) = 22.90, p < .001, d = 1.17$) revealing better performance for older children.
496 Post-hoc tests revealed a significantly lower performance for KG children compared to 3rd, 4th,
497 and 5th graders ($p < .0029$), significantly lower performance for 1st graders compared to 5th
498 graders ($p < .030$), marginally significant lower performance for 2nd graders compared to 5th
499 graders ($p = .060$). The main effect of Phonological Difficulty was significant ($\chi^2(1) = 61.90, p$
500 $< .001, d = 2.97$) with lower performance in the difficult condition. The main effect of Noise

501 Condition was also significant ($\chi^2(1) = 32.02, p < .001, d = 1.49$) with lower performance for
 502 the unfavorable SNR (-3 dB). No interaction effect was significant ($p > .12$).

503 *Response times.* The model that best fitted our data was the one using an inverse
 504 gaussian distribution with a log-link function, with random effect structure including Difficulty
 505 and SNR as slopes ($response\ times \sim grade*difficulty*noise\ (noise + difficulty | subject),$
 506 $family=inverse.gaussian(link=log)$). The main effect of grade was significant ($\chi^2(5) = 12.90, p$
 507 $= .024, d = .49$). Post-hoc tests only revealed a marginally significantly faster response time for
 508 5th graders compared to KG ($p = .050$) and for 5th graders compared to 2nd graders ($p = .050$).
 509 Interestingly, longer response times were obtained for the easy phonological condition
 510 compared to the difficult one (main effect of Phonological Difficulty ($\chi^2(1) = 9.49, p = .0020,$
 511 $d = .69$)). No effect of Noise Condition or interaction effects were found ($p > .53$).

512




513 *Fig.2: (a) mean performance (% of correct response) and (b) mean correct response*
 514 *times after onset of the sound (in seconds) as a function of Noise Condition and Phonological*
 515 *Difficulty for all grades and for adults. Response times are for words correctly recognized in*
 516 *both easy and difficult conditions (per participant). Error bars represent standard deviations*
 517 *around the mean. Blue color represents the easy condition; orange color represents the difficult*
 518 *condition. Solid lines represent the favorable noise condition, dashed lines the unfavorable*
 519 *noise condition. KG: Kindergarten. Adult's average response times for the easy and difficult*
 520 *condition in the favorable noise condition were similar, so that in the figure (b), the triangle*
 521 *for the easy condition hides the one for the difficult condition.*

522

523 ***Comparison between children and adults' performance***

524 ***Comparison of children's performance and adult's psychometric parameters***

525 In order to obtain a representative adult psychometric curve, we fitted a psychometric
 526 curve to accuracy averaged across adult data separately for the easy and the difficult
 527 phonological conditions (respectively left and right panels in Figure 3). We then projected
 528 children's performance on the overall adult psychometric curve as a function of their grade and
 529 of the SNR used (-3 dB and +3 dB, respectively top and bottom panels in Figure 3). We thus
 530 obtained the value of the SNR at which an adult would have to be tested to obtain the same
 531 performance as children. This representation allows to highlight the relative deficit of speech
 532 in noise of children in comparison to adults, using the same task and stimuli. Furthermore, it
 533 allows a better comparison across studies that use different languages and parameters, where
 534 the young normally-hearing adult group acts as a reference. Indeed, the « SNR gap » (i.e., the
 535 difference between the SNR used to test children and the SNR corresponding to the same
 536 performance in adult data) goes up to 10 dB for the youngest children and the minimal one is 1
 537 dB for the oldest children in the easy condition and for a testing SNR of -3 dB.



538

539 *Fig.3: projection of children's performance by grade on the adult's psychometric function*
 540 *(derived from the average data of the 19 adult participants). Left panels and blue lines for the*
 541 *easy condition; right panels and orange lines for the difficult condition. Top panels: the*
 542 *projected performance of children when they underwent the -3 dB SNR condition. Bottom*
 543 *panels: the projected performance of children when they underwent the +3 dB SNR condition.*

544 *Filled circles represent the average performance for adults in the corresponding SNR*
 545 *condition.*

546 *Comparison between performance of oldest children and performance of adults*

547 To compare performance of the oldest children (5th grade) with performance of adults,
 548 we performed GLM(M)s on the same measures as for children (accuracy and response times)
 549 with Group (adults ($n = 19$) and children (5th graders, $n = 15$)), Phonological Difficulty
 550 (easy/difficult), and Noise Condition (unfavorable/favorable) as factors. Mean accuracy and
 551 response times are shown in Figure 2a and 2b. Complete statistical results are shown in Table
 552 3 and we present significant effects, interactions, and their follow-up post-hoc tests below.

553

Table 3: Type II Wald chi-square analysis of deviance results on the GLM with binomial modeling for the accuracy measure and on the GLMM with inverse gaussian modeling for the correct response times measure, to compare data in the oldest children with adult data. All main effects were tested for each measure: Group (two levels: 5th grade vs. adults), Phonological Difficulty (two levels: easy/difficult), Noise Condition (two levels: unfavorable/favorable) and their interactions (see text for details). Significant effects are depicted in bold font (p -values $< .05$). df : degrees of freedom. X^2 : chi-square test. P : p -value. Cohen's d : standardized mean difference.

<i>Measure</i>	<i>Factor</i>	<i>Df</i>	χ^2	<i>p</i>	<i>Cohen's d</i>
Accuracy	Group	1	6.16	.013	0.94
	Phonological Difficulty	1	9.55	.0020	1.25
	Noise Condition	1	4.04	.044	0.74
	Group:Difficulty	1	4.08	.043	0.74
	Group:Noise	1	1.19	.28	0.38
	Difficulty: Noise	1	1.93	.17	0.49
	Group:Difficulty: Noise	1	0.32	.57	0.20
Response times	Group	1	26.92	2.1e- 07	3.9
	Phonological Difficulty	1	3.20	.074	0.53
	Noise Condition	1	2.24	.13	0.65

Group:Difficulty	1	2.31	.13	0.54
Group:Noise	1	12.63	3.8 -04	1.54
Difficulty:Noise	1	1.13	.29	0.37
Group:Difficulty:Noise	1	0.0020	.97	0.015

554

555 *Accuracy.* The following model was used for accuracy: *correct responses/number of*
556 *trials ~ group*difficulty*noise, family = binomial(link=logit)*. All main effects were
557 significant: Group ($\chi^2(1) = 6.16, p = .013, d = 0.94$ with adults having better performance than
558 5th graders, Phonological Difficulty ($\chi^2(1) = 9.55, p = .0020; d = 1.25$) with better performance
559 in the easy condition, and Noise Condition ($\chi^2(1) = 4.04, p = .044, d = .74$) with better
560 performance for the favorable noise condition. In addition, a significant Group-by-Difficulty
561 interaction was found ($\chi^2(1) = 4.08, p = .043, d = 0.74$) with significantly better performance
562 in the easy condition compared to the difficult one for 5th graders ($p < .001$) but not for adults
563 ($p = .29$). Note that the absence of difficulty effect in the adult group was expected as in the
564 present analysis, we are comparing children and adults' data recorded at exactly the same
565 SNR levels, and therefore, there is a ceiling effect for the adult's data.

566 *Response times.* The model that best fitted our data was the one using an inverse
567 gaussian distribution with a log-link function, and a random effect structure including Difficulty
568 and SNR as slopes (*response times ~ group*difficulty*noise + (noise + difficulty | subject),*
569 *family = inverse.gaussian(link=log)*). There was a marginally significant effect of Phonological
570 Difficulty ($\chi^2(1) = 3.20, p = .074, d = 0.65$) for which longer response times were found for the
571 easy condition compared to the difficult condition. Critically, the main effect of Group was
572 significant ($\chi^2(1) = 26.92, p < .001, d = 3.90$) as well as the Group-by-Noise interaction ($\chi^2(1)$
573 $= 12.63, p < .001, d = 1.54$), that revealed longer response times for the unfavorable noise
574 condition (-3 dB) for children ($p = .018$) but not for adults ($p = .66$).

575 *Impact of learning, familiarity, and psycholinguistic parameters*

576 *Learning in the course of a block: Influence of the repetition of the trials*

577 The model used in this analysis of accuracy is the following: *correct responses/number*
 578 *of trials* ~ *grade*difficulty*noise*series, family = binomial(link=logit)*. As expected from the
 579 developmental trajectory analysis above, effects of Grade ($\chi^2(5) = 22.90, p < .001, d = 1.17$),
 580 Phonological Difficulty ($\chi^2(5) = 62.19, p < .001, d = 2.99$) and Noise Condition ($\chi^2(5) = 32.18,$
 581 $p < .001, d = 1.49$) were found. No effect of series was observed ($\chi^2(3) = 0.32, p = .96, d = 0.12$)
 582 indicating an absence of significant increase or decrease of performance within a block. No
 583 interaction effect was found (all $p > .45$). Reproducibility was good with an interclass
 584 correlation coefficient of .60 ($F = 2.57, p < .001$) for the unfavorable noise condition and of .68
 585 ($F = 3.14, p < .001$) for the favorable noise condition.

586 *Word picture familiarity*

587 Ranking of performance per word were correlated between groups, and all pairwise
 588 correlations were significant ($p < .001, r$ ranging from .69 to .87). Correlations between the
 589 ranking of each group with the overall ranking were also significant ($p < .001, r$ ranging from
 590 .83 (KG) to .92 (2nd grade)).

591 *Influence of psycholinguistic parameters*

592 *Frequency of occurrence*

593 The effect of the Frequency of occurrence of target words, Grade (kindergarten to 5th
 594 grade), Phonological Difficulty (easy/difficult), and Noise Condition (unfavorable/favorable)
 595 factors was tested on accuracy. Except for the expected Grade ($\chi^2(5) = 24.30, p < .001, d =$
 596 0.86), Phonological Difficulty ($\chi^2(1) = 66.74, p < .001, d = 3.39$), and Noise Condition ($\chi^2(1)$
 597 $= 34.61, p < .001, d = 1.58$) effects (already found in the previous analysis), no other significant

598 effect or interaction was found (all $p > .11$), with, in particular, no significant effect of the
599 frequency of occurrence.

600 *Phonological neighbors*

601 The effect of the Number of unique phonological neighbors within each 4-images set,
602 Grade (kindergarten to 5th grade), and Noise Condition (unfavorable/favorable) factors were
603 tested on accuracy. One model was fitted for each Phonological Difficulty (easy/difficult). The
604 expected effect of Grade was found for the easy ($\chi^2(5) = 11.16, p = .048, d = 0.43$) and the
605 difficult conditions ($\chi^2(5) = 12.49, p = .029, d = 0.47$), as well as the Noise Condition effect for
606 the easy ($\chi^2(1) = 9.55, p = .0020, d = 0.69$) and the difficult ($\chi^2(1) = 26, p < .001, d = 1.27$)
607 conditions. All other effects or interactions were non-significant ($p > .11$).

608 **Discussion**

609 The present study depicts the developmental aspects of speech perception in multi-talker
610 babble noise, in French language, in primary school children, using a newly developed French
611 closed-set test with a 4AFC paradigm. Additionally, the exploration of children's response
612 times allowed to unveil an effect of competition between phonologically similar words within
613 the lexicon. Finally, we explored the impact on performance of several factors that have
614 implications for the design of central hearing screening test: reproducibility, word familiarity,
615 and psycholinguistic factors.


616 *Developmental trajectory of speech-in-babble-noise perception*

617 The main results show definite steps into the development, with significant differences
618 in accuracy between 5-year-olds (KG) and 7-, 9-, and 10-year-olds (3rd, 4th and 5th grade),
619 between 6-year-olds (1st grade) and 9-10-year-olds (4th and 5th grade), and between 7- and 10-
620 year-olds (2nd and 5th grade), revealing a major maturation step around 7 years of age (2nd
621 grade). The accuracy obtained by the oldest children (10-year-olds, 5th grade) were equivalent

622 to a difference of more than 3 dB SNR on average with normally hearing young adults, showing
623 that speech perception in one of the most common, but challenging situation (i.e., speech-in-
624 babble noise), is far from mature during nearly the entirety of primary and secondary school
625 attendance years. Those results are in agreement with Wilson et al. (2010)'s large normative
626 study, using an open set task in babble noise (Word in Noise test). Indeed, the authors showed
627 that the largest improvement in recognition performance occurred between 6- and 7-year-olds,
628 then a slight improvement between 7 and 9-year-olds, and performance remained fairly constant
629 between 9- and 12-year-olds, with a significant difference between 12-year-olds and young
630 adults.

631 It could be argued that phonological awareness maturation could account for the lower
632 scores of the youngest children, especially for Kindergarten children. For instance, using an
633 oddity non-sense syllable recognition task (presented with 3 auditory non-sense syllables - two
634 identical and one odd), Hnath-Chisolm et al. (1998) obtained significantly lower accuracy for
635 5- to 7-year-olds compared to 7- to-9-, or 9-to-11-year-olds. Those differences could possibly
636 be attributed to cognitive and phonological development for children under 7 years (Hnath-
637 Chisolm et al., 1998). In the present study, even though we obtained the expected greater
638 accuracy for phonologically distant words than for close words, we did not observe any
639 interaction between grade and difficulty in the children's data analysis. This fairly parallel
640 evolution of accuracy in easy versus difficult phonological conditions with age (see also Figure
641 4) argues against an important influence of phonological awareness maturation in the evolution
642 of accuracy as a function of age, at least between 5 and 10 years of age. However, we observed
643 a weakly significant interaction between difficulty and group when comparing children's data
644 with adults' data (with a difference between difficult and easy conditions in children only).
645 Therefore, we cannot dismiss the potential impact of later maturation of phonological
646 awareness on the current test. However, the lack of a difficulty effect in adults could

647 alternatively be attributed to a ceiling effect, particularly influenced by the favorable SNR
 648 condition (+3 dB SNR). Additionally, there might be a potential ceiling effect on response times
 649 for older children, as indicated by the statistical weakness of the interaction.



651 *Fig.4: comparison between the current study and Buss et al. (2016). For the current*
 652 *study a SNR threshold was derived for each child and each condition from the 2 points obtained*
 653 *along the psychometric curve and adding 2 other points: one at 100% correct score for a 15*
 654 *dB SNR and one at 25% chance level for a -15 dB SNR. We then fitted a linear regression as a*
 655 *function of the decimal logarithm of age in years, as in Buss et al. (2022). For the Buss et al.*
 656 *(2016)'s figures, data were recovered from their Figure 2 using the Engauge Digitizer software*
 657 *(<https://markummittell.github.io/engauge-digitizer/>). For the current study, the effect of age*
 658 *was significant ($p < .001$) and slopes were -0.53 and -0.75 for easy and difficult conditions*
 659 *respectively. For Buss et al.'s speech-shaped noise, the effect of age was significant ($p < .014$)*
 660 *and slopes were -0.64 and -0.47 for the easy and difficulty conditions respectively. For the two-*
 661 *talker noise, the effect of age was significant only for the easy condition ($p < .001$) but not for*
 662 *the difficult condition ($p = .14$). Slopes were -1.43 and -0.03. The objective of this comparison*
 663 *with the existing literature is to evaluate the contribution of the informational/energetic*
 664 *masking component within our 16-talker babble noise. This is in contrast to the study we are*
 665 *comparing with (Buss et al., 2016), which employed a two-talker masker with a pronounced*
 666 *informational masking component and speech-shaped noise with a strong energetic masking*
 667 *component.*

668 The difference between phonologically similar and dissimilar words obtained here (as
 669 measured with the procedure described in Figure 4, e.g., a threshold difference of 3.8 dB SNR
 670 in children and 4.3 dB in adults) appears to be lower than the one obtained by Buss et al. (2016)
 671 in a 4AFC task, in a speech shaped noise (about 5 dB in children and 7 dB in adults). Differences
 672 in language (French versus English), in the degree of similarity between the target words and
 673 the foils, and in the type of noise used, could account for that. In addition, Buss et al. (2016)
 674 used the 2 conditions in 2 different blocks, determining the SNR threshold using an adaptative

675 method, whereas we used a fixed SNR method, where easy target words and difficult trials were
676 randomly intermixed. Both in the present study and in Buss et al.'s data (Figure 4), performance
677 improved with age for both easy and difficult phonological conditions. It is interesting to note
678 though, that our results match better with Buss et al. (2016)'s results in speech shaped noise
679 than with their results in 2-talker masker. Indeed, in the 2-talker masker, they obtained a
680 significantly sharper improvement of accuracy with age in the easy condition than in the
681 difficult condition, with a difference of almost 10 dB SNR between both conditions in adults.
682 This similarity between Buss et al. (2016)'s results in speech shaped noise and ours can be
683 easily attributed to the 16-talker babble noise we used, that involves both energetic and
684 informational masking, and is closer, in performance, to speech shaped noise than the strong
685 informational masking of a 2 talker noise (Rosen et al., 2013). The 16-talker babble noise thus
686 offers a good mix between energetic and informational masking. The definite existence of an
687 informational masking component, involving central auditory processing, in this 16-talker
688 babble noise, has been shown in a case of venous cerebral infarct in an adult patient (Bourgeois-
689 Vionnet et al., 2020). Indeed, this patient showed a specific deficit in the 16-talker babble noise
690 speech perception in the ear contralateral to the infarct, without any deficit in a speech shaped
691 noise of the same spectrum, using the same test as in the present study.

692 The informational masking component could explain the absence of mature
693 performance at 10 years of age, as already pointed out by Elliott (1979), who observed
694 significant differences between 13-year-olds and 17-year-olds in speech-in-babble-noise
695 perception. Indeed, speech-in-babble-noise perception depends on a wide range of cognitive
696 abilities, whose development extend well beyond 11 years of age, such as sustained attention
697 (Betts et al., 2006; Hoyer et al., 2021; Thillay et al., 2015), distractibility (Hoyer et al., 2021;
698 Wetzell et al., 2009), impulsivity and motor control (Booth et al., 2003; Wright et al., 2003),
699 verbal short-term memory (Alloway et al., 2006; Gathercole et al., 2004), and the ability to use

700 pitch cues to focus on the target, including pitch short-term memory (Ginzburg et al., 2022).
701 This explains the great difference in the amount of noise that children and young adults can
702 sustain to achieve the same performance, with more than 3 dB of difference in SNR between
703 10-year-olds and young adults, and more than 9 dB between 5-year-olds and adults in the
704 difficult condition (see Figure 3). The latter result is very close to the 10 dB threshold difference
705 reported between 5-year-olds and young adults, in a 2-talker masker, in Buss et al. (2019). This
706 emphasizes the relevance of using a babble-noise in a speech perception screening test as speech
707 in babble noise is often associated with several deficits, such as attention or short-term memory
708 impairments, and would be more sensitive to all those deficits than perception in speech-shaped
709 noise.

710 *Response times reveals competition processes within the lexicon*

711 Response times reflected the expected slower speed of processing from the youngest to
712 the eldest children (with about 500 ms of difference in response times observed here), due to
713 processing speed maturation (Kail, 1991), in addition with slower response times for the oldest
714 children as compared to adults. In a similar 4AFC task, Rigler et al. (2015) obtained a 400 ms
715 difference between 16-year-olds and 9-year-olds having to click on the picture of a target word
716 amongst 4 pictures, revealing that processing speed, as in the present study, undergoes
717 maturation well beyond 10 years of age. Unexpectedly, we observed longer response times for
718 the easy condition compared to the difficult condition, whereas for accuracy, the expected
719 higher accuracy for the easy condition over the difficult one was observed in both children and
720 adults.

721 One mechanism that could explain these counterintuitive results is the organisation of
722 mental lexicon with distant words engendering a greater lexical competition within the mental
723 lexicon than phonologically close words. Using eye tracking with the visual world paradigm
724 (introduced by Huettig et al., 2011), Rigler et al. (2015) showed more activation for competition

725 words for 9-year-olds than for 16-year-olds. Indeed, according to most models for spoken word
726 recognition (e.g., Frauenfelder & Floccia, 1999, for a review), words are organised within the
727 mental lexicon as clusters of phonologically similar words (e.g., the neighbourhood activation
728 model, Luce & Pisoni, 1998) and words sharing the same onset sounds (e.g. the Cohort model
729 Marslen-Wilson, 1987; Marslen-Wilson & Zwitserlood, 1989). As a word begins to be heard,
730 several potential candidates are activated within the mental lexicon and the process of selecting
731 the correct target involves competition within the several activated candidates (e.g., the trace
732 model, Dahan et al., 2001; McClelland & Elman, 1986). Several studies, using open response
733 tests, have shown that words with numerous phonological neighbours are more difficult to
734 recognize and take a longer time to process, especially when those phonological neighbours are
735 of higher occurrence frequency than the target word, than words with a low number of
736 phonological neighbours (Dirks et al., 2001; Kirk et al., 1995; Krull et al., 2010). We would
737 thus expect a greater response time when more numerous candidates are activated. In the
738 present task, candidates are not only activated by the word heard, but, as well, by the three other
739 words displayed in the pictures (i.e., the target and the three foils). Here, in the difficult
740 condition, the words are phonologically close, so the phonological patterns that are activated
741 by the target words and the foils are similar, and are less numerous than the phonological
742 patterns activated in the easy condition. Indeed, because the words, in the easy condition, are
743 phonologically dissimilar, they would activate four different clusters of phonological patterns,
744 demanding a greater number of comparisons between the target word and all the phonological
745 similar words to the foils. In other words, a difficult set (e.g., /ku/, /kul/, /ru/, /ruʒ/) would
746 activate fewer candidates in the participant's lexicon because all of these words share common
747 phonological neighbors. Conversely, an easy set (e.g., /bã/, /flœR/, /mẽ/, /trwa/) would activate
748 a larger number of candidates due to the dissimilarity between each word in the set. Thus, the

749 greater number of activated candidates could account for the longer response times in the ‘easy’
750 condition, even if this condition shows a higher accuracy.

751 *Central hearing screening test design: impact of reproducibility, word familiarity, and*
752 *psycholinguistic factors*

753 Another major result obtained here is the absence of short-term practice effects. Indeed,
754 by using a fixed SNR ratio method, rather than an adaptative one, we were able to assess the
755 potential changes from the beginning of a block (12 first sets) and the end (12 last sets). Indeed,
756 although we avoided the major source of practice effect by not using trial per trial visual
757 feedback (Munro & Lutman, 2005), there are still a wide range of reasons why substantial
758 modifications could occur. Firstly, the same sets of 4 pictures are used throughout, so more
759 familiarity with the specific phonetic contrasts to listen to would be expected as the sets repeat
760 themselves, especially in the difficult condition. Secondly, as the same feminine voice for the
761 target words is used throughout, better identification and separation of the talker’s voice from
762 the babble background could be expected with practice. Those factors could account for a
763 significant improvement of the accuracy with time. Conversely, a deterioration of accuracy
764 could account for lack of maintenance of the necessary listening effort, involving a strong
765 focused attention component, during the entire 2-minute-long blocks, for such a challenging
766 task. We did not observe any significant difference in accuracy between series, whether the
767 whole population was considered, or only the youngest ones (i.e., first quartile, < 7.3 years old).
768 This shows the absence of learning during the course of one block with the task used here, and,
769 as well, the absence of loss of motivation during the course of a block, which is particularly
770 important when addressing young children. Indeed, to maintain motivation, Fallon et al., (2000)
771 used a “game-like” automated visual feedback for correct and incorrect responses in a 4AFC
772 task in babble noise. The authors obtained a significant improvement between the first 20 trials
773 (82%) and the last 20 trials (86%) of a 40-word block at constant SNR in their children and

774 adult groups. In order to be able to use the test later in an adaptative manner, a stability of
775 scores, at least within one block, is recommended.

776 Performance is likely to be influenced by familiarity with the target words depending
777 on children's age (G. D. Brown & Watson, 1987), as contextual influences on those words
778 develop. To assess this possibility, Wilson et al. (2010) checked the performance of their
779 youngest groups in quiet, and obtained significant lower performance in the 6-year-olds versus
780 the older children. They concluded that part of the 6-year-olds increased threshold was probably
781 due to less familiarity for the target words. Although the pictures we used were designed so
782 that 5-year-olds could recognize them easily, a growing familiarity as age increases is difficult
783 to rule out. To address this question, we ranked each target word according to its score and
784 correlated the ranking in each age group with the overall ranking, as in Fallon et al. (2000), who
785 used a 4AFC task as well. We obtained quite similar results (r ranging from .83 to .92) as in
786 their study ($r > .90$), showing that the relative difficulty of each target word didn't depend on
787 age, and that the improvement of accuracy observed with age was not likely due to an increase
788 in familiarity of the target words with age.

789 Although the experimental design was not aimed at testing specifically the influence of
790 psycholinguistic factors on the test accuracy, it is important to ascertain whether they have
791 substantial influence on the accuracy at a test aimed at screening for speech in babble noise
792 perception deficits. Indeed, the influence of such psycholinguistic parameters is rarely
793 considered when developing auditory perception tests. Yet, those parameters can introduce
794 unwanted variability in audiological speech perception accuracy (Moulin et al., 2017),
795 especially in interaction with cognitive status and lexical knowledge (Braza et al., 2022; Moulin
796 & Richard, 2015). Elliot et al. (1983) specifically designed a 4AFC task to test the potential
797 influence of occurrence frequency. By comparing scores for the same target words presented
798 with foils of high and low occurrence frequencies, the authors observed a word frequency effect

799 that was significantly greater in adults than in 3- to 7-year-olds. Furthermore, this frequency
800 effect, in children, increased significantly with receptive vocabulary knowledge, which argues
801 for the need to control for that effect when developing audiological tests of speech perception
802 (Elliott et al., 1983). In the present test, no effect of target-word occurrence frequency was
803 found, nor any effect of phonological neighbourhood density. To ensure that scores reflect more
804 auditory perception (Kosky & Boothroyd, 2003; Mendel, 2008) than mental lexicon
805 development, it is important, for audiological assessment, to ensure that there is no significant
806 influence of lexical factors (such as occurrence frequencies and/or phonological density) of the
807 stimuli chosen on the scores.

808 The limitations of this study lie in the relatively moderate sample size in each grade, the
809 limited age range (focusing on primary school years) and the absence of other thorough clinical
810 investigations (e.g., pure-tone audiometry, including high frequencies audiometry, lexical
811 knowledge and other cognitive investigations). However, these investigations would go beyond
812 the aims of the present study. Indeed, to establish normative data with the aim of a screening
813 test, we would need at least 40 children per age group. Several refinements in the words used
814 could be made as well, in order to further decrease the duration of the test, and being able to
815 use it in children younger than 5 years of age.

816 Be that as it may, this study represents a stepping stone towards a screening test. Indeed,
817 this study showed the developmental course of speech perception in a babble noise that
818 combines both energetic and informational masking, in primary school children, and confirms
819 that processes involved in speech perception in babble noise (in particular informational
820 masking) show a definite maturation step around 7 years of age and that they are far from mature
821 at 10 years of age. We also unveiled the effect of lexicon competition processes on speed
822 processing and checked for crucial factors that can influence such central hearing screening
823 tests, such as word occurrence frequency and phonological neighborhood density. This resulted

824 in a preliminary version of a screening test, that is quick (around 2 minutes for one block),
 825 reproducible, engaging for children, devoid of short-term learning effect and that has therefore
 826 good potential for screening speech perception in babble noise deficits.

827 **References**

- 828 Alloway, T. P., Gathercole, S. E., & Pickering, S. J. (2006). Verbal and Visuospatial Short-Term and
 829 Working Memory in Children : Are They Separable? *Child Development*, *77*(6), 1698-1716.
 830 <https://doi.org/10.1111/j.1467-8624.2006.00968.x>
- 831 Anderson, K. (2004). The Problem of Classroom Acoustics : The Typical Classroom Soundscape Is a
 832 Barrier to Learning. *Seminars in Hearing*, *25*(02), 117-129. [https://doi.org/10.1055/s-2004-](https://doi.org/10.1055/s-2004-828663)
 833 [828663](https://doi.org/10.1055/s-2004-828663)
- 834 Anthony, J. L., & Lonigan, C. J. (2004). The Nature of Phonological Awareness : Converging Evidence
 835 From Four Studies of Preschool and Early Grade School Children. *Journal of Educational*
 836 *Psychology*, *96*(1), 43-55. <https://doi.org/10.1037/0022-0663.96.1.43>
- 837 Bamiou, D., Musiek, F., & Luxon, L. (2001). Aetiology and clinical presentations of auditory processing
 838 disorders—A review. *Archives of Disease in Childhood*, *85*(5), 361-365.
 839 <https://doi.org/10.1136/adc.85.5.361>
- 840 Bates, D., Kliegl, R., Vasishth, S., & Baayen, H. (2015). *Parsimonious Mixed Models*.
 841 <https://doi.org/10.48550/ARXIV.1506.04967>
- 842 Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4.
 843 *arXiv preprint arXiv:1406.5823*.
- 844 Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate : A practical and powerful
 845 approach to multiple testing. *Journal of the Royal statistical society: series B*
 846 *(Methodological)*, *57*(1), 289-300.
- 847 Betts, J., Mckay, J., Maruff, P., & Anderson, V. (2006). The development of sustained attention in
 848 children : The effect of age and task load. *Child Neuropsychology*, *12*(3), 205-221.

- 849 Boets, B., Vandermosten, M., Poelmans, H., Luts, H., Wouters, J., & Ghesquière, P. (2011). Preschool
850 impairments in auditory processing and speech perception uniquely predict future reading
851 problems. *Research in Developmental Disabilities, 32*(2), 560-570.
852 <https://doi.org/10.1016/j.ridd.2010.12.020>
- 853 Booth, J. R., Burman, D. D., Meyer, J. R., Lei, Z., Trommer, B. L., Davenport, N. D., Li, W., Parrish, T. B.,
854 Gitelman, D. R., & Mesulam, M. M. (2003). Neural development of selective attention and
855 response inhibition. *Neuroimage, 20*(2), 737-751.
- 856 Bourgeois-Vionnet, J., Moulin, A., Hermier, M., Pralus, A., & Nighoghossian, N. (2020). A case of
857 verbal and emotional prosody processing dissociation after a right temporal venous infarct.
858 *Neurological Sciences: Official Journal of the Italian Neurological Society and of the Italian*
859 *Society of Clinical Neurophysiology, 41*(6), 1615-1618. [https://doi.org/10.1007/s10072-019-](https://doi.org/10.1007/s10072-019-04175-w)
860 [04175-w](https://doi.org/10.1007/s10072-019-04175-w)
- 861 Bradlow, A. R., Kraus, N., & Hayes, E. (2003). *Speaking Clearly for Children With Learning*
862 *Disabilities Sentence Perception in Noise*. 18.
- 863 Braza, M. D., Porter, H. L., Buss, E., Calandruccio, L., McCreery, R. W., & Leibold, L. J. (2022). Effects of
864 word familiarity and receptive vocabulary size on speech-in-noise recognition among young
865 adults with normal hearing. *PLOS ONE, 17*(3), e0264581.
866 <https://doi.org/10.1371/journal.pone.0264581>
- 867 Bregman, A. S. (1994). *Auditory scene analysis : The perceptual organization of sound*. MIT press.
- 868 Brown, D. K., Cameron, S., Martin, J. S., Watson, C., & Dillon, H. (2010). The North American Listening
869 in Spatialized Noise—Sentences Test (NA LiSN-S) : Normative Data and Test-Retest Reliability
870 Studies for Adolescents and Young Adults. *Journal of the American Academy of Audiology,*
871 *21*(10), 629-641. <https://doi.org/10.3766/jaaa.21.10.3>
- 872 Brown, G. D., & Watson, F. L. (1987). First in, first out : Word learning age and spoken word
873 frequency as predictors of word familiarity and word naming latency. *Memory & cognition,*
874 *15*, 208-216.

- 875 Buss, E., Felder, J., Miller, M. K., Leibold, L. J., & Calandruccio, L. (2022). Can Closed-Set Word
876 Recognition Differentially Assess Vowel and Consonant Perception for School-Age Children
877 With and Without Hearing Loss? *Journal of Speech, Language, and Hearing Research, 65*(10),
878 3934-3950. https://doi.org/10.1044/2022_JSLHR-20-00749
- 879 Buss, E., Hodge, S. E., Calandruccio, L., Leibold, L. J., & Grose, J. H. (2019). Masked Sentence
880 Recognition in Children, Young Adults, and Older Adults : Age-Dependent Effects of Semantic
881 Context and Masker Type. *Ear and Hearing, 40*(5), 1117-1126.
882 <https://doi.org/10.1097/AUD.0000000000000692>
- 883 Buss, E., Leibold, L. J., & Hall, J. W. (2016). Effect of response context and masker type on word
884 recognition in school-age children and adults. *The Journal of the Acoustical Society of*
885 *America, 140*(2), 968-977. <https://doi.org/10.1121/1.4960587>
- 886 Buss, E., Leibold, L. J., Porter, H. L., & Grose, J. H. (2017). Speech recognition in one- and two-talker
887 maskers in school-age children and adults : Development of perceptual masking and
888 glimpsing. *The Journal of the Acoustical Society of America, 141*(4), 2650-2660.
889 <https://doi.org/10.1121/1.4979936>
- 890 Cabbage, K. L., & Hitchcock, E. R. (2022). Clinical Considerations for Speech Perception in School-Age
891 Children With Speech Sound Disorders : A Review of the Current Literature. *Language,*
892 *Speech & Hearing Services in Schools, 53*(3), 768-785. [https://doi.org/10.1044/2022_LSHSS-](https://doi.org/10.1044/2022_LSHSS-21-00120)
893 [21-00120](https://doi.org/10.1044/2022_LSHSS-21-00120)
- 894 Calandruccio, L., Gomez, B., Buss, E., & Leibold, L. J. (2014). Development and Preliminary Evaluation
895 of a Pediatric Spanish–English Speech Perception Task. *American Journal of Audiology, 23*(2),
896 158-172. https://doi.org/10.1044/2014_AJA-13-0055
- 897 Calandruccio, L., Leibold, L. J., & Buss, E. (2016). Linguistic Masking Release in School-Age Children
898 and Adults. *American Journal of Audiology, 25*(1), 34-40. [https://doi.org/10.1044/2015_AJA-](https://doi.org/10.1044/2015_AJA-15-0053)
899 [15-0053](https://doi.org/10.1044/2015_AJA-15-0053)

- 900 Calandruccio, L., & Zhou, H. (2014). Increase in Speech Recognition due to Linguistic Mismatch
901 Between Target and Masker Speech : Monolingual and Simultaneous Bilingual Performance.
902 *Journal of Speech Language and Hearing Research*, 1089.
903 https://doi.org/10.1044/2013_JSLHR-H-12-0378
- 904 Clopper, C. G., Pisoni, D. B., & Tierney, A. T. (2006). Effects of Open-Set and Closed-Set Task Demands
905 on Spoken Word Recognition. *Journal of the American Academy of Audiology*, 17(05),
906 331-349. <https://doi.org/10.3766/jaaa.17.5.4>
- 907 Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112(1), 155-159.
908 <https://doi.org/10.1037/0033-2909.112.1.155>
- 909 Corbin, N. E., Bonino, A. Y., Buss, E., & Leibold, L. J. (2016). Development of Open-Set Word
910 Recognition in Children : Speech-Shaped Noise and Two-Talker Speech Maskers. *Ear &*
911 *Hearing*, 37(1), 55-63. <https://doi.org/10.1097/AUD.0000000000000201>
- 912 Crandell, C. C., & Smaldino, J. J. (2000). Classroom Acoustics for Children With Normal Hearing and
913 With Hearing Impairment. *Language, Speech, and Hearing Services in Schools*, 31(4),
914 362-370. <https://doi.org/10.1044/0161-1461.3104.362>
- 915 Dahan, D., Magnuson, J. S., Tanenhaus, M. K., & Hogan, E. M. (2001). Subcategorical mismatches and
916 the time course of lexical access : Evidence for lexical competition. *Language and Cognitive*
917 *Processes*, 16(5-6), 507-534. <https://doi.org/10.1080/01690960143000074>
- 918 Devous, M. D., Altuna, D., Furl, N., Cooper, W., Gabbert, G., Ngai, W. T., Chiu, S., Scott, J. M., Harris, T.
919 S., Payne, J. K., & Tobey, E. A. (2006). Maturation of Speech and Language Functional
920 Neuroanatomy in Pediatric Normal Controls. *Journal of Speech, Language, and Hearing*
921 *Research*, 49(4), 856-866. [https://doi.org/10.1044/1092-4388\(2006/061\)](https://doi.org/10.1044/1092-4388(2006/061))
- 922 Dirks, D. D., Takayanagi, S., Moshfegh, A., Noffsinger, P. D., & Fausti, S. A. (2001). Examination of the
923 neighborhood activation theory in normal and hearing-impaired listeners. *Ear and Hearing*,
924 22(1), 1-13.

- 925 Ellen Peng, Z., & Litovsky, R. Y. (2021). The Role of Interaural Differences, Head Shadow, and Binaural
926 Redundancy in Binaural Intelligibility Benefits Among School-Aged Children. *Trends in*
927 *Hearing, 25*, 233121652110453. <https://doi.org/10.1177/23312165211045313>
- 928 Elliott, L. L. (1979). Performance of children aged 9 to 17 years on a test of speech intelligibility in
929 noise using sentence material with controlled word predictability. *The Journal of the*
930 *Acoustical Society of America, 66*(3), 651-653. <https://doi.org/10.1121/1.383691>
- 931 Elliott, L. L., Clifton, L. A., & Servi, D. G. (1983). Word frequency effects for a closed-set word
932 identification task. *Audiology, 22*(3), 229-240. <https://doi.org/10.3109/00206098309072787>
- 933 Fallon, M., Trehub, S. E., & Schneider, B. A. (2000). Children's perception of speech in multitalker
934 babble. *The Journal of the Acoustical Society of America, 108*(6), 3023-3029.
935 <https://doi.org/10.1121/1.1323233>
- 936 Flaherty, M. M., Buss, E., & Leibold, L. J. (2019). Developmental Effects in Children's Ability to Benefit
937 From F0 Differences Between Target and Masker Speech. *Ear & Hearing, 40*(4), 927-937.
938 <https://doi.org/10.1097/AUD.0000000000000673>
- 939 Fox, J., & Weisberg, S. (2018). *An R companion to applied regression*. Sage publications.
- 940 Frauenfelder, U. H., & Floccia, C. (1999). The Recognition of Spoken Words. In A. D. Friederici,
941 *Language Comprehension : A Biological Perspective* (p. 1-40). Springer Berlin Heidelberg.
942 https://doi.org/10.1007/978-3-642-59967-5_1
- 943 Freyman, R. L., Balakrishnan, U., & Helfer, K. S. (2004). Effect of number of masking talkers and
944 auditory priming on informational masking in speech recognition. *The Journal of the*
945 *Acoustical Society of America, 115*(5), 2246-2256. <https://doi.org/10.1121/1.1689343>
- 946 Gathercole, S. E., Pickering, S. J., Ambridge, B., & Wearing, H. (2004). The Structure of Working
947 Memory From 4 to 15 Years of Age. *Developmental Psychology, 40*(2), 177-190.
948 <https://doi.org/10.1037/0012-1649.40.2.177>
- 949 Gelfand, S. A. (1998). Optimizing the Reliability of Speech Recognition Scores. *Journal of Speech,*
950 *Language, and Hearing Research, 41*(5), 1088-1102. <https://doi.org/10.1044/jslhr.4105.1088>

- 951 Gelfand, S. A. (2003). Tri-Word Presentations With Phonemic Scoring for Practical High-Reliability
952 Speech Recognition Assessment. *Journal of Speech, Language, and Hearing Research, 46*(2),
953 405-412. [https://doi.org/10.1044/1092-4388\(2003/033\)](https://doi.org/10.1044/1092-4388(2003/033))
- 954 Gijbels, L., Burkhardt, A., Ma, W. A., & Yeatman, J. D. (2024). Rapid online assessment of reading and
955 phonological awareness (ROAR-PA). *Scientific Reports, 14*(1), 10249.
956 <https://doi.org/10.1038/s41598-024-60834-9>
- 957 Ginzburg, J., Moulin, A., Fornoni, L., Talamini, F., Tillmann, B., & Caclin, A. (2022). Development of
958 auditory cognition in 5- to 10-year-old children : Focus on musical and verbal short-term
959 memory. *Developmental Science, 25*(3). <https://doi.org/10.1111/desc.13188>
- 960 Guzek, A., & Iwanicka-Pronicka, K. (2022). Analysis of the auditory processing skills in 1,012 children
961 aged 6–9 confirms the adequacy of APD testing in 6-year-olds. *PLOS ONE, 17*(8), e0272723.
962 <https://doi.org/10.1371/journal.pone.0272723>
- 963 Hall, J. W., Grose, J. H., Buss, E., & Dev, M. B. (2002). Spondee Recognition in a Two-Talker Masker
964 and a Speech-Shaped Noise Masker in Adults and Children: *Ear and Hearing, 23*(2), 159-165.
965 <https://doi.org/10.1097/00003446-200204000-00008>
- 966 Hnath-Chisolm, T. E., Laipply, E., & Boothroyd, A. (1998). Age-related changes on a children's test of
967 sensory-level speech perception capacity. *Journal of Speech, Language, and Hearing*
968 *Research: JSLHR, 41*(1), 94-106. <https://doi.org/10.1044/jslhr.4101.94>
- 969 Hoyer, R. S., Elshafei, H., Hemmerlin, J., Bouet, R., & Bidet-Caulet, A. (2021). Why Are Children So
970 Distractible? Development of Attention and Motor Control From Childhood to Adulthood.
971 *Child Development, 92*(4). <https://doi.org/10.1111/cdev.13561>
- 972 Huettig, F., Rommers, J., & Meyer, A. S. (2011). Using the visual world paradigm to study language
973 processing : A review and critical evaluation. *Acta Psychologica, 137*(2), 151-171.
974 <https://doi.org/10.1016/j.actpsy.2010.11.003>
- 975 Jerger, J., Speaks, C., & Trammell, J. L. (1968). A New Approach to Speech Audiometry. *Journal of*
976 *Speech and Hearing Disorders, 33*(4), 318-328. <https://doi.org/10.1044/jshd.3304.318>

- 977 Jerger, S., Jerger, J., & Abrams, S. (1983). Speech Audiometry in the Young Child: *Ear and Hearing*,
978 4(1), 56-66. <https://doi.org/10.1097/00003446-198301000-00010>
- 979 Jerger, S., Jerger, J., & Lewis, S. (1981). Pediatric Speech Intelligibility test. II. Effect of receptive
980 language age and chronological age. *International Journal of Pediatric Otorhinolaryngology*,
981 3(2), 101-118. [https://doi.org/10.1016/0165-5876\(81\)90026-4](https://doi.org/10.1016/0165-5876(81)90026-4)
- 982 Kail, R. (1991). Developmental change in speed of processing during childhood and adolescence.
983 *Psychological Bulletin*, 109(3), 490-501. <https://doi.org/10.1037/0033-2909.109.3.490>
- 984 Kidd, G., & Colburn, H. S. (2017). Informational Masking in Speech Recognition. In J. C. Middlebrooks,
985 J. Z. Simon, A. N. Popper, & R. R. Fay (Éds.), *The Auditory System at the Cocktail Party* (Vol.
986 60, p. 75-109). Springer International Publishing. [https://doi.org/10.1007/978-3-319-51662-](https://doi.org/10.1007/978-3-319-51662-2_4)
987 2_4
- 988 Killion, M. C., Niquette, P. A., & Gudmundsen, G. I. (2004). Development of a quick speech-in-noise
989 test for measuring signal-to-noise ratio loss in normal-hearing and hearing-impaired
990 listeners). *J. Acoust. Soc. Am.*, 116(4).
- 991 Kirk, K. I., Pisoni, D. B., & Osberger, M. J. (1995). Lexical effects on spoken word recognition by
992 pediatric cochlear implant users. *Ear and Hearing*, 16(5), 470-481.
- 993 Kolk, S. M., & Rakic, P. (2022). Development of prefrontal cortex. *Neuropsychopharmacology*, 47(1),
994 41-57. <https://doi.org/10.1038/s41386-021-01137-9>
- 995 Kosky, C., & Boothroyd, A. (2003). Validation of an On-line Implementation of the Imitative Test of
996 Speech Pattern Contrast Perception (IMSPAC). *Journal of the American Academy of*
997 *Audiology*, 14(02), 072-083. <https://doi.org/10.3766/jaaa.14.2.3>
- 998 Krull, V., Choi, S., Kirk, K. I., Prusick, L., & French, B. (2010). Lexical effects on spoken-word
999 recognition in children with normal hearing. *Ear and Hearing*, 31(1), 102-114.
1000 <https://doi.org/10.1097/AUD.0b013e3181b7892f>

- 1001 Leibold, L. J., & Buss, E. (2013). Children's Identification of Consonants in a Speech-Shaped Noise or a
1002 Two-Talker Masker. *Journal of Speech, Language, and Hearing Research, 56*(4), 1144-1155.
1003 [https://doi.org/10.1044/1092-4388\(2012/12-0011\)](https://doi.org/10.1044/1092-4388(2012/12-0011))
- 1004 Leibold, L. J., & Buss, E. (2019). Masked Speech Recognition in School-Age Children. *Frontiers in*
1005 *Psychology, 10*, 1981. <https://doi.org/10.3389/fpsyg.2019.01981>
- 1006 Leibold, L. J., Buss, E., & Calandruccio, L. (2018). Developmental Effects in Masking Release for
1007 Speech-in-Speech Perception Due to a Target/Masker Sex Mismatch. *Ear & Hearing, 39*(5),
1008 935-945. <https://doi.org/10.1097/AUD.0000000000000554>
- 1009 Leibold, L. J., Buss, E., Miller, M. K., Cowan, T., McCreery, R. W., Oleson, J., Rodriguez, B., &
1010 Calandruccio, L. (2024). Development of the Children's English and Spanish Speech
1011 Recognition Test : Psychometric Properties, Feasibility, Reliability, and Normative Data. *Ear &*
1012 *Hearing, 45*(4), 860-877. <https://doi.org/10.1097/AUD.0000000000001480>
- 1013 Lenth, R. V. (2021). *emmeans : Estimated Marginal Means, aka Least-Squares Means*.
1014 <https://CRAN.R-project.org/package=emmeans>
- 1015 Lévêque, Y., Teyssier, P., Bouchet, P., Bigand, E., Caclin, A., & Tillmann, B. (2018). Musical emotions in
1016 congenital amusia : Impaired recognition, but preserved emotional intensity.
1017 *Neuropsychology, 32*(7), 880-894. <https://doi.org/10.1037/neu0000461>
- 1018 Lewis, H. D., Benignus, V. A., Muller, K. E., Malott, C. M., & Barton, C. N. (1988). Babble and Random-
1019 Noise Masking of Speech in High and Low Context Cue Conditions. *Journal of Speech,*
1020 *Language, and Hearing Research, 31*(1), 108-114. <https://doi.org/10.1044/jshr.3101.108>
- 1021 Linares, D., & López-Moliner, J. (2016). quickpsy : An R Package to Fit Psychometric Functions for
1022 Multiple Groups. *The R Journal, 8*(1), 122-131. <https://doi.org/10.32614/RJ-2016-008>
- 1023 Litovsky, R. Y. (2005). Speech intelligibility and spatial release from masking in young children. *The*
1024 *Journal of the Acoustical Society of America, 117*(5), 3091-3099.
1025 <https://doi.org/10.1121/1.1873913>

- 1026 Lo, S., & Andrews, S. (2015). To transform or not to transform : Using generalized linear mixed
1027 models to analyse reaction time data. *Frontiers in Psychology, 6*, 1171.
1028 <https://doi.org/10.3389/fpsyg.2015.01171>
- 1029 Luce, P. A., & Pisoni, D. B. (1998). Recognizing spoken words : The neighborhood activation model.
1030 *Ear and hearing, 19*(1), 1.
- 1031 Lüdtke, D. (2019). *esc : Effect Size Computation for Meta Analysis (Version 0.5.1)*.
1032 <https://doi.org/10.5281/zenodo.1249218>
- 1033 Majerus, S., & Cowan, N. (2016). The Nature of Verbal Short-Term Impairment in Dyslexia : The
1034 Importance of Serial Order. *Frontiers in Psychology, 7*.
1035 <https://doi.org/10.3389/fpsyg.2016.01522>
- 1036 Marslen-Wilson, W. D. (1987). Functional parallelism in spoken word-recognition. *Cognition, 25*(1-2),
1037 71-102.
- 1038 Marslen-Wilson, W. D., & Zwitserlood, P. (1989). Accessing spoken words : The importance of word
1039 onsets. *Journal of Experimental Psychology: Human perception and performance, 15*(3), 576.
- 1040 McClelland, J. L., & Elman, J. L. (1986). The TRACE model of speech perception. *Cognitive Psychology,*
1041 *18*(1), 1-86.
- 1042 McCreery, R. W., Miller, M. K., Buss, E., & Leibold, L. J. (2020). Cognitive and Linguistic Contributions
1043 to Masked Speech Recognition in Children. *Journal of Speech, Language, and Hearing*
1044 *Research, 63*(10), 3525-3538. https://doi.org/10.1044/2020_JSLHR-20-00030
- 1045 Mendel, L. L. (2008). Current considerations in pediatric speech audiometry. *International Journal of*
1046 *Audiology, 47*(9), 546-553. <https://doi.org/10.1080/14992020802252261>
- 1047 Moulin, A., Bernard, A., Tordella, L., Vergne, J., Gisbert, A., Martin, C., & Richard, C. (2017). Variability
1048 of word discrimination scores in clinical practice and consequences on their sensitivity to
1049 hearing loss. *European Archives of Oto-Rhino-Laryngology: Official Journal of the European*
1050 *Federation of Oto-Rhino-Laryngological Societies (EUFOS): Affiliated with the German Society*

- 1051 *for Oto-Rhino-Laryngology - Head and Neck Surgery*, 274(5), 2117-2124.
- 1052 <https://doi.org/10.1007/s00405-016-4439-x>
- 1053 Moulin, A., Favier, C., Richard, C., & Farget, V. (2013). French Babble Noises for Speech Perception
- 1054 Tests. Development and Spectro-temporal analysis. *Annales D'oto-Laryngologie Et De*
- 1055 *Chirurgie Cervico Faciale*.
- 1056 Moulin, A., & Richard, C. (2015). Lexical Influences on Spoken Spondaic Word Recognition in Hearing-
- 1057 Impaired Patients. *Auditory Cognitive Neuroscience*, 476.
- 1058 <https://doi.org/10.3389/fnins.2015.00476>
- 1059 Munro, K. J., & Lutman, M. E. (2005). The influence of visual feedback on closed-set word test
- 1060 performance over time. *International Journal of Audiology*, 44(12), 701-705.
- 1061 <https://doi.org/10.1080/14992020500271795>
- 1062 New, B., Brysbaert, M., Veronis, J., & Pallier, C. (2007). The use of film subtitles to estimate word
- 1063 frequencies. *Applied Psycholinguistics*, 28(4), 661-677.
- 1064 <https://doi.org/10.1017/S014271640707035X>
- 1065 New, B., Pallier, C., Brysbaert, M., & Ferrand, L. (2004). Lexique 2 : A new French lexical database.
- 1066 *Behavior Research Methods, Instruments, & Computers*, 36(3), 516-524.
- 1067 <https://doi.org/10.3758/BF03195598>
- 1068 Plaza, M., Cohen, H., & Chevrie-Muller, C. (2002). Oral Language Deficits in Dyslexic Children :
- 1069 Weaknesses in Working Memory and Verbal Planning. *Brain and Cognition*, 48(2-3), 505-512.
- 1070 <https://doi.org/10.1006/brcg.2001.1407>
- 1071 Pujol, J., Soriano-Mas, C., Ortiz, H., Sebastian-Galles, N., Losilla, J. M., & Deus, J. (2006). Myelination
- 1072 of language-related areas in the developing brain. *Neurology*, 66(3), 339-343.
- 1073 <https://doi.org/10.1212/01.wnl.0000201049.66073.8d>
- 1074 R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. R Foundation for
- 1075 Statistical Computing. <https://www.R-project.org/>

- 1076 Rigler, H., Farris-Trimble, A., Greiner, L., Walker, J., Tomblin, J. B., & McMurray, B. (2015). The slow
1077 developmental time course of real-time spoken word recognition. *Developmental*
1078 *Psychology, 51*, 1690-1703. <https://doi.org/10.1037/dev0000044>
- 1079 Rosen, S., Souza, P., Ekelund, C., & Majeed, A. A. (2013). Listening to speech in a background of other
1080 talkers : Effects of talker number and noise vocoding. *The Journal of the Acoustical Society of*
1081 *America, 133*(4), 2431-2443. <https://doi.org/10.1121/1.4794379>
- 1082 Simpson, S. A., & Cooke, M. (2005). Consonant identification in N-talker babble is a nonmonotonic
1083 function of N. *The Journal of the Acoustical Society of America, 118*(5), 2775-2778.
1084 <https://doi.org/10.1121/1.2062650>
- 1085 Skarżyński, H., & Piotrowska, A. (2012). Screening for pre-school and school-age hearing problems :
1086 European Consensus Statement. *International Journal of Pediatric Otorhinolaryngology,*
1087 *76*(1), 120-121. <https://doi.org/10.1016/j.ijporl.2011.10.016>
- 1088 Skarzynski, P. H., Włodarczyk, A. W., Kochanek, K., Pilka, A., Jędrzejczak, W., Olszewski, L., Bruski, L.,
1089 Niedzielski, A., & Skarzynski, H. (2015). Central auditory processing disorder (CAPD) tests in a
1090 school-age hearing screening programme – analysis of 76,429 children. *Annals of Agricultural*
1091 *and Environmental Medicine, 22*(1), 90-95. <https://doi.org/10.5604/12321966.1141375>
- 1092 Snowling, M. J., & Hulme, C. (1994). The development of phonological skills. *Philosophical*
1093 *Transactions of the Royal Society of London. Series B: Biological Sciences, 346*(1315), 21-27.
1094 <https://doi.org/10.1098/rstb.1994.0124>
- 1095 Sommers, M. S., Kirk, K. I., & Pisoni, D. B. (1997). Some considerations in evaluating spoken word
1096 recognition by normal-hearing, noise-masked normal-hearing, and cochlear implant listeners.
1097 I: The effects of response format. *Ear and Hearing, 18*(2), 89.
- 1098 Stahl, S. A., & Murray, B. A. (1994). Defining phonological awareness and its relationship to early
1099 reading. *Journal of Educational Psychology, 86*(2), 221-234. <https://doi.org/10.1037/0022->
1100 [0663.86.2.221](https://doi.org/10.1037/0022-0663.86.2.221)

- 1101 Stekić, K., Ilić, O., Ković, V., & Savić, A. M. (2023). ERP Indicators of Phonological Awareness
1102 Development in Children : A Systematic Review. *Brain Sciences*, 13(2), 290.
1103 <https://doi.org/10.3390/brainsci13020290>
- 1104 Thillay, A., Roux, S., Gissot, V., Carreau-Martin, I., Knight, R. T., Bonnet-Brilhault, F., & Bidet-Caulet, A.
1105 (2015). Sustained attention and prediction : Distinct brain maturation trajectories during
1106 adolescence. *Frontiers in Human Neuroscience*, 9, 519.
- 1107 Tillmann, B., Albouy, P., Caclin, A., & Bigand, E. (2014). Musical familiarity in congenital amusia :
1108 Evidence from a gating paradigm. *Cortex*, 59, 84-94.
1109 <https://doi.org/10.1016/j.cortex.2014.07.012>
- 1110 Vickers, D. A., Moore, B. C. J., Majeed, A., Stephenson, N., Alferaih, H., Baer, T., & Marriage, J. E.
1111 (2018). Closed-Set Speech Discrimination Tests for Assessing Young Children. *Ear & Hearing*,
1112 39(1), 32-41. <https://doi.org/10.1097/AUD.0000000000000528>
- 1113 Watson, B. U., & Miller, T. K. (1993). Auditory Perception, Phonological Processing, and Reading
1114 Ability/Disability. *Journal of Speech, Language, and Hearing Research*, 36(4), 850-863.
1115 <https://doi.org/10.1044/jshr.3604.850>
- 1116 Werner, L., Fay, R. R., & Popper, A. N. (Éds.). (2012). *Human Auditory Development* (Vol. 42). Springer
1117 New York. <https://doi.org/10.1007/978-1-4614-1421-6>
- 1118 Wetzel, N., Widmann, A., & Schröger, E. (2009). The cognitive control of distraction by novelty in
1119 children aged 7–8 and adults. *Psychophysiology*, 46(3), 607-616.
- 1120 Wightman, F., & Allen, P. (1992). Individual differences in auditory capability among preschool
1121 children. In *Developmental psychoacoustics* (p. 113-133). American Psychological
1122 Association. <https://doi.org/10.1037/10119-004>
- 1123 Wilson, R. H., Farmer, N. M., Gandhi, A., Shelburne, E., & Weaver, J. (2010). Normative Data for the
1124 Words-in-Noise Test for 6- to 12-Year-Old Children. *Journal of Speech, Language, and*
1125 *Hearing Research*, 53(5), 1111-1121. [https://doi.org/10.1044/1092-4388\(2010/09-0270\)](https://doi.org/10.1044/1092-4388(2010/09-0270))

- 1126 Wilson, W. J., & Arnott, W. (2013). Using Different Criteria to Diagnose (Central) Auditory Processing
1127 Disorder : How Big a Difference Does It Make? *Journal of Speech, Language, and Hearing*
1128 *Research, 56*(1), 63-70. [https://doi.org/10.1044/1092-4388\(2012/11-0352\)](https://doi.org/10.1044/1092-4388(2012/11-0352))
- 1129 Wright, I., Waterman, M., Prescott, H., & Murdoch-Eaton, D. (2003). A new Stroop-like measure of
1130 inhibitory function development : Typical developmental trends. *Journal of Child Psychology*
1131 *and Psychiatry, 44*(4), 561-575.
- 1132 Wróblewski, M., Lewis, D. E., Valente, D. L., & Stelmachowicz, P. G. (2012). Effects of reverberation
1133 on speech recognition in stationary and modulated noise by school-aged children and young
1134 adults. *Ear and hearing, 33*(6), 731-744. <https://doi.org/10.1097/AUD.0b013e31825aead>
- 1135 Ziegler, J. C., Pech-Georgel, C., George, F., Alario, F.-X., & Lorenzi, C. (2005). Deficits in speech
1136 perception predict language learning impairment. *Proceedings of the National Academy of*
1137 *Sciences, 102*(39), 14110-14115. <https://doi.org/10.1073/pnas.0504446102>
- 1138 Ziegler, J. C., Pech-Georgel, C., George, F., & Lorenzi, C. (2009). Speech-perception-in-noise deficits in
1139 dyslexia. *Developmental Science, 12*(5), 732-745. [https://doi.org/10.1111/j.1467-](https://doi.org/10.1111/j.1467-7687.2009.00817.x)
1140 [7687.2009.00817.x](https://doi.org/10.1111/j.1467-7687.2009.00817.x)
- 1141