

LARGE-SCALE LABORATORY EXPERIMENTS OF THE GIBRALTAR EXCHANGE FLOW, INCLUDING GULF OF CADIZ AND ALBORAN SEA

Maria Eletta Negretti, Axel Tassigny, Louis Gostiaux, Lucie Bordois, Xavier

J. Carton

► To cite this version:

Maria Eletta Negretti, Axel Tassigny, Louis Gostiaux, Lucie Bordois, Xavier J. Carton. LARGE-SCALE LABORATORY EXPERIMENTS OF THE GIBRALTAR EXCHANGE FLOW, INCLUD-ING GULF OF CADIZ AND ALBORAN SEA. CIESM - Mediterranean Science Commission, Oct 2024, Palerme, Italy. hal-04770477

HAL Id: hal-04770477 https://hal.science/hal-04770477v1

Submitted on 7 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LARGE-SCALE LABORATORY EXPERIMENTS OF THE GIBRALTAR EXCHANGE FLOW, INCLUDING GULF OF CADIZ AND ALBORAN SEA

Maria Eletta Negretti ^{1*}, Axel Tassigny ¹, Louis Gostiaux ², Lucie Bordois ³ and Xavier Carton ⁴

¹ CNRS, LEGI/CORIOLIS, Grenoble, 38000, France - eletta.negretti@legi.cnrs.fr

² Univ Lyon, Ecole Centrale de Lyon, CNRS, Univ Claude Bernard Lyon 1, INSA Lyon, LMFA, UMR5509, 69130, Ecully,

France

³ Service Hydrographique et Océanographique de la Marine (SHOM), Brest, France

⁴ University Brest, CNRS, Ifremer, IRD, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, Plouzané, France

Abstract

Results from large scale laboratory experiments of a realistic physical model of the Gibraltar Strait and adjacent Gulf of Cadiz and Alboran Sea are presented. The physical model includes all forcings: the baroclinic exchange flow, the barotropic tide, the realistic bathymetry and the Earth's rotation. Results show that experiments are able to model and correctly reproduce such a large area of 250km x 150km with different dynamics, including turbulent mixing at the hydraulic control areas, the Mediterranean outflow propagation in the Gulf of Cadiz and the formation of internal solitary waves propagating in the Alboran Sea, revealing intriguing insights in the reach dynamics of the area.

Keywords: Gibraltar Strait, Straits and channels

Results from large scale laboratory experiments of a realistic physical model of the Gibraltar Strait and adjacent Gulf of Cadiz and Alboran Sea are presented, including all forcings: the baroclinic exchange flow, the barotropic tide, the realistic bathymetry and the Earth's rotation. The experiments were conducted at the Coriolis platform at LEGI, Grenoble (France), the largest rotating tank in the world. The baroclinic exchange flow was realized using a lock-exchange configuration with two different salinities at each side. Scaled semi-diurnal (M2) tides for both spring tide (ST) and neap tide (NT) regimes were generated, including the transient with the superposition of the M2+S2 tide. The experimental duration was of 13 rotational days. Similarity has been achieved for the Froude, Rossby, Burger numbers, and the Reynolds number is 5,000. Variation parameters in the experiments were the density difference and the different barotropic flows. The results show that experiments are able to model such a large area of 250km x 150km with different dynamics, including turbulent mixing at the hydraulic control areas, the Mediterranean outflow propagation in the Gulf of Cadiz and the formation of internal solitary waves propagating in the Alboran Sea.



Fig. 1. Instantaneous PIV/PLIF image downstream CS (0,-3). Vertical scale 1/2,500, horizontal scale 1/25,000, velocity scale 1/20, density scale 6.25. Bottom - Fluorescent dye highlighting the propagation of the MOW in the different channels of the Gulf of Cadiz.

Measurements techniques were based on PIV in all regions at different horizontal and vertical sections, including combined PIV/PLIF in high mixing regions (Camarinal/Espartell Sills) which deliver the full turbulent fluxes crucial for parametrization of small-scale turbulent processes (see Figure 1). They appear to be very different during the different phase of the tide. Intrusive devices (ADV, ADV profilers, SSH, CTD) have been used as well, especially to track the Mediterranean outflow in the Gulf of Cadiz. The dense water transport reveals to be crucially determined by the bathymetric features (see Figure 1).



Fig. 2. Hovmöller diagram of the Mediterranean interface during two tidal cycles highlighting the internal tide, the soliton and small-scale instabilities propagating in the Alboran Sea. (Bottom) Dye image with the evident soliton with superposed characteristic velocity field. Speed: 8cm/s, amplitude: 2cm, wavelength 20cm. Scales as in Fig.1.

Solitons are produced during spring tide (ST) at CS and propagate toward the Alboran Sea (see figure 2), at each tidal inversion (West-Est). Preliminary results suggest hence that they are not related to the relaxation of the internal hydraulic jump downstream of Camarinal Sill, but rather from an internal bore generated by the internal tide at CS at the tidal inversion (West-Est).

References

1 - Negretti ME, Tucciarone FL and Wirth A (2021) Intruding gravity currents and recirculation in a rotating frame: Laboratory experiments Phys. Fluids 10.1063/5.0058629.

2 - J.-B. Roustan, L. Bordois, F. Dumas, F. Auclair, and X. Carton (2023) In situ observations of the small-scale dynamics at Camarinal Sill-Strait of Gibraltar, J. Geophys. Res. Oceans 128, e2023JC019738

3 - Hilt, M., Roblou, L., Nguyen, C., Marchesiello, P., Lemarié, F., Jullien, S., Dumas, F., Debreu, L., Capet, X., Bordois, L., Benshila, R., Auclair, F., 2020. Numerical modeling of hydraulic control, solitary waves and primary instabilities in the Strait of Gibraltar. Ocean Modelling 155, 101642. https://doi.org/10.1016/j.ocemod.2020.101642