



HAL
open science

Mindfulness Meditation–Based Interventions in Parents of Autistic Children: a Systematic Review of Effects on Children

Jérôme Lichtlé, Marco Sperduti, Jean-Louis Monestès, Emilie Cappe

► To cite this version:

Jérôme Lichtlé, Marco Sperduti, Jean-Louis Monestès, Emilie Cappe. Mindfulness Meditation–Based Interventions in Parents of Autistic Children: a Systematic Review of Effects on Children. Review Journal of Autism and Developmental Disorders, 2024, 10.1007/s40489-024-00483-w . hal-04774170

HAL Id: hal-04774170

<https://hal.science/hal-04774170v1>

Submitted on 19 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Review Journal of Autism and Developmental Disorders

Mindfulness meditation–based interventions in parents of autistic children: A systematic review of effects on children --Manuscript Draft--

Manuscript Number:	RJAD-D-24-00058R2
Full Title:	Mindfulness meditation–based interventions in parents of autistic children: A systematic review of effects on children
Article Type:	Review Paper
Funding Information:	
Abstract:	<p>The quality of life (QoL) of autistic children and their parents is a significant clinical concern and an important focus of research in the field. The well-being of autistic children and their parents can be improved through interventions based on mindfulness meditation. However, when implemented with autistic children, mindfulness-based interventions (MBIs) have certain limitations. These limitations can be overcome through the indirect positive effects on autistic children resulting from their parents' engagement in MBIs. We conducted a systematic review of the literature in this field, targeting studies on the effects of MBIs on autistic children through interventions being offered exclusively to their parents. A total of 14 studies were identified for inclusion in the review. Overall, autistic children showed a significant reduction in challenging behaviours following the implementation of MBIs aimed at their parents. Given this reduction, MBIs for parents of autistic children could be a viable option for improving the QoL of families affected by autism.</p>
Corresponding Author:	Emilie Cappe, Ph.D. Université Paris Cité Faculté Sociétés et Humanités: Université Paris Cité Faculté Sociétés et Humanités Boulogne-Billancourt, FRANCE
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Université Paris Cité Faculté Sociétés et Humanités: Université Paris Cité Faculté Sociétés et Humanités
Corresponding Author's Secondary Institution:	
First Author:	Jérôme Lichtlé
First Author Secondary Information:	
Order of Authors:	Jérôme Lichtlé Marco Sperduti Jean-Louis Monestès Emilie Cappe, Ph.D.
Order of Authors Secondary Information:	
Author Comments:	
Suggested Reviewers:	

**Mindfulness meditation-based interventions in parents of autistic children:
A systematic review of effects on children**

Jérôme Lichtlé¹ (jeromelichtle@gmail.com)  <https://orcid.org/0000-0001-5147-559X>

Marco Sperduti² (marco.sperduti@u-paris.fr)

Jean-Louis Monestès³ (jean-louis.monestes@univ-grenoble-alpes.fr)

Emilie Cappe^{1,4} (emilie.cappe@u-paris.fr)  <https://orcid.org/0000-0002-4792-8653>

¹ Université Paris Cité, Laboratoire de Psychopathologie et Processus de Santé, F-92100
Boulogne-Billancourt, France

² Université Paris Cité, LMC2, F-92100 Boulogne-Billancourt, France

³ LIP/PC2S Lab, Univ. Grenoble Alpes, France

⁴ Institut universitaire de France (IUF)

Abstract

The quality of life (QoL) of autistic children and their parents is a significant clinical concern and an important focus of research in the field. The well-being of autistic children and their parents can be improved through interventions based on mindfulness meditation. However, when implemented with autistic children, mindfulness-based interventions (MBIs) have certain limitations. These limitations can be overcome through the indirect positive effects on autistic children resulting from their parents' engagement in MBIs. We conducted a systematic review of the literature in this field, targeting studies on the effects of MBIs on autistic children through interventions being offered exclusively to their parents. A total of 14 studies were identified for inclusion in the review. Overall, autistic children showed a significant reduction in challenging behaviours following the implementation of MBIs aimed at their parents. Given this reduction, MBIs for parents of autistic children could be a viable option for improving the QoL of families affected by autism.

Keywords: autism, parent, mindfulness, child outcomes, challenging behaviours, systematic review

Correspondance concerning this article should be addressed to Emilie Cappe,
Laboratoire de Psychopathologie et Processus de Santé EA4057, Institut de Psychologie,
Université de Paris, 71 avenue Edouard Vaillant, 92100 Boulogne-Billancourt, France.

Email: emilie.cappe@u-paris.fr

Mindfulness meditation–based interventions in parents of autistic children:

A systematic review of effects on children

Abstract

The quality of life (QoL) of autistic children and their parents is a significant clinical concern and an important focus of research in the field. The well-being of autistic children and their parents can be improved through interventions based on mindfulness meditation. However, when implemented with autistic children, mindfulness-based interventions (MBIs) have certain limitations. These limitations can be overcome through the indirect positive effects on autistic children resulting from their parents' engagement in MBIs. We conducted a systematic review of the literature in this field, targeting studies on the effects of MBIs on autistic children through interventions being offered exclusively to their parents. A total of 14 studies were identified for inclusion in the review. Overall, autistic children showed a significant reduction in challenging behaviours following the implementation of MBIs aimed at their parents. Given this reduction, MBIs for parents of autistic children could be a viable option for improving the QoL of families affected by autism.

Keywords: autism, parent, mindfulness, child outcomes, challenging behaviours, systematic review

Introduction

1
2
3
4
5 Regardless of age, the quality of life (QoL) of autistic people is lower than that of people
6
7 with typical development (Arias et al., 2018; van Heijst & Geurts, 2015). QoL is a
8
9 multidimensional and subjective concept (Schalock, 2004), encompassing several dimensions,
10
11 including physical, social, and psychological functioning (World Health Organization [WHO],
12
13 1946). [Autism](#) can also impact the well-being of the family (Dückert et al., 2023; National
14
15 Institute for Health and Care Excellence [NICE], 2013). For example, parents of autistic
16
17 children have higher stress scores (Hayes & Watson, 2013), and lower QoL levels
18
19 (Vasilopoulou & Nisbet, 2016) compared to parents of typically developing children. However,
20
21 the QoL (Vernhet et al., 2022) and stress levels (Hastings et al., 2005; McStay et al., 2014) of
22
23 parents of autistic children are less dependent on the fundamental characteristics of their child's
24
25 autism than on its clinical specifiers, such as mental, neurodevelopmental, or behavioural
26
27 conditions (APA, 2013). For example, research has identified an association between autistic
28
29 children's behavioural and emotional problems and their parent's psychological distress (Yorke
30
31 et al., 2018). Similarly, the QoL of autistic people is less dependent on the fundamental
32
33 characteristics of autism than on its clinical specifiers (van Heijst & Geurts, 2015; Kim &
34
35 Bottema-Beutel, 2019).

36
37
38 Mindfulness-based interventions (MBIs) are interventions that teach various meditation
39
40 techniques to continuously redirect attention to the present moment (Kabat Zinn, 2003). MBIs
41
42 are the interventions of choice for improving QoL (Jones & Drummond et al., 2023). Despite
43
44 the lack of evidence for interventions aimed at reducing the burden on parents with autistic
45
46 children (Dückert et al., 2023), MBIs have been shown to improve QoL for these parents
47
48 (Cachia et al., 2016a; Juvín et al., 2021). Furthermore, intervention programmes that include
49
50 mindfulness meditation practices for parents of autistic children are more effective in reducing
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 their stress levels than programmes that do not include such practices (Anonymous et al., 2020;
2 Li et al., 2024). MBIs can be implemented with autistic people and have been shown to improve
3 their QoL (Benevides et al., 2020; Byrne & O’Mahony, 2020; Cachia et al., 2016b; Menezes et
4 al., 2020; Semple, 2019). Given the heterogeneity of the autistic phenotype and co-occurring
5 conditions, MBIs aimed directly at autistic people raise questions of adaptability (Beck,
6 Connor, et al., 2020; Poquérusse et al., 2021). For example, autistic participants who are offered
7 these interventions are sometimes excluded from studies for comorbid intellectual disability
8 diagnoses (Beck, Greco, et al., 2020; Spek et al., 2013).

19 Some research has demonstrated that MBIs targeted at caregivers can have indirect
20 effects on autistic people. Singh et al. (2004) and Singh, Lancioni, Winton, Curtis, et al. (2006),
21 for example, showed that a predisposition to mindfulness among carers was positively
22 associated with the well-being of the disabled person they support. In the same vein, Harmony
23 and Woodard (2020) showed that mindfulness training for staff members in schools with
24 autistic pupils was negatively correlated with pupils’ challenging behaviours. [Challenging
25 behaviours are defined as a range of behaviours that people with severe learning disabilities
26 may display to get their needs met, such as hurting others, destructive behaviours or self-
27 injurious behaviours, which](#) can have a negative impact on the autistic person’s well-being
28 (Carter Leno et al., 2019). The literature has identified a high prevalence rate of these
29 behaviours among autistic people (Hattier et al., 2011; Matson & Nebel-Schwalm, 2007).
30 Several studies have also found that children’s challenging behaviours is negatively associated
31 with parents’ ability to be present in their relationship with their autistic child, otherwise known
32 as “dispositional mindfulness” (Aydin, 2023; Beer et al., 2013; Chan et al., 2022; Raulston et
33 al., 2021). This association has not been found with other parenting practices (Suvarna et al.,
34 2024). Utilising emotional co-regulation strategies (Ting & Weiss, 2017) could also help
35 parents mitigate challenging behaviours in their autistic children. Finally, studies have

1 identified a significant positive association between the emotional openness (or acceptance
2 skills) of parents of preschool-aged autistic children and their child's QoL (Anonymous et al.,
3
4 2023), which is considered the "gold standard" outcome for autism interventions (Ikeda et al.,
5
6 2014).
7

8
9
10 Ultimately, the adaptability challenges associated with implementing MBIs with autistic
11 individuals could be partially mitigated by the indirect effects of implementing MBIs with their
12 parents. The aim of the present research was therefore to (1) identify studies examining the
13 effects of MBIs on autistic children when these interventions are offered exclusively to their
14 parents and (2) explore the variables examined in autistic individuals, especially the clinical
15 specifiers. To the best of our knowledge, there are no other systematic reviews addressing this
16 topic.
17
18
19
20
21
22
23
24
25

26 **Methods**

27
28
29 A systematic literature review was conducted in accordance with the Preferred
30 Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Moher et
31 al., 2009; Page et al., 2021). No meta-analysis was planned.
32
33
34

35 *Information sources and search strategies*

36
37
38
39 An electronic search was conducted for the period 2006 to April 2024 by one of the
40 authors (JL). To our knowledge, Singh, Lancioni, Winton, Fisher, et al. (2006) were the first
41 to examine the indirect effects of MBIs offered to parents on their autistic children; therefore,
42 the search was limited to the period between 2006 and April 2024. A comprehensive query of
43 PubMed, PsycINFO via the EBSCOhost access provider, Education Resources Information
44 Center (ERIC), PubPsych, and the Science Direct databases was conducted. We used the
45 following keywords: autism, ASD, mindfulness, mindful, parent, and caregiver. These
46 keywords were combined with Boolean operators and used with truncation; the algorithm
47 used in all but the Science Direct database was: (autism* OR ASD) AND (parent* OR
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 caregiver) AND mindful*. Because Boolean operators and truncation are not available in
2 Science Direct, the keywords entered into this database were “ASD”, “Mindfulness”, and
3 “parent”. We applied this query to titles and abstracts in PubMed and PsycINFO, and full text
4 in ERIC, PubPsych, and Science Direct. The reference lists of the articles reviewed were also
5 consulted to identify further studies.
6
7
8
9
10

11 *Eligibility criteria*

12 We selected studies published in English or French in peer-reviewed scientific journals.
13
14 The studies had to propose an MBI for parents of autistic children, utilising a protocol that
15 included either a formal or informal mindfulness meditation practice. The former refers to a
16 period of time, whether short or extended, dedicated to a specific meditation practice (e.g.
17 breathing, sitting, body scan), while the latter refers to a mindfulness practice that can be
18 flexible and integrated into everyday activities (e.g. when eating, doing the dishes, playing with
19 one’s child). This meditation could be the focal point of the intervention or an additional
20 component of a broader programme that includes other interventions. Children were only
21 included if they had been diagnosed with autism according to international classification
22 criteria: autism spectrum disorder or pervasive developmental disorder (ICD-10, DSM-IV-TR
23 or DSM-5). The diagnosis had to be either explicitly confirmed during the research procedure,
24 taken from the child’s medical records, or obtained from other reliable sources (e.g. parents’
25 reports). Included studies also had to report quantitative or qualitative measures of the indirect
26 effects of MBI on the child, with no age limit, given that the well-being of parents of autistic
27 adults is as much a concern as that of parents of autistic children (Herrema et al., 2017). Studies
28 that offered MBI to both parents and children were excluded, as this study targeted the effects
29 of mindfulness meditation practised solely by the parent.
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Study selection process

After removing duplicates via the Excel software program, one of the authors (JL) reviewed the titles and abstracts of the studies identified through the database search to determine which ones met the inclusion criteria. The full texts of these studies were then reviewed independently by two of the authors (JL and MS) to confirm whether they met the inclusion criteria. This resulted in a 100% agreement rate.

Data collection process

The study characteristics and results were extracted by the lead author (JL) and verified by two co-authors (EC and MS). Omissions or discrepancies were resolved by consensus between these three authors. Data collected included the name of the first author, year of publication, country, characteristics of participating parents and their children, details of the MBI and any additional interventions, study design, measures concerning the child, any follow-up, main results concerning the effect of MBI on the child, and methodological quality and level of evidence.

Assessing the methodological quality and level of evidence of the studies

The methodological quality and level of evidence of each study were assessed independently by two of the authors (JL and MS). This resulted in agreement rates of 83% and 62% for methodological quality and level of evidence, respectively. Any discrepancies were discussed with a third author (EC) to reach a consensus.

The STrengthening the Reporting of OBservational Studies in Epidemiology (STROBE) guidelines (von Elm et al., 2008) were used to assess the methodological quality of each study, ensuring a clear presentation of what was planned, done, and found. STROBE reporting reveals the strengths and weaknesses of studies and facilitates sound interpretation and application of the results. The STROBE statement includes a checklist of 22 items related to the title, abstract, introduction, methods, results, and discussion sections of articles. The

1 scores were assigned as follows: 1 = all the item's criteria were met, 0.5 = the criteria were
2 partially met, and 0 = none of the criteria were found in the study. The overall evaluation of
3 each study was obtained by transforming the total score into the percentage of criteria met.
4
5
6
7 Studies with a quality assessment below 50% should be excluded (da Costa et al., 2011).
8

9
10 The level of evidence was determined using the hierarchy of evidence described in the
11 National Health and Medical Research Council (NHMRC; 1999) guidelines, with six levels
12 ranging from the highest level of evidence (Level I) to the lowest level of evidence (Level IV):
13
14 Level I (evidence obtained from a systematic review of all relevant randomised controlled trials
15 [RCTs]), Level II (evidence obtained from at least one properly designed RCT), Level III-1
16 (evidence obtained from well-designed pseudo-RCTs), Level III-2 (evidence obtained from
17 comparative studies with concurrent controls and allocation not randomised, case-control
18 studies, or interrupted time series with a control group), Level III-3 (evidence obtained from
19 comparative studies with historical control, two or more single-arm studies, or interrupted time
20 series without a parallel control group), and Level IV (evidence obtained from case series, either
21 post-test or pre-test and post-test). The most appropriate study design to answer the intervention
22 question is Level II evidence. Level I studies are systematic reviews of the appropriate Level II
23 studies. Study designs that are progressively less robust for answering the intervention question
24 are shown at Levels III and IV.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 *Risk of bias inherent in randomised controlled trials*

44
45 Two of the authors (JL and JLM) independently assessed the risk of bias inherent in
46 each RCT using the risk of bias tool recommended by Munder and Barth (2017), which includes
47 seven domains of bias: (1) random sequence generation, (2) allocation concealment, (3)
48 performance bias, (4) detection bias, (5) attrition bias, (6) reporting bias, and (7) treatment
49 implementation. For each domain, the risk of bias was assessed as low (+), high (-) or uncertain
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results

Selection of studies

The PRISMA flow chart for identifying the studies is shown in Figure 1. Of the 607 abstracts identified after removing duplicates, 549 were excluded and 58 studies were evaluated in their entirety, including three additional references found through other sources and one study from updated database searches (Rojas-Torres et al. 2023). Of these 58, 44 did not meet the inclusion criteria and were excluded; therefore, 14 studies were retained for inclusion. The studies by Chan and Neece (2018) and Neece et al. (2019) were based on results obtained from the same sample, so they were combined for reporting and discussion purposes. A subsample of participants from the Chan and Neece (2018) study was also used in the Neece (2014) study, but we were unable to treat them separately.

Characteristics of included studies

All the studies included in this review measured the effects of MBIs on both the child and their parents, except Weitlauf et al. (2022), which targeted only child-related variables. Regarding the parent variables, most of the studies showed a significant reduction in parents' perceived stress following intervention (Andrews et al., 2022; Chan & Neece, 2018; Jones et al., 2018; Neece, 2014; Neece et al., 2019; Raulston et al., 2019; Rojas-Torres et al. 2023; Schwartzman et al., 2022; Singh et al., 2014; Singh et al., 2019; Singh et al., 2021). The characteristics of the included studies are presented in Table 1.

Four of the 14 studies selected for this review included not only parents of autistic children but also parents of children with other diagnoses. Neece (2014) and Neece et al. (2019) did not specify these other diagnoses, indicating only that the children had developmental delays. In addition to autism, Chan and Neece (2018) reported the following diagnoses, but without specifying the number of children involved: Down's syndrome, Prader-Willi syndrome, Fragile X syndrome, intellectual disability, learning disability, language delay,

1 cerebral palsy, and microcephaly. Jones et al. (2018) included 16 autistic children, one child
2 with Down's syndrome, two children with cerebral palsy, and two children with intellectual
3 disability. The autistic children in Singh et al. (2014) had all been diagnosed with Asperger's
4 (autism without intellectual and language impairment).
5
6
7

8
9 The mean age of the children included in the studies ranged from 2.3 years (Weitlauf et
10 al., 2022) to 16.7 years (Singh et al., 2014). For 10 of the 14 studies, the mean age of the children
11 was less than 7 years. The remaining four studies included children with an average age of over
12 10 years (Jones et al., 2018; Singh et al., 2014; Singh et al., 2019; Singh et al., 2021).
13
14
15
16
17
18

19 In total, there were 636 parents across the 14 studies. The number of parents included
20 in the studies ranged from three (Andrews et al., 2022; Raulston et al., 2019; Singh, Lancioni,
21 Winton, Fisher, et al., 2006; Singh et al., 2014) to 80 (Chan & Neece, 2018; Neece et al., 2019).
22 Five studies included only mothers (Raulston et al., 2019; Singh, Lancioni, Winton, Fisher, et
23 al., 2006; Singh et al., 2014; Singh et al., 2019; Singh et al., 2021). The remaining studies
24 included a smaller number of fathers compared to mothers. McGregor et al. (2020) did not
25 specify the ratio of mothers to fathers.
26
27
28
29
30
31
32
33
34
35

36 Five of the included studies offered parents MBI without any additional intervention
37 (Chan and Neece, 2018; Jones et al., 2018; McGregor et al., 2020; Neece, 2014; Neece et al.,
38 2019). In seven of the included studies, parents received training in cognitive-behavioural
39 educational strategies in addition to MBI (Andrews et al., 2022; Raulston et al., 2019;
40 Schwartzman et al., 2022; Singh et al., 2014; Singh et al., 2019; Singh et al., 2021; Weitlauf et
41 al., 2022). The parents in the Rojas-Torres et al. study (2023) received psychoeducation
42 sessions on autism. The parents in the Singh, Lancioni, Winton, Fisher, et al. study (2006) had
43 access to additional interventions not specified in the article.
44
45
46
47
48
49
50
51
52
53
54

55 The 14 studies included in this review proposed different types of MBI: training based
56 on acceptance and commitment therapy (ACT; Andrews et al., 2022), an intervention inspired
57
58
59
60
61
62
63
64
65

1 by the mindfulness-based stress reduction (MBSR) protocol (Chan & Neece, 2018; McGregor
2 et al., 2020; Neece, 2014; Neece et al., 2019; Schwartzman et al., 2022; Weitlauf et al., 2022),
3
4 an intervention called mindfulness-based well-being for parents (MBW-P; Jones et al., 2018),
5
6 mindfulness-based positive behaviour support (MBPBS; Singh et al., 2014; Singh et al., 2019;
7
8 Singh et al., 2021), the mindfulness parenting programme (Rojas-Torres et al., 2023), and a
9
10 formal meditation practice whose protocol was specially developed for the study (Raulston et
11
12 al., 2019; Singh, Lancioni, Winton, Fisher, et al., 2006). All 14 included studies were
13
14 manualized, and eight proposed an assessment of implementation fidelity (Andrews et al., 2022;
15
16 Chan & Neece, 2018; Neece et al., 2019; Raulston et al., 2019; Singh et al., 2014; Singh et al.,
17
18 2019; Singh et al., 2021; Weitlauf et al., 2022). The number of MBI sessions used in the 14
19
20 studies ranged from four (Andrews et al., 2022) to 12 (Singh, Lancioni, Winton, Fisher, et al.,
21
22 2006), with durations from two days (Andrews et al., 2022) to 12 weeks (Singh, Lancioni,
23
24 Winton, Fisher, et al., 2006). The sessions lasted from five minutes (Andrews et al., 2022) to
25
26 one day (Singh et al., 2014; Singh et al., 2019; Singh et al., 2021) and were offered either
27
28 individually (Andrews et al., 2022; Raulston et al., 2019; Singh, Lancioni, Winton, Fisher, et
29
30 al., 2006; Singh et al., 2014; Weitlauf et al., 2022) or in a parent group (Chan & Neece, 2018;
31
32 Jones et al., 2018; McGregor et al., 2020; Neece, 2014; Neece et al., 2019; Rojas-Torres et al.,
33
34 2023; Schwartzman et al., 2022; Singh et al., 2014; Singh et al., 2019; Singh et al., 2021). Three
35
36 of the MBIs were delivered remotely, either for all participants (Andrews et al., 2022), some of
37
38 the participants (Singh et al., 2021), or all participants for only one session (Rojas-Torres et al.,
39
40 2023). Finally, all the MBIs included a home meditation practice for parents.

41
42 Of the 14 included studies, three involved an RCT (Schwartzman et al., 2022; Singh et
43
44 al., 2021; Weitlauf et al., 2022). The remaining involved a multiple baseline design (Andrews
45
46 et al., 2022; Raulston et al., 2019; Singh, Lancioni, Winton, Fisher, et al., 2006; Singh et al.,
47
48 2014), a wait-list control group design (Chan & Neece, 2018; McGregor et al., 2020; Neece,
49
50

1 2014; Neece et al., 2019; Rojas-Torres et al., 2023), a pre/post-test design (Jones et al., 2018),
2 or an equivalent diagnostic two-group design (Singh et al., 2019). Seven studies included
3
4 follow-ups from one month (Andrews et al., 2022) to three years (Singh et al., 2021), including
5
6 one in which the follow-up data were not processed (McGregor et al., 2020).
7

8 9 *Variables studied*

10 All the studies included in this review focused on assessing challenging behaviours in
11
12 autistic children, except for those by Rojas-Torres et al. (2023) and Weitlauf et al. (2022). The
13
14 former assessed the perceived severity of autism signs using the autism impact measure (AIM)
15
16 while the latter targeted the core features of autism, cognitive level, and adaptive behaviour,
17
18 assessed using the Autism Diagnostic Observation Schedule, Second Edition (ADOS-2),
19
20 Mullen Scales of Early Learning, and the Vineland-II, respectively. The challenging behaviours
21
22 of autistic children were assessed using observations of behaviours targeted by the authors
23
24 (Andrews et al., 2022; Singh, Lancioni, Winton, Fisher, et al., 2006; Singh et al., 2014; Singh
25
26 et al., 2019; Singh et al., 2021), or using standardised tools, such as the Child Behaviour
27
28 Checklist (CBCL) (Chan & Neece, 2018; McGregor et al., 2020; Neece, 2014; Neece et al.,
29
30 2019; Raulston et al., 2019), the Strengths and Difficulties Questionnaire (SDQ; Jones et al.,
31
32 2018; Schwartzman et al., 2022), or the Aberrant Behaviour Checklist, Second Edition (ABC-
33
34 2; Schwartzman et al., 2022). All measurements were conducted solely by the parents, except
35
36 in the case of Andrews et al. (2022), where the experimenter also participated in the assessment.
37
38 Four studies also used the father as a second independent assessor in addition to the mother
39
40 (Singh, Lancioni, Winton, Fisher, et al., 2006; Singh et al., 2014; Singh et al., 2019; Singh et
41
42 al., 2021). Finally, using the Vineland-II, Jones et al. (2018) also assessed the adaptive
43
44 behaviour of the children in their study, and Schwartzman et al. (2022) also assessed the
45
46 children's social responsiveness, using the Social Responsiveness Scale, Second Edition (SRS-
47
48 2).
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Intervention-related children's outcomes

1
2 The children's behaviour showed significant improvement (see Table 3). Overall,
3
4 Andrews et al. (2022) observed a decrease in the frequency and average severity of challenging
5
6 behaviours between baseline, intervention, and follow-up for most children in their study. Chan
7
8 and Neece (2018) and Neece et al. (2019) identified a significant decrease in withdrawal and
9
10 attention problems between baseline and second assessment for children in the treatment group.
11
12 In addition, the scores for somatic complaints, withdrawal, attention problems, and aggressive
13
14 behaviour at the six-month follow-up were significantly lower than at baseline. In McGregor et
15
16 al. (2020), children whose parents were in the treatment group showed a significantly greater
17
18 reduction in challenging behaviours compared to children in the control group. In Neece (2014),
19
20 at the second assessment, the children in the treatment group showed significantly fewer signs
21
22 of hyperactivity compared to those in the control group. In their study, Raulston et al. (2019)
23
24 observed a moderate association between the intervention and a decrease in children's
25
26 challenging behaviours. The treatment group in Schwartzman et al. (2022) showed a significant
27
28 decrease in children's total behavioural difficulties, emotional problems, and inappropriate
29
30 speech compared with the control group; in addition, treatment-related gains in parents'
31
32 perceptions of their children's behaviours were maintained at eight weeks.
33
34
35
36
37
38
39
40

41 In studies by Singh, Lancioni, Winton, Fisher, et al. (2006), Singh et al. (2014), and
42
43 Singh et al. (2019), the average occurrence of three target behaviours decreased significantly
44
45 following intervention. In all three conditions of Singh et al.'s (2021) study, the occurrence of
46
47 children's aggressive and disruptive behaviours decreased significantly. Additionally,
48
49 compliance increased significantly following intervention, with the most significant reductions
50
51 being in the MBPBS condition, then the mindfulness (MB) and positive behaviour support
52
53 (PBS) conditions. These changes were maintained three years after the intervention. Finally,
54
55 following intervention, only Jones et al. (2018) found no significant reduction in behaviour
56
57
58
59
60
61
62
63
64
65

1 problems or increase in prosocial behaviours in the children in their study. Similarly, Weitlauf
2 et al. (2022) observed no significant differences between the two treatment groups (P-ESDM
3 condition and P-ESDM + MBSR condition) regarding autism signs, cognitive level, and
4 adaptive behaviour in the children in their study. Rojas-Torres et al. (2023) did not observe
5 changes in the perceived severity of signs across the measures.
6
7
8
9
10

11 *Methodological quality and levels of evidence*

12 Methodological quality scores and levels of evidence are presented in Table 2. The
13 assessment of methodological quality using STROBE guidelines ranged from 71% (Rojas-
14 Torres et al., 2023) to 95% (Schwartzman et al., 2022). The level of evidence, determined using
15 NHMRC guidelines, ranged from Level II for the three RCTs (Schwartzman et al., 2022; Singh
16 et al., 2021; Weitlauf et al., 2022) to Level III-3 (Andrews et al., 2022; Jones et al., 2018;
17 Raulston et al., 2019; Singh, Lancioni, Winton, Fisher, et al., 2006; Singh et al., 2014; Singh et
18 al., 2019).
19
20
21
22
23
24
25
26
27
28
29
30

31 *Risk of bias assessment of RCT studies*

32 Finally, the risk of bias inherent in the RCT studies is presented in Table 4. Weitlauf et
33 al. (2022) had the lowest risk of bias concerns while the highest risk was found in Schwartzman
34 et al. (2022) and Singh et al. (2021).
35
36
37
38
39
40
41
42

43 **Discussion**

44 This systematic review provides preliminary data on how MBIs offered exclusively to
45 parents indirectly affect their autistic children.
46
47
48
49
50

51 Most studies included in this review (12 out of 14) targeted a particular clinical specifier
52 of autism in children: challenging behaviours. Of these 12 studies, 11 showed a significant
53 reduction in challenging behaviours in autistic children following the implementation of MBIs
54 for their parents. The literature notes a high prevalence rate of challenging behaviours among
55
56
57
58
59
60
61
62
63
64
65

1 autistic individuals (Hattier et al., 2011; Matson & Nebel-Schwalm, 2007). Challenging
2 behaviours are also negatively associated with QoL in both adults (Chiang & Wineman, 2014)
3
4 and autistic children (Chuang et al., 2014; Delahaye et al., 2014; Kuhlthau et al., 2010; Kuhlthau
5
6 et al., 2018). Challenging behaviours have also been identified as a factor in low QoL among
7
8 parents of autistic children (Vasilopoulou & Nisbet, 2016). Such behaviours exacerbate parents'
9
10 perceived stress (Barroso et al., 2017), which in turn impacts their educational abilities (Bonis,
11
12 2016) and increases the child's challenging behaviours (Hastings, 2003). For the parents of
13
14 autistic children, challenging behaviours are considered the most stressful element, even more
15
16 than the core signs of autism (Hastings et al., 2005). It is therefore crucial to offer interventions
17
18 to families affected by autism to help parents better manage their children's challenging
19
20 behaviours. However, the effectiveness of the currently available interventions remains unclear
21
22 (Zwilling et al., 2022). The results of the present review suggest that MBIs aimed at the parents
23
24 of autistic children provide a viable alternative that could help reduce their children's
25
26 challenging behaviours and thus improve QoL for children and parents alike. The results align
27
28 with those of previous studies showing the relevance of MBIs for the well-being of parents of
29
30 autistic children (Chua and Shorey, 2022; Merriman et al., 2020; Yu et al., 2019).

31
32
33
34
35
36
37
38
39 The results of this review also complement those of previous studies showing a link
40
41 between MBIs offered to both parents and their autistic children and a significant reduction in
42
43 challenging behaviours in these children, as measured using the CBCL (Beck et al., 2022;
44
45 Bögels et al., 2008; Hwang et al., 2015; Ridderinkhof et al., 2018; Shaffer et al., 2019) or the
46
47 ABC (Shaffer et al., 2019; Shaffer et al., 2023). It would therefore seem that MBIs offered
48
49 solely to parents could achieve similar results to MBIs offered to both parents and their autistic
50
51 children. Comparative studies would shed further light on the added value of MBIs addressed
52
53 to the parent-child dyad, compared with those offered solely to parents.
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

One question raised by the results of the present review concerns the appropriateness of using interventions in addition to the MBIs addressed to parents (which are aimed at decreasing their children’s challenging behaviours). Five studies included in this review offered parents an MBI without any other intervention (Chan & Neece, 2018; Jones et al., 2018; McGregor et al., 2020; Neece, 2014; Neece et al., 2019) and one of these studies (Jones et al., 2018) showed no significant relationship between the MBI addressed to parents and their children’s challenging behaviours. Rayan and Ahmad (2017) suggested that in addition to MBIs, parents of autistic children may need other interventions, such as skills-based training on how to deal with their child’s challenging behaviours. However, other authors note that these interventions, which are mainly behavioural, are laborious to implement (Koerting et al., 2013). As a result, parents may struggle to apply them consistently, potentially leading to intervention failure and increased parental stress (Singh et al., 2014). Furthermore, these behavioural interventions, unlike MBIs, are less likely to produce substantial personal transformations in the parent (Singh, Lancioni, Winton, Fisher, et al., 2006).

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Finally, the findings of this research suggest that the positive effects of MBIs on the stress levels of parents of autistic children, as measured in the included studies, could be transferred to their children by helping them reduce their challenging behaviours. These findings align with those of similar studies involving children with other conditions, such as attention deficit hyperactivity disorder (ADHD; Lee et al., 2022), conduct disorder, oppositional defiant disorder, anxiety disorder, or learning disability (Burgdorf et al., 2019). Although the interpersonal mechanisms through which the positive effects of mindfulness are transferred from parents to their children remain poorly understood (Khoury et al., 2023), several hypotheses have been proposed to explain such effects (Singh, Lancioni, Winton, Fisher, et al., 2006). For instance, the nonjudgmental acceptance of children’s behaviours transforms the therapeutic environment into one that promotes positive behavioural changes

1 without directly targeting specific behaviours. It has been hypothesised that in typically
2 developing children, the parents' emotional management skills may influence the children's
3 emotional regulation abilities, notably through an observational learning mechanism (Morris et
4 al., 2007). On the other hand, in autistic children, a supportive parenting style and less excessive
5 involvement are associated with better emotion regulation (Ostfeld-Etzion et al., 2016).
6
7
8
9
10

11 The literature highlights the difficulties experienced by autistic children in regulating
12 their emotions (Cai et al., 2018), which involves controlling, evaluating, and modifying their
13 emotional responses to environmental demands (Thompson, 1994). This emotional
14 dysregulation, which is considered more prevalent in autistic children than in typically
15 developing children (Mallise et al., 2020), plays a central role in the adjustment difficulties
16 faced by autistic children (Mazefsky & White, 2014). It is a key factor in triggering challenging
17 behaviours (Korbut et al., 2020) regardless of intelligence level (Berkovits et al., 2017). These
18 emotional difficulties provide a clearer explanation for anger, loss of control, aggression or self-
19 mutilation than do associated psychiatric diagnoses (Mazefsky & White, 2014) or the severity
20 of autism signs (Korbut et al., 2020). For example, the self-injury behaviours often observed in
21 autistic people are thought to have a regulatory function for negatively valenced affective states
22 (Moseley et al., 2019). In other words, these behaviours enable autistic people to escape certain
23 emotional states (Chapman et al., 2006).
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 Better emotional regulation is associated with a reduction in challenging behaviours in
45 autistic children (Wilson et al., 2013), leading to higher QoL (Oakley et al., 2021). Therefore,
46 some authors recommend supporting the parents of autistic children by providing emotional
47 scaffolding strategies to help them assist their child in better regulating their emotions (Crowell
48 et al., 2019). This type of parental support is increasingly relevant, as emotional support is one
49 of the most important needs expressed by parents of autistic children (Derguy et al., 2015;
50 Boshoff et al., 2019). When confronted with challenging behaviour, parents may experience
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 negatively valenced emotional states that can act as antecedents to coercive reactions towards
2 their child (Chang et al., 2003; Lindsey & Barry, 2018). This may inadvertently escalate the
3
4 child’s negative affect (Omer, 2017), thus influencing new escape behaviours (Maljaars et al.
5
6 2014). Mindfulness meditation, as a tool for acceptance defined as openness to emotions (Hayes
7
8 et al., 2004), can help parents mitigate overreactions to their emotional states. This could
9
10 potentially decrease negative parent–child interactions and reduce the risk of escalation (Singh
11
12 et al., 2010). Furthermore, developing acceptance could help parents more effectively manage
13
14 their apprehensions when implementing extinction procedures, such as ignoring consequence-
15
16 maintained challenging behaviour (Raulston et al., 2019).
17
18
19
20
21

22 As Thomas et al. pointed out as early as 1968, an extreme temperament in a child only
23
24 leads to adjustment problems when parents respond with energetic confrontation (overreaction),
25
26 or conversely, capitulation (extinction failure). Finally, parental acceptance may also enable
27
28 them to respond more effectively to their child’s needs (Duncan et al., 2009; Parent et al., 2016).
29
30 Mindfulness practice, by giving the parent a “beginner’s eye” on their child’s behaviour, could
31
32 allow them to spontaneously intuit possibilities they had not previously envisaged (Suzuki,
33
34 1970). This could prompt them to adjust their conditioned parenting habits and engage in novel
35
36 responses. Parents can come to perceive autism less as a clinical construct and more as a human
37
38 variant. This shift may lead them to let go of some of their expectations of normality
39
40 (Poquérousse et al., 2021).
41
42
43
44
45

46 *Limitations*

47
48 While the studies included in this review indicate a positive trend, these results cannot
49
50 be generalised due to their varied methodological quality and low to intermediate levels of
51
52 evidence. Inadequate reporting makes it difficult to synthesise data. Likewise, the risks of bias
53
54 for the three RCT studies in this review are mostly high or uncertain, resulting in unconvincing
55
56 evidence. The scope of this systematic review is also limited by the small number of participants
57
58
59
60
61
62
63
64
65

1 in some of the included studies, the diverse research methodologies and study designs
2 employed, the variety of tools used to assess child-related variables, the presence of
3 interventions additional to MBIs in more than half of the included studies, and the wide
4 variations between the different MBIs offered to parents, which may partly explain the
5 variability between the studies. It should also be noted that in eight studies, there is a marked
6 absence of any implementation fidelity assessment. All 14 studies included in this review are
7 represented by just eight research teams. Furthermore, the collection of data by the parents
8 themselves in most of the included studies could call into question the reliability of the
9 assessments, since parental stress can influence evaluations (Schwartzman et al., 2021).
10 Nevertheless, this review sheds light on the quality of the child–parent relationship and the risk
11 of dysfunctional interactions between child and parent (Duthu et al., 2008). This implies that
12 measures taken by parents and clinicians can be complementary.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 This systematic review also has several methodological limitations. Firstly, a limited
30 number of studies were identified, all of which are written in English or French, which limits
31 the generalisability of the results to the target population. Secondly, although the protocol
32 registration prior to review starting is done for transparency of research methods used, the
33 protocol was not registered, as one of the authors had already initiated literature searches.
34 Thirdly, only one author screened the titles and abstracts during the initial phase. This
35 shortcoming may have partially influenced the identification of articles. However, this issue is
36 mitigated by the small number of available articles, which limits the risk of false negatives.
37 Additionally, all the selected articles were fully read by two of the authors. Finally, although
38 the discrepancy concerning the evaluation of the level of evidence was discussed with a third
39 author to reach a consensus, the low rate of agreement (62%) was not improved beforehand by
40 a reviewer’s training.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Conclusion and perspectives

1
2 Mindfulness and acceptance are important parenting support processes that clinicians
3
4 can offer to autistic children and their parents (Jones et al. 2014). However, clinicians must
5
6 remain vigilant for the possible adverse effects (such as transient distress) of programmes
7
8 drawing on these processes (Britton et al., 2021).
9
10

11
12 Further research is needed to better understand the mechanisms underlying the indirect
13
14 effects on autistic children of MBIs addressed to their parents. One approach to understanding
15
16 these mechanisms could involve using behavioural analytic methods to capture changes in the
17
18 behaviours of both parents and children (Singh et al., 2023). More experimental studies are
19
20 needed on the association between MBIs for the parents of autistic children and child outcomes.
21
22 In particular, these studies should focus on the impact of MBIs on children’s QoL, which is the
23
24 “gold standard” outcome for autism interventions. Researchers should also be alert to the
25
26 homogeneity (or lack of homogeneity) of MBIs across studies while isolating MBIs from other
27
28 types of interventions. The use of challenging behaviour assessments taken by parents as well
29
30 as clinicians would enable us to better understand whether the behavioural problems are
31
32 objectively less significant or whether it is only the parents’ perceptions of their child’s
33
34 challenging behaviour that are changing. Finally, researchers should also focus on measuring
35
36 and adapting their intervention’s implementation processes to increase the reliability of its
37
38 results.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- 1
2 American Psychiatric Association (2013). *Diagnostic and statistical manual of mental*
3 *disorders* (5th ed.). American Psychiatric Association.
4
5
6 Andrews, M. L., Garcia, Y. A., Catagnus, R. M., & Gould, E. R. (2022). Effects of acceptance
7 and commitment training plus behavior parent training on parental implementation of
8 autism treatment. *The Psychological Record*, 72, 601–617. doi: 10.1007/s40732-021-
9 00496-5
10
11
12
13 Arias, V. B., Gomez, L. E., Moran, M. L., Alcedo, M. A., Monsalve, A., & Fontanil, Y. (2018).
14 Does quality of life differ for children with autism spectrum disorder and intellectual
15 disability compared to peers without autism? *Journal of Autism and Developmental*
16 *Disorders*, 48(1), 123–136. doi: 10.1007/s10803-017-3289-8
17
18
19
20 Aydin, A. (2023). Examining the mediating role of mindful parenting: A study
21 on the relationship between parental emotion regulation difficulties and problem
22 behaviors of children with ASD. *Journal of Autism and Developmental Disorders*,
23 53(5), 1873–1883. <https://doi.org/10.1007/s10803-022-05455-9>
24
25
26
27 Barroso, N. E., Mendez, L., Graziano, P. A., & Bagner, D. M. (2017). Parenting stress through
28 the lens of different clinical groups: A systematic review & meta-analysis. *Journal of*
29 *Abnormal Child Psychology*, 46, 449–461. doi: 10.1007/s10802-017-0313-6
30
31
32
33 Beck, K. B., Conner, C. M., White, S. W., & Mazefsky, C. A. (2020). Mindfulness “here and
34 now”: Strategies for helping adolescents with autism. *Journal of the American Academy*
35 *of Child & Adolescent Psychiatry*, 59(10), 1125–1127. doi: 10.1016/j.jaac.2020.07.004
36
37
38
39 Beck, K. B., Greco, C. M., Terhorst, L. A., Skidmore, E. R., Kulzer, J. L., & McCue, M. P.
40 (2020). Mindfulness-based stress reduction for adults with autism spectrum disorder:
41 Feasibility and estimated effects. *Mindfulness*, 11, 1286–1297.
42 <https://doi.org/10.1007/s12671-020-01345-2>
43
44
45
46 Beck, K. B., Northrup, J. B., Breitenfeldt, K. E., Porton, S., Day, T. N., MacKenzie, K. T.,
47 Conner, C. M., & Mazefsky, C. A. (2022). Stakeholder informed development of the
48 Emotion Awareness and Skills Enhancement team-based program (EASE-Teams).
49 *Autism*, 26(3), 586–600. doi: 10.1177/13623613211061936
50
51
52
53 Beer, M., Ward, L., & Moas, K. (2013). The relationship between mindful parenting and
54 distress in parents of children with an autism spectrum disorder. *Mindfulness*, 4, 102–
55 112. <https://doi.org/10.1007/s12671-012-0192-4>
56
57
58
59
60
61
62
63
64
65

- 1 Benevides, T. W., Shore, S. M., Andresen, M. L., Caplan, R., Cook, B., Gassner, D. L., Erves,
2 J. M., Hazlewood, T. M., King, M. C., Morgan, L., Murphy, L. E., Purkis, Y.,
3 Rankowski, B., Rutledge, S. M., Welch, S. P., & Wittig, K. (2020). Interventions to
4 address health outcomes among autistic adults: A systematic review. *Autism*, 24(6),
5 1345–1359. doi: 10.1177/1362361320913664
6
7
8
9 Bögels, S., Hoogstad, B., van Dun, L., de Schutter, S., & Restifo, K. (2008). Mindfulness
10 training for adolescents with externalizing disorders and their parents. *Behavioural and*
11 *Cognitive Psychotherapy*, 36(2), 193–209.
12 https://doi.org/10.1017/S1352465808004190
13
14
15
16 Bonis, S. (2016). Stress and parents of children with autism: A review of literature. *Issues in*
17 *Mental Health Nursing*, 37(3), 153–163.
18 https://doi.org/10.3109/01612840.2015.1116030
19
20
21
22 Boshoff, K., Gibbs, D., Phillips, R. L., Wiles, L., & Porter, L. (2019). A meta-synthesis of how
23 parents of children with autism describe their experience of advocating for their children
24 during the process of diagnosis. *Health and Social Care in the Community*, 27, e143–
25 e157. doi: 10.1111/hsc.12691
26
27
28
29 Britton, W. B., Lindahl, J. R., Cooper, D. J., Canby, N. K., & Palitsky, R. (2021). Defining and
30 measuring meditation-related adverse effects in mindfulness-based programs. *Clinical*
31 *Psychological Science*, 9(6), 1185–1204. https://doi.org/10.1177/2167702621996340
32
33
34
35 Burgdorf, V., Szabo, M., & Abbott, M. J. (2019). The effect of mindfulness interventions for
36 parents on parenting stress and youth psychological outcomes: A systematic review and
37 meta-analysis. *Frontiers in Psychology*, 10. https://doi.org/10.3389/fpsyg.2019.01336
38
39
40
41 Byrne, G., & O'Mahony, T. (2020). Acceptance and commitment therapy (ACT) for adults
42 with intellectual disabilities and/or autism spectrum conditions (ASC): A systematic
43 review. *Journal of Contextual Behavioral Science*, 18, 247–255.
44 https://doi.org/10.1016/j.jcbs.2020.10.001
45
46
47
48 Cachia, R. L., Anderson, A., & Moore, D. W. (2016a). Mindfulness, stress and well-being in
49 parents of children with autism spectrum disorder: A systematic review. *Journal of*
50 *Child and Family Studies*, 25, 1–14. https://doi.org/10.1007/s10826-015-0193-8
51
52
53
54 Cachia, R. L., Anderson, A., & Moore, D. W. (2016b). Mindfulness in individuals with autism
55 spectrum disorder: A systematic review and narrative analysis. *Review Journal of*
56 *Autism and Developmental Disorders*, 3, 165–178. https://doi.org/10.1007/s40489-016-
57 0074-0
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Cai, R. Y., Richdale, A. L., Uljarevic, M., Dissanayake, C., & Samson, A. C. (2018). Emotion regulation in autism spectrum disorder: Where we are and where we need to go. *Autism Research, 11*, 962–978. doi: 10.1002/aur.1968
- Carter Leno, V., Vitoratou, S., Kent, R., Charman, T., Chandler, S., Jones, C. R., Happe, F. Baird, G., Pickles, A., & Simonoff, E. (2019). Exploring the neurocognitive correlates of challenging behaviours in young people with autism spectrum disorder. *Autism, 23*(5), 1152–1164. <https://doi.org/10.1177/1362361318769176>
- Chan, N., & Neece, C. L. (2018). Mindfulness-based stress reduction for parents of children with developmental delays: A follow-up study. *Evidence-Based Practice in Child and Adolescent Mental Health, 3*(1), 16–29. <https://doi.org/10.1080/23794925.2017.1399484>
- Chan, K. K. S., Wang, Z., & Li, H. (2022). Longitudinal impact of mindful parenting on internalizing and externalizing symptoms among children with autism spectrum disorder. *Mindfulness, 13*(8), 1984–1993. <https://doi.org/10.1007/s12671-022-01933-4>
- Chang, L., Schwartz, D., Dodge, K. A., & McBride-Chang, C. (2003). Harsh parenting in relation to child emotion regulation and aggression. *Journal of Family Psychology, 17*(4), 598–606. doi: 10.1037/0893-3200.17.4.598
- Chapman, A. L., Gratz, K. L., & Brown, M. Z. (2006). Solving the puzzle of deliberate self-harm: The experiential avoidance model. *Behaviour Research and Therapy, 44*, 371–394. <https://doi.org/10.1016/j.brat.2005.03.005>
- Chiang, H. M., & Wineman, I. (2014). Factors associated with quality of life in individuals with autism spectrum disorders: A review of literature. *Research in Autism Spectrum Disorders, 8*(8), 974–986. <https://doi.org/10.1016/j.rasd.2014.05.003>
- Chua, J. Y. X., & Shorey, S. (2022). The effect of mindfulness-based and acceptance commitment therapy-based interventions to improve the mental well-being among parents of children with developmental disabilities: A systematic review and meta-analysis. *Journal of Autism and Developmental Disorders, 52*(6), 2770–2783. doi: 10.1007/s10803-021-04893-1
- Chuang, I. C., Tseng, M. H., Lu, L., Shieh, J. Y., & Cermak, S. A. (2014). Predictors of the health-related quality of life in preschool children with autism spectrum disorders. *Research in Autism Spectrum Disorder, 8*, 1062–1070. <https://doi.org/10.1016/j.rasd.2014.05.015>

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Crowell, J. A., Keluskar, J., & Gorecki, A. (2019). Parenting behavior and the development of children with autism spectrum disorder. *Comprehensive Psychiatry*, *90*, 21–29. doi: 10.1016/j.comppsy.2018.11.007
- da Costa, B. R., Cevallos, M., Altman, D. G., Rutjes, A. W. S., & Egger, M. (2011). Uses and misuses of the STROBE statement: Bibliographic study. *BMJ Open*, *1*, e000048. doi: 10.1136/bmjopen-2010-000048
- Delahaye, J., Kovacs, E., Sikora, D., Hall, T. A., Orlich, F., Clemons, T. E., van der Weerd, E., Glick, L., & Kuhlthau, K. (2014). The relationship between health-related quality of life and sleep problems in children with autism spectrum disorder. *Research in Autism Spectrum Disorder*, *8*(3), 292–303. <https://doi.org/10.1016/j.rasd.2013.12.015>
- Derguy, C., Michel, G., M'Bailara, K., Roux, S., & Bouvard, M. (2015). Assessing needs in parents of children with autism spectrum disorder: A crucial preliminary step to target relevant issues for support programs. *Journal of Intellectual and Developmental Disability*, *40*(2), 1–11. <https://doi.org/10.3109/13668250.2015.1023707>
- Dückert, S., Gewohn, P., König, H., Schöttle, D., Konnopka, A., Rahlff, P., Vogeley, K., Schulz, H., David, N., & Peth, J. (2023). Multidimensional burden on family caregivers of adults with autism spectrum disorder: A scoping review. *Review Journal of Autism and Developmental Disorders*. <https://doi.org/10.1007/s40489-023-00414-1>
- Duncan, L. G., Coatsworth, J. D., & Greenberg, M. T. (2009). A model of mindful parenting: Implications for parent-child relationships and prevention research. *Clinical Child and Family Psychology Review*, *12*(3), 255–270. <https://doi.org/10.1007/s10567-009-0046-3>
- Duthu, S., Blicharski, T., Bouchet, G., & Bourdet-Loubère, S. (2008). Caractéristiques parentales et représentations du tempérament de l'enfant: Influence des traits de personnalité, des symptômes psychiatriques et de l'écologie familiale. *Neuropsychiatrie de l'enfance et de l'adolescence*, *56*, 155–164. <https://doi.org/10.1016/j.neurenf.2008.01.014>
- Harmony, C., & Woodard, C. R. (2020). Mindfulness training for staff in a school for children with autism and other developmental disabilities: Effects on staff mindfulness and student behavior. *Advances in Neurodevelopmental Disorders*, *4*, 113–121. <https://doi.org/10.1007/s41252-020-00148-9>
- Hastings, R. P. (2003). Child behaviour problems and partner mental health as correlates of stress in mothers and fathers of children with autism. *Journal of Intellectual Disability Research*, *47*, 231–237. doi: 10.1046/j.1365-2788.2003.00485.x

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Hastings, R. P., Kovshoff, H., Ward, N. J., Degli Espinosa, F., Brown, T., & Remington, B. (2005). Systems analysis of stress and positive perceptions in mothers and fathers of pre-school children with autism. *Journal of Autism and Developmental Disorders*, 35(5), 635–644. <https://doi.org/10.1007/s10803-005-0007-8>
- Hattier, M. A., Matson, J. L., Belva, B. C., & Horovitz, M. (2011). The occurrence of challenging behaviours in children with autism spectrum disorders and atypical development. *Developmental Neurorehabilitation*, 14(4), 221–229. <https://doi.org/10.3109/17518423.2011.573836>
- Hayes, S. C., Follette, V. M., & Linehan, M. M. (2004). *Mindfulness and acceptance*. The Guilford Press.
- Hayes, S. A., & Watson, S. L. (2013). The impact of parenting stress: A meta-analysis of studies comparing the experience of parenting stress in parents of children with and without autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 43, 629–642. doi: 10.1007/s10803-012-1604-y
- Herrema, R., Garland, D., Osborne, M., Freeston, M., Honey, E. and Rodgers, J. (2017). Mental wellbeing of family members of autistic adults. *Journal of Autism and Developmental Disorders*, 47(11), 3589–3599. doi: 10.1007/s10803-017-3269-z
- Hwang, Y.-S., Kearney, P., Klieve, H., Lang, W., & Roberts, J. (2015). Cultivating mind: Mindfulness interventions for children with autism spectrum disorder and problem behaviours, and their mothers. *Journal of Child and Family Studies*. 24(10), 3093–3106. <https://doi.org/10.1007/s10826-015-0114-x>
- Ikeda, E., Hinckson, E., & Krägeloh, C. (2014). Assessment of quality of life in children and youth with autism spectrum disorder: A critical review. *Quality of Life Research*, 23(4), 1069–1085. doi: 10.1007/s11136-013-0591-6
- Jones, P., & Drummond, P. (2023). Construction and evaluation of a mindfulness-based quality of life and well-being program (MQW) in a randomized trial. *Current Psychology*, 42, 14782–14803. <https://doi.org/10.1007/s12144-021-02672-w>
- Jones, L., Gold, E., Totsika, V., Hastings, R. P., Jones, M., Griffiths, A., & Silverton, S. (2018). A mindfulness parent well-being course: Evaluation of outcomes for parents of children with autism and related disabilities recruited through special schools. *European Journal of Special Needs Education*, 33(1), 16–30. <https://doi.org/10.1080/08856257.2017.1297571>

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Jones, L., Hastings, R. P., Totsika, V., Keane, L., & Rhule, N. (2014) Child behavior problems and parental well-being in families of children with autism: The mediating role of mindfulness and acceptance. *American Journal on Intellectual and Developmental Disabilities*, 119(2), 171–85. doi: 10.1352/1944-7558-119.2.171
- Juvin, J., Sadeg, S., Julien-Sweerts, S., & Zebdi, R. (2021). A systematic review: Acceptance and commitment therapy for the parents of children and adolescents with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 52(1), 124–141.
- Kabat-Zinn J. (2003). Mindfulness-based interventions in context: Past, present, and future. *Clinical Psychology: Science and Practice*, 10(2), 144–156. doi: 10.1007/s10803-021-04923-y
- Khoury, B., Manova, V., Adel, L., Dumas, G., Lifshitz, M., Vergara, R. C., Sekhon, H., & Rej, S. (2023). Tri-process model of interpersonal mindfulness: Theoretical framework and study protocol. *Frontiers in Psychology*, 14. <https://doi.org/10.3389/fpsyg.2023.1130959>
- Kim, S. Y., & Bottema-Beutel, K. (2019). A meta regression analysis of quality of life correlates in adults with ASD. *Research in Autism Spectrum Disorders* 63, 23–33. <https://doi.org/10.1016/j.rasd.2018.11.004>
- Koerting J., Smith E., Knowles M. M., Latter S., Elsey H., McCann D. C., Thompson, M., & Sonuga-Barke E. J. (2013). Barriers to, and facilitators of, parenting programmes for childhood behavior problems: A qualitative synthesis of studies of parents’ and professionals’ perceptions. *European Child & Adolescent Psychiatry*, 22, 653–670. <https://doi.org/10.1007/s00787-013-0401-2>
- Korbut, S., Hedley, D., Chetcuti, L., Sahin, E., & Nuske, H. J. (2020). Temperament predicts challenging behavior in children with autism spectrum disorder at age 5. *Research in Autism Spectrum Disorders*, 71. <https://doi.org/10.1016/j.rasd.2019.101492>
- Kuhlthau, K. A., McDonnell, E., Cury, D. L., Payakachat, N., & Macklin, E. (2018). Associations of quality of life with health-related characteristics among children with autism. *Autism*, 22(7), 804–813. doi: 10.1177/1362361317704420
- Kuhlthau, K., Orlich, F., Hall, T. A., Sikora, D., Kovacs, E. A., Delahaye, J., & Clemons, T. E. (2010). Health-related quality of life in children with autism spectrum disorders: Results from the autism treatment network. *Journal of Autism and Developmental Disorders*, 40, 721–729. doi: 10.1007/s10803-009-0921-2
- Lee, C. S. C., Ng, K. H., Chan, P. C. K., & Peng, X. (2022). Effectiveness of mindfulness parent training on parenting stress and children’s ADHD-related behaviors: A systematic

review and meta-analysis. *Hong Kong Journal of Occupational Therapy*, 35(1), 3–24.
doi: 10.1177/15691861211073826

Anonymous et al. (2023).

Anonymous et al. (2020).

Li, S. N., Chien, W. T., Lam, S. K. K., Chen, Z. Y., & Ma, X. (2024). Effectiveness of parent-focused interventions for improving the mental health of parents and their children with autism spectrum disorder: A systematic review and meta-analysis. *Research in Autism Spectrum Disorders*, 114, 102389. <https://doi.org/10.1016/j.rasd.2024.102389>

Lindsey, R. A., & Barry, T. D. (2018). Protective factors against distress for caregivers of a child with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 48(4), 1092–1107. doi: 10.1007/s10803-017-3372-1

Maljaars, J., Boonen, H., Lambrechts, G., Van Leeuwen, K., & Noens, I. (2014). Maternal parenting behavior and child behavior problems in families of children and adolescents with autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 44(3), 501–512. <https://doi.org/10.1007/s10803-013-1894-8>

Mallise, C. A., Lane, A. E., Woolard, A. J., Whalen, O. M., Murphy, V. E., Karayanidis, F., & Campbell, L. E. (2020). The temperament features associated with autism spectrum disorder in childhood: A systematic review. *Research in Developmental Disabilities*, 104. <https://doi.org/10.1016/j.ridd.2020.103711>

Matson, J. L., & Nebel-Schwalm, M. S. (2007). Comorbid psychopathology with autism spectrum disorder in children: An overview. *Research in Developmental Disabilities*, 28, 341–352. doi: 10.1016/j.ridd.2005.12.004

Mazefsky, C. A., & White, S. W. (2014). Emotion regulation: Concepts & practice in autism spectrum disorder. *Child and Adolescent Psychiatric Clinics of North America*, 23(1), 15–24. doi: 10.1016/j.chc.2013.07.002

McGregor, H. A., Sanner, C. M., & Neece, C. L. (2020). Effects of MBSR parent intervention on internalizing problems in children: ASD status as a moderator. *Journal of Mental Health Research in Intellectual Disabilities*, 13(4), 343–363. <https://doi.org/10.1080/19315864.2020.1815913>

McStay, R. L., Trembath, D., & Dissanayake, C. (2014). Stress and family quality of life in parents of children with autism spectrum disorder: Parent gender and the double ABCX model. *Journal of Autism and Developmental Disorders*, 44(12), 3101–3118. <https://doi.org/10.1007/s10803-014-2178-7>

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Menezes, M., Harkins, C., Robinson, M. F., & Mazurek, M. O. (2020). Treatment of depression in individuals with autism spectrum disorder: A systematic review. *Research in Autism Spectrum Disorders*, 78. <https://doi.org/10.1016/j.rasd.2020.101639>
- Merriman, K., Burke, T., & O'Reilly, G. (2020). A systematic review of the effectiveness and efficacy of clinician-led psychological interventions for parents of children with ASD. *Research in Autism Spectrum Disorders*, 76(4). <https://doi.org/10.1016/j.rasd.2020.101584>
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine*, 6(7), e1000097. <https://doi.org/10.1371/journal.pmed.1000097>
- Morris, A. S., Silk, J. S., Steinberg, L., Myers, S. S., & Robinson, L. R. (2007). The role of the family context in the development of emotion regulation. *Social Development*, 16(2), 361–388. doi: 10.1111/j.1467-9507.2007.00389.x
- Moseley, R. L., Gregory, N. J., Smith, P., Allison, C., & Baron-Cohen, S. (2019). A ‘choice’, an ‘addiction’, a way ‘out of the lost’: Exploring self-injury in autistic people without intellectual disability. *Molecular Autism*, 10, 20–23. <https://doi.org/10.1186/s13229-019-0267-3>
- Munder, T., & Barth, J. (2017). Cochrane’s risk of bias tool in the context of psychotherapy outcome research. *Psychotherapy Research*. <https://doi.org/10.1080/10503307.2017.1411628>
- National Health Medical Research Council. (1999). *A guide to the development, implementation and evaluation of clinical practice guidelines*. Commonwealth of Australia.
- National Institute for Health and Care Excellence (NICE) (2013). *Autism: The management and support of children and young people on the autism spectrum*. National Institute for Health and Care Excellence.
- Neece, C. L. (2014). Mindfulness-based stress reduction for parents of young children with developmental delays: Implications for parental mental health and child behavior problems. *Journal of Applied Research in Intellectual Disabilities*, 27, 174–186. doi: 10.1111/jar.12064
- Neece, C. L., Chan, N., Klein, K., Roberts, L., & Fenning, R. M. (2019). Mindfulness-based stress reduction for parents of children with developmental delays: Understanding the experiences of Latino families. *Mindfulness*, 10, 1017–1030. doi: 10.1007/s12671-018-1011-3

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Oakley, B. F. M., Tillmann, J., Ahmad, J., Crawley, D., Caceres, A. S. J., Holt, R., Charman, T., Banaschewski, T., Buitelaar, J., Simonoff, E., Murphy, D., & Loth, E. (2021). How do core autism traits and associated symptoms relate to quality of life? Findings from the Longitudinal European Autism Project. *Autism*, 25(2), 389–404. <https://doi.org/10.1177/1362361320959959>
- Omer, H. (2017). *La résistance non violente*. De Boeck.
- Ostfeld-Etzion, S., Feldman, R., Hirschler-Guttenberg, Y., Laor, N., & Golan, O. (2016). Self-regulated compliance in preschoolers with autism spectrum disorder: The role of temperament and parental disciplinary style. *Autism*, 20(7), 868–878. <https://doi.org/10.1177/1362361315615467>
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E. Chou, R., Glanville, J., Grimshaw, J. M., Hrobjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E., McDonald, S., ..., & Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*, 372(71). <https://doi.org/10.1136/bmj.n71>
- Parent, J., McKee, L. G., Rough, J. N., & Forehand, R. (2016). The association of parent mindfulness with parenting and youth psychopathology across three developmental stages. *Journal of Abnormal Child Psychology*, 44(1), 191–202. doi: 10.1007/s10802-015-9978-x
- Poquérusse, J., Pagnini, F., & Langer, E. J. (2021). Mindfulness for autism. *Advances in Neurodevelopmental Disorders*, 5, 77–84. <https://doi.org/10.1007/s41252-020-00180-9>
- Raulston, T. J., Kosty, D., & McIntyre, L. L. (2021). Mindful parenting, caregiver distress, and conduct problems in children with autism. *American Journal on Intellectual and Developmental Disabilities*, 126(5), 396–408. doi: 10.1352/1944-7558-126.5.396
- Raulston, T. J., Zemantic, P. K., Machalicek, W., Hieneman, M., Kurtz-Nelson, E., Barton, H., Hansen, S. G., & Frantz, R. J. (2019). Effects of a brief mindfulness-infused behavioral parent training for mothers of children with autism spectrum disorder. *Journal of Contextual Behavioral Science*, 13, 42–51. <https://doi.org/10.1016/j.jcbs.2019.05.001>
- Rayan, A., & Ahmad, M. (2017). Effectiveness of mindfulness-based intervention on perceived stress, anxiety, and depression among parents of children with autism spectrum disorder. *Mindfulness*, 8, 677–690. <https://doi.org/10.1007/s12671-016-0595-8>

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Ridderinkhof, A., de Bruin, E. I., Blom, R., & Bögels, S. M. (2018). Mindfulness-based program for children with autism spectrum disorder and their parents: Direct and long-term improvements. *Mindfulness*, 9, 773–791. doi: 10.1007/s12671-017-0815-x
- Rojas-Torres, L., Alonso-Esteban, Y., & Alcantud-Marin, F. (2023). Mindfulness parenting and childish play: A clinical trial with parents of children with autism spectrum disorders. *Psicothema*, 35(3), 259–270. doi: 10.7334/psicothema2022.339
- Schalock, R. L. (2004). The concept of quality of life: What we know and do not know. *Journal of Intellectual Disability Research*, 48(3), 203–216. <https://doi.org/10.1111/j.1365-2788.2003.00558.x>
- Schwartzman, J. M., Hardan, A. Y., & Gengoux, G. W. (2021). Parenting stress in autism spectrum disorder may account for discrepancies in parent and clinician ratings of child functioning. *Autism*, 25(6), 1601–1614. doi: 10.1177/1362361321998560
- Schwartzman, J. M., Millan, M. E., Uljarevic, M., & Gengoux, G. W. (2022). Resilience intervention for parents of children with autism: Findings from a randomized controlled trial of the AMOR method. *Journal of Autism and Developmental Disorders*, 52, 738–757. <https://doi.org/10.1007/s10803-021-04977-y>
- Semple, R. J. (2019). Review: Yoga and mindfulness for youth with autism spectrum disorder: review of the current evidence. *Child and Adolescent Mental Health*, 24(1), 12–18. <https://doi.org/10.1111/camh.12295>
- Shaffer, R. C., Schmitt, L. M., Reisinger, D. L., Coffman, M., Horn, P., Goodwin, M. S., Mazefsky, C., Randall, S., & Erickson C. (2023). Regulating together: Emotion dysregulation group treatment for ASD youth and their caregivers. *Journal of Autism and Developmental Disorders*, 53(5), 1942–1962. doi: 10.1007/s10803-022-05461-x
- Shaffer, R. C., Wink, L. K., Ruberg, J., Pittenger, A., Adams, R., Sorter, M., Manning, P., & Erickson, C. A. (2019). Emotion regulation intensive outpatient programming: Development, feasibility, and acceptability. *Journal of Autism and Developmental Disorders*, 49, 495–508. doi: 10.1007/s10803-018-3727-2
- Singh, N. N., Lancioni, G. E., Felver, J. C., Myers, R. E., Hwang, Y. S., Chan, J., & Medvedev, O. N. (2023). Effects of mindful engagement and attention on reciprocal caregiver and client interactions: A behavioral analysis of moment-to-moment changes during mindfulness practice. *Mindfulness* 14, 1893–1907. <https://doi.org/10.1007/s12671-023-02190-9>
- Singh, N. N., Lancioni, G. E., Karazsia, B. T., Myers, R. E., Hwang, Y. S., & Analayo, B. (2019). Effects of mindfulness-based positive behavior support (MBPBS) training are

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

equally beneficial for mothers and their children with autism spectrum disorder or with intellectual disabilities. *Frontiers in Psychology*, *10*.
<https://doi.org/10.3389/fpsyg.2019.00385>

Singh, N. N., Lancioni, G. E., Medvedev, O. N., Hwang, Y. S., & Myers, R. E. (2021). A component analysis of the mindfulness-based positive behavior support (MBPBS) program for mindful parenting by mothers of children with autism spectrum disorder. *Mindfulness*, *12*, 463–475. doi: 10.1007/s12671-020-01376-9

Singh, N. N., Lancioni, G. E., Winton, A. S. W., Curtis, W. J., Wahler, R. G., Sabaawi, M., Singh, J., & McAleavey, K. (2006). Mindful staff increase learning and reduce aggression in adults with developmental disabilities. *Research in Developmental Disabilities*, *27*, 545–558. doi: 10.1016/j.ridd.2005.07.002

Singh, N. N., Lancioni, G. E., Winton, A. S. W., Fisher, B. C., Wahler, R. G., McAleavey, K., Singh, J., & Sabaawi, M. (2006). Mindful parenting decreases aggression, noncompliance and self-injury in children with autism. *Journal of Emotional and Behavioral Disorders*, *14*, 169–177. <https://doi.org/10.1177/10634266060140030401>

Singh, N. N., Lancioni, G. E., Winton, A. S. W., Karazsia, B. T., Myers, R. E., Latham, L. L., & Singh, J. (2014). Mindfulness-based positive behavior support (MBPBS) for mothers of adolescents with autism spectrum disorder: Effects on adolescents' behavior and parental stress. *Mindfulness*, *5*, 646–657. <https://doi.org/10.1007/s12671-014-0321-3>

Singh, N. N., Lancioni, G. E., Winton, A. S. W., Wahler, R. G., Singh, J., & Sage, M. (2004). Mindful caregiving increases happiness among individuals with profound multiple disabilities. *Research in Developmental Disabilities*, *25*, 207–218. doi: 10.1016/j.ridd.2003.05.001

Singh, N. N., Singh, A. N., Lancioni, G. E., Singh, J., Winton, A. S. W., & Adkins, A. D. (2010). Mindfulness training for parents and their children with ADHD increases the children's compliance. *Journal of Child and Family Studies* *19*, 157–166. <https://doi.org/10.1007/s10826-009-9272-z>

Spek, A. A., van Ham, N. C., & Nyklıcek, I. (2013). Mindfulness-based therapy in adults with an autism spectrum disorder: A randomized controlled trial. *Research in Developmental Disabilities*, *34*, 246–253. doi: 10.1016/j.ridd.2012.08.009

Suvarna, V., Farrell, L., Adams, D., Emerson, L.-M., & Paynter, J. (2024). Parenting practices and externalizing behaviors in autistic children: A systematic literature review. *Clinical Child and Family Psychology Review*, *27*, 235–256. doi: 10.1007/s10567-024-00467-

- 1 Suzuki, S. (1970). *Zen mind, beginner's mind*. Weatherhill.
- 2 Thomas, A., Chess, S., & Birch, H. G. (1968). *Temperament and behaviour disorders in*
3 *children*. New York University Press.
- 4
- 5 Thompson, R. A. (1994). Emotion regulation: A theme in search of definition. *Monographs of*
6 *the Society for Research in Child Development*, 59(2–3), 25–52. PMID: 7984164
- 7
- 8
- 9 Ting, V., & Weiss, J. A. (2017). Emotion regulation and parent co-regulation in children with
10 autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 47, 680–
11 689. doi: 10.1007/s10803-016-3009-9
- 12
- 13
- 14 Van Heijst, B. F. C., & Geurts, H. M. (2015). Quality of life in autism across the lifespan: A
15 meta-analysis. *Autism*, 19, 158–167. doi: 10.1177/1362361313517053
- 16
- 17
- 18 Vasilopoulou, E., & Nisbet, J. (2016). The quality of life of parents of children with autism
19 spectrum disorder: A systematic review. *Research in Autism Spectrum Disorders*, 23,
20 36–49. <https://doi.org/10.1016/j.rasd.2015.11.008>
- 21
- 22
- 23 Vernhet, C., Michelon, C., Dellapiazza, F., Rattaz, C., Geoffray, M. M., Roeyers, H., Picot, M.
24 C., & Baghdadli, A. (2022). Perceptions of parents of the impact of autism spectrum
25 disorder on their quality of life and correlates: Comparison between mothers and
26 fathers. *Quality of Life Research*, 31, 1499–1508. doi: 10.1007/s11136-021-03045-3
- 27
- 28
- 29 von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gotsche, P. C., & Vandenbroucke, J. P.
30 (2007). The strengthening the reporting of observational studies in epidemiology
31 (STROBE) statement: Guidelines for reporting observational studies. *Journal of*
32 *Clinical Epidemiology*, 61(4), 344–349. doi: 10.1136/bmj.39335.541782
- 33
- 34
- 35
- 36
- 37
- 38 Weitlauf, A. S., Broderick, N., Alacia Stainbrook, J., Slaughter, J. C., Taylor, J. L., Herrington,
39 C. G., Nicholson, A. G., Santulli, M., Dorris, K., Garrett, L. J., Hopton, M., Kinsman,
40 A., Morton, M., Vogel, A., Dykens, E. M., Juarez, A. P., & Warren, Z. E. (2022). A
41 longitudinal RCT of P-ESDM with and without parental mindfulness based stress
42 reduction: Impact on child outcomes. *Journal of Autism and Developmental Disorders*,
43 52, 5403–5413. doi: 10.1007/s10803-021-05399-6
- 44
- 45
- 46
- 47
- 48
- 49 Wilson, B. J., Berg, J. L., Zurawski, M. E., & King, K. A. (2013). Autism and externalizing
50 behaviors: Buffering effects of parental emotion coaching. *Research in Autism*
51 *Spectrum Disorders*, 7, 767–776. <https://doi.org/10.1016/j.rasd.2013.02.005>
- 52
- 53
- 54
- 55 World Health Organization. (1946). *Constitution*. WHO.
- 56
- 57 Yorke, I., White, P., Weston, A., Rafla, M., Charman, T., & Simonoff, E. (2018). The
58 association between emotional and behavioral problems in children with autism
59 spectrum disorder and psychological distress in their parents: A systematic review and
60
- 61
- 62
- 63
- 64
- 65

meta-analysis. *Journal of Autism and Developmental Disorders*, 48, 3393–3415. doi:
10.1007/s10803-018-3605-y

1
2
3 Yu, Y., McGrew, J. H., & Bolor, J. (2019). Effects of caregiver-focused programs on
4 psychosocial outcomes in caregivers of individuals with ASD: A meta-analysis. *Journal*
5 *of Autism and Developmental Disorders*, 49(12), 4761–4779. doi: 10.1007/s10803-019-
6 04181-z
7
8
9

10 Zwillling, M., Romano, A., Hoffman, H., Lotan, M., & Tesler, R. (2022). Development and
11 validation of a system for the prediction of challenging behaviors of people with autism
12 spectrum disorder based on a smart wearable shirt: A mixed-methods design. *Frontiers*
13 *in Behavioral Neuroscience*. <https://doi.org/10.3389/fnbeh.2022.948184>
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 1. PRISMA 2020 flow diagram (*updated database searches limiting to the period between January 2023 and April 2024)

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Study characteristics

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Andrews et al. (2022) USA	3 mothers and 1 transgender father 35–51 years	AS ¹ (n = 4) 1 girl and 3 boys 6–11 years	4 sessions of 5 minutes each over 2 days. The mindfulness activities were part of the larger overall ACT+ RUBI ² treatment package (2-hr online individual training).	Behaviour parent training (the RUBI Autism Network).	Concurrent multiple baseline across participants (participants were randomly assigned in three tiers).	Responsiveness to ACT ³ was measured by the percentage of demonstrated objectives during the ACT portion of the training. This was scored as either a 2 (the parent correctly demonstrated the learning objective in full), 1 (the parent partially demonstrated the learning objective), 0 (the parent did not demonstrate the learning objective), or not applicable (the objective was not covered in the session). All parents demonstrated 100% responsiveness to ACT and ACT homework.

¹ Autism Spectrum.
² Research Units in Behavioural Intervention.
³ Acceptance and Commitment Therapy.

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Chan & Neece (2018) Neece et al. (2019) USA	80 parents 96.3% mothers 37.21 (7.22)	Developmental delay 63.6% on the AS. 71.25% boys 2.5–5 years 4.18 (1.01)	MBSR ⁴ . 8 weekly 2-hr sessions, a day-long 6-hr meditation retreat after the sixth session, and daily home practice based on audio CDs with instructions.	/	Waitlist-control design (39 and 41 parents were randomly assigned respectively to the treatment or waitlist-control group).	Two trained research assistants assessed treatment fidelity each session using a treatment fidelity checklist developed for this project. Inter-rater reliability was high with 95.04% agreement between the two raters. In the treatment group, 73.27% (SD = 16.60) of the treatment content items were covered, compared to 78.03% in the control group (SD = 9.93), $t(34) = -1.046$, $p = .305$.
Jones et al. (2018) UK	18 mothers and 3 fathers (including 1 mother–father couple) 45 (6.48)	AS (n = 16) Down’s syndrome (n = 1) Cerebral palsy (n = 2) Intellectual disorder (n = 2) 8 girls and 13 boys 4–16 years 10.53 (3.16)	MBW-P ⁵ . 8 weekly 2-hr sessions and daily home short practice from 5 to 20 min based on audio CDs instructions.	/	A pilot pre-post comparison of treatment.	/

⁴ Mindfulness-Based Stress Reduction.

⁵ Mindfulness Based Well-Being for Parents.

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
McGregor et al. (2020) USA	Treatment group: 39 parents 37.71 (8.38) Control group: 41 parents 36.76 (6.06)	80 children with developmental delays and AS Treatment group: n = 39 (56.40% AS) 66.70% male 4.01 (0.94) Control group: n = 41 (65.90% AS) 75.60% male 4.34 (1.05)	MBSR. 8 weekly 2-hr sessions, a day-long meditation retreat, and daily home practice based on audio CDs with instruction.	/	Waitlist-control design (39 and 41 parents were randomly assigned to the immediate treatment or waitlist-control group, respectively).	/
Neece (2014) USA	46 parents 78.3% mothers Treatment group: 21 parents 34.15 (8.71) Control group: 25 parents 36.40 (8.41)	Children with developmental delays, 84.8% on the AS Treatment group: 66.70% boys 3.59 (0.88) Control group: 76.50% boys 4.12 (0.90)	MBSR. 8 weekly 2-hr sessions, a day-long 6-hr meditation retreat after the sixth session, and daily home practice based on audio CDs with instructions.	/	Waitlist-control design (21 and 25 parents were randomly assigned to the immediate treatment or waitlist-control group, respectively)	/
Raulston et al. (2019) USA	3 mothers 31–34 years	AS (n = 3) 1 girl and 2 boys 4 years 4 months – 8 years 7 months	1 weekly 1.5-hr training sessions with homework in between. Formal mindfulness meditation practice (non-judgemental awareness or attention to internal and external experiences) + homework (listening to sound meditations on a mobile app).	Behaviour parent training.	Single-case concurrent multiple baseline across mother–child dyad design.	An independent observer viewed screencasts to assess treatment fidelity using a task analysis for 33% of sessions. Mean treatment fidelity was 100%.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Rojas-Torres et al. (2023) Spain	<p>Group A: 7 parents Fathers: 48-51 49 (1.73) Mothers: 41-47 43.5 (2.64)</p> <p>Group B: 7 parents Fathers: 30-48 40.25 (7.5) Mothers: 30-48 39.57 (7.5)</p>	<p>AS (n = 8) 1 girl and 7 boys</p> <p>Group A: 36– 46 months 41 (6)</p> <p>Group B: 48– 58 months 51 (1.5)</p>	<p>8 weekly 1.5-hr Mindfulness Parenting Programme sessions with homework in between (15-30 minutes per day).</p> <p>Each session was divided into two parts: development and training in Mindfulness exercises, and developing psychoeducation.</p>	<p>Psychoeducation sessions on AS.</p> <p>Children were cared for by a group of volunteer students guided by a member of the research team. During this period of time the children engaged in non-directive play activities on the floor with the intention of improving communication and social interaction.</p>	<p>Waitlist-control group design (Parents Group A and Group B assigned to the immediate intervention or wait-list-control group, respectively. Allocation of participants to the groups was not random and was determined according to the couple's possibilities of reconciling work and work schedules)</p>	/

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Schwartzman et al. (2022) USA	Treatment group: 17 parents 13 mothers 39.5 (3.7) Control group: 17 parents 13 mothers 42.5 (5.4)	Treatment group: AS (n = 17) 14 male 6.6 (1.8) Control group: AS (n = 17) 14 male 6.7 (1.9)	AMOR ⁶ Method. 8 weekly 1.5-hr group sessions (6–9 parents). The content for 2 of 8 sessions was based on treatments, including mindfulness from MBSR and values and committed actions from ACT. At the start of each session, parents engaged in a brief meditation exercise as a group and exercises included: (a) self-compassion break, (b) mindful breathing from MBSR, or (c) <i>Leaves on a Stream</i> from ACT.	The AMOR method also includes acceptance, optimism and resilience. The majority of sessions utilised principles of CBT ⁷ , such as psychoeducation, behavioural activation, identifying cognitive distortions, cognitive restructuring, and weekly homework assignments and review.	RCT ⁸ of two parallel groups: immediate treatment (AMOR) and DTG ⁹	/

⁶ Acceptance, Mindfulness, Optimism, Resilience.

⁷ Cognitive Behavioural Therapy.

⁸ Randomised Controlled Trial.

⁹ Delayed Treatment Group.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Singh et al. (2006b) USA	3 mothers 24–33 years	AS (n = 3) 4 years 5 months – 6 years 1 month	Mindfulness Parent Training Programme. 12 individual 2-hr sessions (3 sessions per week at Weeks 3, 6, 9 and 12). During the initial session, the senior investigator explained the details of the training and the philosophy of mindfulness. The mothers were taught meditation methods to enhance their mindfulness, and exercises to help them practice mindfulness during their interactions with their children. Mindfulness Practice Phase: This phase immediately followed the mindfulness training phase and lasted 52 weeks. The mothers were requested to continue with all the mindfulness exercises and use all their mindfulness skills in interactions with their children (homework).	/	Multiple baseline across subjects (parent–child dyad design).	/

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Singh et al. (2014) USA	3 mothers 37–43 years	Asperger (n = 3) 3 boys 15–19 years	<p>MB component of the MBPBS¹⁰. A month before the beginning of baseline, mothers were taught basic Samatha meditation during a 2-hr individual pre-training session. In addition, the mothers were taught Kinhin and insight meditations. Following the pre-training, they were instructed to develop a personal meditation practice, beginning with a few minutes each day and incrementally increasing their practice until they reached 20–30 min of daily practice.</p> <p>By the beginning of the baseline phase, each mother had practised the three foundational meditations daily for 4 weeks and was required to continue with these meditations during this phase.</p> <p>MB component training: 8 weekly 1-day individual training sessions. The training included instruction on the four immeasurables (loving-kindness, compassion, joy, and equanimity), the three poisons (attachment, anger, and ignorance), Shenpa and compassionate abiding, and meditation on the soles of the feet.</p> <p>MB component practice: Mothers were requested to mindfully utilise the PBS techniques in their interactions with their children.</p>	<p>PBS components of the MBPBS: Throughout the MBPBS training, mothers were taught how to use PBS within the context of mindfulness practices (the mothers had been taught the basics of PBS procedures when their children were between 6 and 8 years of age).</p>	Multiple baseline design across participants, with three phases: baseline, mindfulness training, and mindfulness practice.	<p>Teaching sessions were videotaped, and 6 randomly chosen sessions, stratified by mothers, were assessed for fidelity of training by another qualified mindfulness trainer. The fidelity of mindfulness training was rated at 100%. Training adherence: The daily logs showed that the three mothers were highly diligent in their meditation practice. On average, they engaged in 5–10 min of meditation practice, first thing in the morning, every day during the first week of pre-training. They gradually increased their meditation sessions to 30–40 min a day by the end of the MBPBS training phase. They were able to maintain and increase their daily meditations for up to an hour each day by the end of the MBPBS practice phase. On average, the mothers sat for 93% of the days during the MBPBS practice phase, for from 45–60 min.</p>

¹⁰ Mindfulness-Based Positive Behavior Support.

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Singh et al. (2019) USA	AS group: 47 mothers 39–59 years 47.91 ID group: 45 mothers 38–55 years 48.82	AS (n = 47) 66% male 13–17 years 15.15 ID (n = 45) 67% male 13–17 years 15.56	3-day MBPBS protocol in the intervention phase. The first day was devoted to training and practice in mindfulness procedures that included 3 basic meditations, 4 immeasurables, 5 hindrances, 3 poisons, beginner’s mind, and ethical precepts. It also covered issues related to home practice, daily logs of meditation, journaling, and data collection. The second day was devoted to developing PBIS ¹¹ plans. The third day was devoted to a review and practice of daily meditation practices for the mothers, day-to-day implementation of an integrated MB positive behavioural intervention with their child, living mindfully with their family on a daily basis, informal mindfulness practices, collection and interpretation of outcome data, and frequently asked questions. Following the 3-day training, the mothers implemented the MBPBS procedures in daily life.	PBIS plans.	Equivalent diagnostic 2-group design, with a control and intervention condition. In a 40-wk study, the first 10 weeks constituted a control phase during which no additional procedures were instituted, followed by a 3-day training in MBPBS for both groups, and 30 weeks of implementation. The training was the same for both groups.	The training was videotaped for 10–12 min per hour per day during random segments of each hour of training. 24 videotaped segments were rated for fidelity of MBPBS training by two independent raters, one a mindfulness expert and the other an expert in PBS. The fidelity of MBPBS training was rated at 100% for meditation instructions and principles.

¹¹ Positive Behavior Intervention and Support.

Citation and location	Number, gender and age range of parents, mean (SD)	Diagnostic, number, gender and age range of children, mean (SD)	Mindfulness-based interventions	Additional interventions	Study design	Implementation process assessment
Singh et al. (2021) USA	<p>MBPBS condition: 65 mothers 33–49 years 40.03 (4.08)</p> <p>MB condition: 65 mothers 29–46 years 36.74 (4.63)</p> <p>PBS condition: 65 mothers 28–48 years 38.09 (4.83)</p>	<p>MBPBS condition: AS (n = 65) 76.92% male 9–14 years 12.17 (1.62)</p> <p>MB condition: AS (n = 65) 72.31% male 9–14 years 11.80 (1.65)</p> <p>PBS condition: AS (n = 65) 73.85% male 9–14 years 11.95 (1.68)</p>	<p>MBPBS condition: 3-day stepped-care version of the MBPBS protocol for teaching mindfulness, related meditation practices, and guiding principles of PBS (the training was provided in small groups at locations close to the participants or via telemedicine).</p> <p>MB component of the MBPBS: mothers were taught mindfulness and related meditation practices from the 3-day stepped-care version of the MBPBS protocol.</p> <p>Mothers were encouraged to engage in formal meditation for at least 20 min a day.</p>	<p>PBS components of the MBPBS: Parents learned the PBS principles by working through case examples, and by developing PBS plans for their own children.</p>	<p>3-arm RCT, with a 10-week pre-intervention control condition. The experimental condition included a 3-day training in each active experimental condition (i.e. MBPBS, MB, and PBS). The training in each experimental condition was followed by 30 weeks of implementation.</p>	<p>Training provided in all 3 experimental conditions was videotaped for 15 min each hour per day during random segments of the training. Two independent raters, trained in both mindfulness and PBS, rated 24 videotaped segments for fidelity assessment across the 3 experimental conditions. Two aspects of fidelity were assessed: structural fidelity (i.e. what is being taught) and process fidelity (i.e. how the contents are being taught). Both aspects of fidelity were assessed at 100% for all three experimental conditions.</p>
Weitlauf et al. (2022) USA	<p>P-ESDM¹² condition: 31 parents 87% mothers 33.79 (5.53)</p> <p>P-ESDM + MBSR condition: 30 parents 87% mothers 33.27 (6.24)</p>	<p>P-ESDM condition: AS (n = 31) 81% male 2.46 (1.64)</p> <p>P-ESDM + MBSR condition: AS (n = 30) 83% male 2.30 (0.45)</p>	<p>MBSR. 6 sessions, 1 hr per week.</p>	<p>P-ESDM. 12 consecutive weekly clinic-based sessions. Each lasted approximately 1 hr and included the child, one parent, and a therapist. Speech, occupational and ABA¹³ therapy in a variable way depending on the child.</p>	<p>Longitudinal RCT of P-ESDM with and without MBSR.</p>	<p>Fidelity of the MBSR intervention was tracked using fidelity checklists that therapists completed at the end of each visit. The fidelity of covering the required topics across all completed sessions was 97.9%.</p>

¹² Parent-Implemented Early Start Denver Model.

¹³ Applied Behaviour Analysis.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 2. Methodological quality scores and levels of evidence

Citation and location	Child outcomes (measures)	Follow-up	Results	STROBE rating (%)	NHMRC level of evidence
Andrews et al. (2022) USA	Children’s rate per minute of challenging behaviours (measured during target routines by the experimenter). Frequency and severity of challenging behaviours (4-point rating scale measured by parents each day).	1 week and 1 month	<p>Children’s rate per minute of CB¹⁴ for all 4 children. During baseline: ($M = 0.1, SD = 0.2$); ($M = 0.5, SD = 0.5$); ($M = 1.4, SD = 0.9$); ($M = 0.4, SD = 0.5$). During intervention: ($M = 0.8, SD = 0.5$); ($M = 0.7, SD = 1.1$); ($M = 0.2, SD = 0.3$); ($M = 0.2, SD = 0.2$). During the 1-wk FU¹⁵: increased CB (Child 1); at the low end of baseline range (Child 2 and 4); zero responses (Child 3). During the 1-mth FU: zero CB (Child 1); same level of CB (Child 2 and Child 4); increase CB (Child 3).</p> <p>Frequency of CB. Between baseline and intervention: CB decreased for each child. Between baseline and 1-wk FU: CB decreased (Child 1, 2, 3); increased CB (Child 4). At 1-mth FU: CB decreased (Child 1, 2, 4); increased CB return to baseline (Child 3).</p> <p>Severity of CB. Between baseline and intervention: CB decreased (Child 2, 3, 4); CB slight increase (Child 1) At 1-wk FU: CB decreased (Child 1, 2); no change for Child 3; slight increase returning to baseline (Child 4). At 1-mth FU: CB decreased (Child 1, 2, 4); CB increased (Child 3)</p>	89	III-3

¹⁴ Challenging Behavior.

¹⁵ Follow-up.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Citation and location	Child outcomes (measures)	Follow-up	Results	STROBE rating (%)	NHMRC level of evidence
Chan & Neece (2018); Neece et al. (2019) USA	Parents rated child behavioural problems (CBCL ¹⁶ for ages 1.5–5 years).	6 months	Withdrawn significantly predicted by TG ¹⁷ × TI ¹⁸ , $F(1, 65) = 4.51, p < .05$, partial $\eta^2 = .07$. Decrease in being withdrawn from baseline ($M = 6.19, SD = 2.62$) to second assessment ($M = 4.84, SD = 2.81$) for TG. AP ¹⁹ significantly predicted by TG × TI, $F(1, 65) = 10.67, p < .01$, partial $\eta^2 = .14$. Decrease in AP from baseline ($M = 5.41, SD = 2.60$) to second assessment ($M = 4.41, SD = 2.28$) for TG. TG × TI effect was marginally significant for emotionally reactive, $F(1, 65) = 2.88, p < .095$, partial $\eta^2 = .04$. Anxious/depressed, somatic complaints and aggressive behaviour not significantly predicted by the TG × TI ($p > .05$).	93 & 91	III-1
Jones et al. (2018) UK	Child’s adaptive functioning (VABS-II ²⁰ , administered as a semi-structured interview). Children’s prosocial behaviour and behavioural difficulties (SDQ ²¹).	/	No significant change in child’s behavioural problems or prosocial behaviour.	86	III-3
McGregor et al. (2020) USA	Parents rated child’s behavioural problems. (Internalising Scale of the CBCL for ages 1.5–5 years).	6 months (unprocessed data)	Greater reductions in internalising problems in the MBSR TG compared to the waitlist-control group ($b = -5.71, 95\% \text{ CI } [-10.55, -.87], sr^2 = .04, p < .05$).	89	III-1

¹⁶ Child Behavior Checklist.

¹⁷ Treatment Group.

¹⁸ Time interaction.

¹⁹ Attention Problems.

²⁰ Vineland Adaptive Behavior Scales, Second Edition.

²¹ Strengths and Difficulties Questionnaire.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Citation and location	Child outcomes (measures)	Follow-up	Results	STROBE rating (%)	NHMRC level of evidence
Neece (2014) USA	Child behavioural problems (CBCL for ages 1.5–5 years).	/	Significantly fewer ADHD ²² signs in the TG in the control group at the second assessment ($t = 2.27, p < 0.05, d = 0.85$), and marginally significant difference in attention signs ($t = 1.89, p = 0.69, d = 0.71$). Clinically significant reduction in ADHD signs for 28.9% of parents. Meaningful change in attention problems for 18.4% of parents.	93	III-1
Raulston et al. (2019) USA	Interview with parent about child's challenging behaviour (Externalising subscale from the CBCL).	/	Reduction in behavioural problems for the 3 children during the intervention (Hedges' $g = 0.24$).	86	III-3
Rojas-Torres et al. (2023) Spain	Parent perception of autism signs. (AIM ²³ . It is a measure of frequency and impact of behaviors and symptoms characteristic of autism)	8-week follow-up	Parents' assessment of sign severity unchanged across the measures.	71	III-1
Schwartzman et al. (2022) USA	Parent perceptions of children: Strengths/prosocial behaviours and difficulties/problem behaviours of children (SDQ). SRS -2 ²⁴ . Type and severity of problem behaviour (ABC-2 ²⁵).	8-week follow-up for AMOR parents only	More positive ratings of child's total difficulties (SDQ; $d = 1.06, p = 0.032, 95\% \text{ CI } [-3.871, 0.203]$) and emotional problems (SDQ; $d = 1.55, p = 0.003, 95\% \text{ CI } [-2.190, -0.475]$), social communication deficits (SRS-2; $d = 0.88, p = 0.025, 95\% \text{ CI } [-6.771, -1.479]$), and inappropriate speech (ABC-2; $d = 0.76, p = 0.043, 95\% \text{ CI } [-1.264, 0.514]$) for AMOR group relative to the TG. No significant changes in the remaining subscales of the SDQ, SRS-2, or ABC-2 by TG. Maintenance of treatment gains of parent perceptions of children with AS on all measures. Significant decrease in parent ratings of child hyperactivity (SDQ).	95	II

²² Attention Deficit Hyperactivity Disorder.

²³ Autism Impact Measure.

²⁴ Social Responsiveness Scale, Second Edition.

²⁵ Aberrant Behavior Checklist, Second Edition.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Citation and location	Child outcomes (measures)	Follow-up	Results	STROBE rating (%)	NHMRC level of evidence
Singh et al. (2006b) USA	3 behaviours were targeted for observation: aggression, noncompliance, and self-injury. Data were collected during the waking hours of each child when at home with mother, between 8 and 14 hrs each day. Each child's mother recorded the occurrence of each target behaviour, in real-time, on a Palm PDA ²⁶ . Fathers collected data on a separate Palm PDA for 5 to 10 hrs each week, typically in the evenings and on weekends, to establish inter-rater agreement.	/	First parent-child dyad: Aggressive behaviours decreased from baseline (8.8) to training (7.4) to practice (0.9). Noncompliance behaviours decreased from baseline (53.6) to training (35.7) to practice (11.2). Second dyad: Aggressive behaviours decreased from baseline (17.1) to training (16.0) to practice (4.9). Noncompliance behaviours decreased from baseline (37.2) to training (32.9) to practice (11.8). Third dyad: Aggressive behaviours decreased from baseline (8.0) to training (7.2) to practice (1.1). Self-injurious behaviours decreased from baseline (10.4) to training (8.7) to practice (4.2).	75	III-3
Singh et al. (2014) USA	Three behaviours were targeted: incidents of aggression, incidents of disruptive behaviour, and compliance with mother's requests. (Data were collected at all times when the adolescent was at home and in the presence of their mother. Each mother collected data using an iPhone app during the three phases: baseline, MBPBS training, and MBPBS practice. The fathers served as reliability raters, collecting data using the same system as the mothers, but on their own iPhones, for an average of 3 hrs each week, typically in the evenings or weekends).	/	Decreased aggressive incidents and disruptive behaviours ($\Phi = 0.76, p < 0.001$ and $\Phi = 0.93, p < 0.001$, respectively). Increased in compliance percentage ($\Phi = 0.86, p < 0.001$).	89	III-3

²⁶ Personal Digital Assistant.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Citation and location	Child outcomes (measures)	Follow-up	Results	STROBE rating (%)	NHMRC level of evidence
Singh et al. (2019) USA	Aggressive behaviour, disruptive behaviour and compliance with mother's requests. (The mothers collected data on their adolescent's behaviour using an app that allowed recording of multiple events in real-time. On average, the mothers collected data for about 5 hrs during weekdays and about 8 hrs during weekends. To establish inter-rater agreement, fathers also collected data using the same system but independently of the mothers, for an average of 2 hrs each week, usually during evenings and weekends).	/	Decreased aggressive and disruptive behaviour from control to intervention phase (aggressive behaviour: $\phi = 0.71, p < 0.001$; disruptive behaviour: $\phi = 0.86, p < 0.001$). Aggressive behaviour decreased from control ($M = 14.00, SD = 2.31$) to intervention ($M = 3.33, SD = 3.92$). Disruptive behaviour decreased from control ($M = 38.80, SD = 7.47$) to intervention ($M = 9.20, SD = 8.25$). Percent compliance increased ($\phi = 0.86, p < 0.001$) from control ($M = 20.10, SD = 2.85$) to intervention ($M = 55.83, SD = 22.50$).	86	III-3
Singh et al. (2021) USA	Aggressive behaviour, disruptive behaviour and compliance with mother's requests. Data were collected by the mothers using an app that enabled real-time recording of multiple events. The mothers collected event data when their children were at home or in community settings and in their presence. On average, data collection occurred for 4–6 hrs during the week and 7–9 hrs during weekends. Fathers independently collected data for an average of 2 hrs per week, during evenings and weekends to establish interobserver agreement.	3 years of post-intervention follow-up (variables monitored each year for 3 years)	Aggression and disruptive behaviour most to least significant reductions in MBPBS, MB, and PBS, respectively. Significant increases in compliance largest in the MBPBS, followed by MB, and PBS. Changes were maintained for 3 years post-intervention. When accounting for the effects of time and MBPBS, both MB and PBS were not significant predictors. Meditation time explained 29% of the reduction in the children's aggressive behaviour ($R^2 = 0.29; \beta = -0.88; p < 0.001$). When controlling for the effects of time and type of intervention, meditation practice explained 24% of disruptive behaviour decrease ($R^2 = 0.24; \beta = -0.79; p < 0.001$). When accounting for the effects of time and MBPBS, MB made a significant contribution of an additional 5% ($R^2 = 0.05; \beta = 0.26; p = 0.002$). When controlling for time and type of intervention, meditation made a significant contribution ($R^2 = 0.27; \beta = 0.85; p < 0.001$).	91	II

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Citation and location	Child outcomes (measures)	Follow-up	Results	STROBE rating (%)	NHMRC level of evidence
Weitlauf et al. (2022) USA	Autism signs: ADOS-2 ²⁷ . Cognitive assessment: MSEL ²⁸ . Adaptative functioning: VABS-II.	6 months	<p>Scores on the ADOS-2 decreased from T1 to T2 (− 0.82 points; $p < 0.01$), but did not change from T2 to T3 (0.05 points; $p > 0.10$). Early learning composite scores of MSEL did not change from T1 to T2 (0.51 points, $p > 0.10$), but increased from T2 to T3 (3.94 points, $p < 0.01$).</p> <p>Raw scores increased for visual reception (T1–T2, 3.12 points, $p < 0.001$; T2–T3, 1.6 points, $p < 0.001$), receptive language (T1–T2, 6.00 points, $p < 0.001$; T2 to T3, 2.39 points, $p < 0.01$), expressive language (T1 to T2, 4.41 points, $p < 0.001$; T2 to T3, 3.5 points, $p < 0.001$), and fine motor (T1–T2, 2.09 points, $p < 0.001$; T2–T3, 2.49 points, $p < 0.001$).</p> <p>No significant change occurred in adaptive behaviour composite (T1–T2, 2.02 points, $p = 0.052$; T2–T3, − 0.40 points, $p > 0.10$).</p>	93	II

²⁷ Autism Diagnostic Observation Schedule, Second Edition.

²⁸ Mullen Scales of Early Learning.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 3. Significant improvement in children’s behaviour

Children’s behaviour	n*
Reduced frequency and/or severity of challenging behaviours	10
Reduced withdrawal, attention problems and somatic complaints	2
Hyperactivity signs	1
Emotional problems and inappropriate speech	1

* Number of studies.

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 4. Risk of bias assessment for the RCT studies. The risk of bias was assessed as low (+), high (-) or uncertain (?)

Citation	Generation of sequence (selection bias)	Concealment of allocation sequence (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessors (detection bias)	Incomplete outcome data (attrition bias)	Selective outcome reporting (reporting bias)	Treatment implementation
Schwartzman et al. (2022)	+	+	?	-	+	?	-
Singh et al. (2021)	+	?	?	-	?	?	-
Weitlauf et al. (2022)	+	+	?	?	+	?	+

Author Note

This submission is an original systematic review paper. We certify that this manuscript will not be submitted elsewhere during the *Review Journal of Autism and Developmental Disorders* review process. We confirm that all named authors provided substantial contributions to this study and have reviewed and approved the final version of this manuscript. We also all agreed to the listing-order for authors.

Funding: This research was conducted for a university degree and received no funding.

Acknowledgements: The first author (JL) would like to thank Dr Marine Colombel and Dr Jean Sixou for the wealth of their theoretical contributions.

Conflicts of interest: The authors declare that they have no conflicts of interest.

Dear Editor,

We would like to thank the reviewers for their latest comments on our manuscript entitled "Mindfulness meditation-based interventions in parents of autistic children: A systematic review of effects on children". Their comments helped us improving our manuscript. We hope the proposed changes meet your/their expectations.

In this document, we have replied to each comment separately (our answers appear in blue). We have also made changes to the original manuscript, which are also highlighted in blue.

REVIEWER 1

1. Given that the review period was extended from 17 December 2022 to April 2024, this should stated on page 4 as: "An electronic search was conducted for the period 2006 to April 2024." We added this information (see page 4).

This statement on page 5 should be omitted to avoid confusion: "We updated the database searches since the initial search date was more than 12 months from the intended publication date. The results of the updated searches, limited to the period between January 2023 and April 2024, were screened for potentially eligible studies." This statement will only make sense to the reviewers because new readers will not know that the extended date was requested by the reviewers.

We removed this sentence from the manuscript (see page 5).

2. In the Reference section, some of the article include the DOI but the majority of them do not. Be consistent.

We added the DOI.

REVIEWER 3

1. PP. 2, line 14 - In my opinion the phase "The particularities of autism" is not appropriate and should be revised.

We revised this phrase: "Autism can also impact the well-being of the family" (see page 2).

2. Pp. 3, lines 32-39 - The added definition of challenging behaviours is not clear. Please define externalising behaviours. See website of The Challenging Behaviours Foundation for definition that is accepted by family carers and people with challenging behaviours. <https://www.challengingbehaviour.org.uk/>

We revised the definition of challenging behaviours: "Challenging behaviours are defined as a range of behaviours that people with severe learning disabilities may display to get their needs met, such as hurting others, destructive behaviours or self-injurious behaviours" (see page 3).

3. Pp. 28, line 34-51 - Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines outline how systematic reviews should be reported, not conducted. Therefore, explanation provided by authors to justify only one person screening titles and abstracts is incorrect. For example, see Cochrane Handbook for Systematic Reviews of Interventions for guidelines on how systematic reviews should be conducted - <https://community.cochrane.org/mecir-manual/standards-conduct-new-cochrane-intervention-reviews-c1-c75/performing-review-c24-c75/selecting-studies-include-review-c39-c42> Since authors have done full text review in duplicate, the paper is still publishable. However, they need to describe their rationale for only one person doing initial screening

which is against good practice guidelines for systematic review (irrespective of systematic review reporting guidelines). The rationale provided on page 28 is not correct.

We concur that the screening by a single author represents a deviation from the guidelines; it is only justifiable by our misreading of the guidelines. Therefore, we added the following paragraph to the limitation sections:

“Thirdly, only one author screened the titles and abstracts during the initial phase. This shortcoming may have partially influenced the identification of articles. However, this issue is mitigated by the small number of available articles, which limits the risk of false negatives. Additionally, all the selected articles were fully read by two of the authors.” (see page 18).

4. Pp. 28, line 34-51 - Protocol registration prior to review starting is done for transparency of research methods used.

This was rephrased as follows: “Secondly, although the protocol registration prior to review starting is done for transparency of research methods used, the protocol was not registered, as one of the authors had already initiated literature searches” (see page 18).