



# The earliest stylophorans (Echinodermata) from South America: new cornutes from the Early Tremadocian (Lower Ordovician) of Jujuy Province, Argentina

Bertrand Lefebvre, Ninon Allaire, Romain Vaucher, N. Emilio Vaccari,  
Beatriz Waisfeld

## ► To cite this version:

Bertrand Lefebvre, Ninon Allaire, Romain Vaucher, N. Emilio Vaccari, Beatriz Waisfeld. The earliest stylophorans (Echinodermata) from South America: new cornutes from the Early Tremadocian (Lower Ordovician) of Jujuy Province, Argentina. Rocks and the Rise of Ordovician Life, 4th annual meeting IGCP 735, IGCP 735, Oct 2024, Cordoba, Argentine, Argentina. hal-04776048


HAL Id: hal-04776048

<https://hal.science/hal-04776048v1>

Submitted on 11 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Rocks and the Rise of Ordovician Life

---

4th Annual Meeting  
IGCP 735

---

Córdoba, Argentina  
2024

Book of Abstracts



exhaustive search for brachiopods is needed in order to confirm the occurrence of the widespread deep-water *Foliomena* Fauna in South America.

## **THE EARLIEST STYLOPHORANS (ECHINODERMATA) FROM SOUTH AMERICA: NEW CORNUTES FROM THE EARLY TREMADOCIAN (LOWER ORDOVICIAN) OF JUJUY PROVINCE, ARGENTINA**

BERTRAND LEFEBVRE<sup>1</sup>, NINON ALLAIRE<sup>2,3</sup>, ROMAIN VAUCHER<sup>4</sup>, N. EMILIO VACCARI<sup>2,3,5</sup> and BEATRIZ G. WAISFELD<sup>2,3</sup>

<sup>1</sup>Laboratoire de Géologie de Lyon: Terre, Planètes, Environnement (LGLTPE), Université Claude Bernard Lyon 1, 2 rue Raphaël Dubois, 69622 Villeurbanne cedex, France. [bertrand.lefebvre@univ-lyon1.fr](mailto:bertrand.lefebvre@univ-lyon1.fr)

<sup>2</sup>Facultad de Ciencias Exactas, Físicas y Naturales, Universidad Nacional de Córdoba, Avenida Vélez Sarsfield 1611, CP X5016GCA Córdoba, Argentina. [ninon.allaire@gmail.com](mailto:ninon.allaire@gmail.com), [evaccari@unc.edu.ar](mailto:evaccari@unc.edu.ar), [bwaisfeld@unc.edu.ar](mailto:bwaisfeld@unc.edu.ar)

<sup>3</sup>Centro de Investigaciones en Ciencias de la Tierra (CICTERRA), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Avenida Vélez Sarsfield 1611, CP X5016GCA Córdoba, Argentina. [ninon.allaire@gmail.com](mailto:ninon.allaire@gmail.com), [evaccari@unc.edu.ar](mailto:evaccari@unc.edu.ar), [bwaisfeld@unc.edu.ar](mailto:bwaisfeld@unc.edu.ar)

<sup>4</sup>Department of Earth Sciences, Université de Genève, 13 rue des Maraîchers, 1205 Genève, Switzerland. [romain.vaucher@unige.ch](mailto:romain.vaucher@unige.ch)

<sup>5</sup>Universidad Nacional de La Rioja, Avenida Luis M. de la Fuente s/n, F5300, La Rioja, Argentina. [evaccari@unc.edu.ar](mailto:evaccari@unc.edu.ar)

Stylophorans are a class of Palaeozoic echinoderms (Miaolingian–Bashkirian) characterized by the possession of a single feeding appendage (aulacophore) and a flattened, asymmetrical body (theca). Along with eocrinoids, glyptocystitid rhombiferans, and solutans, stylophorans were a major component of the low diversity cosmopolitan echinoderm assemblages spanning the Furongian–mid Tremadocian interval, before the diversification of crinoids. In northwestern Argentina (eastern Cordillera of Jujuy Province, Angosto del Moreno locality), the Pupusa Formation (Upper Member of the Guayoc Chico Group) reflects deposition in an offshore environment ending in a shoreface environment. The Pupusa Formation yielded a remarkably well-preserved assemblage of early Tremadocian (Tr1) echinoderms comprising eocrinoids and cornute stylophorans, found at the base of storm-induced sandstone beds in offshore settings. The cornute can be assigned to the scotiaecystid genus *Thoralicyctis*, which is also present in the Lower Ordovician of the Montagne Noire (France) and the Anti-Atlas (Morocco). Several mitrate stylophorans have already been described in the Lower and Middle Devonian of South America (Argentina, Bolivia and Brazil). Therefore, the new Ordovician cornute reported here from the Pupusa Formation (Jujuy Province, Argentina) represent the earliest record of the class Stylophora in South America.