



HAL
open science

Evaluation of Nutritional Composition of *Ocimum gratissimum* Leaf Extract

Moneme, E. C, Nwaka, A. C, Ajakpofo, F. O, Anyanwu, R. O, Onuegbu, M. E

► **To cite this version:**

Moneme, E. C, Nwaka, A. C, Ajakpofo, F. O, Anyanwu, R. O, Onuegbu, M. E. Evaluation of Nutritional Composition of *Ocimum gratissimum* Leaf Extract. *International Journal of Biochemistry Research & Review*, 2024, 33 (6), pp.415 - 419. <https://doi.org/10.9734/ijbcr/2024/v33i6923> . hal-04776090

HAL Id: hal-04776090

<https://hal.science/hal-04776090v1>

Submitted on 11 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Evaluation of Nutritional Composition of *Ocimum gratissimum* Leaf Extract

Moneme, E. C. ^{a*}, Nwaka, A. C. ^a, Ajakpofo, F. O. ^b,
Anyanwu, R. O. ^b and Onuegbu, M. E ^b

^a Department of Biochemistry, Chukwuemeka Odumegwu Ojukwu University, Uli, Nigeria.

^b Department of Applied Biochemistry, Nnamdi Azikiwe University, Awka, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijbcrr/2024/v33i6923>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126300>

Original Research Article

Received: 02/09/2024

Accepted: 05/11/2024

Published: 09/11/2024

ABSTRACT

Vegetables are known to be vital sources of nutritive foods. This study evaluates the nutritional composition of *Ocimum gratissimum* leaf extract vis-a-vis proximate, minerals and vitamins analyses. Fresh leaves of *O. gratissimum* were purchased from a local market in Awka North Local Government Area of Anambra State, Nigeria and were washed, destalked, oven-dried at 40 °C, pulverized, and sieved through 2.0 mm mesh prior to analysis. Proximate, minerals, and vitamin analyses were carried out, using standard methods. Results indicated that *O. gratissimum* contained high levels of carbohydrate (51.78 ± 0.50 %), protein (10.50 ± 0.04 %), and fiber (9.38 ± 0.05 %), with minute quantity of crude fat (4.94 ± 0.03 %). Analysis of the mineral compositions showed that the vegetable contained considerable quantity of minerals that is sufficient to improve general well-being, while the vitamin analysis revealed that the leaves of *O. gratissimum* was sufficiently rich in Vitamin E and could, therefore, act as antioxidant and consequently help to improve health status of its consumers.

*Corresponding author: E-mail: emekamoneme5@gmail.com;

Cite as: E. C., Moneme, Nwaka, A. C., Ajakpofo, F. O., Anyanwu, R. O., and Onuegbu, M. E. 2024. "Evaluation of Nutritional Composition of *Ocimum Gratissimum* Leaf Extract". *International Journal of Biochemistry Research & Review* 33 (6):415-19. <https://doi.org/10.9734/ijbcrr/2024/v33i6923>.

Keywords: Green leafy vegetables; *Ocimum gratissimum*; nutrition; health.

1. INTRODUCTION

“Green leafy vegetables are widely succulent plants parts grown in gardens and consumed as a side dish or soup with starchy staples among the tribes in Nigeria” [1]. “The importance of dietary components of leafy vegetables is significant in African population since they are comparatively rich in fiber, whereas cereals, root vegetables, and other food are relatively poor sources” [2]. “Similar to other African countries, a typical Nigerian diet is dominated by starchy staple foods, and vegetables are considered a cheap source of energy. Vegetables are major sources of nutrients such as carotene, protein, vitamins, calcium, iron, and ascorbic acid, and they also contain tangible concentration of trace minerals” [3]. “Many species of vegetables abound in the world, and they are utilized in addition of flavor, taste, and color to diet which would otherwise be monotonous. The invaluable food nutrients contained in vegetables can be used for body growth, protective, and regulatory roles as well as sources of energy for the body” [2].

“Vegetables also contain plants secondary metabolites which act as anti-nutrients” [4]. “These secondary metabolites include: tannins, phytate, oxalate, and cyanide; all of which have various detrimental effects to nutrition and health” [5]. A good leafy vegetable is the one that contains high amount of these beneficial nutrients and a very minute amount of anti-nutrients. One of the promising vegetables that has been acclaimed ‘good’ is *Ocimum gratissimum*.

“*O. gratissimum* Linn (Lamiaceae) is a perennial herbaceous shrub notably found in tropical countries, including Nigeria” [6], where it is commonly called Clove basil, Sweet basil, tea-bush, Scent leaf or fever plant. It is also popularly known with different local names in Nigeria as “*Daidoya*” (hausa), “*Effirin*” (yoruba), and “*Nchuanwu*” (igbo). Due to its strong aromatic smell, it is used as a natural flavoring agent, condiment, or vegetable in the preparation of fish, meat, soup, and stew.

“Scientific reports have shown that *O. gratissimum* has a wide range of bio-active compounds such as flavonoids, polyphenols” [7] and essential oils with several beneficial effects [8]. Since the nutrient contents vary from one

leafy vegetable species to another, there is a need to investigate these variations within the vegetables. Hence, this study aimed to evaluate the nutritional composition of *O. gratissimum*.

2. MATERIALS AND METHODS

2.1 Plant Materials

Fresh leaves of *O. gratissimum* were purchased from a daily market, Mgbakwu in Awka North Local Government Area of Anambra State, Nigeria and were authenticated by a Taxonomist, Mr Iroka Finan, from the Department of Botany, Nnamdi Azikiwe University, Awka. A voucher specimen was also deposited with Herbarium no NAUH 35B. The leaves were then detached from the stalk, rinsed with distilled water and were oven-dried at 40 °C and weighed (300 g). The weighed powdered sample (300 g) was then used for the extraction with a solvent combination of ethanol and water (7:3) (2500 ml) for 48 hr via maceration in an unheated medium. The mixture was decanted and filtered using sterile Whatman paper No. 1. The filtrate was there after evaporated to dryness with the aid of a rotary evaporator set at 50 °C to obtain crude ethanol extract (25.2 g) which was carefully preserved for further analysis.

2.2 Proximate Analysis

Moisture, ash, crude fat, and crude fiber were analyzed in the fresh samples in accordance with the methods described by Association of Official Analytical Chemist [9], while the percentage of nitrogen was determined by the micro-Kjeldahl method (1976) and was converted to crude protein by multiplying by 6.25. Carbohydrate was determined by difference [10].

2.3 Vitamin Analysis

Vitamin A and C were determined by the Futtarmayer colorimetric method of Kirk and Sawyer [11], while Vitamin E was determined using the Emmerie-Engel reaction for Tocopherol estimation as reported by Rosenberg [12].

2.4 Mineral Analysis

Wet ashing method was used to release the investigated metals in the samples from biological complexes before evaluating them with atomic absorption spectrophotometer

(PinAAcle™ 900H) and flame photometer (Gulfex FP6410). Wet ashing was achieved by heating the sample with a digestion mixture containing concentrated nitric, perchloric, and sulphuric acids in a Kjeldahl flask [9]. “Calcium, Iron, Manganese, Magnesium, and Zinc were determined by atomic absorption spectrophotometer; Sodium and Potassium by Flame photometer while Phosphorus was determined by Molybdate method using hydroquinone as a reducing agent” [9].

2.5 Data Analysis

The mean values obtained and the significance between the treated and control group was analyzed by one-way ANOVA using the SPSS version 17 and $P < 0.05$ was considered to be statistically significant.

3. RESULTS AND DISCUSSION

Table 1 showed the result for proximate analysis of the leaf extract of *O. gratissimum*. Result indicated highest content of Carbohydrate ($51.78 \pm 0.50\%$), with minimal content of crude fat ($4.94 \pm 0.03\%$).

Table 1. Percentage proximate composition of the leaf extract of *O. gratissimum*

Proximate components	Percentage composition (%)
Moisture	11.92 ± 0.02
Ash	11.48 ± 0.10
Fiber	9.38 ± 0.05
Crude fat	4.94 ± 0.03
Protein	10.50 ± 0.04
Carbohydrates	51.78 ± 0.50

Values are expressed in mean±SEM

Table 2 showed the result for vitamin analysis of the leaf extract of *O. gratissimum*. Result indicated that the sample was rich in vitamin E ($35.85 \pm 1.56 \mu\text{g/g}$), with minimal content of vitamin A ($1.86 \pm 0.07 \mu\text{g/g}$).

Table 2. Vitamin composition of the leaf extract of *O. gratissimum*

Vitamins	Concentration ($\mu\text{g/g}$)
A	1.86 ± 0.07
C	3.40 ± 0.00
E	35.85 ± 1.56

Values are expressed in mean±SEM

Table 3 showed the result for mineral composition of the leaf extract of *O. gratissimum*.

Result indicated that the sample contained considerable quantity of iron ($1.40 \pm 0.50 \text{ ppm}$) with minute quantity of copper ($0.29 \pm 0.17 \text{ ppm}$).

Table 3. Mineral composition of the leaf extract of *O. gratissimum*

Minerals	Quantity (ppm)
Copper	0.29 ± 0.17
Iron	1.40 ± 0.50
Manganese	0.31 ± 0.08
Zinc	0.79 ± 0.10
Selenium	0.37 ± 0.02

Values are expressed in mean±SEM

The results of the proximate analysis of the leafy vegetable was represented in Table 1. Generally, the results indicated that the vegetable was a good source of carbohydrates protein, and crude fiber. Carbohydrate content was much higher ($51.78 \pm 0.50 \%$) compared to other macronutrients. The high Carbohydrate content of *O. gratissimum* was in accordance with the reports of Hussain et al. [13] and Oladejo [14]. *O. gratissimum* met the requirement by providing more than 12% of its calorific value from protein and could, thus, be considered a good source of protein according to Ajayi et al. [1]. However, the quality of a protein is dependent on its ability to supply all the essential amino acids in the required amount [15]; hence, protein quality assessment is necessary to ascertain the ability of *O. gratissimum* to meet dietary protein requirements.

“The microelements: copper, iron, manganese, and zinc were also well represented in the vegetable. Although the Dietary Reference Intake (DRI) of 20.5 mg of iron cannot be met by the vegetable at 100g per day serving, the vegetable is nevertheless a good source of iron as supplementation with other food components will make up for the iron need. Epidemiological studies have indicated that diets high in iron can help to fight infections, boost brain health, and promote systemic circulation of oxygen” [14]. “Manganese is an essential metal because it is required for proper immune function, regulation of blood sugar and cellular energy, reproduction, digestion, bone growth, blood coagulation, and homeostasis as well as defense against reactive oxygen species” [16]. The beneficial effects of manganese are due to the incorporation of the metal into metalloproteins, thus aiding biochemical functions. The minute quantity of manganese ($0.31 \pm 0.08 \text{ ppm}$) present in this leafy vegetable indicates that it could provide the

body with the minimum concentrations of manganese required for basic enzyme activities.

Similarly, the zinc concentrations present in the vegetable as observed in this study were found to be comparable with previous works [1,14]. "Zinc is highly recommended in diets as it is required for proper sexual organ functioning and enzyme activity. High levels of zinc are usually associated with high-protein food stuff, whereas low levels are obtained from food rich in carbohydrates" [17].

"Vitamins are organic compounds that are essential to an organism in small quantities for proper metabolic function. As depicted in this study, *O. gratissimum* contained considerable quantity of vitamins, especially vitamins A, C, and E. Vitamin A is a fat-soluble vitamin that is naturally present in many foods. It is important for normal vision, immune system, reproduction, growth, and development. It has also been reported to help in the proper functioning of the heart and lungs" [16]. The vitamin A content as reported in this study was consistent with the findings of Oladejo [14] and Enemor et al., [17]. In the same vein, the researched vegetable was found to be rich source of vitamin E (35.85 ± 1.56 µg/g). Vitamin E is a fat-soluble vitamin that act as antioxidant and consequently help to protect the cells from damages occasioned by free radicals. By implication, consumption of this vegetable could aid the body's antioxidant defense mechanism and consequently improve health.

4. CONCLUSION

This study indicated that the investigated vegetable was a rich source of carbohydrate, protein, fiber, and essential minerals, as well as vitamins A, C, and E. *O. gratissimum* presented a nutritive profile capable of promoting health and well-being. The anti-nutrients in the leafy vegetable were much lower than the safe limits and the reports for other plant sourced foods, including nuts, roots, beans and berries. A serving of 100 g per day of the investigated vegetables, complemented by other nutrient sources, will provide the RDA of the essential minerals investigated. While a serving of up to 2 kg per day of the vegetables does not contribute anti-nutrients higher than the safe limits.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models

(ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ajayi OB, Bamidele TJ, Malachi OI, Oladejo AA. Comparative Proximate, Minerals and Antinutrient Analysis of Selected Nigerian Leafy Vegetables. *Journal of Applied Life Sciences International*. 2018;16(1):1-8.
2. Okonwu K, Egerton NA, Ajayi AE, Agogbua JU. Nutritional Composition of Some of the Most Consumed Cereals and Legumes in Nigeria. *European Journal of Nutrition and Food Safety*. 2022;14(2):53-58.
3. Nakano MR, Umar BD, Nuhu AL, Adamu IT. Proximate and mineral composition of some wild leafy vegetables consumed in Katsina State Nigeria. *World Journal of Advanced Engineering Technology and Sciences*. 2023;09(01):084–088.
4. Baiyeri SO, Chimaluka CA, Okorie ON. Proximate, Mineral, Vitamin, And Anti-Nutrient Contents of the Leaves of *Senecio bialfrae*. *Journal of Agricultural Sciences*. 2023;68(1):67-79.
5. Ojmelukwe PC, Amaechi N. Composition of *Vernonia amygdalina* and its potential health benefits. *International Journal of Environment, Agriculture and Biotechnology*. 2019;4 (6):1836-1848.
6. Akara EU, Okezie E, Ude VC, Uche-Ikonke C, Eke G, Ugbogu AE. *Ocimum gratissimum* leaf extract ameliorates phenylhydrazine-induced anaemia and toxicity in Wistar rats. *Drug Metab. Pers. Ther.* 2021;15(1):20-35.
7. Irondi EA, Agboola SO, Oboh G, Boligon AA. Inhibitory effect of leaves extracts of *Ocimum basilicum* and *Ocimum gratissimum* on two key enzymes involved in obesity and hypertension *In vitro*. *J. Intercult. Ethnopharmacol.* 2016;5: 396–402.
8. Melo RS, Azevedo AMA, Pereira AMG, Rocha RR, Cavalcante RMB, Matos MNC, Lopes PHR, Gomes GA, Rodrigues THS, Dos Santos HS, Ponte IL, Costa RA, Brito GS, Júnior FEAC, Carneiro VA. Chemical

- composition and antimicrobial effectiveness of ocimum gratissimum L. Essential oil against multidrug-resistant isolates of Staphylococcus aureus and Escherichia coli. *Molecules*. 2019;3(8) 64:2–17.
9. AOAC. Official method of analysis. Association of Official Analytical Chemist. 1990;15:910-28.
 10. McDonald P, Edwards RA, Greenhalgh JFD. *Animal Nutrition*. Edinburgh: T and A Constable Ltd. 1973;2-5.
 11. Kirk RS, Sawyer. *Pearsons Food Composition and Analysis*. 4th Edition. Macmillan Publisher Company, United Kingdom; 1991.
 12. Rosenberg HR. *Chemistry and physiology of the vitamins*. Interscience Publisher, New York. 1992;452-3.
 13. Hussain J, Rehman NU, Khan AL, Hamayun M, Hussain SM, Shinwari ZK. Proximate and essential nutrients evaluation of selected vegetables species from Kohat region, Pakistan. *Pak. J. Bot.* 2010;42(4):2847-2855.
 14. Oladejo AA. Comparative Vitamin Analysis of Some Selected Nigerian Green Leafy Vegetables from Two Different Zones. *Acta Scientific Nutritional Health*. 2019;3(11):1-7.
 15. Pellett PL, Young VR. Nutritional evaluation of protein foods. *Food and Nutrition Bulletin*. 2018;4(1):1-7.
 16. Nnenne SK, Ubaoji KI, Ogbodo UC, Enemor VHA, Oladejo AA. Comparative Study on the Nutritional and Antioxidant Components of Fruit Parts of Citrullus lanatus. *European Journal of Nutrition and Food Safety*. 2020;12(11):39-51.
 17. Enemor VHA, Oguazu CE, Odiakosa AU, Okafor SC. Evaluation of the medicinal properties and possible nutrient composition of Citrullus lanatus (watermelon) seed. *Res J Med Plants*. 2019;13:129-135.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/126300>