



HAL
open science

Influence of process parameters on nutritional profile and physical stability of innovative infant follow-on formulas based on animal and plant oil bodies

Mathilde Cancalon, Claire Bourlieu-Lacanal, Gaëlle Tanguy, Éric Lacroux, Romain Valentin, Françoise Boissel, Olivia Ménard, Lucie Rault, Pascal Lecouillard, Yann Le Gouar, et al.

► To cite this version:

Mathilde Cancalon, Claire Bourlieu-Lacanal, Gaëlle Tanguy, Éric Lacroux, Romain Valentin, et al.. Influence of process parameters on nutritional profile and physical stability of innovative infant follow-on formulas based on animal and plant oil bodies. 6. Food Structure and Functionality Symposium: Meeting the sustainability challenge, Oct 2024, Bruges, Belgium. <hal-04777019>

HAL Id: hal-04777019

<https://hal.science/hal-04777019v1>

Submitted on 12 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Influence of process parameters on nutritional profile and physical stability of innovative infant follow-on formulas based on animal and plant oil bodies

Mathilde Cancalon¹, Claire Bourlieu-Lacanal⁴, Gaëlle Tanguy¹, Éric Lacroux², Romain Valentin², Françoise Boissel¹, Olivia Ménard¹, Lucie Rault¹, Pascal Lecouillard¹, Yann Le Gouar¹, Didier Dupont¹, Romain Jeantet¹, Jeehyun Lee^{1*} and Amélie Deglaire^{1*}

¹ STLO, INRAE, L'Institut Agro, Rennes, France ; ² Laboratoire de Chimie Agro-industrielle, LCA, Université de Toulouse, INRAE, Toulouse, France ; ³ UMR IATE, UM Montpellier, INRAE, Institut Agro, Montpellier, France

Introduction

Only 34% of infant worldwide are still breastfed at 6 months despite WHO recommendations

→ Majority of formula-fed infants WHO, 2022

Lipids mainly from plant oils derived from extraction processes employing organic solvents posing environmental issues and unstructured

(only 14% of IFF include dairy fat)

Cancalon et al., Recherche Data Gov, 2023
Cancalon et al., Data in brief, 2023

Interest of oil bodies (OB)

- Structured lipids (fat globules, oleosomes)
- From more sustainable extraction processes
- Profile of fatty acids of nutritional interest

AIM OF THE STUDY

Introduction of animal and plant OB obtained from more sustainable procedure into infant follow-on formulas (IFF) and assess their stability under different processing conditions

Materials and methods

Introduction of animal-derived OB (aOB) with dairy cream or plant-based OB from the grinding of sunflower, oleic Sunflower, and rapeseed seeds

Formulation of three IFF (40% dry matter*) differing in lipid origin

*To avoid evaporation before spray-drying



PO - based on plant oils



PO/aOB - based on plant oils and animal OB



pOB/aOB - based on animal and plant OB

Production under different operating conditions

High-pressure homogenization (HPH)

80/20 bars

OR

140/40 bars

Heat treatment (HT)

73°C, 1 min


OR

80°C, 1 min

What impact do process conditions have?

Results

Confocal microscopy




Replacement of phospholipid-stabilized interfaces by protein-stabilized interfaces after HPH
Aggregation for pOB/aOB

Fluorescent probes

- Lipids, Lipidtox®
- Amphiphiles, Rd-DOPE®
- Proteins, Fast green FCF

Physical stability




Phase separation for pOB/aOB

Increasing HT temperature limits phase separation

- 80/20 bars, no HT (A)
- 80/20 bars, 73°C/1 min (A1)
- 80/20 bars, 80°C/1 min (A2)
- 140/40 bars, no HT (B)
- 140/40 bars, 73°C/1 min (B1)
- 140/40 bars, 80°C/1 min (B2)

Droplet size distribution




Bimodal distribution with mode between 1.2 and 1.4 µm or 0.7 and 1.3 µm for HPH of 80/20 and 140/40 bars

No impact of HT on size distribution

Significant aggregation in pOB/aOB (larger size for measurements in water than in SDS)

Apparent viscosity



Conclusions

These results demonstrated the feasibility of homogenization and heat treatment on innovative IFF. The influence of the addition of minerals and of the spray-drying of these IFF will be evaluated, before investigating the nutrient bioaccessibility using *in vitro* digestion models.

The authors thank Carnot Qualiment and 3Bcar for funding the OBEINN (Oil Bodies for INNOvative food products: From plant seed processing to digestive fate) project.