



**HAL**  
open science

## Understanding ketone hydrogenation catalysis with anionic iridium(III) complexes: The crucial role of counterion and solvation

Paven Kisten, Sandrine Vincendeau, Eric Manoury, Jason Martin Lynam, John M Slattery, Simon B Duckett, Agusti Lledos, Rinaldo Poli

### ► To cite this version:

Paven Kisten, Sandrine Vincendeau, Eric Manoury, Jason Martin Lynam, John M Slattery, et al.. Understanding ketone hydrogenation catalysis with anionic iridium(III) complexes: The crucial role of counterion and solvation. *Chemical Science*, 2024, 10.1039/d4sc04629c . hal-04777200

HAL Id: hal-04777200

<https://hal.science/hal-04777200v1>

Submitted on 12 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Chemical Science

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: P. Kisten, S. Vincendeau, E. Manoury, J. M. Lynam, J. M. Slattery, S. B. Duckett, A. Lledos and R. Poli, *Chem. Sci.*, 2024, DOI: 10.1039/D4SC04629C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

# Understanding ketone hydrogenation catalysis with anionic iridium(III) complexes: The crucial role of counterion and solvation

View Article Online  
DOI: 10.1039/D4SC04629C

Paven Kisten,<sup>a,b</sup> Sandrine Vincendeau,<sup>a</sup> Eric Manoury,<sup>a</sup> Jason Lynam,<sup>b</sup> John Slattery,<sup>\*b</sup> Simon B. Duckett,<sup>b</sup> Agustí Lledós<sup>\*c</sup> and Rinaldo Poli<sup>\*a,d</sup>

<sup>a</sup>CNRS, LCC (Laboratoire de Chimie de Coordination), Université de Toulouse, UPS, INPT, 205 route de Narbonne, BP 44099, F-31077 Toulouse Cedex 4, France CNRS. Fax: +33-561553003; Tel: +33-561333174; E-mail: [rinaldo.poli@lcc-toulouse.fr](mailto:rinaldo.poli@lcc-toulouse.fr).

<sup>b</sup> Department of Chemistry, University of York, Heslington, York, YO10 5DD U.K. E-mail: [john.slattery@york.ac.uk](mailto:john.slattery@york.ac.uk).

<sup>c</sup> Departament de Química, Universitat Autònoma de Barcelona, 08193 Cerdanyola del Vallès, Catalonia, Spain. E-mail: [Agusti.Lledos@uab.cat](mailto:Agusti.Lledos@uab.cat).

<sup>d</sup> Institut Universitaire de France, 1, rue Descartes, 75231 Paris Cedex 05, France.

**Keywords:** Iridium, pre-catalyst activation, transfer hydrogenation, cyclooctadiene, phosphine ligands, hydride ligands, alkali-metal cation, solvent effects



## Abstract

View Article Online  
DOI: 10.1039/D4SC04629C

Catalytic asymmetric hydrogenation of ketones is an important approach to prepare valuable chiral alcohols. Understanding how transition metals promote these reactions is key to the rational design of more active, selective and sustainable catalysts. A highly unusual mechanism for asymmetric hydrogenation of benzophenone catalysed by an anionic Ir<sup>III</sup> hydride system, including a strong counterion dependence on catalyst activity, is explored and rationalised here. The active catalyst, generated *in situ* from [IrCl(COD)]<sub>2</sub> and a bidentate ligand (P,S<sup>R</sup>) under H<sub>2</sub> in the presence of a strong base (M<sup>+</sup>PrO<sup>-</sup> in isopropanol, M = Li, Na, K), is the solvated M<sup>+</sup>[Ir(H)<sub>4</sub>(P,S<sup>R</sup>)] salt (P,S<sup>R</sup> = CpFe[1,2-C<sub>5</sub>H<sub>3</sub>(PPh<sub>2</sub>)(CH<sub>2</sub>S<sup>R</sup>)], with R = <sup>i</sup>Pr, Ph, Bz and Cy). Catalyst activity increases, for all R derivatives, significantly as the counterion is varied in the order Li < Na < K. For the most active K system, the addition of 18-crown-6 drastically reduces the activity. While the cation strongly affects catalyst activity, it does not significantly affect enantioselectivity. DFT calculations explored these effects in detail and showed that the solvation model used in the calculations was critical. Only a hybrid implicit/explicit solvent model including sufficient explicit solvent molecules to properly describe the first solvation shell of the cation was able to reproduce the experimental observations. This model revealed the fundamental importance of the alkali-metal cation coordination sphere in understanding the counterion effects. The turnover-determining states in the catalytic cycle are those involved in outer-sphere hydride transfer to the substrate. This step leads to coordination of the alkoxide product to the alkali-metal cation, with a significant rearrangement of the coordination sphere of M, whereas there is little change in the geometrical parameters around Ir or the alkoxide. The DFT calculations also



pinpointed the major enantio-discriminating interactions and rationalised the insensitivity of the enantioselectivity on the alkali metal cation placement. View Article Online  
DOI: 10.1039/D4SC04629C


## Introduction

The asymmetric hydrogenation of polar prochiral substrates, particularly aryl ketones, is a highly important route to valuable chiral alcohols.<sup>1-6</sup> Transition metal catalysis can prove effective for these reactions, but catalyst activation, speciation in solution and catalytic reaction mechanisms are not always well understood. A detailed understanding of how metals can promote these transformations is important to allow the rational design of more active, selective and sustainable catalysts. In many reactions a pre-catalyst, for example  $[\text{IrCl}(\text{COD})]_2$ , is used to deliver the metal to the system. For most pre-catalysts, the presence of a strong base is required to afford high activities. The role of the base is usually attributed to the stabilisation of a more active neutral hydride species, derived from the pre-catalyst, relative to cationic species that may otherwise form in the protic reaction solvents (typically alcohols) that are used.<sup>7</sup> When a chloride-containing complex is used as a pre-catalyst, as is often the case, the base is also invoked in the activation step, where substitution of the chloride by an alkoxide produced by deprotonation of the alcohol solvent (e.g. isopropoxide when the reactions are conducted in isopropanol) is followed by  $\beta$ -H elimination.<sup>8</sup> However, in some of our laboratories we have demonstrated that these rationalisations are insufficient,<sup>9</sup> at least for ketone hydrogenations catalysed by  $[\text{IrCl}(\text{COD})]_2/\text{LL}'$  (COD = 1,5-cyclooctadiene), where LL' is a planar chiral 1,2-disubstituted ferrocene ligand containing diphenylphosphino and thioether donor groups,  $\text{CpFe}[1,2\text{-C}_5\text{H}_3(\text{PPh}_2)(\text{CH}_2\text{SR})]$  (R = Et, <sup>i</sup>Pr, <sup>t</sup>Bu, Bz, Ph, *etc.*),<sup>10</sup> henceforth abbreviated as (P,S<sup>R</sup>).



The stoichiometric reaction between  $[\text{IrCl}(\text{COD})]_2$  and  $((S)\text{-P}, S^R)$  was shown to yield well-defined  $[\text{IrCl}(\text{COD})((S)\text{-P}, S^R)]$  complexes with either a 5-coordinate square pyramidal<sup>11</sup> or a 4-coordinate square planar<sup>12</sup> (for  $R = t\text{Bu}$ ) geometry. These complexes (either isolated or made *in situ*) proved highly active catalysts and led to excellent enantioselectivity in aromatic ketone hydrogenation, though only in the presence of an excess of a strong base (e.g. 5 equivalents of NaOMe per Ir).<sup>13</sup> However, the use of an alternative precursor,  $[\text{Ir}(\text{OMe})(\text{COD})]_2$ , which contains an internal methoxide base, alongside the  $(P, S^R)$  ligand still required the presence of additional strong base to yield an active catalyst, suggesting an anionic active catalyst is formed under the reaction conditions.<sup>9</sup> A parallel DFT investigation revealed that the most stable species generated by a sequence of COD hydrogenation and  $\text{H}_2$  oxidative addition steps is the anionic tetrahydrido complex  $[\text{Ir}(\text{H})_4((S)\text{-P}, S^R)]^-$ . The latter is therefore a strong candidate for the catalyst resting state. This complex does not contain mobile protons for a classical outer-sphere bifunctional (Noyori-type)<sup>7</sup> mechanism, nor a vacant coordination site for an inner-sphere mechanism. Instead, the DFT calculations rationalised the catalytic action through a novel mechanistic variant of the outer-sphere bifunctional mechanism, which is summarised in Scheme 1. This involves a cooperative action between the alkali-metal cation for ketone activation and the iridium centre for hydride transfer to the carbonyl carbon atom, thereby generating an alkoxide intermediate. This step is followed by transfer of a proton originating from heterolytic cleavage of  $\text{H}_2$  by the alkoxide and the Ir centre. The calculated Gibbs energy span of the resulting catalytic cycle,  $18.2 \text{ kcal mol}^{-1}$ , was consistent with the experimental activity.<sup>9</sup> However, the effect of the  $[\text{Ir}(\text{H})_4((S)\text{-P}, S^R)]^-$  counterion was not explored and yet has proved to be a highly important feature of this system, significantly affecting its activity, as described below.





Scheme 1. Catalytic cycle for ketone hydrogenation catalysed by  $[\text{IrCl}(\text{COD})((\text{S})\text{-P,S}^{\text{Ph}})]/\text{NaOMe}$ , according to a DFT investigation.<sup>9</sup>

The extreme lability of the  $[\text{Ir}(\text{H})_4((\text{S})\text{-P,S}^{\text{R}})]^-$  complex has so far not allowed us to isolate or indeed spectroscopically identify it. However, we have obtained a related complex with the chelating diphenylphosphinoethane (dppe) ligand,  $[\text{Ir}(\text{H})_4(\text{dppe})]^-$ , under identical conditions to those used for hydrogenation catalysis (from  $[\text{IrCl}(\text{COD})]_2/\text{dppe}$  in basic isopropanol under  $\text{H}_2$  at room temperature) and also under transfer hydrogenation conditions (no  $\text{H}_2$  gas added), even though the latter required warming to  $80^\circ\text{C}$ .<sup>14</sup> Analogous anionic tetrahydrido complexes of  $\text{Ir}^{\text{III}}$  were previously reported in the chemistry of monodentate phosphine ligands,  $[\text{Ir}(\text{H})_4\text{L}_2]^-$  ( $\text{L} = \text{P}^i\text{Pr}_3$ ,  $\text{PPh}_3$ ), obtained by reaction of  $[\text{Ir}(\text{H})_5\text{L}_2]$  with  $\text{KH}$ , but not under catalytic conditions.<sup>15</sup> <sup>16</sup> This gives further credence to the implication of anionic  $[\text{Ir}(\text{H})_4((\text{S})\text{-P,S}^{\text{R}})]^-$  as the active species in ketone hydrogenation with  $[\text{IrCl}(\text{COD})]_2/((\text{S})\text{-P,S}^{\text{R}})$ .



The action of anionic hydride complexes as (transfer) hydrogenation catalysts and their generation by strong bases, has not been widely acknowledged or studied. In early work, G. P. Pez *et al.* suggested the involvement of anionic Ru hydride complexes, e.g.  $[K(L_n)][RuH_2(\kappa^2\text{-}C,P\text{-}o\text{-}C_6H_4PPh_2)(PPh_3)_2]$  and  $[K(L_n)][(PPh_3)_3(PPh_2)RuH_2]$  ( $L_n$  = solvent or crown ether), as active species for the catalytic reduction of aromatic compounds as well as aldehydes, ketones, esters and nitriles.<sup>17-21</sup> Of special note is the fact that the addition of either a crown ether or cryptand was shown to dramatically reduce the catalytic activity.<sup>19</sup> However, subsequent work by J. Halpern *et al.* demonstrated that the anionic  $[Ru(H)_3(PPh_3)_3]^-$  complex, which is generated from the Pez systems under  $H_2$ ,<sup>22, 23</sup> is reversibly protonated in alcohol solvents to form  $[Ru(H)_2(H_2)(PPh_3)_3]$  and that the latter “tetrahydride” complex is the real catalytically active species.<sup>24</sup> After these debated contributions, the role of anionic hydride species in transition metal catalysed hydrogenation was not considered again, until recently.

Just prior to our above-mentioned contribution, Dub *et al.* suggested that the active species of Noyori's  $[Ru(H)_2(\text{diphosphine})(NH_2CHPhCHPhNH_2)]$  catalyst is, in fact, the deprotonated amine-amido complex  $[K(L_n)[Ru(H)_2(\text{diphosphine})(NHCHPhCHPhNH_2)]$ .<sup>25</sup> DFT calculations carried out for that system underlined the activating action of the  $N-K^+$  function in the rate-determining hydride transfer step to the outer-sphere substrate, though the potassium ion was introduced in the calculation as a naked ion, which does not reflect the solvation of this ion under catalytic conditions. Beller's Ru PNP pincer complex  $[RuH(CO)Cl\{HN(C_2H_4P^iPr_2)_2\}]$ , which catalyses the aqueous dehydrogenation of methanol under highly basic conditions,<sup>26</sup> was suggested to operate *via* an anionic deprotonated pincer hydride resting state on the basis of DFT calculations. However,





these were carried out on the free anion, without consideration of a possible cation effect.<sup>27</sup> The ester hydrogenation catalysed by  $[\text{MnBr}(\text{CO})_3(\text{Ph}_2\text{CH}_2\text{CH}_2\text{NH}_2)]$  under strongly basic conditions<sup>28</sup> was also proposed by Pidko *et al.*, on the basis of a DFT investigation,<sup>29</sup> to proceed *via* an anionic species with a deprotonated ligand,  $[\text{MnH}(\text{CO})_3(\text{Ph}_2\text{CH}_2\text{CH}_2\text{NH})]^-$ . These calculations, like the above-mentioned ones by Dub *et al.*, included only the naked cation and a comparison of the NH and N-M<sup>+</sup> (M = Li, Na, K) systems suggested that, while the barrier of the rate-determining hydride transfer step is reduced on going from the neutral to the anionic system, the cation effect is only minor. A cation effect, however, was not experimentally probed for this manganese catalyst. The  $[\text{IrCl}(\text{COD})]_2/\text{HL}$  system with HL = tridentate ferrocene-based amino-phosphine-sulphonamide (f-Amphamide), used as an asymmetric catalyst for ketone transfer hydrogenation, was also shown by Dang, Zhang *et al.* to operate *via* an anionic hydride complex as a resting state, *i.e.*  $[\text{Ir}(\text{H})_3\text{L}]^-$ .<sup>30</sup> For the latter system, DFT calculations suggested an activating role of Li<sup>+</sup>, which was again introduced in the calculations as a naked ion. Finally, a most recent contribution by Bai, Lang, Zhang *et al.* has shown an extremely active and enantioselective anionic Ir<sup>III</sup> hydride complex with a tetradentate PNNO ligand (f-phamidol), which hydrogenates acetophenone with a TOF of 224 s<sup>-1</sup> up to a TON of over 13 million and up to 99% ee.<sup>31</sup> In this contribution, the cation (Na<sup>+</sup>) was explicitly introduced in the calculations and the effect of the cation solvation (up to two isopropanol molecules per Na<sup>+</sup>) was considered, but a possible effect of the cation nature was not explored, neither experimental nor computationally.

For an anionic active catalyst, the activity may somewhat be affected by the nature of the counter-cation. In addition to the above-mentioned investigation by Pez *et al.*,<sup>19</sup> the effect of an alkali-metal counterion on the activity (and/or selectivity) of a



(transfer) hydrogenation catalyst has been pointed out in a few cases but has generally be attributed to cooperation with a neutral hydride system. A notable example is the  $[[\text{RuCl}_2(\textit{p}\text{-cymene})]_2]$ -pseudodipeptide-catalysed enantioselective transfer hydrogenation of ketones, where the alkali-metal cation was suggested to act in a Meerwein–Ponndorf–Verley fashion in combination with a neutral  $\text{Ru}^{\text{II}}$  hydride in the rate-limiting hydride transfer step.<sup>32, 33</sup> In a series of papers by Hazari *et al.*,<sup>34–36</sup> alkali-metal cation salts were shown to affect the activity of Fe pincer hydride complexes in methanol dehydrogenation. In this case, the DFT studies suggested that the cation accelerates the formate product extrusion, but again in cooperation with a neutral active species and a similar conclusion was reached for the related Ru catalyst.<sup>37</sup> Very recently, it was found that counteranions can exert a remarkable influence on the ability of anionic cobaltate salts to catalyze alkene hydrogenations.<sup>38</sup> Experimental and computational studies suggested an active co-catalytic role of the counterion ( $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Li}^+$ ,  $\text{Mg}^{2+}$ ) in the hydrogenation reaction through coordination to cobalt hydride intermediates.<sup>38</sup> However, despite the importance of these effects, detailed studies and understanding of counteranion effects in catalysis are still very rare. In contrast, anion effects in homogeneous catalysis using cationic metal complexes, e.g.  $\text{Au}^+$  systems,<sup>39, 40</sup> olefin polymerisation catalysis,<sup>41</sup> or in asymmetric counteranion-directed catalysis,<sup>42, 43</sup> have been more widely studied.

In the present contribution, we show the results of catalytic studies for the acetophenone hydrogenation carried out in the presence of strong bases with different alkali metal cations, where pronounced alkali-metal ion effects are apparent. Extensive DFT investigations pinpoint the role of the alkali-metal cation in ketone activation and its assistance in the rate- and enantio-determining hydride-transfer step. A key aspect that emerged during the DFT study was the importance of the solvent model used for



the correct modelling of the catalytic cycle. Quite surprisingly, a major effect was revealed for the rearrangement of the alkali-metal coordination sphere, which must be described by an appropriate number of explicit solvent molecules. These observations are important not only for the present work, but to inform the development of computational methodologies for studying many homogeneous catalytic reactions where solvent plays an underappreciated role.

View Article Online  
DOI: 10.1039/D4SC04629C

## Results and discussion

### (a) Catalytic investigations

The effect of the alkali-metal cation on the catalytic activity of the system generated from  $[\text{IrCl}(\text{COD})((\text{S})\text{-P,S}^{\text{R}})]/\text{MO}^t\text{Pr}/\text{H}_2$  ( $\text{M} = \text{Li}, \text{Na}, \text{K}$ ) was probed for four different  $((\text{S})\text{-P,S}^{\text{R}})$  ligands ( $\text{R} = ^i\text{Pr}, \text{Ph}, \text{Bz}, \text{Cy}$ ), under similar conditions to those of our initial investigation, namely hydrogenation under 30 bar of  $\text{H}_2$  pressure at 3 °C with a 0.2% catalyst loading and with activation by 1% strong base (5 equiv. relative to catalyst) in isopropanol.<sup>13</sup> It is useful to recall here that no activity is obtained unless the neutral  $\text{Ir}^{\text{I}}$  precatalyst is activated by a strong alkali metal base, which is rationalised by the conversion to the active anionic  $[\text{IrH}_4((\text{S})\text{-P,S}^{\text{R}})]^-$  complex.<sup>9</sup> Alkoxide bases ( $\text{NaOMe}$ ,  $\text{KO}^t\text{Bu}$ ) and even  $\text{KOH}$  were found able to activate the precatalyst, whereas no activity resulted in the presence of  $\text{NEt}_3$ .<sup>13</sup> The Ir complexes with Ph- and Bz-substituted ligands were previously reported,<sup>12, 13</sup> whereas those with the  $^i\text{Pr}$ - and Cy-substituted ligands have been used here as hydrogenation catalysts for the first time. The operating base in entries 1-5 (Table 1) is isopropoxide, because the employed  $\text{KO}^t\text{Bu}$  in the experiments of entries 3 and 4 is a stronger base and thus the equilibrium position in  $\text{HO}^i\text{Pr}$  strongly favours the formation of isopropoxide. The results of entries



1-3 indicate a significant effect of the alkali-metal counterion associated with the base on the catalytic activity, confirming the proposition that there is an explicit role for the alkali metal cation in the catalytic cycle. Notably, all four ((S)-P,S<sup>R</sup>) systems show the same trend of activity ( $\text{Li}^+ < \text{Na}^+ < \text{K}^+$ ). When using the same cation, the isopropyl system consistently gives the highest catalytic activity. The addition of 18-crown-6 to the potassium system (entry 4) results in a dramatic quenching of the catalytic activity. Once again, a comparable effect is shown by all four ((S)-P,S<sup>R</sup>) systems. This result is also in perfect agreement with the proposed catalytic cycle (Scheme 1), because the high affinity of this particular crown ether for  $\text{K}^+$  is expected to weaken the ability of the cation to ion pair with the anionic catalyst. In this respect, it is interesting to note that the previously reported  $[\text{Ir}(\text{H})_4\text{L}_2]^-$  ( $\text{L} = \text{P}^i\text{Pr}_3, \text{PPh}_3$ ) complexes were crystallised and structurally investigated as crown-ether adducts of the  $\text{K}^+$  salts.<sup>15, 16</sup> Furthermore, all structures that feature *cis* L ligands also reveal  $\text{K} \cdots \text{H}$  interactions with three *mer* hydride ligands, as found by the DFT calculations for the  $[\text{Na}(\text{MeOH})_3][\text{Ir}(\text{H})_4((\text{S})\text{-P,S}^{\text{Ph}})]$  model (**1** in Scheme 1). The deliberate addition of degassed water to the reaction mixtures has little or no effect on the catalytic activities (*cf.* entries 3 and 5). Finally, using the tetramethylammonium cation (entry 6) also resulted in reduced activity. This result is interesting when compared with the identical activities reported (for the Et-substituted ligand system) in the presence of  $\text{KO}^t\text{Bu}$  and  $\text{KOH}$ .<sup>13</sup> Thus, the activity drop for the Cy-substituted ligand system from entry 3 to entry 6 cannot be attributed to the weaker basicity of  $\text{OH}^-$  in isopropanol, but rather to the replacement of  $\text{K}^+$  with  $\text{Me}_4\text{N}^+$ .

The observed enantioselectivity trends are also interesting. The highest *e.e.* values were obtained for the Bz system, in agreement with the previous report.<sup>13</sup> For this system, as well as for the Ph and Cy systems, all three alkali-metal cations yield



essentially the same e.e., except for a significant drop for the Cy system when using  $\text{Me}_4\text{N}^+$  instead of an alkali metal cation. For the  $^i\text{Pr}$  system, however, the nature of the cation significantly affects the e.e., which decreases in the order  $\text{K}^+ > \text{Na}^+ > \text{Li}^+$ . It is particularly interesting to note that, for all ((S)-P, $S^R$ ) systems, the presence of the crown ether, although strongly quenching the activity, does not significantly affect the e.e. Conversely, the presence of water, although not significantly changing the activity, results in an e.e. decrease of approximately a factor of 2 for the  $^i\text{Pr}$  and Bz systems. The Ph and Cy systems, on the other hand, yield e.e. similar to those of entries 1-3.

Table 1. Results of the acetophenone hydrogenation using different (S-P, $S^R$ ) ligands and bases.<sup>a</sup>

| Entry | Base (+ additive)                                | Conversion/% <sup>b</sup> (e.e. <sup>c</sup> /%) |                     |                     |                     |
|-------|--|--|---------------------|---------------------|---------------------|
|       |  | R = $^i\text{Pr}^d$                              | R = Ph <sup>e</sup> | R = Bz <sup>d</sup> | R = Cy <sup>e</sup> |
| 1     | LiO $^i\text{Pr}$                                | 20 (34)  | 12 (65)             | 8 (78)              | 2 (69)              |
| 2     | NaO $^i\text{Pr}$                                | 61 (51)  | 26 (60)             | 19 (80)             | 60 (72)             |
| 3     | KO $^t\text{Bu}$                                 | 90 (58)  | 34 (66)             | 21 (78)             | 70 (75)             |
| 4     | KO $^t\text{Bu}$ + 18-crown-6 <sup>f</sup>       | 14 (60)  | 12 (60)             | 4 (81)              | 14 (67)             |
| 5     | KO $^t\text{Bu}$ + H <sub>2</sub> O <sup>g</sup> | 93 (29)  | 30 (67)             | 23 (50)             | 84 (75)             |
| 6     | Me <sub>4</sub> NOH                              | -  | -                   | -                   | 10 (23)             |

<sup>a</sup> Conditions:  $p(\text{H}_2) = 30$  bar;  $t = 4$  h;  $T = 3^\circ\text{C}$ ;  $[\text{PhCOMe}] = 1.6$  M in isopropanol;  $[\text{Ir}] = 3.2 \cdot 10^{-3}$  M;  $[\text{Base}] = 1.6 \cdot 10^{-2}$  M. <sup>b</sup> Determined from the area of the residual acetophenone in the gas-chromatogram in comparison with calibration curves (estimated error =  $\pm 1.7$  %). <sup>c</sup> Determined from the relative areas of the two phenylethanol enantiomers separated on a chiral column (estimated error =  $\pm 1.5$  %); the S enantiomer is the major one. <sup>d</sup> Pre-catalyst prepared *in situ* from  $[\text{IrCl}(\text{COD})]_2$  and ((S)-P, $S^R$ ). <sup>e</sup> With isolated  $[\text{IrCl}(\text{COD})(\text{PS}^R)]$ . <sup>f</sup> 18-crown-6/K = 1.5. <sup>g</sup> H<sub>2</sub>O/K = 1.5.

### (b) DFT investigations of the cation effect

We have performed DFT calculations to disclose the influence of the cation on the catalyst activity, using a hybrid implicit-explicit (cluster-continuum) solvent model in which, in addition to the continuum model, five explicit solvent molecules have been introduced. The goal of the calculations is, after finding a model that reproduces the experimental trend, to identify its origin. We are aware that to obtain more quantitative




agreements would require the use of very expensive time-consuming DFT-based Molecular Dynamics (DFT-MD) simulations, placing the reactive system inside a box with a huge number of solvent molecules. This is out of our scope, and we will show that, with a careful description of the solvent, static DFT calculations are able to capture the main features without performing these highly demanding DFT-MD calculations. As we will show in what follows the main point is that a complete description of the first solvation sphere of the cation is mandatory to reproduce the experimental trends.

Using the cluster-continuum solvent model with five solvent molecules, the (S)-P,S<sup>Ph</sup> ligand without any truncation, and acetone as the model substrate molecule, we have computed all the steps of the catalytic cycle of ketone hydrogenation depicted in

Scheme 2 for the three alkali cations, initially modelling the isopropanol solvent used experimentally as methanol. The computed Gibbs energy profiles are shown in Figure 1. However, once the rate determining step (hydride transfer) was identified, we reoptimised all the structures along this step using isopropanol in the hybrid continuum explicit (five solvent molecules) solvent model, in order to provide a solvent description more akin to the experimental system. This approach has allowed the reproduction of the experimental activity trend:  $K^+ > Na^+ > Li^+$ . As we will discuss later, an accurate description of the first solvation sphere of the cation is crucial if DFT is to match the experimental trend. This was already achieved using  $[M(\text{MeOH})_5]^+$ . Contrarily, using the previously reported model<sup>9</sup> with only three methanol molecules ( $[M(\text{MeOH})_3]^+$ ), we found that the experimental trend was not mirrored in the computational study, *i.e.*  $Li^+$  would be expected to show the highest activity. We will discuss this issue later on.





Scheme 2 Catalytic cycle for the hydrogenation of acetone with  $[M(\text{MeOH})_5][\text{Ir}(\text{H})_4((\text{S})\text{-P}, \text{S}^{\text{Ph}})]$  ( $M = \text{Li}^+, \text{Na}^+, \text{K}^+$ ).






Figure 1. Computed Gibbs energy profiles for ketone hydrogenation catalysed by  $[M(\text{MeOH})_5]^+[\text{Ir}(\text{H})_4((\text{S})\text{-P}, \text{S}^{\text{Ph}})]^-$  ( $M = \text{Li}, \text{Na}, \text{K}$ ). The profile follows the catalytic cycle of

Scheme 2, with 5 MeOH molecules explicitly included in the calculations.

As can be seen from Figure 1, the cation has an effect on the energy span of the catalytic cycle. This energy span corresponds to the difference between the transition state of the outer-sphere hydride transfer ( $\text{TS}_{2-3}$ ) and the initial state, **1**, which is the  $\{[\text{Ir}(\text{H})_4((\text{S})\text{-P}, \text{S}^{\text{R}})]^-; [M(\text{MeOH})_5]^+\}$  ion pair plus an acetone molecule infinitely apart. The barrier of the hydride transfer step changes appreciably with the cation, following the experimental trend:  $\text{K}$  ( $9.8 \text{ kcal mol}^{-1}$ ) <  $\text{Na}$  ( $14.1 \text{ kcal mol}^{-1}$ ) <  $\text{Li}$  ( $16.6 \text{ kcal mol}^{-1}$ ). After this step, which generates an O-coordinated alkoxide **4** via a less stable isomeric  $\sigma$ -complex **3**, the cation influence is much smaller. From **4**,  $\text{H}_2$  enters into play, first displacing the alkoxide ( $\text{TS}_{5-6}$ ) to form a  $\eta^2\text{-H}_2$  complex (**6**). The latter is able to protonate the displaced alkoxide *via*  $\text{TS}_{6-7}$ , yielding the alcohol (**7**) and regenerating complex **1**. The proton transfer TS ( $\text{TS}_{6-7}$ ) is the highest point in this pathway, and it is





found 7.9, 10.9 and 9.9 kcal mol<sup>-1</sup> above **4**, for Li, Na and K cations, respectively, thus the barrier to proton transfer from **4** is lower than the hydride transfer barrier for Li<sup>+</sup> and Na<sup>+</sup>, whereas the two barriers are effectively the same for K<sup>+</sup>.

We will present selected structures to discuss the changes in the cation solvation sphere along the catalytic pathway for the three ions. 3D plots of all the optimised structures are collected in Figures S6-S9 in the SI. Figure 2 displays the optimised structures of the iridium tetrahydride complex/cation ion pair before the arrival of the ketone molecule.




Figure 2. Optimised structures of the iridium complex/cation ion pairs (**1**) with the (MeOH)<sub>5</sub> solvent model. Distances in Å. C-H hydrogen atoms have been omitted for clarity.

The cations are placed in less crowded space opposite to the ((S)-P,S<sup>Ph</sup>) ligand and have a well-organised solvation sphere, involving the five MeOH molecules plus interactions with the hydride ligands of the Ir<sup>III</sup> anion. The M-O distances (M = Li, Na, K) involving the five solvent molecules are 2.03-2.16 Å for Li<sup>+</sup>, 2.32-2.40 Å for Na<sup>+</sup> and 2.70-2.82 Å for K<sup>+</sup>. These distances are close to the sum of the covalent radii of the metals and oxygen, particularly for Na<sup>+</sup> and K<sup>+</sup> (1.94, 2.32 and 2.69 Å, for Li, Na and K, respectively),<sup>44</sup>. The additional alkali metal-hydride interactions show a different behaviour for Li<sup>+</sup> vs. the other two cations. For Na<sup>+</sup> and K<sup>+</sup>, one hydride ligand



completes a distorted octahedral solvation sphere, thus bridging the alkali-metal cation close to the Ir<sup>III</sup> centre (Na-H: 2.40 Å; K-H: 2.79 Å), while the hydride ligand is out of the first solvation sphere of Li<sup>+</sup> (Li-H: 2.59 Å), presumably due to its smaller ionic radius and preference for lower coordination numbers. Usually, the promoting effect of alkali cations in hydrogenation reactions involving carbonyl groups have been discussed from DFT calculations in terms of activation of the carbonyl moiety by a direct M<sup>+</sup>··O=C interaction.<sup>29, 30</sup> However, those calculations did not include the solvation sphere of the cations, which leads to non-physical models. Our results highlight that, in order for a direct M<sup>+</sup>··O=C interaction to result in an alcohol solvent, at least one of the solvent molecules needs to be displaced from the cation solvation sphere.

Figure 3 depicts the optimised structures along the hydride transfer step for the three cations: the initial intermediate in which the acetone has been incorporated (**2**), the transition state for the outer-sphere hydride transfer (**TS<sub>2-3</sub>**) and the generated  $\sigma$ -complex where the alkoxide that is produced by hydride transfer interacts with the Ir<sup>III</sup> centre through the formed C-H bond (**3**). In all of the structures, two solvent molecules are H-bonded to the acetone oxygen atom but, notably, the acetone molecule is itself not incorporated in the coordination sphere of the alkali metal cations. These two H-bonds strengthen as the negative charge on the oxygen atom increases. For instance, in the K<sup>+</sup> system the O-H distances are 2.01/1.89 Å in **2**, 1.80/1.90 Å in **TS<sub>2-3</sub>** and 1.61/1.63 Å in **3**. Similar changes were found for the Li<sup>+</sup> and Na<sup>+</sup> systems.






Figure 3. Optimised structures along the hydride transfer step for the three cations in the (MeOH)<sub>5</sub> solvent model. Distances in Å. C-H hydrogen atoms have been omitted for clarity.

Values of the main distances involved in **TS**<sub>2,3</sub> (Ir-H, H-C and C-O), collected in Figure 3, show almost no difference between the three systems. Indeed, they are also very similar to those obtained without cation, just keeping the two solvating methanol molecules, for which the computed hydride transfer barrier is 14.6 kcal mol<sup>-1</sup> (see Figure S4). The same situation is evident with atomic charges, as computed with the



CM5 model (Table 2).<sup>45</sup> Regarding the acetone substrate polarisation in **TS<sub>2,3</sub>**, the charge transferred to the acetone oxygen atom is about the same in the Li<sup>+</sup> and cation-free cases, whereas it is slightly greater for the Na<sup>+</sup> and K<sup>+</sup> systems (Table 2).

The three **TS<sub>2,3</sub>** transition states have markedly different barriers (Figure 1), even though they exhibit very similar geometric and charge parameters for iridium coordination of the acetone substrate. However, the cation solvation sphere changes significantly along the hydride transfer pathway, suggesting that solvation rearrangement is the main cause of the difference in barrier. Alkoxide coordination to the alkali cation stabilises the negative charge that develops during the hydride transfer step, this disrupts the solvation sphere of the cation, leading to different barriers depending on the cation present. Once the alkoxide has formed (**3**), it coordinates to the alkali-metal cation.

Table 2. Main distances (Å) and CM5 charges in the transition states of the hydride transfer (**TS<sub>2,3</sub>**). Values in parentheses have been obtained using isopropanol in the hybrid continuum-explicit (five solvent molecules) solvent model

|                         | Li <sup>+</sup> | Na <sup>+</sup> | K <sup>+</sup> | No cation     |
|-------------------------|-----------------|-----------------|----------------|---------------|
| <b>distances</b>        |                 |                 |                |               |
| <b>Ir-H</b>             | 1.80 (1.76)     | 1.80 (1.78)     | 1.79 (1.78)    | 1.78 (1.78)   |
| <b>H-C</b>              | 1.66 (1.80)     | 1.69 (1.73)     | 1.69 (1.72)    | 1.68 (1.68)   |
| <b>C-O</b>              | 1.27 (1.27)     | 1.27 (1.27)     | 1.27 (1.26)    | 1.27 (1.26)   |
| <b>charges</b>          |                 |                 |                |               |
| <b>O</b>                | -0.38 (-0.39)   | -0.45 (-0.41)   | -0.43 (-0.43)  | -0.39 (-0.37) |
| <b>C</b>                | 0.15 (0.18)     | 0.15 (0.17)     | 0.15 (0.16)    | 0.15 (0.15)   |
| <b>H</b>                | -0.07 (-0.08)   | -0.07 (-0.07)   | -0.07 (-0.07)  | -0.08 (-0.07) |
| <b>Ir</b>               | -0.12 (-0.14)   | -0.12 (-0.13)   | -0.12 (-0.12)  | -0.20 (-0.18) |
| <b>CH<sub>3</sub>CO</b> | -0.19 (-0.10)   | -0.24 (-0.16)   | -0.20 (-0.29)  | -0.19 (-0.19) |
| <b>M</b>                | 0.47 (0.46)     | 0.63 (0.63)     | 0.77 (0.77)    | /             |

In the Li<sup>+</sup> system, two methanol molecules are displaced to produce the tetrahedral solvation sphere of this cation in **3-Li**, formed by the alkoxide and three solvent molecules. The lithium cation does not interact with any hydride ligand throughout this



process. In contrast, for the Na<sup>+</sup> and K<sup>+</sup> systems, one methanol molecule is displaced by alkoxide, thereby maintaining a distorted octahedral solvation sphere for the alkali-metal cation in **3-Na** and **3-K**, which is made up by the alkoxide, four solvent molecules and a bridging hydride ligand. This change in solvation is more favourable for K<sup>+</sup>, the cation with the largest ionic radius and greater effective charge (Table 2). Overall, the calculations using the (MeOH)<sub>5</sub> solvent model reproduce correctly the experimentally observed cation effect on the hydrogenation rate and allow its rationalisation, stressing the importance of the changes in the solvation sphere of the alkali cation along the reaction. However, to assess the validity of this model and to get more reliable results, we have reoptimized all the structures involved in the hydride transfer step (**1-M**, **2-M**, **TS<sub>23</sub>-M**, **3-M** and **4-M**) in the experimentally employed isopropanol solvent, described with the cluster-continuum solvent model.

Table 3. Comparison of relative Gibbs energies (kcal mol<sup>-1</sup>) of the species involved in the hydride transfer step with methanol and isopropanol solvents. Cluster-continuum solvent model with (ROH)<sub>5</sub> explicit solvent molecules + SMD continuum.

|                                      | K <sup>+</sup> |            | Na <sup>+</sup> |             | Li <sup>+</sup> |             | No cation   |             | TMA <sup>+</sup> |
|--------------------------------------|----------------|------------|-----------------|-------------|-----------------|-------------|-------------|-------------|------------------|
|                                      | Me(OH)         | iPr(OH)    | Me(OH)          | iPr(OH)     | Me(OH)          | iPr(OH)     | Me(OH)      | iPr(OH)     | iPr(OH)          |
| <b>1-M + ketone</b>                  | 0.0            | 0.0        | 0.0             | 0.0         | 0.0             | 0.0         | 0.0         | 0.0         | 0.0              |
| <b>2-M</b>                           | 0.4            | 0.9        | 4.5             | 4.7         | 0.6             | 6.6         | 8.1         | 6.7         | 8.0              |
| <b>TS<sub>23</sub>-M<sup>a</sup></b> | <b>9.8</b>     | <b>9.9</b> | <b>14.1</b>     | <b>13.2</b> | <b>16.6</b>     | <b>14.9</b> | <b>14.6</b> | <b>16.9</b> | <b>17.2</b>      |
| <b>3-M</b>                           | 2.2            | 2.3        | 7.9             | 4.5         | 2.1             | 1.3         | 6.3         | 9.0         | 6.2              |
| <b>4-M</b>                           | -4.7           | -9.3       | -0.3            | -5.5        | 0.0             | -1.4        | 0.1         | 1.5         | 3.8              |

<sup>a</sup> These values give the energy span of the catalytic cycle.



The obtained relative Gibbs energies are compared to those in methanol in Table 3. The main geometrical parameters and charges of the hydride-transfer transition states are compared in Table 2. 3D-structures of all the species can be found in the supporting Information (Figures S10-S14). The main conclusion from these calculations using the solvent used experimentally is that they give essentially very similar results to those in methanol. The trend in the barriers for the hydride transfer step ( $K < Na < Li$ ) is preserved, with very close barriers. We have also computed the barriers for the hydride-transfer step to acetophenone leading to the lowest S enantiomer with the three cations and (iPrOH)<sub>5</sub>. The obtained values ( $K^+$ : 9.9 kcal mol<sup>-1</sup>;  $Na^+$ : 12.5 kcal mol<sup>-1</sup>;  $Li^+$ : 14.4 kcal mol<sup>-1</sup>) are very similar to those obtained with acetone (Table 3). However, while with acetone there is no stabilization of the initial intermediates **2-M** and the barriers are computed with respect to the separated species (**1-M** + ketone), with acetophenone the initial intermediates are below the separated species and the energy span is computed as a difference between **2-M** and **TS<sub>2-3-M</sub>**. The absolute barriers are not consistent with the activities seen experimentally (non-quantitative yields in 4 hours). There are also concentration effects at play when considering rates, that would be difficult to address in the calculated barriers.

The most noticeable change when moving from the methanol to the isopropanol solvent model is a higher stabilisation of the alkoxide intermediates **4-M** in isopropanol solvent. Isopropanol has a significant effect on the stability of the alkoxide intermediate **4**, and this may then have an impact on the barriers to the subsequent steps. To address this issue, we have first computed the barrier for the **3** to **4** rearrangement for the three cations and the (iPrOH)<sub>5</sub> model, to discard it as rate-determining step, and



then we have recomputed the full energy profile with the potassium cation and the isopropanol model. View Article Online  
DOI: 10.1039/D4SC04629C

The **3** to **4** step involves a change in the coordination mode of the alkoxide formed in the previous step, from a weak C-H...Ir interaction to a stronger Ir-O. In this rearrangement solvation stabilizes the anionic alkoxide when it has been released from the iridium coordination sphere and the barriers are low, well below those of the hydride transfer for the three cations. However, they display a cation dependence (K: 1.3, Na: 3.0, Li: 6.5 kcal/mol) that follows that of the stability of the O-bound alkoxide (structures **4-M**; K: -9.3, Na: -5.5, Li: -1.4 kcal/mol). See **TS<sub>3,4-M-iso</sub>** in Figures S10-S12.

Regarding the complete profile with potassium cation in the isopropanol solvent model, comparing with the same profile with methanol, a strong stabilization of all the structures involving an alkoxide is observed (from **4-K**). Isopropanol solvent stabilises the alkoxide better than methanol. However, the barriers remain unchanged: 9.9 kcal mol<sup>-1</sup> for the hydride transfer step and 9.9 kcal mol<sup>-1</sup> for the second part of the reaction (Figure 4, optimized structures in Figure S12). Overall, these results substantiate the choice of acetophenone as a model to validate the catalytic mechanism.






Figure 4. Comparison of the computed Gibbs energy profiles for ketone hydrogenation catalysed by  $[\text{K}(\text{solvent})_5]^+[\text{Ir}(\text{H})_4((\text{S})\text{-P}, \text{S}^{\text{Ph}})]^-$  with 5 MeOH or 5 iPrOH molecules explicitly included in the calculations.

With the isopropanol solvent model, the highest barrier is found in the cation-free system, in agreement with the quenching of the catalytic activity by addition an 18-crown-6 to the potassium system, which effectively sequesters the cation. This barrier displays consistent values of 16.9 kcal mol<sup>-1</sup> when keeping two solvating isopropanol molecules, 17.1 kcal mol<sup>-1</sup> with three solvent molecules and 17.6 kcal mol<sup>-1</sup> when isopropanol is only modelled as a continuum (no explicit solvent).








Figure 5. Favoured solvation spheres for  $[M(\text{iPrOH})_6]^+$  systems. Distances in Å.

From our analysis, the solvation sphere of the cation plays an important role. We have always found ordered and strongly oriented first solvation shells around the spherical alkali cations, with oxygen atoms pointing toward the cation. We have further analysed this issue by determining the preferred conformation of lithium, sodium and potassium solvated cations in the presence of six solvent (isopropanol) molecules, in isopropanol (continuum). Figure 5 displays these structures. Lithium shows a clear preference to accommodate four isopropanol molecules in the first solvation sphere. In presence of 6 isopropanol molecules a 4+2 conformation, with four solvent molecules in the first solvation sphere in a tetrahedral-like arrangement is 5.3 kcal mol<sup>-1</sup> more stable than a 5+1 and 7.7 kcal mol<sup>-1</sup> than having the 6 isopropanol molecules interacting with the cation. The situation is reversed for potassium, for which an octahedral-like coordination of six alcohols molecules is favoured over the 5+1 and 4+2 configurations (3.1 and 2.7 kcal mol<sup>-1</sup> above). An intermediate situation is found for sodium. The 4+2 is the preferred conformation, but six-coordination is only 1.6 kcal mol<sup>-1</sup> less favourable. Both the favoured coordination numbers and the optimised M-O distances agree with experimental studies and molecular dynamics simulations



studying structural characteristics of the solvation environment of alkali metal ions in various electron-donor oxygen containing solvents, with methanol among them.<sup>46, 47</sup>

These results raise the question of the influence in the ketone hydrogenation of a cation like tetramethylammonium (TMA). TMA is a large cation with a relatively low charge density that has been described as one of the simplest and most spherical representatives of the so-called “hydrophobic ions” and for which minor polarization effects can be expected.<sup>48</sup> A loose solvation can be expected for TMA, and indeed that is what we have found when optimising a  $[(\text{TMA})(\text{iPrOH})_6]^+$  cluster in isopropanol (Figure 6a).




Figure 6. a) Solvation sphere for  $[(\text{TMA})(\text{iPrOH})_6]^+$  system. Transition states for the hydride transfer in the  $[(\text{TMA})(\text{iPrOH})_5]^+$  (b) and cation free (c) systems. Distances in Å.

From this solvation scenario, a behaviour similar to the cation free system can be foreseen in the ketone hydrogenation reaction. We have computed the hydride transfer step in isopropanol with a  $[(\text{TMA})(\text{iPrOH})_5]^+$  cluster. The results, collected in Table 3, agree with this similarity with the cation-free case (barriers of 16.9 and 17.2 kcal mol<sup>-1</sup> for the cation-free and TMA systems, respectively). Indeed, this result was found in the experimental study using Me<sub>4</sub>N<sup>+</sup> instead of an alkali-metal cation (Table 1, entry 6). Comparison of both transition states (**TS<sub>23-iso</sub>** and **TS<sub>23-TMA</sub>**, Figure 6)



highlight the structural likeness of the transition states, in which the incipient alkoxide is interacting with only two solvent molecules with no influence of the cation.

### (c) The importance of a proper solvent model

As commented above, the hydride transfer goes along with a reorganisation of the cation solvation sphere. It can be expected that an appropriate description of the first solvation shell of the cation is a requirement for a reliable description of the process. Indeed, in the initial solvent model we only included three methanol molecules, as we did in a previous study of this reaction mechanism.<sup>9</sup> We found, however, that this model was not able to reproduce the experimental activity trend (see Figure S5). We will only comment on the main features of this inaccurate model. First, the order of the Gibbs energy barriers for the hydride transfer step is reversed:  $\text{Li}^+$  (9.7 kcal mol<sup>-1</sup>) <  $\text{Na}^+$  (13.1 kcal mol<sup>-1</sup>) <  $\text{K}^+$  (13.6 kcal mol<sup>-1</sup>). Secondly, the alkoxide/H<sub>2</sub> exchange (**TS**<sub>5-6</sub>) is associated with remarkably higher barriers, in such a way that the cycle energy span ( $\text{Li}^+$ : 11.6 kcal mol<sup>-1</sup>;  $\text{Na}^+$ : 18.5 kcal mol<sup>-1</sup>;  $\text{K}^+$ : 20.6 kcal mol<sup>-1</sup>) is determined by the relative energy of this ligand exchange transition state and **4-M**, rather than by the hydride transfer step.

Figure 7 collects the optimised structures of the **2-M-3MeOH** intermediates and **TS**<sub>2-3-M-3MeOH</sub> transition states. With this model, the carbonyl substrate is already in contact with the cation in **2**, hence there is no need to displace a solvent molecule from the cation during the hydride transfer process. Moreover, the  $\text{Na}^+$  and  $\text{K}^+$  cations cannot reach their optimal distorted octahedral solvation shell in any of the states.






Figure 7. Optimised structures along the hydride transfer step for the three cations with the  $[M(\text{MeOH})_3]^+$  solvent model. Distances in Å. C-H hydrogen atoms have been omitted for clarity.

After formation of the alkoxide intermediate **4**, the next step involves the exchange of a strongly bound, negatively charged alkoxide ligand by a neutral, weakly donating  $\text{H}_2$  molecule. It is apparent that, without the help of solvent molecules to stabilise the departing alkoxide ligand *via* H-bonding, ligand exchange will not be favoured. Figure 8 compares the transition states of the alkoxide/ $\text{H}_2$  exchange (**TS<sub>5-6</sub>**), optimised with the  $[M(\text{MeOH})_3]^+$  and  $[M(\text{MeOH})_5]^+$  solvent models, for the three cations. The stabilizing H-bond interaction between two  $\text{M}^+$ -coordinated MeOH molecules and the leaving alkoxide can be appreciated in all **TS<sub>5-6</sub>-M** structures, while it is absent in all **TS<sub>5-6</sub>-M-3MeOH** structures. Thanks to the assistance by these interactions, the barriers for this step are lowered to 5.4 ( $\text{Li}^+$ ), 9.8 ( $\text{Na}^+$ ) and 6.9 ( $\text{K}^+$ )  $\text{kcal mol}^{-1}$ . The corresponding barriers for the 3MeOH model are 11.6 ( $\text{Li}^+$ ), 13.7 ( $\text{Na}^+$ ) and 14.1 ( $\text{K}^+$ )  $\text{kcal mol}^{-1}$ .







Figure 8. Transition states of the alkoxide by H<sub>2</sub> ligand exchange with the [M(MeOH)<sub>3</sub>]<sup>+</sup> (top) and [M(MeOH)<sub>5</sub>]<sup>+</sup> (bottom) solvent models. Distances in Å. C-H hydrogen atoms have been omitted for clarity.

#### (d) DFT investigations of the cation effect on enantioselectivity and limitations of the solvent model

Contrarily to the strong alkali-metal cation effect on the catalytic activity, its influence on enantioselectivity is small (Table 1). To understand the origin of this behaviour, we have performed DFT calculations on the enantio-determining hydride-transfer step using the real prochiral acetophenone substrate while employing the ((S)-P,S<sup>Ph</sup>) ligand. Preliminary results revealed that calculations dealing with enantioselectivity required a description of the system that is as realistic as possible. For this reason, they were performed using an isopropanol solvent model, with the usual cluster-explicit approach. For this system, the S enantiomer of the ligand leads to a formation preference for the S enantiomer of the 1-phenylethanol product with all three alkali-



metal cations yielding essentially the same *ee*. The *pro-R* and *pro-S* transition states **TS**<sub>2-3</sub> were computed for the three [M-(iPrOH)<sub>5</sub>]<sup>+</sup> cations, as well as for the cation-free catalyst without explicit solvent molecules for comparative purposes. Figure 9 shows the resulting TS geometries, and important distances for the cation-free (top) and K (bottom) systems. Similar outputs result for the other two cations (see Figure S15). The main interaction that discriminates between the two diastereomeric transition states involves a phenyl group of the phosphine ligand. This group interacts with the acetophenone methyl group (C-H- $\pi$  interaction) in the *pro-S* TS and with the acetophenone phenyl group ( $\pi$ - $\pi$  interaction) in the *pro-R* TS. The cation is located in the less crowded space opposite to this key P-phenyl group. Consequently, the nature of the cation does not significantly alter the main interactions that govern the *ee*. Table 4 gathers the calculated  $X_{\text{Ph}} \cdots \text{HC}_{\text{Me}}$  and  $X_{\text{Ph}} \cdots X_{\text{Ph}}$  distances for all computed systems, together with the Gibbs energy difference between both transition states ( $\Delta\Delta G^\ddagger_{\text{R-S}}$ ). The geometrical parameters describing the substrate-ligand interaction are very similar in all cases, in agreement with the limited influence the cation plays on enantioselectivity.






Figure 9. *pro-S* and *pro-R* transition states for hydride transfer to acetophenone, in absence of the cation and with  $[K(iPrOH)_5]^+$ . Distances in Å.

Table 4. Main distances (Å), Gibbs energy differences ( $\text{kcal mol}^{-1}$ ) and computed enantiomeric excesses between the *pro-S* and *pro-R* transition states for hydride transfer to acetophenone with the  $[M(iPrOH)_5]^+$  model.

|                       | $X_{Ph} \cdots HC_{Me}$ | $X_{Ph} \cdots X_{Ph}$ | $\Delta\Delta G_{R-S}^\ddagger$ <sup>a</sup> | e.e. <sup>a</sup> |
|-----------------------|-------------------------|------------------------|--|-------------------|
| <b>Cation free</b>    | 2.767                   | 4.049                  | 3.3 (3.7)                                    | 99 (100)          |
| <b>Li<sup>+</sup></b> | 2.516                   | 3.850                  | 3.8 (2.7)                                    | 100 (98)          |
| <b>Na<sup>+</sup></b> | 2.534                   | 3.867                  | 3.1 (2.1)                                    | 99 (94)           |
| <b>K<sup>+</sup></b>  | 2.615                   | 3.918                  | 1.2 (1.7)                                    | 77 (89)           |

<sup>a</sup> B3LYP-D3/BS2 values in parentheses.

In each case, the *pro-S* transition state is lower than that of *pro-R*, in qualitative agreement with the experimental results. However, the computed  $\Delta\Delta G_{R-S}^\ddagger$  overestimate these differences, as comparison of experimental and computed e.e.s showcases.<sup>49</sup> We have recomputed the  $\Delta\Delta G_{R-S}$  with the B3LYP-D3 functional at the fixed M06-optimised geometries, with very similar outcome (Table 4). The calculations correctly predict that the *S* isomer is preferred and rationalizes the little influence of the cation nature on the e.e.s. as a consequence of the cation placement, but also



reveal the difficulties in quantitatively reproducing the experimental e.e.s in such a complex system. Several factors can be invoked to justify this discrepancy. On one hand, the energies are very sensitive to the position of isopropanol molecules and slightly different arrangements of the five solvent molecules appreciably modifies the value of  $\Delta\Delta G^\ddagger_{R-S}$ . On the other hand, the balance between the relative strength of the C-H- $\pi$  and  $\pi$ - $\pi$  interactions may be functional-dependent. Finally, our model of five solvent molecules satisfies the coordination requirements of the alkali cation but does not satisfy the van der Waals requirements of the substrate. The  $\Delta\Delta G^\ddagger$  reflects the different TS stabilization when docking the substrate with the two different enantiofaces, but this could be overestimated by the lack of explicit van der Waals interactions between the substrate and the solvent. The stronger interaction in the TS leading to the preferred product enantiomer may be somewhat buffered by stronger interactions with the solvent as well. To prove that, a much more expensive full explicit solvent model would be needed.

## Conclusions

The role of anionic hydride complexes as active catalysts in hydrogenation and hydrogen-transfer reactions is now becoming well-established. When such species operate in solvents such as *i*-PrOH, the cation may associate with the anion in addition to coordination of the ion by solvent molecules. The present contribution has illustrated one such case for the active catalyst produced *in situ* by activation of [IrCl(COD)((S)-P,S<sup>R</sup>)] under H<sub>2</sub> in the presence of a strong base. The stark dependence of the catalytic activity on the nature of the base alkali cation (K<sup>+</sup> > Na<sup>+</sup> > Li<sup>+</sup>) and the severe activity dampening observed when an 18-crown-6 is added to the K<sup>+</sup> system confirm our previous suggestion of an anionic active complex, with ion pairing between the





anion and cation.<sup>9</sup> The proposition, made solely on the basis of DFT calculations in the same contribution,<sup>9</sup> that such a species is the Ir<sup>III</sup> tetrahydrido complex  $M^+[\text{Ir}(\text{H})_4((\text{S})\text{-P}, \text{S}^{\text{R}})]$ , has recently found strong support by the observed formation of the related (and more stable)  $\text{K}^+[\text{Ir}(\text{H})_4(\text{dppe})]^-$  from  $[\text{IrCl}(\text{COD})(\text{dppe})]$  in isopropanol and  $\text{KO}^t\text{Bu}$  under hydrogenation conditions ( $\text{H}_2$  gas at 25 °C) and even under transfer hydrogenation conditions (without  $\text{H}_2$  at 80 °C).<sup>14</sup> This hypothesis is further supported by the DFT calculations carried out in the present contribution, where the experimental trend of alkali metal cation-dependent activity has not only been reproduced but also rationalised, at the condition of using a fully explicit cation coordination sphere,  $M(\text{ROH})_5^+$ . Most intriguingly, the activity does not appear to depend on any activating effect by the cation on either the ketone substrate or the iridium hydride species, but rather on the coordination sphere rearrangement at the alkali metal cation when, in the rate-determining transition state, the incipient alkoxide is formed and concurrently establishes a bond with the alkali metal cation, replacing one or more alcohol molecules in its coordination sphere. These results indicate that the inclusion of explicit solvent molecules (and a sufficient number of them) is of paramount importance for the appropriate interpretation of certain effects that may be observed in catalytic cycles that involve anionic active species.

The calculations reported here were also able to pinpoint the main enantio-discriminating interaction in the rate-determining transition state, rationalizing the relatively low effect of the alkali metal nature on the e.e. and further supporting the proposed catalyst structure. However, they are not able to quantitatively reproduce the experimental e.e.s, pointing out the need of a full explicit solvent model, involving very expensive DFT-based Molecular Dynamics simulations with a solvent box, for a more quantitative agreement. Despite this shortcoming, static DFT calculations with a



careful choice of the solvent model can reproduce the experimental trends and identify its origin. On the basis of the work presented in this contribution, the DFT tool may now be applied with confidence to further explorations of the ligand space in order to achieve improved activities and enantioselectivities for any prochiral substrate of interest.

### Supporting Information

General experimental details. NMR characterisation data. Full details of the computational methodology, computed energies and xyz coordinates.

### Acknowledgments

This work has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860322, and was granted access to the resources of the CICT (Centre Interuniversitaire de Calcul de Toulouse, project CALMIP). Calculations on this project were carried out with the use of CSUC supercomputing resources. We are also grateful to the CNRS (Centre National de la Recherche Scientifique) for additional funding. JML is supported by a Royal Society Industrial Fellowship (INF/R1\221057).

### References

1. W. Tang and X. Zhang, New Chiral Phosphorus Ligands for Enantioselective Hydrogenation, *Chem. Rev.*, 2003, **103**, 3029-3069.
2. J. H. Xie, S. F. Zhu and Q. L. Zhou, Transition Metal-Catalyzed Enantioselective Hydrogenation of Enamines and Imines, *Chem. Rev.*, 2011, **111**, 1713-1760.
3. H.-U. Blaser, B. Pugin and F. Spindler, in *Organometallics as Catalysts in the Fine Chemical Industry*, eds. M. Beller and H. U. Blaser, 2012, vol. 42, pp. 65-102.
4. C. S. G. Seo and R. H. Morris, Catalytic Homogeneous Asymmetric Hydrogenation: Successes and Opportunities, *Organometallics*, 2019, **38**, 47-65.



5. D.-S. Wang, Q.-A. Chen, S.-M. Lu and Y.-G. Zhou, Asymmetric Hydrogenation of Heteroarenes and Arenes, *Chem. Rev.*, 2012, **112**, 2557-2590. View Article Online  
DOI: 10.1039/D4SC04629C
6. J. L. Wen, F. Y. Wang and X. M. Zhang, Asymmetric hydrogenation catalyzed by first-row transition metal complexes, *Chem. Soc. Rev.*, 2021, **50**, 3211-3237.
7. C. A. Sandoval, T. Ohkuma, K. Muniz and R. Noyori, Mechanism of asymmetric hydrogenation of ketones catalyzed by BINAP/1,2-diamine-ruthenium(II) complexes, *J. Am. Chem. Soc.*, 2003, **125**, 13490-13503.
8. M. Yamakawa, H. Ito and R. Noyori, The metal-ligand bifunctional catalysis: A theoretical study on the ruthenium(II)-catalyzed hydrogen transfer between alcohols and carbonyl compounds, *J. Am. Chem. Soc.*, 2000, **122**, 1466-1478.
9. J. M. Hayes, E. Deydier, G. Ujaque, A. Lledós, R. Malacea-Kabbara, E. Manoury, S. Vincendeau and R. Poli, Ketone Hydrogenation with Iridium Complexes with “non N–H” Ligands: The Key Role of the Strong Base, *ACS Catal.*, 2015, **5**, 4368–4376.
10. L. Routaboul, S. Vincendeau, J.-C. Daran and E. Manoury, New ferrocenyl P,S and S,S ligands for asymmetric catalysis, *Tetrahedron: Asymmetry*, 2005, **16**, 2685-2690.
11. R. Malacea, J.-C. Daran, S. B. Duckett, J. P. Dunne, C. Godard, E. Manoury, R. Poli and A. C. Whitwood, Parahydrogen studies of H<sub>2</sub> addition to an Ir(I) complex containing chiral phosphine- thioether ligands: Implications for catalysis, *Dalton Trans.*, 2006, 3350-3359.
12. R. Malacea, E. Manoury, L. Routaboul, J.-C. Daran, R. Poli, J. P. Dunne, A. C. Withwood, C. Godard and S. B. Duckett, Coordination chemistry and diphenylacetylene hydrogenation catalysis of planar chiral ferrocenylphosphine-thioether ligands with cyclooctadieneiridium(I), *Eur. J. Inorg. Chem.*, 2006, 1803–1816.
13. E. Le Roux, R. Malacea, E. Manoury, R. Poli, L. Gonsalvi and M. Peruzzini, Highly efficient asymmetric hydrogenation of alkyl aryl ketones catalyzed by iridium complexes with chiral planar ferrocenyl phosphino-thioether ligands, *Adv. Synth. & Catal.*, 2007, **349**, 309-313.
14. P. Kisten, E. Manoury, A. Lledós, A. C. Whitwood, J. Lynam, J. Slattery, S. B. Duckett and R. Poli, Ir( $\eta^4$ -diene) precatalyst activation by strong bases: formation of an anionic Ir<sup>III</sup> tetrahydride, *Dalton*, 2023, **52**, 2495-2505.
15. K. Abdur-Rashid, D. G. Gusev, S. E. Landau, A. J. Lough and R. H. Morris, Organizing chain structures by use of proton-hydride bonding. The single-crystal X-ray diffraction structures of K(Q) Os(H)(5)((PPr3)-Pr-i)(2) and K(Q) Ir(H)(4)((PPr3)-Pr-i)(2), Q = 18-crown-6 and 1,10-diaza-18-crown-6, *J. Am. Chem. Soc.*, 1998, **120**, 11826-11827.
16. S. E. Landau, K. E. Groh, A. J. Lough and R. H. Morris, Large effects of ion pairing and protonic-hydridic bonding on the stereochemistry and basicity of crown- ; azacrown- ; and cryptand-222-potassium salts of anionic tetrahydride complexes of iridium(III), *Inorg. Chem.*, 2002, **41**, 2995-3007.



17. R. A. Grey, G. P. Pez and A. Wallo, Selective Homogeneous Catalytic Hydrogenation of Polynuclear Aromatics, *J. Am. Chem. Soc.*, 1980, **102**, 5948-5949. View Article Online  
DOI: 10.1039/D4SC04629C
18. R. A. Grey, G. P. Pez, A. Wallo and J. Corsi, Homogeneous Catalytic Hydrogenation of Carboxylic Acid Esters to Alcohols, *Chem. Commun.*, 1980, 783-784.
19. R. A. Grey, G. P. Pez and A. Wallo, Anionic Metal Hydride Catalysts. 2. Application to the Hydrogenation of Ketones, Aldehydes, Carboxylic Acid Esters, and Nitriles, *J. Am. Chem. Soc.*, 1981, **103**, 7536-7542.
20. G. P. Pez, R. A. Grey and J. Corsi, Anionic Metal Hydride Catalysts. 1. Synthesis of Potassium Hydrido(phosphine)ruthenate Complexes, *J. Am. Chem. Soc.*, 1981, **103**, 7528-7535.
21. R. A. Grey, G. P. Pez, A. Wallo and J. Corsi, The hydrogenation of carbonyl compounds catalyzed by anionic ruthenium hydride complexes and alkali-doped supported group-VIII metals, *Ann. NY Acad. Sci.*, 1983, **415**, 235-243.
22. R. Wilczynski, W. A. Fordyce and J. Halpern, Coordination Chemistry and Catalytic Properties of Hydrido(phosphine)ruthenate Complexes, *J. Am. Chem. Soc.*, 1983, **105**, 2066-2068.
23. W. A. Fordyce, R. Wilczynski and J. Halpern, Hydrido(phosphine)ruthenate complexes and their role in the catalytic hydrogenation of arenes *J. Organomet. Chem.*, 1985, **296**, 115-125.
24. D. E. Linn and J. Halpern, Roles of Neutral and Anionic Ruthenium Polyhydrides in the Catalytic Hydrogenation of Ketones and Arenes, *J. Am. Chem. Soc.*, 1987, **109**, 2969-2974.
25. P. A. Dub, N. J. Henson, R. L. Martin and J. C. Gordon, Unravelling the Mechanism of the Asymmetric Hydrogenation of Acetophenone by [RuX<sub>2</sub>(diphosphine)(1,2-diamine)] Catalysts, *J. Am. Chem. Soc.*, 2014, **136**, 3505-3521.
26. M. Nielsen, E. Alberico, W. Baumann, H. J. Drexler, H. Junge, S. Gladiali and M. Beller, Low-temperature aqueous-phase methanol dehydrogenation to hydrogen and carbon dioxide, *Nature*, 2013, **495**, 85-89.
27. E. Alberico, A. J. J. Lennox, L. K. Vogt, H. Jiao, W. Baumann, H.-J. Drexler, M. Nielsen, A. Spannenberg, M. P. Checinski, H. Junge and M. Beller, Unravelling the Mechanism of Basic Aqueous Methanol Dehydrogenation Catalyzed by Ru-PNP Pincer Complexes, *J. Am. Chem. Soc.*, 2016, **138**, 14890-14904.
28. R. van Putten, E. A. Uslamin, M. Garbe, C. Liu, A. Gonzalez-de-Castro, M. Lutz, K. Junge, E. J. M. Hensen, M. Beller, L. Lefort and E. A. Pidko, Non-Pincer-Type Manganese Complexes as Efficient Catalysts for the Hydrogenation of Esters, *Angew. Chem. Int. Ed.*, 2017, **56**, 7531-7534.



29. C. Liu, R. van Putten, P. O. Kulyaev, G. A. Filonenko and E. A. Pidko, Computational insights into the catalytic role of the base promoters in ester hydrogenation with homogeneous non-pincer-based Mn-P,N catalyst, *J. Catal.*, 2018, **363**, 136-143. [View Article Online](#)  
DOI: 10.1039/D4SC04629C
30. Z. Q. Liang, T. L. Yang, G. X. Gu, L. Dang and X. M. Zhang, Scope and Mechanism on Iridium-f-Amphamide Catalyzed Asymmetric Hydrogenation of Ketones, *Chin. J. Chem.*, 2018, **36**, 851-856.
31. C. Yin, Y.-F. Jiang, F. Huang, C.-Q. Xu, Y. Pan, S. Gao, G.-Q. Chen, X. Ding, S.-T. Bai, Q. Lang, J. Li and X. Zhang, A 13-million turnover-number anionic Ir-catalyst for a selective industrial route to chiral nicotine, *Nat. Commun.*, 2023, **14**, 3718.
32. P. Västilä, A. B. Zaitsev, J. Wettergren, T. Privalov and H. Adolfsson, The importance of alkali cations in the {RuCl<sub>2</sub>(p-cymene)}<sub>2</sub>-pseudodipeptide-catalyzed enantioselective transfer hydrogenation of ketones, *Chem. Eur. J.*, 2006, **12**, 3218-3225.
33. T. Slagbrand, T. Kivijarvi and H. Adolfsson, Bimetallic Catalysis: Asymmetric Transfer Hydrogenation of Sterically Hindered Ketones Catalyzed by Ruthenium and Potassium, *ChemCatChem*, 2015, **7**, 3445-3449.
34. E. A. Bielinski, P. O. Lagaditis, Y. Y. Zhang, B. Q. Mercado, C. Wurtele, W. H. Bernskoetter, N. Hazari and S. Schneider, Lewis Acid-Assisted Formic Acid Dehydrogenation Using a Pincer-Supported Iron Catalyst, *J. Am. Chem. Soc.*, 2014, **136**, 10234-10237.
35. E. A. Bielinski, M. Forster, Y. Zhang, W. H. Bernskoetter, N. Hazari and M. C. Holthausen, Base-Free Methanol Dehydrogenation Using a Pincer-Supported Iron Compound and Lewis Acid Co-catalyst, *ACS Catal.*, 2015, **5**, 2404-2415.
36. Y. Zhang, A. D. MacIntosh, J. L. Wong, E. A. Bielinski, P. G. Williard, B. Q. Mercado, N. Hazari and W. H. Bernskoetter, Iron catalyzed CO<sub>2</sub> hydrogenation to formate enhanced by Lewis acid co-catalysts, *Chem. Sci.*, 2015, **6**, 4291-4299.
37. N. Govindarajan and E. J. Meijer, Elucidating cation effects in homogeneously catalyzed formic acid dehydrogenation, *Faraday Discuss.*, 2019, **220**, 404-413.
38. M. Gawron, F. Gilch, D. Schmidhuber, J. A. Kelly, T. M. Horsley Downie, A. Jacobi von Wangelin, J. Rehbein and R. Wolf, Counterion Effect in Cobaltate-Catalyzed Alkene Hydrogenation, *Angew. Chem. Int. Ed.*, 2024, **63**, e202315381.
39. M. Jia and M. Bandini, Counterion Effects in Homogeneous Gold Catalysis, *ACS Catal.*, 2015, **5**, 1638-1652.
40. R. G. Epton, W. P. Unsworth and J. M. Lynam, DFT Studies of Au(I) Catalysed Reactions: Anion Effects and Reaction Selectivity, *Israel Journal of Chemistry*, 2023, **63**, e202200033.
41. F. Zaccaria, L. Sian, C. Zuccaccia and A. Macchioni, in *Advances in Organometallic Chemistry*, ed. P. J. Pérez, Academic Press, 2020, vol. 73, pp. 1-78.



42. R. J. Phipps, G. L. Hamilton and F. D. Toste, The progression of chiral anions from concepts to applications in asymmetric catalysis, *Nature Chemistry*, 2012, **4**, 603-614. View Article Online  
DOI: 10.1039/D4SC04629C
43. M. Mahlau and B. List, Asymmetric Counteranion-Directed Catalysis: Concept, Definition, and Applications, *Angew. Chem. Int. Ed.*, 2013, **52**, 518-533.
44. B. Cordero, V. Gomez, A. E. Platero-Prats, M. Reves, J. Echeverria, E. Cremades, F. Barragan and S. Alvarez, Covalent radii revisited, *Dalton Trans.*, 2008, 2832-2838.
45. A. V. Marenich, S. V. Jerome, C. J. Cramer and D. G. Truhlar, Charge Model 5: An Extension of Hirshfeld Population Analysis for the Accurate Description of Molecular Interactions in Gaseous and Condensed Phases, *J. Chem. Theory Comput.*, 2012, **8**, 527-541.
46. P. R. Smirnov, Structure of the Li<sup>+</sup> Ion Close Environment in Various Solvents, *Russ. J. Gen. Chem.*, 2019, **89**, 2443-2452.
47. P. R. Smirnov, Structure of the Nearest Environment of Na<sup>+</sup>, K<sup>+</sup>, Rb<sup>+</sup>, and Cs<sup>+</sup> Ions in Oxygen-Containing Solvents, *Russ. J. Gen. Chem.*, 2020, **90**, 1693-1702.
48. P. E. Mason, T. Martinek, B. Fábrián, M. Vazdar, P. Jungwirth, O. Tichacek, E. Duboué-Dijon and H. Martinez-Seara, Hydration of biologically relevant tetramethylammonium cation by neutron scattering and molecular dynamics, *Phys. Chem. Chem. Phys.*, 2024, **26**, 3208-3218.
49. J. Wahlers, J. Margalef, E. Hansen, A. Bayesteh, P. Helquist, M. Diéguez, O. Pàmies, O. Wiest and P.-O. Norrby, Proofreading experimentally assigned stereochemistry through Q2MM predictions in Pd-catalyzed allylic aminations, *Nat. Commun.*, 2021, **12**, 6719.



The experimental and computational data, including xyz coordinates for all states, have been included as part of the Supplementary Information.

[View Article Online](#)

DOI: 10.1039/D4SC04629C

