



HAL
open science

Toeplitz operators, submultiplicative filtrations and weighted Bergman kernels

Siarhei Finski

► **To cite this version:**

Siarhei Finski. Toeplitz operators, submultiplicative filtrations and weighted Bergman kernels. 2024.
hal-04778364

HAL Id: hal-04778364

<https://hal.science/hal-04778364v1>

Preprint submitted on 12 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Toeplitz operators, submultiplicative filtrations and weighted Bergman kernels

Siarhei Finski

Abstract. We demonstrate that the weight operator associated with a submultiplicative filtration on the section ring of a polarized complex projective manifold is a Toeplitz operator. We further analyze the asymptotics of the associated weighted Bergman kernel, presenting the local refinement of earlier results on the convergence of jumping measures for submultiplicative filtrations towards the pushforward measure defined by the corresponding geodesic ray.

Table of contents

1	Introduction	1
2	The spaces of norms, Kähler forms and geodesic rays	5
3	Asymptotics of weight operators, proofs of Theorems 1.4, 1.6	10
4	Transfer operator between L^2 -norms, a proof of Theorem 3.2	14
5	Functional calculus for Toeplitz operators	18
6	Superadditivity of weighted Bergman kernels, a proof of Theorem 1.1	24

1 Introduction

The main goal of this paper is to make a connection between the theory of Toeplitz operators, Bergman kernels and submultiplicative filtrations.

Throughout the whole article, we fix a complex projective manifold X , $\dim X = n$, and an ample line bundle L over it. Recall that a decreasing graded filtration $\mathbb{R} \ni \lambda \mapsto \mathcal{F}^\lambda R(X, L) := \bigoplus_{k=0}^{+\infty} \mathcal{F}_k^\lambda H^0(X, L^{\otimes k})$, $\mathcal{F}_k^\lambda H^0(X, L^{\otimes k}) \subset H^0(X, L^{\otimes k})$, on the section ring,

$$R(X, L) := \bigoplus_{k=0}^{+\infty} H^0(X, L^{\otimes k}), \quad (1.1)$$

is called *submultiplicative* if the multiplication map on $R(X, L)$ factors through

$$\mathcal{F}_k^\lambda H^0(X, L^{\otimes k}) \otimes \mathcal{F}_l^\mu H^0(X, L^{\otimes l}) \rightarrow \mathcal{F}_{k+l}^{\lambda+\mu} H^0(X, L^{\otimes(k+l)}), \quad \text{for any } \lambda, \mu \in \mathbb{R}, k, l \in \mathbb{N}. \quad (1.2)$$

The most geometrically natural example is the filtration given by the order of vanishing along a fixed divisor. Other examples include filtrations associated with the weight of a \mathbb{C}^* -action on the pair (X, L) , filtrations given by the restriction of the Harder-Narasimhan filtration on direct image vector bundles associated with tensor powers of a polarization on a family of manifolds [17], [43], filtrations associated with valuations or graded ideals [62], or finitely generated filtrations induced by an arbitrary filtration on $H^0(X, L^{\otimes k})$, for $k \in \mathbb{N}$ big enough.

Remark that any (decreasing) filtration \mathcal{F} on the Hermitian vector space (V, H) induces the *weight operator* $A(\mathcal{F}, H) \in \text{End}(V)$, defined as

$$A(\mathcal{F}, H)e_i = w_{\mathcal{F}}(e_i) \cdot e_i, \quad \text{where } w_{\mathcal{F}}(e) := \sup\{\lambda \in \mathbb{R} : e \in \mathcal{F}^\lambda V\}, e \in V, \quad (1.3)$$

and e_1, \dots, e_r , $r := \dim V$, is an orthonormal basis of (V, H) adapted to the filtration \mathcal{F} in the sense that e_1 has the maximal weight, e_2 has the maximal weight among vectors orthogonal to e_1 ,

and so on. We call $w_{\mathcal{F}} : V \rightarrow]-\infty, +\infty]$ the weight function of \mathcal{F} . The main purpose of this article is to study the weight operator associated with submultiplicative filtrations.

More precisely, we fix a positive Hermitian metric h^L on L , and denote by $\text{Hilb}_k(h^L)$ the L^2 -metric on $H^0(X, L^{\otimes k})$ induced by h^L , see (1.10). For a continuous function $g : \mathbb{R} \rightarrow \mathbb{R}$, we consider the *weighted Bergman kernel*, $B_k^{\mathcal{F},g}(x) \in \mathbb{R}$, $k \in \mathbb{N}$, $x \in X$, defined as

$$B_k^{\mathcal{F},g}(x) := \sum_{i=1}^{N_k} \left\langle g\left(\frac{A(\mathcal{F}_k, \text{Hilb}_k(h^L))}{k}\right) s_{i,k}(x), s_{i,k}(x) \right\rangle_{h^{L^{\otimes k}}}, \quad (1.4)$$

where $N_k := \dim H^0(X, L^{\otimes k})$ and $s_{i,k}$, $i = 1, \dots, N_k$, is an orthonormal basis of $(H^0(X, L^{\otimes k}), \text{Hilb}_k(h^L))$. The reader will check that (1.4) doesn't depend on the choice of an orthonormal basis, and if the basis is adapted to \mathcal{F}_k , then $B_k^{\mathcal{F},g}(x) = \sum_{i=1}^{N_k} g(w_{\mathcal{F}_k}(s_{i,k})/k) |s_{i,k}(x)|_{h^{L^{\otimes k}}}^2$. When \mathcal{F} coincides with the grading on $R(X, L)$ (we then say \mathcal{F} is trivial), $B_k^{\mathcal{F},g}$ coincides with the usual Bergman kernel up to a constant.

Our first goal is to study the behavior of $B_k^{\mathcal{F},g}$, as $k \rightarrow \infty$. As we shall see, this study depends largely on the algebraic properties of \mathcal{F} .

We say that a submultiplicative filtration \mathcal{F} on $R(X, L)$ is *bounded* if there is $C > 0$, such that for any $k \in \mathbb{N}^*$, $\mathcal{F}^{Ck} H^0(X, L^{\otimes k}) = \{0\}$. We say that \mathcal{F} is *finitely generated* if it has integral weights and the associated $\mathbb{C}[\tau]$ -algebra $\text{Rees}(\mathcal{F}) := \sum_{(\lambda,k) \in \mathbb{Z} \times \mathbb{N}} \tau^{-\lambda} \mathcal{F}^{\lambda} H^0(X, L^{\otimes k})$, also called the *Rees algebra*, is finitely generated. Finitely generated submultiplicative filtrations are clearly automatically bounded. As the section ring $R(X, L)$ is finitely generated, cf. [49, Example 2.1.30], the set of finitely generated submultiplicative filtrations is non-empty, and for an arbitrary submultiplicative filtration, there is $C > 0$, such that for any $k \in \mathbb{N}^*$, $\mathcal{F}^{-Ck} H^0(X, L^{\otimes k}) = H^0(X, L^{\otimes k})$.

Recall that Phong-Sturm [60, Theorem 3] and Ross-Witt Nyström [64] associated for an arbitrary bounded submultiplicative filtration on $R(X, L)$ the *geodesic ray* $h_t^{\mathcal{F}}$, $t \in [0, +\infty[$, of Hermitian metrics on L , emanating from h^L . In general $h_t^{\mathcal{F}}$ is not smooth, however, due to convexity properties in t -variable, one can always define its derivative at $t = 0$, $\dot{h}_0^{\mathcal{F}} := (h_0^{\mathcal{F}})^{-1} \frac{d}{dt} h_t^{\mathcal{F}}|_{t=0} : X \rightarrow \mathbb{R}$, and this derivative is bounded, see Section 2 for details. We denote $\phi(h^L, \mathcal{F}) = -\dot{h}_0^{\mathcal{F}}$ for brevity.

Theorem 1.1. *For a bounded submultiplicative filtration \mathcal{F} on $R(X, L)$, the sequence of functions $x \mapsto \frac{1}{k^n} B_k^{\mathcal{F},g}(x)$, $x \in X$, $k \in \mathbb{N}$, is uniformly bounded and converges pointwise to a function which equals $g(\phi(h^L, \mathcal{F}))$ almost everywhere. If, moreover, \mathcal{F} is finitely generated, then the limit coincides with $g(\phi(h^L, \mathcal{F}))$ everywhere, and the convergence is uniform.*

Remark 1.2. a) When \mathcal{F} is trivial, Theorem 1.1 recovers the result of Tian [72] concerning the Bergman kernel asymptotics, and $\phi(h^L, \mathcal{F})$ equals to 1 (we emphasize that our proof builds on Tian's result!). When the filtration is induced by the \mathbb{C}^* -action on (X, L) , so that the induced S^1 -action is isometric for the Kähler form associated with h^L , the result is also well known, [70, §7.3], [55], and $\phi(h^L, \mathcal{F})$ then coincides with the Hamiltonian of the induced S^1 -action. See also [67] for a recent result on the off-diagonal behavior in the related context.

b) In the end of Section 2, we provide several examples indicating that Theorem 1.1 is sharp. Specifically, we show that without the finite-generation assumption, pointwise convergence cannot be strengthened to uniform convergence, and the limit doesn't coincide with $g(\phi(h^L, \mathcal{F}))$ everywhere. We then show that with finite-generation assumption, uniform convergence cannot be improved to \mathcal{C}^1 -convergence.

c) We explain in (2.10) that Theorem 1.1 partially justifies a folklore conjecture on the \mathcal{C}^1 -convergence of quantized geodesic rays towards the geodesic ray associated with the filtration.

We will now describe the context behind Theorem 1.1. Remark the following basic identity

$$\frac{1}{n!} \int B_k^{\mathcal{F},g}(x) c_1(L, h^L)^n = \text{Tr} \left[g \left(\frac{A(\mathcal{F}_k, \text{Hilb}_k(h^L))}{k} \right) \right], \quad (1.5)$$

which in particular implies that Theorem 1.1 guarantees the weak convergence of the spectral measures of the operators $\frac{1}{k} A(\mathcal{F}_k, \text{Hilb}_k(h^L))$ towards the pushforward measure $\phi(h^L, \mathcal{F})_*(c_1(L, h^L)^n / \int c_1(L)^n)$, as $k \rightarrow \infty$. From the definitions, we see that these spectral measures coincide (up to a normalization) with the jumping measures associated with the filtration, cf. [74], [47], where the latter probability measure on \mathbb{R} is defined as

$$\mu_{\mathcal{F},k} := \frac{1}{N_k} \sum_{j=1}^{N_k} \delta \left[\frac{e_{\mathcal{F}}(j, k)}{k} \right], \quad (1.6)$$

where $\delta[x]$ is the Dirac mass at $x \in \mathbb{R}$ and $e_{\mathcal{F}}(j, k)$ are the *jumping numbers*, defined as follows

$$e_{\mathcal{F}}(j, k) := \sup \left\{ t \in \mathbb{R} : \dim \mathcal{F}^t H^0(X, L^{\otimes k}) \geq j \right\}. \quad (1.7)$$

Weak convergence of jumping measures was first established by Chen [17] and Boucksom-Chen [14]. Subsequently, Witt Nyström [74] proved that for filtrations associated with a \mathbb{C}^* -action, the weak limit coincides with the pushforward measure; he also conjectured the analogous relation for finitely generated filtrations. Hisamoto established this in [47], and the author [39, Theorem 5.4] further extended it for bounded submultiplicative filtrations. In light of (1.5), Theorem 1.1 can be interpreted as a local refinement of these statements, showing that *the convergence occurs at the level of functions themselves*, rather than solely at the level of their integrals.

We invite the reader to compare Theorem 1.1 with Berman-Boucksom-Witt Nyström [7, Theorem B] and Darvas-Xia [32, Theorem 1.2], where authors establish the convergence in weak topology of partial Bergman kernels associated with Nadel multiplier ideal sheaves of plurisubharmonic potentials. Remark however that filtrations associated with multiplier ideals are not necessarily submultiplicative (because a product of two L^2 -integrable sections is not necessarily L^2 -integrable), and so it seems that there is no direct connection between our findings.

Theorem 1.1 in particular applies to the filtration associated with the vanishing order along a submanifold. Related study on the partial Bergman kernel was initiated by Berman [4, Theorem 4.3], and then developed by Ross-Singer [65], Coman-Marinescu [24], Zelditch-Zhou [76], Sun [69], and others. As partial Bergman kernel corresponds to indicator functions g in (1.4), which are not continuous, our study doesn't apply directly to the partial Bergman kernel.

Theorem 1.1 is related with a much more refined result concerning the asymptotic properties of the weight operator. To explain this statement, let us recall a version of [51, Definition 7.2.1].

Definition 1.3. A sequence of operators $T_k \in \text{End}(H^0(X, L^{\otimes k}))$, $k \in \mathbb{N}$, is called a *Toeplitz operator* if there is a continuous function $f : X \rightarrow \mathbb{R}$, called the symbol of $\{T_k\}_{k=0}^{+\infty}$, such that for any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that for every $k \geq k_0$, we have

$$\|T_k - T_{f,k}\| \leq \epsilon, \quad (1.8)$$

where $\|\cdot\|$ is the operator norm on $\text{End}(H^0(X, L^{\otimes k}))$, subordinate to $\text{Hilb}_k(h^L)$, $T_{f,k} := B_k \circ M_{f,k}$, and $B_k : L^\infty(X, L^{\otimes k}) \rightarrow H^0(X, L^{\otimes k})$ is the orthogonal (Bergman) projection to $H^0(X, L^{\otimes k})$, and $M_{f,k} : H^0(X, L^{\otimes k}) \rightarrow L^\infty(X, L^{\otimes k})$ is the multiplication map by f , acting as $s \mapsto f \cdot s$.

Our second main result goes as follows.

Theorem 1.4. *For finitely generated submultiplicative filtrations \mathcal{F} , the rescaled weight operator, $\{\frac{1}{k}A(\mathcal{F}_k, \text{Hilb}_k(h^L))\}_{k=0}^{+\infty}$, forms a Toeplitz operator with symbol $\phi(h^L, \mathcal{F})$.*

It turns out that the assumption on finitely generatedness is crucial in Theorem 1.4 and quite surprisingly, it cannot be replaced by the regularity assumption on the geodesic ray associated with the filtration, see the end of Section 2 for an explicit example showing this.

To cover the case of a general bounded submultiplicative filtration, we introduce another definition which, though natural, appears not to have been previously considered.

Definition 1.5. A sequence of operators $T_k \in \text{End}(H^0(X, L^{\otimes k}))$ is called a *Toeplitz operator of Schatten class* if it is uniformly bounded in operator's norm, i.e. there is $C > 0$, such that $\|T_k\| \leq C$, for any $k \in \mathbb{N}$, and there is $f \in L^\infty(X)$, called the symbol of $\{T_k\}_{k=0}^{+\infty}$, so that for any $\epsilon > 0$, $p \in [1, +\infty[$, there is $k_0 \in \mathbb{N}$, such that for every $k \geq k_0$, in the notations of Definition 1.3,

$$\|T_k - T_{f,k}\|_p \leq \epsilon, \quad (1.9)$$

where $\|\cdot\|_p$ is the p -Schatten norm, defined for an operator $A \in \text{End}(V)$, of a finitely-dimensional Hermitian vector space (V, H) as $\|A\|_p = (\frac{1}{\dim V} \text{Tr}[|A|^p])^{\frac{1}{p}}$, $|A| := (AA^*)^{\frac{1}{2}}$.

We can now state the final result of this paper.

Theorem 1.6. *For any bounded submultiplicative filtration \mathcal{F} , the rescaled weight operator, $\{\frac{1}{k}A(\mathcal{F}_k, \text{Hilb}_k(h^L))\}_{k=0}^{+\infty}$, forms a Toeplitz operator of Schatten class with symbol $\phi(h^L, \mathcal{F})$.*

Let us describe the relation between Theorems 1.4, 1.6 and Theorem 1.1. Remark that it is well known that the space of Toeplitz operators forms an algebra, and the symbol map is an algebra morphism, cf. [12], [51, §7]. The analogue of this statement extends to Toeplitz operators of Schatten class, see Proposition 5.8, and it implies that the vector space of Toeplitz operators of Schatten class is closed under the continuous functional calculus. As we shall explain in Section 5, this result along with some basic properties of the diagonal kernel show that Theorem 1.4 implies the second part of Theorem 1.1, and Theorem 1.6 implies the first part of Theorem 1.1, if one replaces the pointwise convergence by the convergence in $L^p(X)$ -spaces, for any $p \in [1, +\infty[$. However, the diagonal kernels of Toeplitz operators of Schatten class do not necessarily converge pointwise (just as convergence in $L^p(X)$ of a sequence of functions doesn't imply the pointwise convergence), and so Theorem 1.1 is not a consequence of Theorems 1.4, 1.6.

This paper is organized as follows. In Section 2, we recall the preliminaries. In Section 3, we prove Theorems 1.4, 1.6 modulo a similar statement concerning the transfer operator between L^2 -norms, which we establish in Section 4. In Section 5, we discuss the convergence properties of the diagonal kernels of Toeplitz operators and as a consequence establish Theorem 1.1 where the pointwise convergence is replaced by the convergence in $L^p(X)$ -spaces, for any $p \in [1, +\infty[$. The pointwise convergence part from Theorem 1.1 is then established in Section 6.

To conclude, we emphasize that we used the results of this paper in [42] to construct approximate solutions to a certain equation arising in relation to the Wess-Zumino-Witten equation.

Notation. A *filtration* \mathcal{F} of a vector space V is a map from \mathbb{R} to vector subspaces of V , $t \mapsto \mathcal{F}^t V$, verifying $\mathcal{F}^t V \subset \mathcal{F}^s V$ for $t > s$, and such that $\mathcal{F}^t V = V$ for sufficiently small t and $\mathcal{F}^t V = \{0\}$ for sufficiently big t . We assume that this map is left-continuous, i.e. for any $t \in \mathbb{R}$, there is $\epsilon_0 > 0$, such that $\mathcal{F}^t V = \mathcal{F}^{t-\epsilon} V$ for any $0 < \epsilon < \epsilon_0$.

For a given function f on a topological space, we denote by f^* (resp. f_*) the upper (resp. lower) semi-continuous regularization of f . The same notations are used for metrics on line bundles.

By a positive Hermitian metric on a line bundle we mean a smooth Hermitian metric with strictly positive curvature. For a positive Hermitian metric h^L on L (resp. and a fixed Kähler form χ), we denote by $\text{Hilb}_k(h^L)$ (resp. $\text{Hilb}_k(h^L, \chi)$) the Hermitian metric on $H^0(X, L^{\otimes k})$ defined for arbitrary $s_1, s_2 \in H^0(X, L^{\otimes k})$ as follows

$$\begin{aligned} \langle s_1, s_2 \rangle_{\text{Hilb}_k(h^L)} &= \frac{1}{n!} \int_X \langle s_1(x), s_2(x) \rangle_{h^L} \cdot c_1(L, h^L)^n, \\ \langle s_1, s_2 \rangle_{\text{Hilb}_k(h^L, \chi)} &= \frac{1}{n!} \int_X \langle s_1(x), s_2(x) \rangle_{h^L} \cdot \chi^n. \end{aligned} \quad (1.10)$$

Let V be a complex vector space, $\dim V = n$, and \mathcal{H}_V be the space of Hermitian norms on V . For $H_0, H_1 \in \mathcal{H}_V$, the *transfer map*, $T \in \text{End}(V)$, between H_0, H_1 , is the Hermitian operator (with respect to both H_0, H_1), defined so that the Hermitian products $\langle \cdot, \cdot \rangle_{H_0}, \langle \cdot, \cdot \rangle_{H_1}$ induced by H_0 and H_1 , are related as $\langle \cdot, \cdot \rangle_{H_1} = \langle \exp(-T) \cdot, \cdot \rangle_{H_0}$. For Hermitian $A_0, A_1 \in \text{End}(V)$ on a Hermitian vector space (V, H) , we say that $A_0 \geq A_1$ if the difference $A_0 - A_1$ is positively-definite. For a \mathcal{C}^1 -path $H_t \in \mathcal{H}_V$, we denote by $\dot{H}_t := H_t^{-1} \frac{d}{dt} H_t$.

We denote by $\mathbb{D}(a, b)$ the complex annuli if inner and outer radiuses a and b respectively. We denote by \mathbb{D} the complex unit disc. For a Kähler form ω on X , we denote by $\text{PSH}(X, \omega)$ the space of ω -quasipsh potentials, consisting of functions $\psi : X \rightarrow [-\infty, +\infty[$, which are locally the sum of a psh function and of a smooth function so that the $(1, 1)$ -current $\omega + \sqrt{-1} \partial \bar{\partial} \psi$ is positive. We denote $\pi : X \times \mathbb{D}(e^{-1}, 1) \rightarrow X$ and $z : X \times \mathbb{D}(e^{-1}, 1) \rightarrow \mathbb{D}(e^{-1}, 1)$ the usual projections.

For $T_k \in \text{End}(H^0(X, L^{\otimes k}))$ and $x \in X$, we denote

$$T_k(x) := \sum_{i=1}^{N_k} \langle T_k s_{i,k}(x), s_{i,k}(x) \rangle_{h^L}, \quad (1.11)$$

in the notations of (1.4), and call it the diagonal kernel of T_k . Recall that the diagonal Bergman kernel $B_k(x)$, $k \in \mathbb{N}$, $x \in X$, defined for an orthonormal basis $s_{i,k}$, $i = 1, \dots, N_k$, $N_k := \dim H^0(X, L^{\otimes k})$, of $(H^0(X, L^{\otimes k}), \text{Hilb}_k(h^L))$, as $B_k(x) = \sum_{i=1}^{N_k} |s_{i,k}(x)|_{h^L}^2$. We also define $B_k(x, y) \in L_x^{\otimes k} \otimes (L_y^{\otimes k})^*$ as $B_k(x, y) = \sum_{i=1}^{N_k} s_{i,k}(x) \otimes s_{i,k}(y)^*$.

Acknowledgement. Author acknowledges the support of CNRS and École Polytechnique.

2 The spaces of norms, Kähler forms and geodesic rays

The primary aim of this section is to revisit some well-known results that will be utilized in the following sections. Specifically, we begin by discussing results related to the geometry of the space of Hermitian norms on a finite-dimensional vector space and complex interpolation. We also recall the connection with the space of filtrations.

Next, we examine the Mabuchi geometry of the space of Kähler potentials and recall the correspondence between test configurations and submultiplicative filtrations. This is followed by a construction of geodesic rays from test configurations and submultiplicative filtrations.

The section concludes with an explicit calculation of a geodesic ray associated with two filtrations, highlighting the sharpness of the paper's main results.

Geodesics in the space of Hermitian norms. The space \mathcal{H}_V of Hermitian norms on a complex vector space V , $\dim V = n$, carries a natural metric, cf. [10, §6]. A path $H_t \in \mathcal{H}_V$, $t \in [0, 1]$, is the *geodesic* between $H_0, H_1 \in \mathcal{H}_V$ with respect to this metric, if for the transfer map, $T \in \text{End}(V)$, between H_0, H_1 , the endomorphism tT , $t \in [0, 1]$ is the transfer map between H_0 and H_t .

It is possible to view the space of filtrations on V as the boundary at the infinity of \mathcal{H}_V , where the latter space is interpreted in terms of geodesic rays. For this, for any filtration \mathcal{F} on V , we associate a geodesic ray in \mathcal{H}_V as follows. We fix $H_V \in \mathcal{H}_V$, and define $H_t^{\mathcal{F}} \in \mathcal{H}_V$, $t \in [0, +\infty[$, through the associated Hermitian product as $\langle \exp(-tA(\mathcal{F}, H_V)) \cdot, \cdot \rangle_{H_V}$, where $A(\mathcal{F}, H_V) \in \text{End}(V)$ is the weight operator associated with \mathcal{F} , defined in the Introduction. It is immediate that $H_t^{\mathcal{F}}$ is a geodesic ray emanating from H_V .

It is well known that geodesics between Hermitian norms can be seen through the lens of complex interpolation theory, cf. [3, Theorem 5.4.1]. Let us recall the following crucial statement concerning the order-preserving properties of complex interpolation theory.

Proposition 2.1 (cf. [3, Theorem 5.4.1] and [41, Proposition 4.12]). For any $H_i \in \mathcal{H}_V$, $i = 0, 1, 2$, verifying $H_1 \leq H_2$, the geodesics H_t^1, H_t^2 , $t \in [0, 1]$, between H_0, H_1 and H_0, H_2 respectively, compare as $H_t^1 \leq H_t^2$. In particular, we have $H_0^{-1} \dot{H}_0^1 \leq H_0^{-1} \dot{H}_0^2$. Similarly, for any filtrations \mathcal{F}_i , $i = 1, 2$, on V , so that their weight functions are related as $w_{\mathcal{F}_1} \geq w_{\mathcal{F}_2}$, the associated geodesic rays $H_t^{\mathcal{F}_1}, H_t^{\mathcal{F}_2}$, $t \in [0, +\infty[$, emanating from H_0 compare as $H_t^{\mathcal{F}_1} \leq H_t^{\mathcal{F}_2}$. In particular, we have $A(\mathcal{F}_1, H_0) \geq A(\mathcal{F}_2, H_0)$.

Mabuchi geodesics. We denote by the \mathcal{H}_L space of positive Hermitian metrics on L . Upon fixing $h_0^L \in \mathcal{H}_L$, one can identify \mathcal{H}_L with the space \mathcal{H}_ω of Kähler potentials of $\omega := 2\pi c_1(L, h_0^L)$, consisting of $u \in \mathcal{C}^\infty(X, \mathbb{R})$, such that $\omega_u := \omega + \sqrt{-1} \partial \bar{\partial} u$ is strictly positive, through the map

$$u \mapsto h^L := e^{-u} \cdot h_0^L. \quad (2.1)$$

Mabuchi in [57] introduced a certain metric on \mathcal{H}_ω , the geodesics of which admit the description as solutions to a certain homogeneous Monge-Ampère equation. To recall this, we identify paths $u_t \in \mathcal{H}_\omega$, $t \in [0, 1]$, with rotationally-invariant $\hat{u} : X \times \mathbb{D}(e^{-1}, 1) \rightarrow \mathbb{R}$, as follows

$$\hat{u}(x, \tau) = u_t(x), \quad \text{where } x \in X \text{ and } t = -\log |\tau|. \quad (2.2)$$

According to [66], [36] smooth geodesic segments in Mabuchi space can be described as the only path $u_t \in \mathcal{H}_\omega$, $t \in [0, 1]$, connecting u_0 to u_1 , so that \hat{u} is the solution of the Dirichlet problem associated with the homogeneous Monge-Ampère equation

$$(\pi^* \omega + \sqrt{-1} \partial \bar{\partial} \hat{u})^{n+1} = 0, \quad (2.3)$$

with boundary conditions $\hat{u}(x, e^{\sqrt{-1}\theta}) = u_0(x)$, $\hat{u}(x, e^{-1+\sqrt{-1}\theta}) = u_1(x)$, $x \in X$, $\theta \in [0, 2\pi]$. By the work of X. Chen [19] and later compliments by Błocki [11] and Chu-Tosatti-Weinkove [22], we now know that $\mathcal{C}^{1,1}$ solutions to (2.3) always exist.

Berndtsson in [8, §2.2] also proved that even for $u_0, u_1 \in \text{PSH}(X, \omega) \cap L^\infty(X)$, *weak solutions* to (2.3) exist, i.e. (2.3) has solutions when the wedge power is interpreted in Bedford-Taylor

sense [2] and the boundary conditions mean that $\|u_\epsilon - u_0\|_{L^\infty(X)} \rightarrow 0$ and $\|u_{1-\epsilon} - u_1\|_{L^\infty(X)} \rightarrow 0$, as $\epsilon \rightarrow 0$. We then have $u_t \in \text{PSH}(X, \omega) \cap L^\infty(X)$, but u_t are highly non-regular in general. However, since $u_t(x)$ is convex in $t \in [0, 1]$, cf. [35, Theorem I.5.13], the one-sided derivatives \dot{u}_t^-, \dot{u}_t^+ of u_t are well-defined for $t \in]0, 1[$ and they increase in t . In particular, one can define the derivative of the geodesic segment in this case as $\dot{u}_0 := \lim_{t \rightarrow 0} \dot{u}_t^- = \lim_{t \rightarrow 0} \dot{u}_t^+$. From [8, §2.2], we know that \dot{u}_0 is bounded and by Darvas [28, Theorem 1], we, moreover, have

$$\sup |\dot{u}_0| \leq \sup |u_1 - u_0|. \quad (2.4)$$

Analogously to Theorem 2.1, we have the following comparison result, which follows directly from the envelope description of the Mabuchi geodesics, cf. [8, (2.1)].

Proposition 2.2. For any $u_0 \in \mathcal{H}_\omega$, $u_i \in \text{PSH}(X, \omega) \cap L^\infty(X)$, $i = 1, 2$, verifying $u_1 \leq u_2$, the geodesics u_t^1, u_t^2 , $t \in [0, 1]$, between u_0, u_1 and u_0, u_2 respectively, compare as $u_t^1 \leq u_t^2$. In particular, the respective derivatives at zero compare as $\dot{u}_0^1 \leq \dot{u}_0^2$.

Test configurations and submultiplicative filtrations. Recall, cf. [73], [37], that a test configuration $\mathcal{T} = (\pi : \mathcal{X} \rightarrow \mathbb{C}, \mathcal{L})$ for (X, L) consists of the following data

1. A scheme \mathcal{X} with a \mathbb{C}^* -action ρ ,
2. A \mathbb{C}^* -equivariant *ample* line bundle \mathcal{L} over \mathcal{X} ,
3. A flat \mathbb{C}^* -equivariant projection $\pi : \mathcal{X} \rightarrow \mathbb{C}$, where \mathbb{C}^* acts on \mathbb{C} by multiplication, such that if we denote its fibers by $X_\tau := \pi^{-1}(\tau)$, $\tau \in \mathbb{C}$, then $(X_1, \mathcal{L}|_{X_1})$ is isomorphic to (X, L) .

Remark that the \mathbb{C}^* -action induces the canonical isomorphisms

$$\mathcal{X} \setminus X_0 \simeq \mathbb{C}^* \times X, \quad \mathcal{L}|_{\mathcal{X} \setminus X_0} \simeq p^*L, \quad (2.5)$$

where $p : \mathbb{C}^* \times X \rightarrow X$ is the natural projection.

Following Witt Nyström [74, Lemma 6.1], one can construct a submultiplicative filtration $\mathcal{F}^{\mathcal{T}}$ on $R(X, L)$ associated with \mathcal{T} as follows. Pick an element $s \in H^0(X, L^k)$, $k \in \mathbb{N}^*$, and consider the section $\tilde{s} \in H^0(\mathcal{X} \setminus X_0, \mathcal{L}^k)$, obtained by the application of the \mathbb{C}^* -action to s . By the flatness of π , the section \tilde{s} extends to a meromorphic section over \mathcal{X} , cf. Witt Nyström [74, Lemma 6.1]. In other words, there is $l \in \mathbb{Z}$, such that for a coordinate τ on \mathbb{C} , we have $\tilde{s} \cdot \tau^l \in H^0(\mathcal{X}, \mathcal{L}^k)$. We define the restriction $\mathcal{F}_k^{\mathcal{T}}$ of the filtration $\mathcal{F}^{\mathcal{T}}$ to $H^0(X, L^k)$ as

$$\mathcal{F}_k^{\mathcal{T}\lambda} H^0(X, L^k) := 2 \cdot \left\{ s \in H^0(X, L^k) : \tau^{-\lceil \lambda \rceil} \cdot \tilde{s} \in H^0(\mathcal{X}, \mathcal{L}^k) \right\}, \quad \lambda \in \mathbb{R}. \quad (2.6)$$

Remark the non-standard normalization by the factor 2 in (2.6), the motivation for which comes from [41, Theorem 1.1]. One can verify that the filtration $\mathcal{F}^{\mathcal{T}}$ is finitely generated and, up to a restriction to $R(X, L^d) \subset R(X, L)$, for some $d \in \mathbb{N}$, an arbitrary finitely generated filtration is produced from a test configuration for $(X, L^{\otimes d})$, cf. [74, (9)] or [15, §A.2].

Construction of geodesic rays in Mabuchi space. Consider the restriction $\pi' : \mathcal{X}'_{\mathbb{D}} \rightarrow \mathbb{D}$ of a resolution of singularities $\mathcal{T}' := (\pi' : \mathcal{X}' \rightarrow \mathbb{C}, \mathcal{L}')$ of a test configuration \mathcal{T} for (X, L) to the unit disc \mathbb{D} and denote $\mathcal{L}'_{\mathbb{D}} := \mathcal{L}'|_{\mathcal{X}'_{\mathbb{D}}}$. Phong-Sturm in [60, Theorem 3] established that for any fixed smooth positive metric h_0^L on L , there is a rotation-invariant bounded psh metric $h_{\mathbb{D}}^{\mathcal{L}'}$ over $\mathcal{L}'_{\mathbb{D}}$, solving weakly the Monge-Ampère equation

$$c_1(\mathcal{L}'_{\mathbb{D}}, h_{\mathbb{D}}^{\mathcal{L}'})^{n+1} = 0, \quad (2.7)$$

and such that its restriction over $\partial\mathcal{X}'_{\mathbb{D}}$ coincides with the rotation-invariant metric obtained from the fixed metric h_0^L on L . Under the identification (2.5), we then construct a ray $h_t^{\mathcal{T}}, t \in [0, +\infty[$, of metrics on L , such that $\dot{h}^{\mathcal{T}} = h_{\mathbb{D}}^{\mathcal{L}'}$ in the notations (2.2). Due to the equation (2.7) and the description of the geodesic ray as in (2.3), we see that the ray of metrics $h_t^{\mathcal{T}}, t \in [0, +\infty[$, is a geodesic ray emanating from h_0^L . This ray of metrics is $\mathcal{C}^{1,1}$, see [61], [22].

More generally, one can construct a geodesic ray from an arbitrary bounded submultiplicative filtration. For this, we first recall the well-known construction which associates for an arbitrary Hermitian norm H_k on $H^0(X, L^{\otimes k})$, for $k \in \mathbb{N}$ so that $L^{\otimes k}$ is very ample, a positive metric $FS(H_k)$ on $L^{\otimes k}$, is constructed for any $l \in L_x^{\otimes k}$, $x \in X$, as follows

$$|l|_{FS(H_k)}^2 := \frac{|l|_{h_0^{L^{\otimes k}}}^2}{\sum_{i=1}^{N_k} |s_{i,k}(x)|_{h_0^{L^{\otimes k}}}^2}, \quad (2.8)$$

where $s_{i,k} \in H^0(X, L^{\otimes k})$, is an orthonormal basis of $(H^0(X, L^{\otimes k}), H_k)$.

Now, let \mathcal{F} be a bounded submultiplicative filtration on $R(X, L)$. We fix a smooth positive metric h_0^L on L , and for any $t \in [0, +\infty[$, $k \in \mathbb{N}$, we define, following Ross-Witt Nyström [64], $H_{t,k}^{\mathcal{F}}$ as the (geodesic) ray of Hermitian norms on $H^0(X, L^k)$ emanating from $\text{Hilb}_k(h_0^L)$ and associated with the restriction \mathcal{F}_k of \mathcal{F} to $H^0(X, L^{\otimes k})$. We denote by $h_t^{\mathcal{F}}, t \in [0, +\infty[$, the ray of metrics on L , constructed as follows

$$h_t^{\mathcal{F}} := \left(\liminf_{k \rightarrow \infty} \liminf_{l \geq k} (FS(H_{t,l}^{\mathcal{F}})^{\frac{1}{k}}) \right)_*. \quad (2.9)$$

It was established in [64], following previous work [59] treating the finitely generated case, that $h_t^{\mathcal{F}}$ is a geodesic ray, i.e. it solves (2.3) weakly, cf. also [40, Theorem 5.1] for an independent proof based on quantization [20] and Mabuchi geometry [30]. In general, however, the metrics $h_t^{\mathcal{F}}$ are only bounded. It was established in [61] that the two constructions of geodesic rays, (2.7) and (2.9), are compatible when the filtration is associated with a test configuration.

It is a well-known open problem to study the precise convergence rate of $FS(H_{t,k}^{\mathcal{F}})^{\frac{1}{k}}$ towards $h_t^{\mathcal{F}}$, see [68] for example. Let us put Theorem 1.1 in this context. Let $x \mapsto B_k^{\mathcal{F}}(x)$ denote the weighted Bergman kernel, as defined in (1.4) for the function $g(x) := x$. Directly from (2.8), we see that

$$FS(H_{0,k}^{\mathcal{F}})^{-\frac{1}{k}} \frac{d}{dt} FS(H_{t,k}^{\mathcal{F}})^{\frac{1}{k}}|_{t=0} = -\frac{1}{B_k(x)} B_k^{\mathcal{F}}(x). \quad (2.10)$$

Recall now that a well-known result of Tian, [72], says

$$\frac{1}{k^n} B_k(x) \quad \text{converges uniformly to 1, as } k \rightarrow \infty, \quad (2.11)$$

see also [75], [16], [13], [51] for more refined results. In particular, Theorem 1.1 can be interpreted as a statement about convergence of the derivative of the initial point of the ray $FS(H_{t,k}^{\mathcal{F}})^{\frac{1}{k}}$ towards $\dot{h}_0^{\mathcal{F}}$, as $k \rightarrow \infty$, giving a partial justification for the \mathcal{C}^1 -convergence.

Example 1. The goal of the following example is to show that the assumption of finite degeneration in Theorem 1.1 is necessary and cannot be replaced by the regularity assumption on the geodesic ray associated with the filtration. The results of Section 5 will also imply that the weight operator associated with the filtration constructed here is not a Toeplitz operator, and so the assumptions of generation in Theorem 1.4 is necessary as well.

We consider the projective space $(X, L) := (\mathbb{P}^1, \mathcal{O}(1))$, and the filtration \mathcal{F} associated the weight function $w_{\mathcal{F}_k}(s) := k \min\{\text{ord}_0(s), 1\}$, where $\text{ord}_0(s)$ is the order of vanishing of $s \in H^0(\mathbb{P}^1, \mathcal{O}(k))$ at the point $0 := [1, 0] \in \mathbb{P}^1$. A straightforward verification reveals that the filtration \mathcal{F} is submultiplicative and bounded, but not finitely generated.

We identify \mathbb{P}^1 to $\mathbb{P}(V^*)$, where V is a vector space generated by two elements: x and y . Let us consider a metric H on V , which makes x and y an orthonormal basis, and denote by h^{FS} the induced Fubini-Study metric on $\mathcal{O}(1)$. For any $k \in \mathbb{N}^*$, $i, j \in \mathbb{N}$, $i + j = k$, under the isomorphism $\text{Sym}^k(V) \rightarrow H^0(\mathbb{P}(V^*), \mathcal{O}(k))$, an easy calculation shows that we have

$$\|x^i \cdot y^j\|_{\text{Hilb}_k(h^{FS})}^2 = \frac{i!j!}{(k+1)!}. \quad (2.12)$$

For the weight operator of \mathcal{F} , we then obtain the following formula

$$A(\mathcal{F}_k, \text{Hilb}_k(h^{FS}))(x^i \cdot y^j) = k(1 - \delta_{i,0})x^i \cdot y^j, \quad (2.13)$$

where the Kronecker symbol $\delta_{i,0}$ is defined so that $\delta_{i,0} = 1$ if $i = 0$, and $\delta_{i,0} = 0$ otherwise.

We then denote by $H_{t,k}^{\mathcal{F}}$ the geodesic ray departing from $\text{Hilb}_k(h^{FS})$ and associated with \mathcal{F}_k . For any $a, b \in \mathbb{C}$, not simultaneously equal to zero, we have

$$\frac{FS(\text{Hilb}_k(h^{FS}))}{FS(H_{t,k}^{\mathcal{F}})}([ax^* + by^*]) = \frac{e^{tk}(|a|^2 + |b|^2)^k + (1 - e^{tk})|a|^{2k}}{(|a|^2 + |b|^2)^k}. \quad (2.14)$$

In particular, for any $t \in [0, +\infty[$, we conclude that

$$\lim_{k \rightarrow \infty} \left(\frac{FS(\text{Hilb}_k(h^{FS}))}{FS(H_{t,k}^{\mathcal{F}})} \right)^{\frac{1}{k}}([ax^* + by^*]) = \begin{cases} e^t, & \text{if } b \neq 0, \\ 1, & \text{otherwise.} \end{cases} \quad (2.15)$$

Lower semicontinuous regularization will give $h_t^{\mathcal{F}} = e^{-t}h^{FS}$, which implies that $\phi(h^{FS}, \mathcal{F}) = 1$.

Let us calculate the weighted Bergman kernel for the function $g(x) = x$. We see directly that

$$B_k^{\mathcal{F},g}([ax^* + by^*]) = \sum_{i=1}^k \frac{(k+1)!}{i!(k-i)!} \frac{|a|^{2i}|b|^{2k-2i}}{(|a|^2 + |b|^2)^k} = (k+1) \frac{(|a|^2 + |b|^2)^k - |a|^{2k}}{(|a|^2 + |b|^2)^k}. \quad (2.16)$$

In particular, we deduce that

$$\lim_{k \rightarrow \infty} \frac{1}{k} B_k^{\mathcal{F},g}([ax^* + by^*]) = \begin{cases} 1, & \text{if } b \neq 0, \\ 0, & \text{otherwise,} \end{cases} \quad (2.17)$$

showing in particular that the convergence towards $\phi(h^{FS}, \mathcal{F})$ is neither uniform nor even pointwise, despite the regularity and simplicity of the associated geodesic ray, $h_t^{\mathcal{F}}$.

Example 2. The goal of the following example is to show that the uniform convergence in Theorem 1.1 cannot be improved to the \mathcal{C}^1 -convergence.

We consider the projective space $(X, L) := (\mathbb{P}^1, \mathcal{O}(2))$, and the filtration \mathcal{F} associated the weight function $w_{\mathcal{F}_k}(s) := \min\{\text{ord}_0(s), k\}$ in the notations of the previous example. The reader will check that \mathcal{F} is finitely generated. Similar calculation to the ones behind (2.14) will reveal that for any $a, b \in \mathbb{C}$, not simultaneously equal to zero, we have

$$\frac{FS(\text{Hilb}_{2k}(h^{FS}))}{FS(H_{t,k}^{\mathcal{F}})}([ax^* + by^*]) = \frac{1}{(|a|^2 + |b|^2)^{2k}} \left(\sum_{i=0}^k e^{ti} |a|^{2i} |b|^{2(2k-i)} \frac{(2k)!}{i!(2k-i)!} + e^{tk} \cdot \sum_{i=k+1}^{2k} |a|^{2i} |b|^{2(2k-i)} \frac{(2k)!}{i!(2k-i)!} \right). \quad (2.18)$$

Cramér's theorem from large deviations theory applied for the binomial distribution yields that for any $x < 1$, we have

$$\lim_{k \rightarrow \infty} \frac{1}{2k} \log \left(\sum_{i=k+1}^{2k} x^i \frac{(2k)!}{i!(2k-i)!} \right) = \frac{1}{2} \log(4x). \quad (2.19)$$

From (2.19) and binomial formula, it is immediate to recover that for any $t \in [0, +\infty[$, we have

$$\frac{(h^{FS})^2}{h_t^{\mathcal{F}}}([ax^* + by^*]) = \begin{cases} \left(\frac{e^t |a|^2 + |b|^2}{|a|^2 + |b|^2} \right)^2, & e^{t/2} |a| < |b|, \\ \left(\frac{2e^{t/2} |a||b|}{|a|^2 + |b|^2} \right)^2, & e^{-t/2} |b| < |a| < |b|, \\ e^t, & |b| < |a|. \end{cases} \quad (2.20)$$

By differentiation, we obtain

$$\phi(h^{FS}, \mathcal{F})([ax^* + by^*]) = \begin{cases} \frac{2|a|^2}{|a|^2 + |b|^2}, & |a| < |b|, \\ 1, & |b| < |a|. \end{cases} \quad (2.21)$$

One can easily see that $\phi(h^{FS}, \mathcal{F})$ is Lipschitz but not \mathcal{C}^1 . As for each $k \in \mathbb{N}^*$, the function $x \mapsto \frac{1}{k^n} B_k^{\mathcal{F},g}(x)$, $x \in X$, is clearly \mathcal{C}^1 , the uniform convergence cannot be improved to the \mathcal{C}^1 -convergence for the above filtration. Our example here is of course related to the well-known phenomena that one can expect at most $\mathcal{C}^{1,1}$ -regularity for the geodesic rays, cf. [5], [21], [68].

3 Asymptotics of weight operators, proofs of Theorems 1.4, 1.6

The primary objective of this section is to demonstrate Theorems 1.4 and 1.6, subject to an auxiliary result that will be established in the subsequent section of the article.

We will fix a positive smooth metric h_0^L on L , a continuous psh metric h_1^L on L , and Kähler forms χ_i , $i = 0, 1$, on X . We denote by $T_k(h_0^L, h_1^L) \in \text{End}(H^0(X, L^{\otimes k}))$ the transfer map between $\text{Hilb}_k(h_0^L, \chi_0)$ and $\text{Hilb}_k(h_1^L, \chi_1)$. Let h_t^L , $t \in [0, 1]$, be the Mabuchi geodesic between h_0^L and h_1^L . We denote $\phi(h_0^L, h_1^L) := -h_0^L$. By Demailly's regularization theorem [34], cf. [46], for any $\epsilon > 0$, we can find a positive Hermitian metric $h_{1,\epsilon}^L$ on L , verifying $h_1^L \exp(-\epsilon) \leq h_{1,\epsilon}^L \leq h_1^L \exp(\epsilon)$.

Proposition 3.1. As $\epsilon \rightarrow 0$, the sequence of functions $\phi(h_0^L, h_{1,\epsilon}^L)$ converges to $\phi(h_0^L, h_1^L)$ uniformly. In particular, $\phi(h_0^L, h_1^L)$ is a continuous function.

Proof. By Proposition 2.2, we have

$$\phi(h_0^L, h_{1,\epsilon}^L) - \epsilon \leq \phi(h_0^L, h_1^L) \leq \phi(h_0^L, h_{1,\epsilon}^L) + \epsilon. \quad (3.1)$$

The result follows directly from (3.1) since $\phi(h_0^L, h_{1,\epsilon}^L)$ is continuous for any $\epsilon > 0$ by [61]. \square

The following result, established in Section 4, is among the main results of this paper.

Theorem 3.2. *The operators $\{\frac{1}{k}T_k(h_0^L, h_1^L)\}_{k=1}^{+\infty}$, form a Toeplitz operator with symbol $\phi(h_0^L, h_1^L)$.*

We establish Theorem 1.4 by using Theorem 3.2. In order to make a connection between the transfer operator and the weight operator, we need to relate the associated rays of norms on $R(X, L)$. Let \mathcal{F} be a finitely generated submultiplicative filtration on $R(X, L)$. We define $h_t^{\mathcal{F}}$, $t \in [0, +\infty[$, and $H_{t,k}^{\mathcal{F}}$, $t \in [0, +\infty[$, $k \in \mathbb{N}$, as in (2.9). We also fix an arbitrary Kähler form χ on X . The following result is at the heart of our approach.

Theorem 3.3 ([41, Theorem 4.1]). *There are $C > 0$, $k_0 \in \mathbb{N}^*$, such that for any $t \in [0, +\infty[$, $k \geq k_0$, we have the following comparison of norms*

$$\exp(-C(t+k)) \cdot \text{Hilb}_k(h_t^{\mathcal{F}}, \chi) \leq H_{t,k}^{\mathcal{F}} \leq \exp(C(t+k)) \cdot \text{Hilb}_k(h_t^{\mathcal{F}}, \chi). \quad (3.2)$$

Remark 3.4. In [41, Theorem 4.1], we only established Theorem 3.3 for filtrations \mathcal{F} associated with a test configuration. Due to the correspondence between test configurations and finitely generated submultiplicative filtrations, recalled in Section 2, it implies that there is $d \in \mathbb{N}^*$, such that (3.2) holds for any k divisible by d . However, the same argument as in [41, proof of Theorem 4.1], based on the isometry properties of the surjective map $H^0(X, L^{\otimes k}) \otimes H^0(X, L^{\otimes l}) \rightarrow H^0(X, L^{\otimes(k+l)})$ then gives (3.2) for all $k \in \mathbb{N}$, see more specifically [41, (4.12), (4.14) and (4.18)].

We can now present the proof of the main result of this paper.

Proof of Theorem 1.4. We conserve the notations from Theorems 1.4 and 3.3. By Theorem 3.3 and Proposition 2.1, for any $k \geq k_0$, $t \in]0, +\infty[$, we have the following relation between the weight operator and the transfer operators

$$\left\| A(\mathcal{F}_k, \text{Hilb}_k(h^L)) - \frac{1}{t}T_k(h_0^{\mathcal{F}}, h_t^{\mathcal{F}}) \right\| \leq C\frac{k}{t} + C. \quad (3.3)$$

Theorem 1.4 follows directly from Theorem 3.2 and the trivial fact $\phi(h_0^{\mathcal{F}}, h_t^{\mathcal{F}}) = t\phi(h_0^{\mathcal{F}}, h_1^{\mathcal{F}})$. \square

We will now deduce Theorem 1.6 from Theorem 1.4 and approximation techniques for submultiplicative filtrations from [71], [6], [15], [39]. For this, we will need an auxillary statement, which will be established in Section 5.

Proposition 3.5. *For an arbitrary sequence $f_k \in L^\infty(X)$, $k \in \mathbb{N}$, of uniformly bounded functions converging, as $k \rightarrow \infty$, in $L^1(X)$ towards $f \in L^\infty(X)$, the sequence of operators $\{T_{f_k, k}\}_{k=0}^{+\infty}$ forms a Toeplitz operator of Schatten class with symbol f .*

Proof of Theorem 1.6. First of all, without loosing the generality we can assume that the filtration \mathcal{F} has integer weights. In order to see this, for an arbitrary filtration \mathcal{F} , we define the filtration

$\lfloor \mathcal{F} \rfloor$ through its weight function as $w_{\lfloor \mathcal{F} \rfloor}(s) := \lfloor w_{\mathcal{F}}(s) \rfloor$, $s \in R(X, L)$. It is easy to see that $\lfloor \mathcal{F} \rfloor$ is submultiplicative and bounded as well. Moreover, by Theorem 2.1, we have

$$A(\mathcal{F}_k, \text{Hilb}_k(h^L)) \geq A(\lfloor \mathcal{F}_k \rfloor, \text{Hilb}_k(h^L)) \geq A(\mathcal{F}_k, \text{Hilb}_k(h^L)) - \text{Id}. \quad (3.4)$$

Which shows that it suffices to establish Theorem 1.6 for $\lfloor \mathcal{F} \rfloor$. Without losing the generality, we assume from now on that the filtration \mathcal{F} has integer weights.

Now, as $R(X, L)$ is finitely generated, there is $k_0 \in \mathbb{N}$, such that $\bigoplus_{i=0}^{k_0} H^0(X, L^{\otimes i})$ generates $R(X, L)$ as a ring. For any $k \geq k_0$, we denote by $\mathcal{F}^{[k]}$ the filtration on $R(X, L)$, generated by the restriction of \mathcal{F} to $\bigoplus_{i=0}^{k_0} H^0(X, L^{\otimes i})$. Clearly, $\mathcal{F}^{[k]}$ has integer weights and is finitely generated. By Theorem 1.4, for any $\epsilon > 0$, $k \geq k_0$, there is $l_0 \in \mathbb{N}$, such that for any $l \geq l_0$, we have

$$\left\| \frac{1}{l} A(\mathcal{F}_l^{[k]}, \text{Hilb}_l(h^L)) - T_{\phi(h^L, \mathcal{F}^{[k]}, l)} \right\| \leq \epsilon. \quad (3.5)$$

Our proof resides on establishing the following two statements. First, we establish that

$$\phi(h^L, \mathcal{F}^{[k]}), k \in \mathbb{N} \text{ increases to } \phi(h^L, \mathcal{F}) \text{ almost everywhere.} \quad (3.6)$$

Second, we establish that for any $\epsilon > 0$, $p \in [1, +\infty[$, there is $k_1 \geq k_0$, such that for any $k \geq k_1$, there is $l_1 \in \mathbb{N}$, so that for any $l \geq l_0$, we have

$$\left\| A(\mathcal{F}_l^{[k]}, \text{Hilb}_l(h^L)) - A(\mathcal{F}_l, \text{Hilb}_l(h^L)) \right\|_p \leq \epsilon l. \quad (3.7)$$

Remark that Theorem 1.6 follows directly from Proposition 3.5, (3.5), (3.6) and (3.7) by Lebesgue dominated convergence theorem. From now on, we concentrate on the proofs of (3.6) and (3.7).

Remark first that as an immediate consequence of submultiplicativity of \mathcal{F} , the weight functions of the filtrations are related for any $k \geq k_0$, $l \in \mathbb{N}$, as follows

$$w_{\mathcal{F}_l^{[k]}} \leq w_{\mathcal{F}_l^{[k+1]}} \leq w_{\mathcal{F}_l}. \quad (3.8)$$

By Theorem 2.1 and (3.8), we deduce that for any $k \geq k_0$, $l \in \mathbb{N}^*$, we have

$$A(\mathcal{F}_l^{[k]}, \text{Hilb}_l(h^L)) \leq A(\mathcal{F}_l^{[k+1]}, \text{Hilb}_l(h^L)) \leq A(\mathcal{F}_l, \text{Hilb}_l(h^L)), \quad (3.9)$$

By Theorem 2.1, (2.9) and (3.8), we deduce that in the notations of (2.9), for any $k \in \mathbb{N}$, $t \in [0, +\infty[$, we have

$$h_t^{\mathcal{F}^{[k]}} \geq h_t^{\mathcal{F}^{[k+1]}} \geq h_t^{\mathcal{F}}. \quad (3.10)$$

By taking derivatives at $t = 0$ in (3.10), for any $k \in \mathbb{N}$, we establish

$$\phi(h^L, \mathcal{F}^{[k]}) \leq \phi(h^L, \mathcal{F}^{[k+1]}) \leq \phi(h^L, \mathcal{F}) \quad (3.11)$$

Let $k_2 \in \mathbb{N}$ is defined so that for any $l_1, l_2 \geq k_2$, the multiplication maps $H^0(X, L^{\otimes l_1}) \otimes H^0(X, L^{\otimes l_2}) \rightarrow H^0(X, L^{\otimes (l_1+l_2)})$ is surjective. Recall, cf. [17, Proposition 3.2.6], that for an arbitrary submultiplicative filtration \mathcal{F} , $l_1, l_2 \geq k_2$, we have

$$\min_{s \in H^0(X, L^{\otimes l_1})} w_{\mathcal{F}_{l_1}}(s) + \min_{s \in H^0(X, L^{\otimes l_2})} w_{\mathcal{F}_{l_2}}(s) \leq \min_{s \in H^0(X, L^{\otimes (l_1+l_2)})} w_{\mathcal{F}_{l_1+l_2}}(s). \quad (3.12)$$

From this and (3.8), we see in particular that the following bound holds

$$\sup_{k \geq \max(k_0, k_2)} \sup_{l \in \mathbb{N}^*} \frac{1}{l} \max_{s \in H^0(X, L^{\otimes l}) \setminus \{0\}} |w_{\mathcal{F}_l^{[k]}}(s)| \leq \sup_{l \in \mathbb{N}^*} \frac{1}{l} \max_{s \in H^0(X, L^{\otimes l}) \setminus \{0\}} |w_{\mathcal{F}_l}(s)|. \quad (3.13)$$

From [41, Lemma 2.4], using the boundness of \mathcal{F} , for $C := \sup_{l \in \mathbb{N}^*} \frac{1}{l} \max |w_{\mathcal{F}_l}|$, we have $\phi(h^L, \mathcal{F}) \leq C$. From this and (3.11), we see that $\phi(h^L, \mathcal{F}^{[k]})$ is uniformly bounded in $k \in \mathbb{N}$. Due to this and (3.11), in order to get (3.6), it suffices to establish that $\phi(h^L, \mathcal{F}^{[k]})$ converges to $\phi(h^L, \mathcal{F})$ in $L^1(X)$, as $k \rightarrow \infty$.

Recall that Darvas-Lu in [30, Theorem 3.1] established the following bound

$$\frac{1}{n! \int c_1(L)^n} \int |\phi(h^L, \mathcal{F}^{[k]}) - \phi(h^L, \mathcal{F})| c_1(L, h^L)^n \leq d_1(h_1^{\mathcal{F}^{[k]}}, h_1^{\mathcal{F}}), \quad (3.14)$$

where d_1 is the Darvas distance, [27]. In particular, to show the convergence of $\phi(h^L, \mathcal{F}^{[k]})$ to $\phi(h^L, \mathcal{F})$ in $L^1(X)$, as $k \rightarrow \infty$, it only suffices to show that $d_1(h_1^{\mathcal{F}^{[k]}}, h_1^{\mathcal{F}}) \rightarrow 0$, as $k \rightarrow \infty$. This was done in [39, Theorem 5.9] for a similar approximation scheme, where instead of the filtration $\mathcal{F}^{[k]}$, we considered the filtration $\mathcal{F}^{(k)}$ on $R(X, L^{\otimes k})$ induced by \mathcal{F}_k . Let us explain why this implies the searched convergence. Indeed, as an immediate consequence of submultiplicativity of \mathcal{F} , similarly to (3.8), the weight functions of the filtrations are related for any $k \geq k_0, l \in \mathbb{N}$, as

$$w_{\mathcal{F}_l^{(k)}} \leq w_{\mathcal{F}_{kl}^{[k]}} \leq w_{\mathcal{F}_{kl}}. \quad (3.15)$$

We denote by $h_t^{\mathcal{F}^{(k)}}$, $t \in [0, +\infty[$, the geodesic ray on $L^{\otimes k}$ emanating from $(h^L)^{\otimes k}$ and induced by $\mathcal{F}^{(k)}$. From (3.15), we deduce in the same way as in (3.10) that

$$(h_t^{\mathcal{F}^{(k)}})^{\frac{1}{k}} \geq h_t^{\mathcal{F}^{[k]}} \geq h_t^{\mathcal{F}}. \quad (3.16)$$

The statement [39, Theorem 5.9] says that $d_1((h_1^{\mathcal{F}^{(k)}})^{\frac{1}{k}}, h_1^{\mathcal{F}}) \rightarrow 0$, as $k \rightarrow \infty$. However, by (3.16) and the usual properties of the Darvas distance, cf. [26, Lemma 4.2], we have $d_1((h_1^{\mathcal{F}^{(k)}})^{\frac{1}{k}}, h_1^{\mathcal{F}}) \geq d_1(h_1^{\mathcal{F}^{[k]}}, h_1^{\mathcal{F}})$, which establishes (3.6).

We will now establish (3.7). Remark that due to (3.13), there is $C > 0$, so that for any $k \geq \max(k_0, k_2), l \in \mathbb{N}^*$, we have

$$\left\| A(\mathcal{F}_l^{[k]}, \text{Hilb}_l(h^L)) - A(\mathcal{F}_l, \text{Hilb}_l(h^L)) \right\| \leq Cl. \quad (3.17)$$

Remark also the classical fact that for an arbitrary operator $A_k \in \text{End}(H^0(X, L^{\otimes k}))$, we have

$$\|A_k\|_p \leq \|A_k\|_1^{\frac{1}{p}} \cdot \|A_k\|^{1-\frac{1}{p}}. \quad (3.18)$$

From (3.17) and (3.18), we see that it suffices to establish (3.7) for $p = 1$.

Directly from (3.9), we have

$$\begin{aligned} & \left\| A(\mathcal{F}_l^{[k]}, \text{Hilb}_l(h^L)) - A(\mathcal{F}_l, \text{Hilb}_l(h^L)) \right\|_1 \\ &= \frac{1}{N_k} \left(\text{Tr}[A(\mathcal{F}_l, \text{Hilb}_l(h^L))] - \text{Tr}[A(\mathcal{F}_l^{[k]}, \text{Hilb}_l(h^L))] \right). \end{aligned} \quad (3.19)$$

Recall that the volume, $\text{vol}(\mathcal{F})$, of a submultiplicative filtration \mathcal{F} is defined as follows

$$\text{vol}(\mathcal{F}) := \lim_{k \rightarrow \infty} \frac{n!}{k^{n+1}} \sum_{i=1}^{N_k} e_{\mathcal{F}}(i, k), \quad (3.20)$$

where $e_{\mathcal{F}}(i, k)$ are the *jumping numbers* of the filtration \mathcal{F}_k on $H^0(X, L^{\otimes k})$, defined in (1.7).

By (1.5), (3.19) and the above discussion, we see that (3.7) would follow if we establish that $\text{vol}(\mathcal{F}^{[k]}) \rightarrow \text{vol}(\mathcal{F})$. For a similar approximation scheme, where instead of the filtration $\mathcal{F}^{[k]}$, we consider the filtration $\mathcal{F}^{(k)}$ defined above, this was done by Boucksom-Jonsson in [15, Theorem 3.18 and (3.14)] and later by the author in [39, Theorem 5.10]. However, by proceeding as in (3.16), we easily see that this implies that the statement also holds for $\mathcal{F}^{[k]}$. \square

4 Transfer operator between L^2 -norms, a proof of Theorem 3.2

In this section, we describe our proof of Theorem 3.2. Our approach is inspired by Berndtsson's previous work [9], in the sense that it also relies on the connection between geodesics in the space of Hermitian metrics and the curvature of the associated Hermitian vector bundles. However, we diverge from it in several important ways.

First, by using the curvature calculations of L^2 -metrics due to Ma-Zhang [56], we demonstrate that L^2 -metrics can be used not only to construct superinterpolating families for the geodesic between two L^2 -metrics, as in [9], but also subinterpolating ones. This enables a much more refined analysis of the transfer operator, which gives an even stronger version of Theorem 3.2 when the Mabuchi geodesic is smooth and passes through positive metrics.

Second, [56] only works for smooth families of metrics, so we need to regularize our geodesic, which is not smooth in general. While a similar regularity issue arises in [9] and is addressed there by selecting an *arbitrary* smooth subgeodesic, the specific choice of subgeodesic is critical in our approach. In particular, the construction of ϵ -geodesics from [19] is crucial to our method. However, due to the lack of non-collapsing estimates for ϵ -geodesics, see the discussion after (4.15), this inevitably introduces additional challenges. Specifically, for non-smooth Mabuchi geodesics, we can only show that L^2 -metrics form subinterpolating families over a small interval that contains an endpoint. Relying on our previous work [39], we nevertheless show that this is enough to establish Theorem 3.2.

We will now proceed as follows. We first establish Theorem 3.2 for smooth endpoints, verifying an additional assumption that the Mabuchi geodesic between them is smooth and passes through positive metrics. Remark that it was established by Lempert-Vivas [50] and Darvas-Lempert [29] that for general endpoints, there is no smooth Mabuchi geodesic relating them. We then establish Theorem 3.2 for arbitrary smooth endpoints. By approximation, we establish Theorem 3.2 in its full generality.

Let us recall first a classical result making a connection between the geodesic construction in the space \mathcal{H}_V of Hermitian norms on a complex vector space V and positivity properties of related vector bundles. For this, similarly to (2.2), we identify a smooth path $H_t \in \mathcal{H}_V$, $t \in [0, 1]$, with the rotationally-invariant Hermitian metric \hat{H} on the (trivial) vector bundle $V \times \mathbb{D}(e^{-1}, 1)$ over $\mathbb{D}(e^{-1}, 1)$, through the formula

$$\hat{H}(\tau) = H_t, \quad \text{where } t = -\log |\tau|. \quad (4.1)$$

Recall that a Hermitian vector bundle (E, h^E) over a Riemann surface S is called positive, cf. [35, §VII.6], if the curvature R^E of the Chern connection of (E, h^E) , for any $s \in S$, can be

written as $R_s^E = dz \wedge d\bar{z} \cdot A(s)$, for a positively definite $A(s) \in \text{End}(E_s)$ and a local holomorphic coordinate z on S , centered at s .

Theorem 4.1 ([63, Theorem 4.2], [23, Theorem 4.1, §15]). *Assume that a smooth path $H_t^0 \in \mathcal{H}_V$, $t \in [0, 1]$, is such that the associated Hermitian metric \hat{H}_0 on $V \times \mathbb{D}(e^{-1}, 1)$ is positive (resp. negative). Then for the geodesic H_t between H_t^0 and H_t^1 , we have $H_t^0 \geq H_t$ (resp. $H_t^0 \leq H_t$). In particular, the following inequality is satisfied $\dot{H}_0^0 \geq \dot{H}_0$ (resp. $\dot{H}_0^0 \leq \dot{H}_0$). The path H_t^0 is then called a superinterpolating (resp. subinterpolating) family.*

We will also need the result of Ma-Zhang [54], [56] on the curvature of direct image bundles. We fix a smooth family of (strictly) positive Hermitian metrics h_τ^L , $\tau \in \mathbb{D}(e^{-1}, 1)$ on L and a smooth family of Kähler forms χ_τ , $\tau \in \mathbb{D}(e^{-1}, 1)$, on X . We denote by $\omega := c_1(L \times \mathbb{D}(e^{-1}, 1), h_\tau^L)$ the curvature of h_τ^L , viewed as a metric on the line bundle $L \times \mathbb{D}(e^{-1}, 1)$ over $X \times \mathbb{D}(e^{-1}, 1)$.

For $\tau \in \mathbb{D}(e^{-1}, 1)$, we define $\omega_H(\tau) \in \mathcal{C}^\infty(X)$ as

$$\omega_H(\tau)(x) := \frac{1}{n+1} \frac{\omega^{n+1}}{\omega^n \wedge \sqrt{-1} dz \wedge d\bar{z}}(x, \tau). \quad (4.2)$$

The denominator above is nonzero, as ω is positive along the fibers.

Now, we denote by R_k the curvature of the Chern connection on the trivial vector bundle $H^0(X, L^{\otimes k}) \times \mathbb{D}(e^{-1}, 1)$ associated with the fiberwise L^2 -metric $\text{Hilb}_k(h_\tau^L, \chi_\tau)$, $\tau \in \mathbb{D}(e^{-1}, 1)$, induced by h_τ^L . We introduce $D_k(\tau) \in \text{End}(H^0(X, L^{\otimes k}))$ through the identity $\frac{\sqrt{-1}}{2\pi} R_{k,\tau} := \sqrt{-1} dz \wedge d\bar{z} \cdot D_k(\tau)$. The following result is the technical backbone of our analysis.

Theorem 4.2 (Ma-Zhang [56, Theorem 0.4]). *There are $C > 0$, $k_0 \in \mathbb{N}$, such that in the notations from Definition 1.3, for any $k \geq k_0$, $\tau \in \mathbb{D}(e^{-1}, 1)$, we have*

$$\left\| D_k(\tau) - kT_{\omega_H(\tau), k} \right\| \leq C, \quad (4.3)$$

where $\|\cdot\|$ is the operator norm subordinate with $\text{Hilb}_k(h_\tau^L, \chi_\tau)$. In particular, $\{\frac{1}{k} D_k(\tau)\}_{k=1}^{+\infty}$ forms a Toeplitz operator with symbol $\omega_H(\tau)$.

We will now establish a refinement of Theorem 3.2 under more restrictive assumptions. We assume that the Mabuchi geodesic h_t^L , $t \in [0, 1]$, between two positive smooth metrics h_i^L , $i = 0, 1$, is smooth (jointly in X and t directions) and h_t^L are (strictly) positive Hermitian metrics. We denote by $T_k(h_0^L, h_1^L) \in \text{End}(H^0(X, L^{\otimes k}))$ the transfer map between $\text{Hilb}_k(h_0^L, \chi_0)$ and $\text{Hilb}_k(h_1^L, \chi_1)$, where χ_0, χ_1 are some Kähler forms on X .

Theorem 4.3. *In the notations of Definition 1.3 and Theorem 3.2, under the described above assumption, there are $C > 0$, $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, we have*

$$\left\| T_k(h_0^L, h_1^L) - kT_{\phi(h_0^L, h_1^L), k} \right\| \leq C. \quad (4.4)$$

Proof. We fix a smooth path of Kähler forms χ_t , $t \in [0, 1]$, on X between χ_0 and χ_1 , and consider the rotationally-invariant Hermitian metric \hat{H}_k^0 on the (trivial) vector bundle $H^0(X, L^{\otimes k}) \times \mathbb{D}(e^{-1}, 1)$ over $\mathbb{D}(e^{-1}, 1)$, constructed from $\text{Hilb}_k(h_t^L, \chi_t)$, $t \in [0, 1]$, as in (4.1). Directly from the fact that smooth Mabuchi geodesics solve the homogeneous Monge-Ampère equation (2.3), by

Theorem 4.2, we deduce that there are $C_0 > 0$, $k_0 \in \mathbb{N}$, such that the curvature R_k of \hat{H}_k^0 satisfies $\|R_k\| \leq C_0$ for any $k \geq k_0$. We denote

$$g : \mathbb{D}(e^{-1}, 1) \rightarrow \mathbb{R}, \quad \tau \mapsto g(\tau) := (2 \log |\tau|^2 - 1)^2 - 1. \quad (4.5)$$

Remark that g is strictly subharmonic and verifies $g(e^{-1+i\theta}) = g(e^{i\theta}) = 0$, for any $\theta \in [0, 2\pi]$. Directly from the bound $\|R_k\| \leq C_0$, and strict subharmonicity of g , there is $C_1 > 0$, such that the curvature of Hermitian metrics $\hat{H}_k^1 = \hat{H}_k^0 \cdot \exp(-C_1 g)$ (resp. $\hat{H}_k^2 = \hat{H}_k^0 \cdot \exp(C_1 g)$) is positive (resp. negative). We denote by $H_{k,t}^1, H_{k,t}^2$, $t \in [0, 1]$, the paths of metrics on $H^0(X, L^{\otimes k})$ induced through (4.1) by \hat{H}_k^1 and \hat{H}_k^2 respectively. Our boundary condition on g implies that $H_{k,0}^1 = \text{Hilb}_k(h_0^L, \chi_0) = H_{k,0}^2$ and $H_{k,1}^1 = \text{Hilb}_k(h_1^L, \chi_1) = H_{k,1}^2$. From this and the above curvature calculation, we deduce by Theorem 4.1 that for any $t \in [0, 1]$, we have

$$H_{k,t}^1 \geq H_{k,t} \geq H_{k,t}^2, \quad (4.6)$$

where $H_{k,t}$ is the geodesic between $\text{Hilb}_k(h_0^L, \chi_0)$ and $\text{Hilb}_k(h_1^L, \chi_1)$. By taking derivatives at $t = 0$ from the above inequality, we deduce that

$$\dot{H}_{k,0}^1 \geq -T_k(h_0^L, h_1^L) \geq \dot{H}_{k,0}^2. \quad (4.7)$$

Directly from the definition of $H_{k,t}^1, H_{k,t}^2$, we deduce that

$$\begin{aligned} \dot{H}_{k,0}^1 &= (\text{Hilb}_k(h_0^L, \chi_0))^{-1} \frac{d}{dt} \text{Hilb}_k(h_t^L, \chi_t)|_{t=0} + C_1 g'(1) \text{Id}, \\ \dot{H}_{k,0}^2 &= (\text{Hilb}_k(h_0^L, \chi_0))^{-1} \frac{d}{dt} \text{Hilb}_k(h_t^L, \chi_t)|_{t=0} - C_1 g'(1) \text{Id}. \end{aligned} \quad (4.8)$$

From the definition of the L^2 -norm, it is direct to see that

$$(\text{Hilb}_k(h_0^L, \chi_0))^{-1} \frac{d}{dt} \text{Hilb}_k(h_t^L, \chi_t)|_{t=0} = -k T_{\phi(h_0^L, h_1^L), k} + T_{\chi_0^{-1} \frac{d}{dt} \chi_t|_{t=0}, k}. \quad (4.9)$$

From (4.7), (4.8) and (4.9), the result follows directly. \square

Now, to drop the additional assumption from Theorem 4.3, we need to regularize the Mabuchi geodesics. Let us recall the approximation scheme, called ϵ -geodesics, introduced by X. Chen [19]. For this, we follow the notations introduced in (2.3). We fix $u_0, u_1 \in \mathcal{H}_\omega$, and for an arbitrary $\epsilon > 0$, consider the Dirichlet problem

$$(\pi^* \omega + \sqrt{-1} \partial \bar{\partial} \hat{u})^{n+1} = \epsilon \pi^* \omega^n \wedge \sqrt{-1} dz \wedge d\bar{z}, \quad (4.10)$$

with boundary conditions $\hat{u}(x, e^{\sqrt{-1}\theta}) = u_0(x)$, $\hat{u}(x, e^{-1+\sqrt{-1}\theta}) = u_1(x)$, $x \in X$, $\theta \in [0, 2\pi]$.

Theorem 4.4 ([19, Lemma 7], [22, §3]). *For any $\epsilon > 0$, there is a unique S^1 -invariant smooth solution \hat{u}_ϵ to (4.10), for which $\pi^* \omega + \sqrt{-1} \partial \bar{\partial} \hat{u}_\epsilon$ is strictly positive. Moreover, as $\epsilon \rightarrow 0$, \hat{u}_ϵ converges to the solution \hat{u} of (2.3) in $\mathcal{C}^1(X \times \mathbb{D}(e^{-1}, 1))$. In particular, as $\epsilon \rightarrow 0$, $\dot{u}_{0,\epsilon} := \frac{d}{dt} u_{t,\epsilon}|_{t=0}$ converges to \dot{u}_0 uniformly.*

We will also need another ingredient from [39, Theorem 4.20], which we recall below.

Theorem 4.5. *We fix continuous psh metrics h_0^L, h_1^L on L , and a smooth path of Kähler forms χ_t , $t \in [0, 1]$, on X . There is a sequence $a_k \in \mathbb{R}$, verifying $a_k/k \rightarrow 0$, as $k \rightarrow \infty$, and $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, $t \in [0, 1]$, the geodesic $H_{k,t}$, $t \in [0, 1]$, between $\text{Hilb}_k(h_0^L, \chi_0)$ and $\text{Hilb}_k(h_1^L, \chi_1)$ is related with the Mabuchi geodesic h_t^L , $t \in [0, 1]$, between h_0^L and h_1^L as follows*

$$\text{Hilb}_k(h_t^L, \chi_t) \cdot \exp(-a_k) \leq H_{k,t} \leq \text{Hilb}_k(h_t^L, \chi_t) \cdot \exp(a_k). \quad (4.11)$$

Remark 4.6. Berndtsson in [9], following a previous work of Phong-Sturm [58], proved the uniform convergence of $FS(H_{k,t})^{\frac{1}{k}}$ to h_t^L , cf. also [31] for a related result. Theorem 4.5 refines this result. Remark, however, that our proof in [39] relies on the results of Berndtsson.

Proof of Theorem 3.2. Let us first reduce our considerations to the case when both endpoints are smooth. We use the notations introduced in Proposition 3.1. Then by Theorem 2.1, we get

$$T_k(h_0^L, h_{1,\epsilon}^L) - \epsilon k \text{Id} \leq T_k(h_0^L, h_1^L) \leq T_k(h_0^L, h_{1,\epsilon}^L) + \epsilon k \text{Id}. \quad (4.12)$$

If we now assume that Theorem 3.2 is valid for smooth endpoints, we deduce that for any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, we have

$$\left\| \frac{1}{k} T_k(h_0^L, h_{1,\epsilon}^L) - T_{\phi(h_0^L, h_{1,\epsilon}^L), k} \right\| \leq \epsilon. \quad (4.13)$$

Directly from (3.1), (4.12) and (4.13), we establish that Theorem 3.2 then holds for continuous endpoint h_1^L . It is, hence, enough to establish Theorem 3.2 when both endpoints are smooth, which we assume from now on.

We denote by $h_{t,\epsilon}^L$, $t \in [0, 1]$, the ϵ -geodesic between h_0^L and h_1^L associated with an auxillary form $\omega := 2\pi c_1(L, h_0^L)$, given by Theorem 4.4. We denote by $\hat{H}_{k,\epsilon}$ the rotationally-invariant Hermitian metric on the (trivial) vector bundle $H^0(X, L^{\otimes k}) \times \mathbb{D}(e^{-1}, 1)$ over $\mathbb{D}(e^{-1}, 1)$, constructed from $\text{Hilb}_k(h_{t,\epsilon}^L)$, $t \in [0, 1]$, as in (4.1). For $\tau \in \mathbb{D}(e^{-1}, 1)$, we define $d_\epsilon(\tau) : X \rightarrow \mathbb{R}$ as follows

$$d_\epsilon(\tau) := \frac{1}{n+1} \frac{c_1(L, h_0^L)^n}{c_1(L, h_{t,\epsilon}^L)^n}, \quad \text{where } t = |\log |\tau||. \quad (4.14)$$

Directly from (4.10), by Theorem 4.2, we deduce that for any $\epsilon > 0$, there are $C > 0$, $k_0 \in \mathbb{N}$, such that the curvature $R_{k,\epsilon}$ of $\hat{H}_{k,\epsilon}$ satisfies

$$\left\| \frac{\sqrt{-1}}{2\pi} R_{k,\epsilon,\tau} - \epsilon k T_{d_\epsilon(\tau), k} \right\| \leq C. \quad (4.15)$$

The major technical problem is that we do not know if $d_\epsilon(\tau)$ stays bounded uniformly in τ , as $\epsilon \rightarrow 0$. In fact, it is even likely that it will not stay bounded as long as the Mabuchi geodesic passes through a non strictly positive Hermitian metric. The absence of this *non-collapsing estimates* makes it unclear if (4.15) implies that $\frac{1}{k} \|R_{k,\epsilon,\tau}\|$ can be made arbitrary small uniformly in $\tau \in \mathbb{D}(e^{-1}, 1)$, for $k \in \mathbb{N}$ big enough and $\epsilon > 0$ small enough. This, however, was crucial in the construction of subinterpolating family of metrics as described in the proof of Theorem 4.3. The rest of the proof will be dedicated to a trick which will overcome this problem.

First, remark that since $d_\epsilon(\tau) > 0$, we deduce from (4.15) that the curvature $R_{k,\epsilon,\tau}$ eventually becomes positive, for k big enough. From Theorem 4.1, we then deduce that

$$\text{Hilb}_k(h_{t,\epsilon}^L) \geq H_{t,k}. \quad (4.16)$$

By proceeding in exactly the same way as we did in the proof of Theorem 4.3, we then see that for any $\epsilon > 0$, there are $k_0 \in \mathbb{N}$, $C > 0$, such that for any $k \geq k_0$, we have

$$kT_{h_{0,\epsilon}^L, k} + C \cdot \text{Id} \geq -T_k(h_0^L, h_1^L). \quad (4.17)$$

We will now show how to modify the argument from Theorem 4.3 to get the lower bound for $T_k(h_0^L, h_1^L)$. We fix a smooth path of Kähler forms χ_t , $t \in [0, 1]$ between χ_0 and χ_1 . By the \mathcal{C}^1 -convergence of ϵ -geodesics to the Mabuchi geodesic from Theorem 4.4, and Theorem 4.5, we conclude that there is a sequence $a_k \in \mathbb{R}$, $k \in \mathbb{N}$, such that $a_k/k \rightarrow 0$, as $k \rightarrow \infty$, and so that for any $\delta > 0$, there are $k_0 \in \mathbb{N}$, $\epsilon_0 > 0$, so that for any $t \in [0, 1]$, $k \geq k_0$, $\epsilon_0 > \epsilon > 0$, we have

$$H_{t,k} \geq \text{Hilb}_k(h_{t,\epsilon}^L, \chi_t) \cdot \exp(-\delta tk - a_k). \quad (4.18)$$

Now, since $h_{t,\epsilon}^L$, $t \in [0, 1]$, is a smooth path, and $d_\epsilon(1) = \frac{1}{n+1}$, there is $t_\epsilon > 0$, such that $d_\epsilon(\tau) \leq 1$ for all $|\tau| = e^{-t}$, $t \in [0, t_\epsilon]$. Then, in the notations (4.5), we deduce that there is $C > 0$, such that for any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that the curvature of the Hermitian vector bundle $\hat{H}_{k,\epsilon}^2 = \hat{H}_{k,\epsilon} \cdot \exp(\epsilon Ckg)$ is negative over $X \times \mathbb{D}(e^{-t_\epsilon}, 1)$ for any $k \geq k_0$. Directly from Theorem 4.1, we deduce that the geodesic $H_{t,k}^0$, $t \in [0, t_\epsilon]$, from $\text{Hilb}_k(h_0^L, \chi_0)$ to $\text{Hilb}_k(h_{t_\epsilon,\epsilon}^L, \chi_{t_\epsilon}) \cdot \exp(\epsilon Ckg(e^{-t_\epsilon}))$, is related with the above path for any $k \geq k_0$, $t \in [0, t_\epsilon]$, as follows

$$H_{t,k}^0 \geq \text{Hilb}_k(h_{t,\epsilon}^L, \chi_t) \cdot \exp(\epsilon Ckg(e^{-t})). \quad (4.19)$$

Remark that we have $H_{0,k} = H_{0,k}^0$, and by the negativity of g , (4.18) and (4.19), we also have $H_{t_\epsilon,k} \geq H_{t_\epsilon,k}^0 \cdot \exp(-t_\epsilon \delta k - a_k)$. From this and Proposition 2.1, we conclude that for any $t \in [0, t_\epsilon]$, $k \geq k_0$, we have

$$H_{t,k} \geq H_{t,k}^0 \cdot \exp(-t\delta k - ta_k/t_\epsilon). \quad (4.20)$$

A combination of (4.19) and (4.20) yields that for any $t \in [0, t_\epsilon]$, $k \geq k_0$, we have

$$H_{t,k} \geq \text{Hilb}_k(h_{t,\epsilon}^L, \chi_t) \cdot \exp(\epsilon Ckg(e^{-t}) - t\delta k - ta_k/t_\epsilon). \quad (4.21)$$

If we then take a derivative at $t = 0$ of (4.21), by (4.9), we immediately get that for any $\delta > 0$, there is $\epsilon_0 > 0$ so that for any $\epsilon_0 > \epsilon > 0$, there are $t_\epsilon > 0$, $k_0 \in \mathbb{N}$, so that for any $k \geq k_0$, we have

$$-T_k(h_0^L, h_1^L) \geq kT_{h_{0,\epsilon}^L, k} - \left(\epsilon Ckg'(1) + \delta k + a_k/t_\epsilon - \inf \left(\chi_0^{-1} \frac{d}{dt} \chi_t \Big|_{t=0} \right) \right) \text{Id}. \quad (4.22)$$

A combination of (4.17) and (4.22) finally yields the result by the uniform convergence of $\hat{h}_{0,\epsilon}^L$ towards $-\phi(h_0^L, h_1^L)$ from Theorem 4.4. \square

5 Functional calculus for Toeplitz operators

The primary objective of this section is to demonstrate that the convergence of operators in various norms leads to the convergence of their diagonal kernels, whereas the converse typically does not hold (unless the sequence of operators forms a Toeplitz operator). As a consequence of our considerations, we show that the vector space of Toeplitz operators of Schatten class form an algebra closed by the functional calculus. Through this, we demonstrate that Theorem 1.1 becomes a consequence of Theorems 1.4, 1.6 if one replaces the pointwise convergence by the convergence in $L^p(X)$ -spaces, for any $p \in [1, +\infty[$.

Throughout the whole section we fix a positive Hermitian metric h^L on L , and a sequence $T_k \in \text{End}(H^0(X, L^{\otimes k}))$ of Hermitian operators (with respect to $\text{Hilb}_k(h^L)$). Recall that $N_k := \dim H^0(X, L^{\otimes k})$. We denote the associated volume form $d\nu := \frac{1}{\int c_1(L)^n} c_1(L, h^L)^n$. We use the notations $\|\cdot\|$, $\|\cdot\|_p$, $p \in [1, +\infty[$, for the norms introduced in Definition 1.3, 1.5.

Let us first discuss the relation between the norm $\|\cdot\|$ and the diagonal kernel.

Proposition 5.1. For any $x \in X$, we have $|T_k(x)| \leq B_k(x) \cdot \|T_k\|$. While this inequality is sharp, already for $(X, L) := (\mathbb{P}^1, \mathcal{O}(1))$, for any $C > 0$, there are $k \in \mathbb{N}$, $T_k \in \text{End}(H^0(X, L^{\otimes k}))$ so that $|T_k(x)| \leq B_k(x)$, for any $x \in X$, but $\|T_k\| \geq C$.

Proof. First of all, by the independence of the expression like in (1.4) on the choice of orthonormal basis, it is immediate that

$$T_k(x) = \langle T_k s_{x,k}, s_{x,k} \rangle_{\text{Hilb}_k(h^L)} \cdot B_k(x), \quad (5.1)$$

where $s_{x,k} \in H^0(X, L^{\otimes k})$ is a peak section at $x \in X$, which is a section of norm 1 with respect to $\text{Hilb}_k(h^L)$, orthogonal to all sections from $H^0(X, L^{\otimes k})$ vanishing at x . This shows the first part of Proposition 5.1 by the trivial inequality $\langle T_k s_{x,k}, s_{x,k} \rangle_{\text{Hilb}_k(h^L)} \leq \|T_k\|$. The sharpness of the established inequality is trivial for $T_k := B_k$.

For the second part of the statement, we assume $(X, L) := (\mathbb{P}^1, \mathcal{O}(1))$ is endowed with the Fubini-Study metric as described in the example in the end of Section 2, from which we also borrow the notation. We consider an operator $T_{2k} := \frac{1}{2}(B_{2k} + \sqrt{k} \cdot P_{2k})$, where P_{2k} is the orthogonal projection from $H^0(X, L^{\otimes 2k})$ to the subspace spanned by $x^k \cdot y^k$ in the notations from (2.12). Clearly, $\|T_{2k}\| = \frac{\sqrt{k+1}}{2}$. Directly from (2.12), we see that

$$T_{2k}([ax^* + by^*]) = \frac{2k+1}{2} + \frac{\sqrt{k}}{2} \left(\frac{|a||b|}{|a|^2 + |b|^2} \right)^{2k} \frac{(2k+1)!}{k!k!}. \quad (5.2)$$

Directly from Stirling's approximation and inequality $|a|^2 + |b|^2 \geq 2|a||b|$, we deduce that for any $x \in \mathbb{P}^1$, $|T_{2k}(x)| \leq 2k+1$, finishing the proof. \square

Let us now discuss the relation between the norms $\|\cdot\|_p$, $p \in [1, +\infty[$, and the diagonal kernel.

Proposition 5.2. For any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, $p \in [1, +\infty[$, we have

$$\sqrt[p]{\int |T_k(x)|^p d\nu(x)} \leq \|T_k\|_p \cdot k^n \cdot (1 + \epsilon). \quad (5.3)$$

While this inequality is sharp, already for $(X, L) := (\mathbb{P}^1, \mathcal{O}(1))$, there are $T_k \in \text{End}(H^0(X, L^{\otimes k}))$, $k \in \mathbb{N}$, so that $\|T_k\| = \|T_k\|_p = 1$, for any $p \in [1, +\infty[$, but for any $\epsilon > 0$, $p \in [1, +\infty[$, there is $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, we have $\int |T_k(x)|^p d\nu(x) \leq \epsilon \cdot k^{np}$.

Proof. Let us first establish the inequality

$$\int |T_k(x)| d\nu(x) \cdot \frac{\int c_1(L)^n}{n!} \leq \|T_k\|_1 \cdot N_k. \quad (5.4)$$

If T_k is positive-definite, (5.4) immediately follows, as we have $\int T_k(x)d\nu(x) \cdot \int c_1(L)^n/n! = \text{Tr}(T_k) = \|T_k\|_1 \cdot N_k$. By decomposing T_k as $T_k := T_k^+ - T_k^-$, where T_k^+, T_k^- is a positive and negative parts of T_k respectively, it is then immediate that (5.4) holds in full generality.

From (2.11) and the asymptotic Riemann-Roch-Hirzebruch theorem, saying that $N_k \sim k^n \cdot \int c_1(L)^n/n!$, we see that (5.4) refines (5.3) for $p = 1$. Moreover, by Proposition 5.1, we see that (5.3) also holds for $p \rightarrow +\infty$. We establish the general case by interpolation.

We consider a map $\pi_k : \text{End}(H^0(X, L^{\otimes k})) \rightarrow L^\infty(X)$, $T_k \mapsto (x \mapsto T_k(x))$. For $p \in [1, +\infty[$, we denote by $\|\pi_k\|_p$ the norm of π_k , viewed as an operator from $(\text{End}(H^0(X, L^{\otimes k})), \|\cdot\|_p)$ to $(L^\infty(X), \|\cdot\|_{L^p(X)})$. Clearly, (5.3) just says that for any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, so that for any $k \geq k_0$, $p \in [1, +\infty[$, we have $\|\pi_k\|_p \leq k^n \cdot (1 + \epsilon)$. Recall that $(\text{End}(H^0(X, L^{\otimes k})), \|\cdot\|_p)$ is a complex interpolation between $(\text{End}(H^0(X, L^{\otimes k})), \|\cdot\|_1)$ and $(\text{End}(H^0(X, L^{\otimes k})), \|\cdot\|)$, cf. [45, Theorem 13.1]. Similarly, $(L^\infty(X), \|\cdot\|_{L^p(X)})$ is a complex interpolation between $(L^\infty(X), \|\cdot\|_{L^1(X)})$ and $(L^\infty(X), \|\cdot\|_{L^\infty(X)})$, cf. [3, Theorem 5.1.1]. The result now follows directly from the fact that complex interpolation is an example of an *exact interpolation functor*, see [3, Theorem 4.1.2] for the proof of this result and [3, (6) on p.27] for the necessary definitions. The sharpness of (5.3) is again easily seen for $T_k := B_k$ by (2.11).

For the second part of the statement, we assume $(X, L) := (\mathbb{P}^1, \mathcal{O}(1))$ is endowed with the Fubini-Study metric as described in the example in the end of Section 2, from which we also borrow the notation. We consider an operator T_k which is diagonal in the monomial basis, and which sends $x^i \cdot y^j$ to $(-1)^i x^i \cdot y^j$ in the notations (2.12). Clearly, we have $\|T_k\| = \|T_k\|_p = 1$ for any $p \in [1, +\infty[$, $k \in \mathbb{N}$. Directly from (2.12) and binomial identity, we see that

$$T_k([ax^* + by^*]) = (k + 1) \cdot \left(\frac{|b|^2 - |a|^2}{|a|^2 + |b|^2} \right)^k. \tag{5.5}$$

It is immediate to see that for any $p \in [1, +\infty[$, we have $\lim_{k \rightarrow \infty} \int \left| \frac{1-|z|^2}{1+|z|^2} \right|^{pk} \frac{\sqrt{-1}dzd\bar{z}}{1+|z|^2} = 0$, which implies that T_k verifies the assumptions from Proposition 5.2. □

Let us now discuss the diagonal kernels for Toeplitz operators and show that the subtleties pointed out in the second parts of Propositions 5.1, 5.2 disappear for these operators. More precisely, we will establish the following two results.

Theorem 5.3. *For any Toeplitz operator (resp. of Schatten class) $\{T_k\}_{k=0}^{+\infty}$ with symbol $f \in \mathcal{C}^0(X)$ (resp. $f \in L^\infty(X)$), the sequence of functions $x \mapsto \frac{1}{k^n} T_k(x)$, $x \in X$, is uniformly bounded and converges uniformly (resp. in $L^p(X)$ for every $p \in [1, +\infty[$) to f .*

Theorem 5.4. *For any Toeplitz operator of Schatten class $\{T_k\}_{k=0}^{+\infty}$, with symbol $f \in L^\infty(X)$, for any $p \in [1, +\infty[$, we have*

$$\lim_{k \rightarrow \infty} \|T_k\|_p = \sqrt[p]{\int |f(x)|^p d\nu(x)} = \lim_{k \rightarrow \infty} \frac{1}{k^n} \sqrt[p]{\int |T_k(x)|^p d\nu(x)}. \tag{5.6}$$

Similarly, for any Toeplitz operator $\{T_k\}_{k=0}^{+\infty}$, with symbol f , we have $\lim_{k \rightarrow \infty} \|T_k\| = \sup|f(x)| = \lim_{k \rightarrow \infty} \frac{1}{k^n} \sup|T_k(x)|$.

In order to prove these statements, we will need several auxiliary results.

Lemma 5.5. For any $p \in [1, +\infty[$, $f \in L^p(X)$, the sequence of functions $k \in \mathbb{N}$, $x \mapsto \frac{1}{k^n} T_{f,k}(x)$, $x \in X$, converges to f in $L^p(X)$, as $k \rightarrow \infty$. If, moreover, f is continuous, then the convergence is uniform.

Remark 5.6. Ma-Marinescu in [52, Theorem 0.1] established the analogous result in realms of Toeplitz operators associated with smooth symbols f . For continuous f , the result was previously established [1, Theorem 3.3].

Proof. By Remark 5.6, it suffices to show the $L^p(X)$ -convergence. Remark that $T_{f,k}(x)$ can be expressed in terms of the Bergman kernel $B_k(x, y) \in L_x^{\otimes k} \otimes (L_y^{\otimes k})^*$, $x, y \in X$, as follows

$$T_{f,k}(x) = \int f(y) |B_k(x, y)|^2 d\nu(y) \cdot \frac{\int c_1(L)^n}{n!}. \quad (5.7)$$

Now, for $x \in X$, we denote by $\exp_x : T_x X \rightarrow X$ the geodesic coordinates, considered with respect to the Kähler form $c_1(L, h^L)$. The main results from Dai-Liu-Ma [25] and Ma-Marinescu [53], cf. [51, Theorem 4.2.1], imply that there are $c, C, \epsilon > 0$, $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, $x \in X$, $Z \in T_x X$, $|Z| < \epsilon$, we have

$$\begin{aligned} \left| |B_k(x, \exp_x(Z))| - k^n \exp\left(-\frac{\pi}{2}k|Z|^2\right) \right| &\leq Ck^{n-\frac{1}{2}} \cdot (\sqrt{k}|Z|)^{2n+1} \exp(-c\sqrt{k}|Z|), \\ |B_k(x, y)| &\leq Ck^n \exp(-c\sqrt{k}\text{dist}(x, y)). \end{aligned} \quad (5.8)$$

Remark that for any $c > 0$, there is $C_0 > 0$, such that for any $k \in \mathbb{N}^*$, we have

$$\int_{Z \in \mathbb{C}^n} k^n \cdot (\sqrt{k}|Z|)^{2n+1} \exp(-c\sqrt{k}|Z|) dZ \leq C_0. \quad (5.9)$$

In particular, if we define the function $G_{f,k} : X \rightarrow \mathbb{R}$ as follows

$$G_{f,k}(x) = k^n \int_{Z \in T_x X} f(\exp_x(Z)) \exp\left(-\pi k|Z|^2\right) dZ, \quad (5.10)$$

then directly from Generalized Young's Inequality, cf. [44, (0.10)], (5.8) and (5.9), we deduce that there is $C_1 > 0$, such that for any $k \geq k_0$, $f \in L^p(X)$, we have

$$\left(\int \left| \frac{1}{k^n} T_{f,k}(x) - G_{f,k}(x) \right|^p d\nu(x) \right)^{\frac{1}{p}} \leq \frac{C_1}{\sqrt{k}} \left(\int |f(x)|^p d\nu(x) \right)^{\frac{1}{p}}. \quad (5.11)$$

Remark, however, that by the usual properties of convolutions, cf. [44, Theorem (0.13)], the sequence of functions $x \mapsto G_{f,k}(x)$ converges to f in $L^p(X)$, as $k \rightarrow \infty$. The result now follows from this and (5.11). \square

Lemma 5.7. For any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that for any $f \in L^1(X)$, $k \geq k_0$, we have

$$\|T_{f,k}\|_1 \leq (1 + \epsilon) \int |f(x)| d\nu(x). \quad (5.12)$$

For any $f \in L^\infty(X)$, $k \geq k_0$, we have $\|T_{f,k}\| \leq \text{esssup}_{x \in X} |f(x)|$, for any $k \in \mathbb{N}$.

Proof. Remark first that if f is positive, then $\|T_{f,k}\|_1 = \frac{1}{N_k} \int T_{f,k}(x) d\nu(x) \cdot \int c_1(L)^n/n!$. As a direct consequence of (5.7) and Tonelli's theorem, we deduce

$$\int T_{f,k}(x) d\nu(x) = \int f(y) |B_k(x, y)|^2 d\nu(x) d\nu(y) \cdot \frac{\int c_1(L)^n}{n!}. \quad (5.13)$$

Remark, however, that $\int |B_k(x, y)|^2 d\nu(y) \cdot \frac{\int c_1(L)^n}{n!} = B_k(x)$, and so we have

$$\int |T_{f,k}(x)| d\nu(x) \leq \int |f(x)| B_k(x) d\nu(x). \quad (5.14)$$

The first part of Lemma 5.7 for positive f now follows from (2.11), (5.14) and the asymptotic Riemann-Roch-Hirzebruch theorem. For general f , The first part of Lemma 5.7 is established by decomposing f as a difference of a positive and a negative part and applying Lemma 5.7 for each of them. The second part follows directly from the trivial bounds $\text{essinf}_{x \in X} f(x) \cdot \text{Id} \leq T_{f,k} \leq \text{esssup}_{x \in X} f(x) \cdot \text{Id}$. \square

Proof of Proposition 3.5. From the uniform bound assumption, the second part of Lemma 5.7 and (3.18), we see that it suffices to establish that for any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, we have

$$\|T_{f_k,k} - T_{f,k}\|_1 \leq \epsilon. \quad (5.15)$$

This follows directly from the first part of Lemma 5.7 and the assumption on the $L^1(X)$ -convergence of f_k towards f . \square

Proof of Theorem 5.3. It follows immediately from Propositions 5.1, 5.2 and Lemma 5.7. \square

We will now study the functional-analytic and algebraic properties of the vector space of Toeplitz operators (resp. of Schatten class).

Proposition 5.8. For any continuous $f, g : X \rightarrow \mathbb{R}$ (resp. $f, g \in L^\infty(X)$) the sequence of operators $T_{f,k} \circ T_{g,k}$ is a Toeplitz operator (resp. of Schatten class) with symbol fg .

Proof. When both f, g are smooth, the result is well known, see [12], [51, §7]. The general case follows easily by approximation and Lemma 5.7. For brevity, we only give the main idea. We fix $C := \text{esssup} \max(|f|, |g|)$ and for any $\epsilon > 0$, consider smooth f_ϵ, g_ϵ , verifying $\sup \max(|f_\epsilon|, |g_\epsilon|) \leq C + 1$ and $\int |f(x) - f_\epsilon(x)| d\eta(x) < \epsilon \int d\eta(x)$, $\int |g(x) - g_\epsilon(x)| d\eta(x) < \epsilon \int d\eta(x)$, the existence of which follows from the usual density statements. It is then a direct that for any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that for any $k \geq k_0$, we have

$$\left\| T_{f,k} \circ T_{g,k} - T_{f_\epsilon,k} \circ T_{g_\epsilon,k} \right\| \leq 4\epsilon(C + 1), \quad \left\| T_{f_\epsilon g_\epsilon,k} - T_{fg,k} \right\| \leq 4\epsilon(C + 1). \quad (5.16)$$

From Lemma 5.7, (5.16) and the validity of Proposition 5.8 for f_ϵ, g_ϵ , we conclude that for any $\epsilon > 0$, there is $k_1 \in \mathbb{N}$, such that for any $k \geq k_1$, we have

$$\left\| T_{f_\epsilon,k} \circ T_{g_\epsilon,k} - T_{f_\epsilon g_\epsilon,k} \right\| \leq 5\epsilon(C + 1). \quad (5.17)$$

A combination of (5.16) and (5.17) yields the proof. \square

Corollary 5.9. The vector space of Toeplitz operators (resp. of Schatten class) forms an algebra, and the symbol map is a algebra morphism.

Proof. Follows directly from Proposition 5.8. \square

Proposition 5.10. For any continuous functions $g : \mathbb{R} \rightarrow \mathbb{R}$, $f : X \rightarrow \mathbb{R}$ (resp. $f \in L^\infty(X)$), the sequence of operators $\{g(T_{f,k})\}_{k=0}^{+\infty}$ form a Toeplitz operator (resp. of Schatten class) with symbol $g(f)$.

Proof. For polynomials g , the result follows from Proposition 5.8. The general case is done by approximation. The details are left to the reader. \square

Corollary 5.11. The vector space of Toeplitz operators (resp. of Schatten class) is closed under functional calculus associated with continuous functions.

Proof. Follows directly from Proposition 5.10. \square

Proof of Theorem 1.1 for $L^p(X)$ -convergence in place of pointwise convergence. By Proposition 5.10, Theorem 1.4 directly implies the second part of Theorem 1.1. Similarly, Proposition 5.10 shows that Theorem 1.6 implies the first part of Theorem 1.1 when pointwise convergence is replaced by the convergence in $L^p(X)$ -spaces for any $p \in [1, +\infty[$. \square

Proof of Theorem 5.4. Let us first establish the first part. The second identity follows from Theorem 5.3. To establish the first identity, we deduce from Proposition 5.10 that $|T_k|^p$, $k \in \mathbb{N}$, is a Toeplitz operator of Schatten class with symbol $|f|^p$. Then for any $\epsilon > 0$, there is $k_0 \in \mathbb{N}$, such that we have

$$\left| \text{Tr} [|T_k|^p] - \text{Tr} [T_{|f|^p,k}] \right| \leq \epsilon N_k. \quad (5.18)$$

However, $\text{Tr} [T_{|f|^p,k}] = \int T_{|f|^p,k}(x) d\nu(x) \cdot \int c_1(L)^n / n!$, and the first identity again follows from Proposition 5.10. The second part of the statement is well known and follows directly from Lemma 5.7 and Theorem 5.3. \square

As a conclusion, we point out that the analogue of the spectral convergence result of Boutet de Monvel-Guillemin [48], also holds in the more general context of Toeplitz operators of Schatten class. More specifically, let us establish the following result.

Proposition 5.12. For any Toeplitz operator of Schatten class $\{T_k\}_{k=0}^{+\infty}$, with symbol $f \in L^\infty(X)$, the spectral measures of T_k converge weakly to the probability measure $f_*(\nu)$ on \mathbb{R} .

Proof. It suffices to prove that for any continuous $g : \mathbb{R} \rightarrow \mathbb{R}$, we have the following convergence

$$\lim_{k \rightarrow \infty} \frac{1}{N_k} \sum_{\lambda \in \text{Spec}(T_k)} g(\lambda) = \int_{x \in \mathbb{R}} g(f(x)) d\nu(x). \quad (5.19)$$

However, we have $\sum_{\lambda \in \text{Spec}(T_k)} g(\lambda) = \text{Tr}[g(T_k)]$, and the result follows directly from Theorem 5.3 and Corollary 5.11. \square

6 Superadditivity of weighted Bergman kernels, a proof of Theorem 1.1

The main goal of this section is to complete the proof of Theorem 1.1 by establishing the pointwise convergence. To clarify our approach, let $x \mapsto B_k^{\mathcal{F}}(x)$ denote the weighted Bergman kernel, as defined in (1.4) for the function $g(x) := x$. The main result of this section, stated below, demonstrates the partial superadditivity of weighted Bergman kernels.

Theorem 6.1. *There are $k_0 \in \mathbb{N}$, $C > 0$, so that for any $k, l \geq k_0$, $l/2 \leq k \leq 2l$, $x \in X$, we have*

$$\frac{B_{k+l}^{\mathcal{F}}(x)}{(k+l)^{n-1}} \geq \frac{B_k^{\mathcal{F}}(x)}{k^{n-1}} + \frac{B_l^{\mathcal{F}}(x)}{l^{n-1}} - C \log(k+l). \quad (6.1)$$

The proof of Theorem 6.1 relies heavily on the multiplicativity properties of L^2 -metrics established by the author in [39], and it will be presented in the end of the section. Before that, let us see how it implies Theorem 1.1.

Proof of Theorem 1.1. By the results of Section 5, it suffices to establish the pointwise convergence of $x \mapsto \frac{1}{k^n} B_k^{\mathcal{F}}(x)$, $x \in X$, as $k \rightarrow \infty$. We will establish it by reducing to the case $g(x) := x$ and using Theorem 6.1.

First, recall that the classical Fekete's Subadditive Lemma says that for any sequence $a_k \in \mathbb{R}$, which is subadditive, i.e. for any $k, l \in \mathbb{N}$, we have $a_{k+l} \leq a_k + a_l$, the limit of $\frac{1}{k} a_k$ exists in $\mathbb{R} \cup \{-\infty\}$. De Bruijn-Erdős further generalized this result in [33, Theorem 23] to sequences, verifying $a_{k+l} \leq a_k + a_l + g_{k+l}$, for any $k, l \in \mathbb{N}$, verifying $l/2 \leq k \leq 2l$, and an auxiliary sequence $g_k \in \mathbb{R}$, so that $\sum \frac{g_k}{k^2} < +\infty$.

Remark that $\sum \frac{\log(k)}{k^2} < +\infty$, and so the result of de Bruijn-Erdős along with Theorem 6.1 imply the pointwise convergence of $x \mapsto \frac{1}{k^n} B_k^{\mathcal{F}}(x)$, $x \in X$, as $k \rightarrow \infty$. The boundness condition on \mathcal{F} implies that the limit is bounded. This establishes Theorem 1.1 for $g(x) := x$.

Let us show that it automatically implies Theorem 1.1 for $g(x) := g_c(x) := \min(x, c)$, for any $c \in \mathbb{R}$. In order to see this, for an arbitrary filtration \mathcal{F} , we define the filtration \mathcal{F}^c through its weight function as $w_{\mathcal{F}^c}(s) := \min(w_{\mathcal{F}}(s), ck)$, $s \in H^0(X, L^{\otimes k})$, $k \in \mathbb{N}$. It is easy to see that \mathcal{F}^c is submultiplicative and bounded whenever \mathcal{F} is. Moreover, it is direct that in the notations of (1.4), we have $g_c(A(\mathcal{F}_k, \text{Hilb}_k(h^L))/k) = A(\mathcal{F}_k^c, \text{Hilb}_k(h^L))/k$. This in particular implies that $B_k^{\mathcal{F}^c}(x) = B_k^{\mathcal{F}, g_c}(x)$. And consequently the already established pointwise convergence result for \mathcal{F}^c , $g(x) := x$, implies the pointwise convergence for \mathcal{F} , $g(x) := g_c(x)$.

We claim that the real vector space spanned by the functions g_c , $c \in \mathbb{R}$, is dense within the set of continuous functions (with respect to the uniform topology over a fixed compact interval in \mathbb{R}). This would complete the proof of Theorem 1.1, as it is straightforward to see that it suffices to verify the theorem's statement for a subset of continuous functions that is dense in the uniform topology among continuous functions with compact support in the interval $[-C, C]$, where $C > 0$ is such that $\max_{s \in H^0(X, L^{\otimes k}) \setminus \{0\}} |w_{\mathcal{F}}(s)| \leq Ck$, for any $k \in \mathbb{N}$.

To see the density of g_c , $c \in \mathbb{R}$, it suffices to establish that an arbitrary smooth function $g : \mathbb{R} \rightarrow \mathbb{R}$ of compact support lies in the closure of this vector space. But this follows directly from the formula $g(x) = \int_{-\infty}^{+\infty} (x+t - g_t(x)) g''(t) dt$, the verification of which is left to the reader. \square

Remark 6.2. Our choice of a dense subset of functions was inspired by Chen-MacLean [18, Proposition 5.1], who used such set in a related context.

From now on, we concentrate on the proof of Theorem 6.1. Recall that a Hermitian norm $H_V = \|\cdot\|_V$ on a finitely dimensional vector space V naturally induces the Hermitian norm $\|\cdot\|_Q$, which we also denote by $[H_V]$, on any quotient $Q, \pi : V \rightarrow Q$ of V through the following identity

$$\|f\|_Q := \inf \{ \|g\|_V; \quad g \in V, \pi(g) = f \}, \quad f \in Q. \quad (6.2)$$

A similar construction associates for an arbitrary filtration \mathcal{F} on V the induced quotient filtration $[\mathcal{F}]$ on Q . Similarly, one can naturally identify V with $Q \oplus \ker \pi$ using the dual to π map $\pi^* : Q \rightarrow V$. Using this identification, for any $A \in \text{End}(V)$, we then can define the operator $A|_Q \in \text{End}(Q)$ by $A|_Q(q) = \pi(A(\pi^*(q)))$.

Proposition 6.3 (cf. [41, Proposition 4.12]). We denote by $H_t^V, t \in [0, +\infty[$, the geodesic ray of Hermitian metrics on V associated with \mathcal{F} , departing from H^V . Similarly, we let $[H^V]_t, t \in [0, +\infty[$, be the geodesic ray of Hermitian metrics on Q associated with $[\mathcal{F}]$, departing from $[H^V]$. Then for any $t \in [0, +\infty[$, we have

$$[H_t^V] \geq [H^V]_t. \quad (6.3)$$

In particular, by taking the derivative at $t = 0$, we obtain

$$A(H^V, \mathcal{F})|_Q \leq A([H^V], [\mathcal{F}]). \quad (6.4)$$

The above result will be applied in the setting of a section ring. More specifically, for any $k, l \in \mathbb{N}$, we denote the multiplication map

$$\text{Mult}_{k,l} : H^0(X, L^{\otimes k}) \otimes H^0(X, L^{\otimes l}) \rightarrow H^0(X, L^{\otimes(k+l)}), \quad s_1 \otimes s_2 \mapsto s_1 \cdot s_2. \quad (6.5)$$

It is classical that there is $k_0 \in \mathbb{N}$, such that $\text{Mult}_{k,l}$ is surjective for any $k, l \geq k_0$. The following statement shows that L^2 -norms respect the algebraic structure of $R(X, L)$ in a certain sense.

Theorem 6.4 ([39, Theorem 1.1]). *There are $C > 0, k_1 \in \mathbb{N}^*$, such that for any $k, l \geq k_1$, for the norms over $H^0(X, L^{k+l})$, the following relation holds*

$$1 - C \left(\frac{1}{k} + \frac{1}{l} \right) \leq \frac{[\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]}{\text{Hilb}_{k+l}(h^L)} \cdot \left(\frac{k \cdot l}{k+l} \right)^{\frac{n}{2}} \leq 1 + C \left(\frac{1}{k} + \frac{1}{l} \right), \quad (6.6)$$

Where the quotient norm $[\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]$ is constructed using (6.5).

Another preliminary result we shall use concerns the stability estimates for the weight operator, which compares the weight operators of a given filtration for two different Hermitian metrics.

Theorem 6.5. *Assume that for a constant $C > 0$, verifying $(1 + 2\lceil \log_2 \dim V \rceil)^2 C < 1$, the Hermitian products H_0, H_1 on V satisfy the bound*

$$1 - C \leq \frac{H_1}{H_0} \leq 1 + C. \quad (6.7)$$

Then for any filtration \mathcal{F} on V , the following bound is satisfied

$$\left\| A(\mathcal{F}, H_0) - A(\mathcal{F}, H_1) \right\| \leq 16C(1 + 2\lceil \log_2 \dim V \rceil) \|\mathcal{F}\|, \quad (6.8)$$

where $\|\cdot\|$ is the operator norm subordinate with H_0 , and $\|\mathcal{F}\| := \sup_{v \in V \setminus \{0\}} |w_{\mathcal{F}}(v)|$.

Proof. Denote $r := \dim V$ and fix a basis f_1, \dots, f_r of V , adapted to \mathcal{F} and H_0 . Let T be a transfer map between H_0 and H_1 , which we view as an $r \times r$ matrix using the above basis. Consider the *Cholesky decomposition* of $\exp(-T)$, i.e. let L be the lower triangular matrix, verifying $\exp(-T) = LL^*$. We claim that

$$A(\mathcal{F}, H_1) = L^{-1}A(\mathcal{F}, H_0)L^*. \quad (6.9)$$

Clearly, (6.9) is equivalent to the statement that $(L^*)^{-1}f_1, \dots, (L^*)^{-1}f_r$ is a basis adapted to \mathcal{F} and H_0 . In order to show this, it suffices to verify that $(L^*)^{-1}f_i \in \mathcal{F}^{w_{\mathcal{F}}(f_i)}$ for $i = 1, \dots, r$, and that $(L^*)^{-1}f_i$ form an orthonormal basis with respect to H_1 . The first claim follows immediately from the fact that $(L^*)^{-1}$ is an upper triangular matrix with positive diagonal entries (the last fact follows from the fact that $\exp(-T)$ is positive definite, and so has positive diagonal entries). The second follows immediately from the definition of L and the transfer map.

Now, recall that Cholesky decomposition is stable with respect to perturbations. More precisely, we denote $B := \exp(-T) - \text{Id}$. Define the matrix $P(B)$ through its entries for $i, j = 1, \dots, r$, as follows $P(B)_{ij} = 0$ if $i > j$, $P(B)_{ij} = \frac{1}{2}B_{ij}$ if $i = j$, $P(B)_{ij} = B_{ij}$ if $i < j$. Recall that in [38, Theorem 2.1 and (25)] authors prove that

$$\|\text{Id} - L\| \leq 2\|P(B)\|, \quad \text{if } 2(1 + 2\lceil \log_2 \dim V \rceil)\|P(B)\| < 1. \quad (6.10)$$

Moreover, in [38, (25)], it is established that

$$\|P(B)\| \leq (1/2 + \lceil \log_2 \dim V \rceil)\|B\|. \quad (6.11)$$

Now, (6.7) can be restated as $\|B\| \leq C$, where $\|\cdot\|$ is the operator norm subordinate with H_0 . This with (6.11) implies that $2(1 + 2\lceil \log_2 \dim V \rceil)\|P(B)\| < 1$ under the stated assumption on C . The result now follows directly from (6.9), (6.10) and the trivial bound $\|A(\mathcal{F}, H_0)\| \leq \|\mathcal{F}\|$. \square

Proof of Theorem 6.1. Recall that in (5.1), we defined the peak sections, $s_{k,x} \in H^0(X, L^{\otimes k})$ up to a multiplication by a unimodular constant. We fix this constant in a compatible way, i.e. so that for any $x \in X$, there is $l \in L_x$, so that for any $k \in \mathbb{N}$, we have $s_{k,x}(x) = c_k l^{\otimes k}$, for some $c_k > 0$. By (2.11) and (5.1), it suffices to establish that there are $k_0 \in \mathbb{N}$, $C > 0$, so that for any $k, l \geq k_0$, $l/2 \leq k \leq 2l$, $x \in X$, we have

$$\begin{aligned} \langle A(\mathcal{F}_{k+l}, \text{Hilb}_{k+l}(h^L))_{s_{x,k+l}, s_{x,k+l}} \rangle_{\text{Hilb}_{k+l}(h^L)} &\geq \langle A(\mathcal{F}_k, \text{Hilb}_k(h^L))_{s_{x,k}, s_{x,k}} \rangle_{\text{Hilb}_k(h^L)} \\ &+ \langle A(\mathcal{F}_l, \text{Hilb}_l(h^L))_{s_{x,l}, s_{x,l}} \rangle_{\text{Hilb}_l(h^L)} - C \log(k+l). \end{aligned} \quad (6.12)$$

From now on, we concentrate on the proof of (6.12).

We denote by $\text{Mult}_{k,l}^*$ the dual of the map (6.5), for which the domain is endowed with the norm $\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)$ and the codomain is endowed with the quotient norm $[\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]$. We claim that

$$\text{Mult}_{k,l}^*(s_{x,k} \cdot s_{x,l}) = s_{x,k} \otimes s_{x,l}. \quad (6.13)$$

To see this, we denote by $\mathcal{J}_x \subset \mathcal{O}_X$ the ideal sheaf of germs of holomorphic functions vanishing at x . It is immediate to see that $\text{Mult}_{k,l}^*(s_{x,k} \cdot s_{x,l})$ has to be the element $h \in H^0(X, L^{\otimes k}) \otimes H^0(X, L^{\otimes l})$ minimizing the norm $\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)$, and such that $\text{Mult}_{k,l}(h) = s_{x,k} \cdot s_{x,l}$. We express h in the basis $s_{x,k} \otimes s_{x,l}, s_{x,k} \otimes s_{x,j,l}, s_{x,i,k} \otimes s_{x,l}, s_{x,i,k} \otimes s_{x,j,l}$, where $s_{x,i,k}$ (resp. $s_{x,j,l}$), $i = 2, \dots, N_k$,

(resp. $j = 2, \dots, N_l$) form the bases for $H^0(X, L^{\otimes k} \otimes \mathcal{J}_x)$ (resp. $H^0(X, L^{\otimes l} \otimes \mathcal{J}_x)$). By comparing the value at x , we see that the coefficient of $s_{x,k} \otimes s_{x,l}$ has to be equal to 1. Also, due to minimality of the norm, all the other components have to vanish. This establishes (6.13).

We also claim that there are $k_0 \in \mathbb{N}$, $C > 0$, so that for any $k, l \geq k_0$, $x \in X$, we have

$$\left\| s_{x,k} \cdot s_{x,l} - \left(\frac{k \cdot l}{k+l} \right)^{\frac{n}{2}} s_{x,k+l} \right\|_{\text{Hilb}_{k+l}(h^L)} \leq C \left(\frac{k \cdot l}{k+l} \right)^{\frac{n}{2}} \cdot \left(\frac{1}{k} + \frac{1}{l} \right). \quad (6.14)$$

To see this, remark that $s_{x,k}(y) \cdot \sqrt{B_k(x)} = B_k(y, x) \cdot l^{\otimes k}$, where the dot stands for the pairing between L_x^* and L_x . Taking this into account, (6.14) follows directly from the estimates of Dai-Liu-Ma [25] and Ma-Marinescu [53], cf. (5.8), using the bound (5.9).

From Proposition 6.3, we see that for $k_0 \in \mathbb{N}$, described after (6.5), for any $k, l \geq k_0$, we have

$$\begin{aligned} A([\mathcal{F}_k \otimes \mathcal{F}_l], [\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]) \\ \geq \left(A(\mathcal{F}_k, \text{Hilb}_k(h^L)) \otimes \text{Id} + \text{Id} \otimes A(\mathcal{F}_l, \text{Hilb}_l(h^L)) \right) \Big|_{H^0(X, L^{\otimes(k+l)})}. \end{aligned} \quad (6.15)$$

We also remark that by the definition of submultiplicativity and Proposition 2.1, we have

$$A(\mathcal{F}_{k+l}, [\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]) \geq A([\mathcal{F}_k \otimes \mathcal{F}_l], [\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]). \quad (6.16)$$

If we evaluate (6.15) at $s_{x,k} \cdot s_{x,l}$ and then use (6.13) along with (6.16), and the fact that both $s_{x,k}$ and $s_{x,l}$ are of unit norm, we get

$$\begin{aligned} \left\langle A(\mathcal{F}_{k+l}, [\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)])(s_{x,k} \cdot s_{x,l}), (s_{x,k} \cdot s_{x,l}) \right\rangle_{[\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]} \\ \geq \left\langle A(\mathcal{F}_k, \text{Hilb}_k(h^L))s_{x,k}, s_{x,k} \right\rangle_{\text{Hilb}_k(h^L)} + \left\langle A(\mathcal{F}_l, \text{Hilb}_l(h^L))s_{x,l}, s_{x,l} \right\rangle_{\text{Hilb}_l(h^L)}. \end{aligned} \quad (6.17)$$

Remark also that in the notations of Theorem 6.5, for an arbitrary $C > 0$, we have $A(\mathcal{F}, H_0) = A(\mathcal{F}, C \cdot H_0)$. Also, by the asymptotic Riemann-Roch-Hirzebruch theorem, for any $\epsilon > 0$, there is $k_2 \in \mathbb{N}$, such that for any $k, l \geq k_2$, $l/2 \leq k \leq 2l$, we have $2(1 + \log N_{k+l}) \left(\frac{1}{k} + \frac{1}{l} \right) \leq \epsilon$. In particular, by Theorem 6.4, the result of Theorem 6.5 applies to $H_0 := [\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)] \cdot \left(\frac{k \cdot l}{k+l} \right)^{n/2}$ and $H_1 := \text{Hilb}_{k+l}(h^L)$ for $k, l \geq k_2$, $l/2 \leq k \leq 2l$. As a conclusion, there are $C > 0$, $k_3 \in \mathbb{N}$, such that for any $k, l \geq k_3$, we have

$$\left\| A(\mathcal{F}_{k+l}, [\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]) - A(\mathcal{F}_{k+l}, \text{Hilb}_{k+l}(h^L)) \right\| \leq C \log(k+l), \quad (6.18)$$

where $\| \cdot \|$ is for the operator norm either with respect to $[\text{Hilb}_k(h^L) \otimes \text{Hilb}_l(h^L)]$ or $\text{Hilb}_{k+l}(h^L)$.

Directly from Theorem 6.4, (6.17) and (6.18), we deduce that there are $k_4 \in \mathbb{N}$, $C > 0$, so that for any $k, l \geq k_4$, $l/2 \leq k \leq 2l$, $x \in X$, we have

$$\begin{aligned} \left\langle A(\mathcal{F}_{k+l}, \text{Hilb}_{k+l}(h^L))(s_{x,k} \cdot s_{x,l}), (s_{x,k} \cdot s_{x,l}) \right\rangle_{\text{Hilb}_{k+l}(h^L)} \cdot \left(\frac{k+l}{k \cdot l} \right)^n \\ \geq \left\langle A(\mathcal{F}_k, \text{Hilb}_k(h^L))s_{x,k}, s_{x,k} \right\rangle_{\text{Hilb}_k(h^L)} \\ + \left\langle A(\mathcal{F}_l, \text{Hilb}_l(h^L))s_{x,l}, s_{x,l} \right\rangle_{\text{Hilb}_l(h^L)} - C \log(k+l). \end{aligned} \quad (6.19)$$

The statement (6.12) now follows directly from (6.14) and (6.19). \square

References

- [1] T. Barron, X. Ma, G. Marinescu, and M. Pinsonnault. Semi-classical properties of Berezin-Toeplitz operators with C^k -symbol. *J. Math. Phys.*, 55(4):042108, 25, 2014.
- [2] E. Bedford and B. A. Taylor. A new capacity for plurisubharmonic functions. *Acta Math.*, 149:1–40, 1982.
- [3] J. Bergh and J. Löfström. *Interpolation spaces. An introduction*, volume 223 of *Grundlehren Math. Wiss.* 1976.
- [4] R. Berman. Bergman kernels and equilibrium measures for ample line bundles, ArXiv: 0704.1640. 2007.
- [5] R. Berman. Bergman kernels and equilibrium measures for line bundles over projective manifolds. *Am. J. Math.*, 131(5):1485–1524, 2009.
- [6] R. Berman, S. Boucksom, and M. Jonsson. A variational approach to the Yau-Tian-Donaldson conjecture. *J. Am. Math. Soc.*, 34(3):605–652, 2021.
- [7] R. Berman, S. Boucksom, and D. Witt Nyström. Fekete points and convergence towards equilibrium measures on complex manifolds. *Acta Math.*, 207(1):1–27, 2011.
- [8] B. Berndtsson. A Brunn-Minkowski type inequality for Fano manifolds and some uniqueness theorems in Kähler geometry. *Invent. Math.*, 200(1):149–200, 2015.
- [9] B. Berndtsson. Probability measures associated to geodesics in the space of Kähler metrics. In *Algebraic and analytic microlocal analysis. AAMA, Evanston, Illinois, USA, May 14–26, 2012 and May 20–24, 2013. Contributions of the workshops*, pages 395–419. Cham: Springer, 2018.
- [10] R. Bhatia. *Positive definite matrices*. Princeton Ser. Appl. Math. Princeton, NJ: Princeton University Press, 2007.
- [11] Z. Błocki. On geodesics in the space of Kähler metrics. In *Advances in geometric analysis. Collected papers of the workshop on geometry in honour of Shing-Tung Yau's 60th birthday, Warsaw, Poland, April 6–8, 2009*, pages 3–19. Somerville, MA: International Press; Beijing: Higher Education Press, 2012.
- [12] M. Bordemann, E. Meinrenken, and M. Schlichenmaier. Toeplitz quantization of Kähler manifolds and $gl(N)$, $N \rightarrow \infty$ limits. *Comm. Math. Phys.*, 165:281–296, 1994.
- [13] T. Bouche. Sur les inégalités de Morse holomorphes lorsque la courbure du fibré en droites est dégénérée. *Ann. Sc. Norm. Super. Pisa*, 18(4):501–523, 1991.
- [14] S. Boucksom and H. Chen. Okounkov bodies of filtered linear series. *Compos. Math.*, 147(4):1205–1229, 2011.
- [15] S. Boucksom and M. Jonsson. A non-Archimedean approach to K-stability, I: Metric geometry of spaces of test configurations and valuations. *Annales de l'Institut Fourier*, 2024.
- [16] D. Catlin. The Bergman kernel and a theorem of Tian. In *Proceedings of the 40th Taniguchi symposium, Katata, Japan, June 23–28, 1997*, pages 1–23. Boston, MA: Birkhäuser, 1999.
- [17] H. Chen. Convergence of Harder-Narasimhan polygons. *Mém. Soc. Math. Fr., Nouv. Sér.*, 120:112, 2010.
- [18] H. Chen and C. Maclean. Distribution of logarithmic spectra of the equilibrium energy.

- Manuscr. Math.*, 146(3-4):365–394, 2015.
- [19] X. Chen. The space of Kähler metrics. *J. Differ. Geom.*, 56(2):189–234, 2000.
- [20] X. Chen and S. Sun. Space of Kähler metrics. V: Kähler quantization. In *Metric and differential geometry. The Jeff Cheeger anniversary volume. Selected papers based on the presentations at the international conference on metric and differential geometry, Tianjin and Beijing, China, May 11–15, 2009*, pages 19–41. Berlin: Springer, 2012.
- [21] X. Chen and Y. Tang. Test configuration and geodesic rays. In *Géométrie différentielle, physique mathématique, mathématiques et société (I). Volume en l’honneur de Jean Pierre Bourguignon*, pages 139–167. Paris: Société Mathématique de France, 2008.
- [22] J. Chu, V. Tosatti, and B. Weinkove. $C^{1,1}$ regularity for degenerate complex Monge-Ampère equations and geodesic rays. *Commun. Partial Differ. Equations*, 43(2):292–312, 2018.
- [23] R. R. Coifman and S. Semmes. Interpolation of Banach spaces, Perron processes, and Yang-Mills. *Am. J. Math.*, 115(2):243–278, 1993.
- [24] D. Coman and G. Marinescu. On the first order asymptotics of partial Bergman kernels. *Ann. Fac. Sci. Toulouse, Math. (6)*, 26(5):1193–1210, 2017.
- [25] X. Dai, K. Liu, and X. Ma. On the asymptotic expansion of bergman kernel. *J. Diff. Geom.*, 72(1):1–41, 2006.
- [26] T. Darvas. The Mabuchi geometry of finite energy classes. *Adv. Math.*, 285:182–219, 2015.
- [27] T. Darvas. The Mabuchi completion of the space of Kähler potentials. *Am. J. Math.*, 139(5):1275–1313, 2017.
- [28] T. Darvas. Weak geodesic rays in the space of Kähler potentials and the class $\mathcal{E}(X, \omega)$. *J. Inst. Math. Jussieu*, 16(4):837–858, 2017.
- [29] T. Darvas and L. Lempert. Weak geodesics in the space of Kähler metrics. *Math. Res. Lett.*, 19(5):1127–1135, 2012.
- [30] T. Darvas and C. H. Lu. Geodesic stability, the space of rays and uniform convexity in Mabuchi geometry. *Geom. Topol.*, 24(4):1907–1967, 2020.
- [31] T. Darvas, C. H. Lu, and Y. A. Rubinstein. Quantization in geometric pluripotential theory. *Commun. Pure Appl. Math.*, 73(5):1100–1138, 2020.
- [32] T. Darvas and M. Xia. The volume of pseudoeffective line bundles and partial equilibrium. *Geom. Topol.*, 28(4):1957–1993, 2024.
- [33] N. G. de Bruijn and P. Erdős. Some linear and some quadratic recursion formulas. II. *Nederl. Akad. Wet., Proc., Ser. A*, 55:152–163, 1952.
- [34] J.-P. Demailly. Regularization of closed positive currents and intersection theory. *J. Algebr. Geom.*, 1(3):361–409, 1992.
- [35] J.-P. Demailly. *Complex Analytic and Differential Geometry*. 2012.
- [36] S. K. Donaldson. Symmetric spaces, Kähler geometry and Hamiltonian dynamics. In *Northern California symplectic geometry seminar*, pages 13–33. Providence, RI: American Mathematical Society, 1999.
- [37] S. K. Donaldson. Scalar curvature and stability of toric varieties. *J. Differ. Geom.*, 62(2):289–349, 2002.

-
- [38] Z. Drmač, M. Omladič, and K. Veselić. On the perturbation of the Cholesky factorization. *SIAM J. Matrix Anal. Appl.*, 15(4):1319–1332, 1994.
- [39] S. Finski. On the metric structure of section ring, arXiv: 2209.03853, 50 p. 2022.
- [40] S. Finski. Submultiplicative norms and filtrations on section rings, arXiv: 2210.03039, 43 p. 2022.
- [41] S. Finski. Geometry at the infinity of the space of positive metrics: test configurations, geodesic rays and chordal distances, arXiv: 2305.15300, 48 p., to appear in *Journal für die reine und angewandte Mathematik*. 2023.
- [42] S. Finski. About Wess-Zumino-Witten equation and Harder-Narasimhan potentials, arXiv: 2407.06034, Second version, 53 p. 2024.
- [43] S. Finski. On Harder-Narasimhan slopes of direct images, arXiv:2402.08554. 2024.
- [44] G. B. Folland. *Introduction to partial differential equations*. Princeton, NJ: Princeton University Press, 2nd ed. edition, 1995.
- [45] I. C. Gohberg and M. G. Krein. *Introduction to the theory of linear nonselfadjoint operators*, volume 18 of *Transl. Math. Monogr.* American Mathematical Society (AMS), Providence, RI, 1969.
- [46] V. Guedj and A. Zeriahi. Intrinsic capacities on compact Kähler manifolds. *J. Geom. Anal.*, 15(4):607–639, 2005.
- [47] T. Hisamoto. On the limit of spectral measures associated to a test configuration of a polarized Kähler manifold. *J. Reine Angew. Math.*, 713:129–148, 2016.
- [48] L. Boutet de Monvel and V. Guillemin. *The Spectral Theory of Toeplitz Operators*. Princeton University Press, 1981.
- [49] R. Lazarsfeld. *Positivity in algebraic geometry. I. Classical setting: line bundles and linear series*, volume 48 of *Ergeb. Math. Grenzgeb., 3. Folge*. Berlin: Springer, 2004.
- [50] L. Lempert and L. Vivas. Geodesics in the space of Kähler metrics. *Duke Math. J.*, 162(7):1369–1381, 2013.
- [51] X. Ma and G. Marinescu. *Holomorphic Morse inequalities and Bergman kernels*, volume 254 of *Progr. Math.* Birkhäuser Verlag Basel, 2007.
- [52] X. Ma and G. Marinescu. Berezin-Toeplitz quantization on Kähler manifolds. *J. Reine Angew. Math.*, 662:1–56, 2012.
- [53] X. Ma and G. Marinescu. Exponential estimate for the asymptotics of Bergman kernels. *Math. Ann.*, 362(3-4):1327–1347, 2015.
- [54] X. Ma and W. Zhang. Superconnection and family Bergman kernels. *C. R. Acad. Sci. Paris*, 344(1):41 – 44, 2007.
- [55] X. Ma and W. Zhang. *Bergman Kernels and symplectic reduction*. Astérisque 318, 2008.
- [56] X. Ma and W. Zhang. Superconnection and family Bergman kernels. *Math. Ann.*, 386(3-4):2207–2253, 2023.
- [57] T. Mabuchi. Some symplectic geometry on compact Kähler manifolds. I. *Osaka J. Math.*, 24:227–252, 1987.

- [58] D. H. Phong and J. Sturm. The Monge-Ampère operator and geodesics in the space of Kähler potentials. *Invent. Math.*, 166(1):125–149, 2006.
- [59] D. H. Phong and J. Sturm. Test configurations for k -stability and geodesic rays. *J. Symplectic Geom.*, 5(2):221–247, 2007.
- [60] D. H. Phong and J. Sturm. The Dirichlet problem for degenerate complex Monge-Ampère equations. *Commun. Anal. Geom.*, 18(1):145–170, 2010.
- [61] D. H. Phong and J. Sturm. Regularity of geodesic rays and Monge-Ampère equations. *Proc. Am. Math. Soc.*, 138(10):3637–3650, 2010.
- [62] D. Rees. *Lectures on the asymptotic theory of ideals*, volume 113 of *Lond. Math. Soc. Lect. Note Ser.* Cambridge: Cambridge University Press, 1988.
- [63] R. Rochberg. Interpolation of Banach spaces and negatively curved vector bundles. *Pac. J. Math.*, 110:355–376, 1984.
- [64] J. Ross and D. W. Nyström. Analytic test configurations and geodesic rays. *J. Symplectic Geom.*, 12(1):125–169, 2014.
- [65] J. Ross and M. Singer. Asymptotics of partial density functions for divisors. *J. Geom. Anal.*, 27(3):1803–1854, 2017.
- [66] S. Semmes. Complex Monge-Ampère and symplectic manifolds. *Am. J. Math.*, 114(3):495–550, 1992.
- [67] O. Shabtai. Off-diagonal estimates of partial Bergman kernels on S^1 -symmetric Kähler manifolds, arXiv:2401.15416. 2024.
- [68] J. Song and S. Zelditch. Bergman metrics and geodesics in the space of Kähler metrics on toric varieties. *Anal. PDE*, 3(3):295–358, 2010.
- [69] J. Sun. Asymptotics of partial density function vanishing along smooth subvariety, arXiv:2010.04951. 2020.
- [70] G. Székelyhidi. *An introduction to extremal Kähler metrics*, volume 152 of *Grad. Stud. Math.* Providence, RI: American Mathematical Society (AMS), 2014.
- [71] G. Székelyhidi. Filtrations and test-configurations. *Math. Ann.*, 362(1-2):451–484, 2015.
- [72] G. Tian. On a set of polarized Kähler metrics on algebraic manifolds. *J. Diff. Geom.*, 32(1):99–130, 1990.
- [73] G. Tian. Kähler-Einstein metrics with positive scalar curvature. *Invent. Math.*, 130(1):1–37, 1997.
- [74] D. Witt Nyström. Test configurations and Okounkov bodies. *Compos. Math.*, 148(6):1736–1756, 2012.
- [75] S. Zelditch. Szegő kernels and a theorem of Tian. *Internat. Math. Res. Notices*, 6:317–331, 1998.
- [76] S. Zelditch and P. Zhou. Interface asymptotics of partial Bergman kernels on S^1 -symmetric Kähler manifolds. *J. Symplectic Geom.*, 17(3):793–856, 2019.