



**HAL**  
open science

# Confirmation of the presence of vaccine DNA in the Pfizer anti-COVID-19 vaccine

Didier Raoult

► **To cite this version:**

Didier Raoult. Confirmation of the presence of vaccine DNA in the Pfizer anti-COVID-19 vaccine. 2024. hal-04778576

**HAL Id: hal-04778576**

**<https://hal.science/hal-04778576v1>**

Preprint submitted on 12 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **TITLE PAGE**

2

3 **Type of article: Short communication**

4

5 **Title: Confirmation of the presence of vaccine DNA in the Pfizer anti-COVID-19 vaccine**

6

7 **Author list: Didier RAOULT<sup>1,2\*</sup>**

8

9 **Affiliations:** <sup>1</sup> IHU Méditerranée Infection, 19-21 boulevard Jean Moulin, 13005 Marseille,  
10 France; <sup>2</sup> Aix-Marseille Univ., Microbes Evolution Phylogeny and Infections (MEPHI), 27  
11 boulevard Jean Moulin, 13005 Marseille, France

12

13 **\* Corresponding author:** Didier Raoult. 19-21 boulevard Jean Moulin, 13005 Marseille,  
14 France. Phone: +33413732401. Email: didier.raoult@gmail.com

15

16 **Keywords:** SARS-CoV-2; Covid-19; vaccine; mRNA; DNA; impurities

17

18

## ABSTRACT

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

The rapid production of messenger RNA (mRNA)-based vaccines has been chosen as the most suitable strategy to fight the COVID-19 pandemic. Three studies reported the presence of DNA in significant amounts in Pfizer mRNA vaccines. We aimed to confirm the presence of this residual DNA. Vaccine plasmid DNA quantification using the Qubit fluorometer on a vaccine vial showed it was 216 ng/dose on average and approximately 24 times greater, reaching 5,160 ng/dose on average, after treatment with Triton-X-100. In addition, we obtained by next-generation sequencing the sequence of the complete plasmid DNA vaccine matrix (7,824 base pairs) with high coverage (98.3%) and sequencing depths (mean, 4,181-4,389 reads), indicating the presence of the plasmid DNA in high copy number. These results calls for an assessment of the copy number and nature of DNA in mRNA vaccines at a larger scale and in multiple batches, notably regarding the putative risk of DNA integration after delivery into cells.

## TEXT

35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58

The SARS-CoV-2 pandemic led to the very fast (<1 year) development of vaccines (Krammer, 2020; Topol, 2021), and more than 13 billion doses have been administered worldwide (including more than 150 million in France) (<https://coronavirus.jhu.edu/map.html>). This was in line with the current framework of thinking that advocate accelerating vaccine development to counter the rapid spread of an emerging infectious disease (Cohen, 2016). The Pfizer-BioNTech COVID-19 mRNA vaccine (or BNT162 Vaccine-19) has been the most used. It is a suspension injected intramuscularly that contains a codon-optimized single-stranded, 5'-capped mRNA encoding a full-length, modified SARS-CoV-2 spike protein once released into target cells (Jackson et al, 2020; Krammer, 2020; Polack et al, 2020). To allow a massive production of the BNT162 Vaccine-19 mRNA, manufacturing included transfection into *Escherichia coli* of the modified spike gene sequence using a plasmid with a final size of 7,824 base pairs (bp) that notably contains an active bacterial origin of replication, an initiation factor of the SV40 virus initiation factor, and a gene of resistance to kanamycin (Rapporteur Rolling Review critical assessment report, 2020). After bacteria lysis, the plasmid DNA matrix was extracted and linearized before its transcription by a T7 RNA polymerase in presence of N-methyl pseudoUridine then its hydrolysis. The Pfizer mRNA vaccine contains in principle extremely minimal quantities of DNA. Nonetheless, at least three studies reported the presence of DNA in Pfizer mRNA vaccine doses in significant amounts (Speicher et al, 2023; McKernan et al, 2023; König and Kirchner, 2024; Hughes, 2024). In a situation such as this, we believed that it was necessary to confirm or deny the presence of DNA without necessarily wanting to draw conclusions about its presence.

59 We carried out DNA quantification using the Qubit fluorometer (Invitrogen, Carlsbad,  
60 CA, United States) on a vaccine vial (batch no. GJ7184) which showed it was 216 ng/dose on  
61 average. However, after treatment with Triton-X-100, DNA quantity was approximately 24  
62 times greater, reaching 5,160 ng/dose on average. As the vaccine dose is reported to contain  
63 30 µg of RNA this means that it contained in this test 17% of DNA.

64 In addition, next-generation sequencing (NGS) was performed as previously described  
65 (Colson et al, 2022) with the Illumina technology using the Nextera XT paired end strategy on  
66 MiSeq instruments (Illumina Inc., San Diego, CA, United States), following the  
67 manufacturer's recommendations. NGS reads were mapped on the vaccine plasmid sequence  
68 deposited in GenBank (<https://www.ncbi.nlm.nih.gov/genbank/>) (Sayers et al, 2023) with the  
69 Accession no. OR134577.1. Reads were processed using bwa-mem2 v.2.2.1  
70 (<https://github.com/bwa-mem2/bwa-mem2>) (Jung et al, 2022) and Samtools v.1.17  
71 (<https://www.htslib.org/>) (Li et al, 2009). Reads shorter than 200 nucleotides and covering  
72 less than 90% of the vaccine plasmid were removed using Samtools v.1.17. Freebayes v1.3.6  
73 (<https://github.com/freebayes/freebayes>) (Garrison and Marth, 2012) and Bcftools v.1.17  
74 (<https://samtools.github.io/bcftools/bcftools.html>) were used for identifying mutations and  
75 generating consensus. NGS in absence of prior reverse transcription step of two vaccine  
76 batches (FP8191 and FP9359), which generated more than 140,000 reads larger than 200  
77 bases, made it possible to retrieve the complete sequences of the DNA plasmid (GenBank  
78 Accession No. PP544445 and PP544446) used to produce the vaccine RNA, with 7.824 base  
79 pairs with high sequencing depth (>4000 reads). Approximately between 63% and 97% of the  
80 sequenced DNA was from the vaccine. Moreover, we observed that DNase treatment  
81 (TURBO DNA-free kit, Invitrogen) of the nucleic acid extracts prevented the obtaining of  
82 NGS reads (only 2 and 1 reads, respectively, were mapped on the reference sequence).

83 Therefore, here we found by several approaches and in different batches of the Pfizer-  
84 BioNTech COVID-19 mRNA vaccine the abundant presence of DNA. In such vaccines, the  
85 maximal residual quantity of DNA not degraded during vaccine purification that are  
86 authorized by the European Medicines Agency and the Food and Drug Administration are  
87 330 ng of DNA per mg of RNA, or 10 ng of DNA per dose (Klinman et al, 2010; Rapporteur  
88 Rolling Review critical assessment report, 2020). Our findings are in line with those of other  
89 (not peer reviewed) studies that reported the presence of DNA in Pfizer-BioNTech COVID-  
90 19 mRNA vaccine vials by qPCR and the obtaining of full-length vaccine plasmid by NGS  
91 (McKernan et al, 2023; Speicher et al, 2023). In the Speicher et al's study, residual DNA  
92 quantities, ranging between 1,896 and 3,720 ng per dose were estimated using fluorometer  
93 based measurements with Qubit, and vaccine spike targeting qPCR obtained cycle threshold  
94 values ranging between 18.0 and 23.8 at 1:10 dilution, and  $16.9 \pm 0.5$  on undiluted vials  
95 contents (Speicher et al, 2023).

96 As a matter of fact, these results of huge quantities of plasmid DNA sequences per  
97 vaccine dose notably raise issues regarding a putative risk of its integration in the human  
98 genome after its entry into cells due to their packaging into cationic lipids (Klinman et al,  
99 2010). In DNA-based gene therapy, it was reported that a proportion of 10- 20% of cells are  
100 usually transfected and about 1-10% of the transiently transfected cells became stably  
101 transfected as a result of subsequent integration likely through crossover events during  
102 nuclear envelope membrane reformation at telophase (Haraguchi et al., 2002; Lim et al, 2023;  
103 Devaux and Camoin-Jau, 2024). Although the potential risk that DNA found in mRNA-based  
104 vaccines integrate and either induce expression of an oncogene or shut down expression of a  
105 tumor suppressor are extremely low (Wang et al, 2004; Klinman et al, 2010), this should  
106 deserve additional investigations.

107 Overall, this work confirms that a significant quantity of vaccine DNA, which is above  
108 recommended levels (Rapporteur Rolling Review critical assessment report, 2020; Klinman et  
109 al, 2010), was present in the tested vaccine batches. Previous findings warrant to be confirmed  
110 at a larger scale, which is technically very easy to perform on a large panel of vaccine vials  
111 from different batches in different laboratories worldwide. There may be variability from one  
112 batch to another, and this requires regular quality controls on the different batches to be  
113 consistent with the requirements of good laboratory practices, and to liberate DNA from lipid  
114 nanoparticles.

115

116

#### 117 **Acknowledgments**

118 I am thankful to Dr Christian Devaux for this help and support.

#### 119 **Data Availability**

120 Sequences of the DNA plasmids recovered from Pfizer mRNA vaccine batches FP8191 and  
121 FP9359 are available from GenBank with Accession No. PP544445 and PP544446,  
122 respectively.

#### 123 **Conflicts of Interest**

124 DR declares grants or contracts and royalties or licenses from Hitachi High-Technologies  
125 Corporation, Tokyo, Japan. He is a scientific board member of Eurofins company, and a  
126 founder and shareholder of a microbial culture company (Culture Top), and two  
127 biotechnology companies (Techno-Jouvence, and Gene and Green TK). Funding sources  
128 played no role in the design and performance of the study, the collection, management,  
129 analysis and interpretation of the data, or the preparation, review and approval of the  
130 manuscript.

#### 131 **Author Contributions**

132 Conceptualization, methodology, formal analysis and interpretation, writing-original draft  
133 preparation, writing—review and editing: DR.

#### 134 **Funding sources**

135 This study was supported by the French Government under the “Investments for the Future”  
136 program managed by the National Agency for Research (ANR) (Méditerranée-Infection 10-  
137 IAHU-03).

#### 138 **Ethics**

139 Not applicable.

140

141

#### 142 **References**

143

144 Cohen J. New vaccine coalition aims to ward off epidemics. *Science*. Available at  
145 [www.sciencemag.org/news/2016/09/new-vaccine-coalition-aims-ward-epidemics](http://www.sciencemag.org/news/2016/09/new-vaccine-coalition-aims-ward-epidemics)  
146 (September 2, 2016); <https://cepi.net/>.

147 Colson P, Fournier PE, Chaudet H, Delerce J, Giraud-Gatineau A, Houhamdi L, Andrieu C,  
148 Brechard L, Bedotto M, Prudent E, Gazin C, Beye M, Burel E, Dudouet P, Tissot-Dupont  
149 H, Gautret P, Lagier JC, Million M, Brouqui P, Parola P, Fenollar F, Drancourt M, La  
150 Scola B, Levasseur A, Raoult D. Analysis of SARS-CoV-2 Variants From 24,181  
151 Patients Exemplifies the Role of Globalization and Zoonosis in Pandemics. *Front*  
152 *Microbiol.* 2022 Feb 7;12:786233. doi: 10.3389/fmicb.2021.786233.

153 Devaux C, Camoin-Jau L. Transient blood release of synthetic SARS-Cov-2 spike protein  
154 after mRNA-based COVID-19 vaccination possibly contributes to ACE2 dysfunction  
155 leading to rare cases of myocarditis. *Cardiol Cardiovascular Med* 2024; 8(4307):307-29.

156 Garrison E and Marth G. Haplotype-based variant detection from short read sequencing.  
157 arXiv 2012. <https://doi.org/10.48550/arXiv.1207.3907>.



158 Haraguchi T, Koujin T, Shindo T, Bilir Ş, Osakada H, Nishimura K, Hirano Y, Asakawa H,  
159 Mori C, Kobayashi S, Okada Y, Chikashige Y, Fukagawa T, Shibata S, Hiraoka Y.  
160 Transfected plasmid DNA is incorporated into the nucleus via nuclear envelope  
161 reformation at telophase. *Commun Biol.* 2022 Jan 20;5(1):78. doi: 10.1038/s42003-022-  
162 03021-8.

163 Hughes DA. Chapter 6: Weaponised Deception. In “Covid-19,” Psychological Operations,  
164 and the War for Technocracy. Part of: Springer Professional "Wirtschaft+Technik" ,  
165 Springer Professional "Wirtschaft". 2024. Publisher: Springer International Publishing.  
166 [https://www.springerprofessional.de/en/covid-19-psychological-operations-and-the-](https://www.springerprofessional.de/en/covid-19-psychological-operations-and-the-war-for-technocracy/26991184)  
167 [war-for-technocracy/26991184.](https://www.springerprofessional.de/en/covid-19-psychological-operations-and-the-war-for-technocracy/26991184)

168 Jackson LA, Anderson EJ, Roupheal NG, Roberts PC, Makhene M, Coler RN, McCullough  
169 MP, Chappell JD, Denison MR, Stevens LJ, Pruijssers AJ, McDermott A, Flach B,  
170 Doria-Rose NA, Corbett KS, Morabito KM, O'Dell S, Schmidt SD, Swanson PA 2nd,  
171 Padilla M, Mascola JR, Neuzil KM, Bennett H, Sun W, Peters E, Makowski M, Albert J,  
172 Cross K, Buchanan W, Pikaart-Tautges R, Ledgerwood JE, Graham BS, Beigel JH;  
173 mRNA-1273 Study Group. An mRNA Vaccine against SARS-CoV-2 - Preliminary  
174 Report. *N Engl J Med.* 2020 Nov 12;383(20):1920-1931. doi: 10.1056/NEJMoa2022483.

175 Jung Y, Han D. BWA-MEME: BWA-MEM emulated with a machine learning approach.  
176 *Bioinformatics.* 2022 Apr 28;38(9):2404-2413. doi: 10.1093/bioinformatics/btac137.

177 Klinman DM, Klaschik S, Tross D, Shirota H, Steinhagen F. FDA guidance on prophylactic  
178 DNA vaccines: analysis and recommendations. *Vaccine.* 2010 Apr 1;28(16):2801-5. doi:  
179 10.1016/j.vaccine.2009.11.025.

180 König B, Kirchner JO. Methodological Considerations Regarding the Quantification of DNA  
181 Impurities in the COVID-19 mRNA Vaccine Comirnaty®. *Methods Protoc.* 2024, 7, 41.  
182 doi: 10.3390/mps7030041.

183 Krammer F. SARS-CoV-2 vaccines in development. *Nature*. 2020 Oct;586(7830):516-527.  
184 doi: 10.1038/s41586-020-2798-3.

185 Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.  
186 *Bioinformatics*. 2009 Jul 15;25(14):1754-60. doi:10.1093/bioinformatics/btp324.

187 Lim S, Yocum RR, Silver PA, Way JC. High spontaneous integration rates of end-modified  
188 linear DNAs upon mammalian cell transfection. *Sci Rep*. 2023 Apr 26;13(1):6835. doi:  
189 10.1038/s41598-023-33862-0.

190 McKernan K, Helbert Y, Kane LT, McLaughlin S. Sequencing of bivalent Moderna and  
191 Pfizer mRNA vaccines reveals nanogram to microgram quantities of expression vector  
192 dsDNA per dose. [https://pflegefueraufklaerung.de/wp-](https://pflegefueraufklaerung.de/wp-content/uploads/2023/05/Sequencing-of-bivalent_4-11-23.pdf)  
193 [content/uploads/2023/05/Sequencing-of-bivalent\\_4-11-23.pdf](https://pflegefueraufklaerung.de/wp-content/uploads/2023/05/Sequencing-of-bivalent_4-11-23.pdf). OSF Preprints 2023.  
194 <https://osf.io/preprints/osf/b9t7m>.

195 Polack FP, Thomas SJ, Kitchin N et al. Safety and Efficacy of the BNT162b2 mRNA Covid-  
196 19 Vaccine. *N Engl J Med* 2020;383:2603-2615.

197 Rapporteur Rolling Review critical assessment report - COVID-19 mRNA Vaccine  
198 BioNTech - Modified mRNA, encoding full length SARS-CoV-2 spike protein. 2020.  
199 [https://factreview.gr/wp-content/uploads/2023/07/Rolling-Review-Report-Quality-](https://factreview.gr/wp-content/uploads/2023/07/Rolling-Review-Report-Quality-COVID-19-mRNA-Vaccine-BioNTech.pdf)  
200 [COVID-19-mRNA-Vaccine-BioNTech.pdf](https://factreview.gr/wp-content/uploads/2023/07/Rolling-Review-Report-Quality-COVID-19-mRNA-Vaccine-BioNTech.pdf).

201 Sayers EW, Cavanaugh M, Clark K, Pruitt KD, Sherry ST, Yankie L, Karsch-Mizrachi I.  
202 GenBank 2023 update. *Nucleic Acids Res*. 2023 Jan 6;51(D1):D141-D144. doi:  
203 10.1093/nar/gkac1012.

204 Speicher DJ, Gutschi JRLM, Wiseman DM, McKernan K. DNA fragments detected in  
205 monovalent and bivalent Pfizer/BioNTech and Moderna modRNA COVID-19 vaccines  
206 from Ontario, Canada: Exploratory dose response relationship with serious adverse  
207 events. OSFPreprints 2023. <https://doi.org/10.31219/osf.io/mjc97>.

208 Topol EJ. Messenger RNA vaccines against SARS-CoV-2. *Cell*. 2021 Mar 18;184(6):1401.

209 doi: 10.1016/j.cell.2020.12.039.

210 Wang Z, Troilo PJ, Wang X, Griffiths TG, Pacchione SJ, Barnum AB, Harper LB, Pauley CJ,

211 Niu Z, Denisova L, Follmer TT, Rizzuto G, Ciliberto G, Fattori E, Monica NL, Manam S,

212 Ledwith BJ. Detection of integration of plasmid DNA into host genomic DNA following

213 intramuscular injection and electroporation. *Gene Ther*. 2004 Apr;11(8):711-21. doi:

214 10.1038/sj.gt.3302213.

215

216