

## Reply to Peng et al.: Chicken tessellation requires more pieces

Joris Peters, Dorian Q Fuller, Evan K Irving-Pease, Ophélie Lebrasseur, Julia Best, Riley Smallman, Greger Larson

## ► To cite this version:

Joris Peters, Dorian Q Fuller, Evan K Irving-Pease, Ophélie Lebrasseur, Julia Best, et al.. Reply to Peng et al.: Chicken tessellation requires more pieces. Proceedings of the National Academy of Sciences of the United States of America, 2022, 119 (44), pp.e2213678119. 10.1073/pnas.2213678119. hal-04782236

## HAL Id: hal-04782236 https://hal.science/hal-04782236v1

Submitted on 14 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



## Reply to Peng et al.: Chicken tessellation requires more pieces

Joris Peters<sup>a,b,c,1</sup>⑩, Dorian Q. Fuller (傅稻镰)<sup>d,e</sup>⑩, Evan K. Irving-Pease<sup>f,g</sup>, Ophélie Lebrasseur<sup>g,h,i</sup>⑩, Julia Best<sup>j,k</sup>⑫, Riley Smallman<sup>l</sup>⑮, and Greger Larson<sup>g,1</sup>⑩

In our PNAS article (1), we analyze the appearance and dispersal of chickens across Eurasia. Numerous factors complicate the confident dating, species determination, and domestic status assignment of archaeological bird remains. To minimize the likelihood of accepting spurious claims for early chickens, we applied strict, conservative criteria. The presence of chickens at the Thai Neolithic site of Ban Non Wat (1650–1250 BCE) thus represents a minimum bound.

In their letter, Peng et al. (2) describe additional Chinese sites with reported chickens. Unfortunately, none of these meet our criteria. At the Inner Mongolian Dadianzi tomb, an earthen-constructed site of the Lower Xiajiadian Culture abandoned by 1500 BCE (3), Yuan (4) attributed the remains to chickens, based on cooccurrence with dogs and pigs. Morphological confirmation and direct dating, however, have yet to be carried out. Similarly, descriptions of remains at Caiyuanzi are not available (5) and thus cannot be evaluated.

At Dadunzi, the single bone was described as "possibly domesticated" (6). Nonskeletal remains have also been excavated, and, although pottery and bronze art objects from Dadunzi and other sites suggest closer relationships between humans and birds resembling galliformes, their species identification and dating are uncertain. The bronze rooster from Sangxingdui is generally accepted, but the specific context of this find (Pit K2) is dated to  $\sim$ 1200 BCE (7). This places the artistic evidence just after our earliest confirmed chicken bones.

Author affiliations: <sup>a</sup>ArchaeoBioCenter, Ludwig Maxmilian University Munich, 80539 Munich, Germany; <sup>b</sup>Institute of Palaeoanatomy, Domestication Research and the History of Veterinary Medicine, Ludwig Maxmilian University Munich, 80539 Munich, Germany; <sup>c</sup>Bavarian Natural History Collections, State Collection of Palaeoanatomy Munich, 80333 Munich, Germany; <sup>d</sup>Institute of Archaeology, University College London, London WC1H 0PY, United Kingdom; <sup>e</sup>School of Cultural Heritage, Northwest University, 710069 Xian, China; <sup>f</sup>Lundbeck Foundation GeoGenetics Centre, GLOBE Institute, University of Copenhagen, 1165 Copenhagen, Denmark; <sup>8</sup>Palaeogenomics & Bio-Archaeology Research Network, School of Archaeology, University of Oxford, Oxford OX1 3TG, United Kingdom; <sup>h</sup>Centre for Anthropobiology and Genomics of Toulouse, Université de Toulouse Paul Sabatier, CNRS, Toulouse, 31000 France; <sup>h</sup>Instituto Nacional de Antropología y Pensamiento Latinoamericano, Ciudad Autonoma de Buenos Aires, C1426BJN, Argentina; <sup>J</sup>Department of Archaeology and Anthropology, Bournemouth University, Poole BH12 5BB, United Kingdom; <sup>\*</sup>School of History, Archaeology and Religion, Cardiff University, Cardiff CF10 3AT, United Kingdom


Author contributions: E.K.I.-P. analyzed data; and J.P., D.Q.F., E.K.I.-P., O.L., J.B., R.S., and G.L. wrote the paper.

The authors declare no competing interest.

Copyright © 2022 the Author(s). Published by PNAS. This article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

<sup>1</sup>To whom correspondence may be addressed. Email: joris.peters@palaeo.vetmed.unimuenchen.de or greger.larson@arch.ox.ac.uk.

Published October 24, 2022.



**Fig. 1.** A map depicting the earliest confidently assigned chicken remains across Eurasia, Africa, and Oceania alongside a spatial kriging interpolation of the timing of the arrival of chickens. The original map in our initial study (1) made use of the most recent end of the age ranges associated with the remains. This figure uses the oldest end of each age range, and this systematic inflation of the date estimates has no material effect on the interpolation.

Peng et al. (2) rightly point to Yunnan as a key region, since, as we note, the chicken ancestral subspecies *Gallus gallus spadiceus* is present. Agriculture was established here in 2600 BCE, based on a mixture of alluvial (wet) rice and rainfed millet (8). Based on our ecological coevolutionary model, the millet fallows would have provided potential commensal habitats for red junglefowl that wet rice did not. The predominance of wet rice over millet in Neolithic Yunnan datasets (8) therefore suggests that opportunities for these commensal pathways were present, but in a more limited fashion relative to peninsular Southeast Asia. Lastly, Peng et al. (2) criticize our decision to use the lower limits of each chicken's age ranges on our map. A revised map using the upper limits of the age ranges (Fig. 1) does not materially affect the pattern of dispersal inferred in our spatial kriging analysis.

Earlier chicken remains will certainly be identified. As Peng et al. (2) suggest, that effort will be facilitated through simultaneous investigations of archaeology, anthropology, ecology, and evolutionary biology, and through international collaborations with experts from Southeast Asia and across the globe.

- 2. M.-S. Peng, J.-L. Han, Y.-P. Zhang, Missing puzzle piece for the origins of domestic chickens. Proc. Natl. Acad. Sci., 10.1073/pnas.2210996119 (2022).
- 3. X. Tang, S. Shen, X. Su, From rammed earth to stone wall: Chronological insight into the settlement change of the Lower Xiajiadian culture. PLoS One 17, e0273161 (2022).
- 4. J. Yuan, Zooarchaeological study on the domestic animals in ancient China [in Chinese]. Quaternary Sci. 30, 298-306 (2010).
- Yunnan Provincial Institute of Cultural Relics and Archaeology, Yunnan Archaeological Team of the Institute of Archaeology CASS, Chengdu Municipal Institute of Cultural Relics and Archaeology, Museum of Chuxiong Autonomous Prefecture, Yongren Country Cultural Center, Excavation of the Caiyuanzi and Mopandi sites in Yongren, Yunnan, in 2001. Acta Geol. Sin., 2, 263–296, 325–328 (2003).
- 6. The Yunnan Provincial Museum, The Neolithic site at Ta-Tun-Tzüin Yuan-Mou County, Yunnan Province. Acta Geol. Sin., 1, 43–72, 169–178 (1977).
- 7. H. Sun, "The Sangxingdui culture of the Sichuan Basin" in A Companion to Chinese Archaeology, A. P. Underhill, Ed. (Wiley-Blakcwell, Oxford, 2013), pp. 147-168.
- 8. R. Dal Martello, The origins of multi-cropping agriculture in Southwestern China: Archaeobotanical insights from third to first millennium BC Yunnan. Asian Archaeol. 6, 65-85 (2022).

<sup>1.</sup> J. Peters et al., The biocultural origins and dispersal of domestic chickens. Proc. Natl. Acad. Sci. U.S.A. 119, e2121978119 (2022).