



HAL
open science

CaV β 1A and CaV β 1E embryonic isoforms in adult skeletal muscle, a Mbnl1 related-expression

Amélie Vergnol, Alain Sureau, Eric Batsché, Eric Allemand, Massiré Traoré, Xavière Lornage, Geneviève Gourdon, Furling D, France Piétri-rouxel, Sestina Falcone

► **To cite this version:**

Amélie Vergnol, Alain Sureau, Eric Batsché, Eric Allemand, Massiré Traoré, et al.. CaV β 1A and CaV β 1E embryonic isoforms in adult skeletal muscle, a Mbnl1 related-expression. *Myology* 2024, Apr 2024, Paris, France. <hal-04782481>

HAL Id: hal-04782481

<https://hal.science/hal-04782481v1>

Submitted on 14 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization


CaVβ1A and CaVβ1E embryonic isoforms in adult skeletal muscle, a Mbn1 related-expression

A. Vergnol¹, A. Sureau¹, E. Batsché², E. Allemand³, M. Traoré¹, X. Lornage¹, G. Gourdon¹, D. Furling¹, F. Pietri-Rouxel^{1*} and S. Falcone^{1*}


¹Sorbonne Université, Inserm, Institut De Myologie, Centre De Recherche En Myologie F-75013 – Paris (France), Sorbonne Université, Institut de Biologie Paris-Seine (IBPS), CNRS Unit of Biological Adaptation and Ageing (B2A), Epigenetics and RNA metabolism in human diseases, 75005, Paris (France.) ²Inserm, U1163, Institut Imagine – Paris (France), ³Equally contributed to this work

INTRODUCTION


CaVβ1-E/GDF5 axis in muscle mass homeostasis



Cacnb1 isoforms in skeletal muscle




Implication of MBNLs in DM1 pathophysiology




Voltage-gated calcium channels (CaVs or VGCCs) are major regulators of calcium-related cellular functions. In skeletal muscle, though the essential component of the pore channel is the CaVα1 subunit, the CaVβ1 subunit is an essential subunit guaranteeing CaV fine-tuning activity. CaVβ1, encoded by *Cacnb1* gene, exists as several variants in skeletal muscle. Our recent work demonstrates CaVβ1D as the constitutive adult isoform, localized with CaVα1 at the triad. On the other hand, our previous and present works show that CaVβ1E and CaVβ1A are expressed at late embryogenesis and peri-natal stages and that these embryonic isoforms are re-expressed after sciatic nerve resection in adult skeletal muscle, both at mRNA and protein level enabling the expression of GDF5 (Growth Differentiation Factor 5) to counteract excessive muscle wasting (Traoré et al. 2019). However, the mechanisms leading to the increase these embryonic isoform expressions are unknown to date.

Our RNAseq data analysis in innervated versus denervated muscles revealed MuscleBlind-Like (MBNL) proteins as potential candidates regulating CaVβ1 expression in skeletal muscle. Interestingly, in a human model of Dystrophy Myotonic 1 (DM1), the sequestration of MBNLs in toxic nuclear aggregates is related to an impaired splicing of CaVβ1 transcript (*CACNB1*) (Arandel et al. 2017). Here, we evaluate the effect of a modulation of MBNLs protein levels on the expression of CaVβ1 isoforms in both *in vitro* and *in vivo* systems as well as in pathological mouse models of DM1.


Diversity of CaVβ1 isoforms in skeletal muscle




Regulation of the expression of CaVβ1 isoforms




Regulation of CaVβ1 isoforms in the physiopathology of muscle disease(s): the possible role of Muscleblind-like proteins (Mbnls)



Cross-talk between Mbnl proteins in the regulation of CaVβ1 isoforms



Cacnb1 isoforms in DM1 models and potential role in the disease



CONCLUSIONS & PERSPECTIVES

- Adult *Cacnb1D* and embryonic *Cacnb1A/E* isoforms are expressed upon specific promoter activation
- A downregulation of *Mbn1*, and the subsequent increase of *Mbn3*, leads to increased *CaVβ1-A* and *CaVβ1-E* and decreased *CaVβ1-D* expression *in vivo*
- Mouse models of DM1 are associated with an increased *Cacnb1-A* and *Cacnb1-E* expression

- Characterization of the splicing events occurring at *Cacnb1* Ex2-3 and Ex13-14
- Deciphering the role of *CaVβ1-E* in DM1 pathophysiology