



CaV β 1A and CaV β 1E embryonic isoforms in adult skeletal muscle, a Mbnl1 related-expression

Amélie Vergnol, Alain Sureau, Eric Batsché, E. Lallemand, Massiré Traoré, Xavière Lornage, Geneviève Gourdon, Furling D, France Piétri-rouxel, Sestina Falcone

► To cite this version:

Amélie Vergnol, Alain Sureau, Eric Batsché, E. Lallemand, Massiré Traoré, et al.. CaV β 1A and CaV β 1E embryonic isoforms in adult skeletal muscle, a Mbnl1 related-expression. Myology 2024, Apr 2024, Paris, France. hal-04782481

HAL Id: hal-04782481

<https://hal.science/hal-04782481v1>

Submitted on 14 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


CaV β 1A and CaV β 1E embryonic isoforms in adult skeletal muscle, a Mbnl1 related-expression

A. Vergnol¹, A. Sureau¹, E. Batsché², E. Allemand³, M. Traoré¹, X. Lornage¹, G. Gourdon¹, D. Furling¹, F. Pietri-Rouxel^{1*} and S. Falcone^{1*}


¹Sorbonne Université, Inserm, Institut De Myologie, Centre De Recherche En Myologie F-75013 – Paris (France), Sorbonne Université, Institut de Biologie Paris-Seine (IBPS), CNRS Unit of Biological Adaptation and Ageing (B2A), Epigenetics and RNA metabolism in human diseases, 75005, Paris (France). ²Inserm, U1163, Institut Imagine – Paris (France), *Equally contributed to this work

INTRODUCTION

Cacnb1 isoforms in skeletal muscle



Implication of MBNLs in DM1 pathophysiology




Voltage-gated calcium channels (CaVs or VGCCs) are major regulators of calcium-related cellular functions. In skeletal muscle, though the essential component of the pore channel is the CaV α 1 subunit, the CaV β 1 subunit is an essential subunit guaranteeing CaV fine-tuning activity. CaV β 1, encoded by Cacnb1 gene, exists as several variants in skeletal muscle. Our recent work demonstrates CaV β 1D as the constitutive adult isoform, localized with CaV α 1 at the triad. On the other hand, our previous and present works show that CaV β 1E and CaV β 1A are expressed at late embryogenesis and peri-natal stages and that these embryonic isoforms are re-expressed after sciatic nerve resection in adult skeletal muscle, both at mRNA and protein level enabling the expression of GDF5 (Growth Differentiation Factor 5) to counteract excessive muscle wasting (Traoré et al. 2019). However, the mechanisms leading to the increase these embryonic isoform expressions are unknown to date.

Our RNAseq data analysis in innervated versus denervated muscles revealed Muscleblind-Like (MBNL) proteins as potential candidates regulating CaV β 1 expression in skeletal muscle. Interestingly, in a human model of Dystrophy Myotonic 1 (DM1), the sequestration of MBNLs in toxic nuclear aggregates is related to an impaired splicing of CaV β 1 transcript (CACNB1) (Arandell et al. 2017).



Here, we evaluate the effect of a modulation of MBNLs protein levels on the expression of CaV β 1 isoforms in both *in vitro* and *in vivo* systems as well as in pathological mouse models of DM1.

Diversity of CaV β 1 isoforms in skeletal muscle




Regulation of the expression of CaV β 1 isoforms


Exon1-Exon2B : Epigenetic marks



Regulation of CaV β 1 isoforms in the physiopathology of muscle disease(s): the possible role of Muscleblind-like proteins (MBNLs)




Cross-talk between Mbni proteins in the regulation of CaV β 1 isoforms





Cacnb1 isoforms in DM1 models and potential role in the disease

Variations of Cacnb1 isoforms at transcriptional level



Fonctional analysis



CONCLUSIONS & PERSPECTIVES

- Adult Cacnb1D and embryonic Cacnb1A/E isoforms are expressed upon specific promoter activation
- A downregulation of Mbni1, and the subsequent increase of Mbni3, leads to increased CaV β 1-A and CaV β 1-E and decreased CaV β 1-D expression *in vivo*
- Mouse models of DM1 are associated with an increased Cacnb1-A and Cacnb1-E expression

- Characterization of the splicing events occurring at Cacnb1 Ex2-3 and Ex13-14
- Deciphering the role of CaV β 1-E in DM1 pathophysiology