



HAL
open science

On curve-flat Lipschitz functions and their linearizations

Gonzalo Flores, Mingu Jung, Gilles Lancien, Colin Petitjean, Antonín Procházka, Andrés Quilis

► **To cite this version:**

Gonzalo Flores, Mingu Jung, Gilles Lancien, Colin Petitjean, Antonín Procházka, et al.. On curve-flat Lipschitz functions and their linearizations. 2024. hal-04784542

HAL Id: hal-04784542

<https://hal.science/hal-04784542v1>

Preprint submitted on 15 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON CURVE-FLAT LIPSCHITZ FUNCTIONS AND THEIR LINEARIZATIONS

GONZALO FLORES, MINGU JUNG, GILLES LANCIEN, COLIN PETITJEAN,
ANTONÍN PROCHAZKA, AND ANDRÉS QUILIS

ABSTRACT. We show that several operator ideals coincide when intersected with the class of linearizations of Lipschitz maps. In particular, we show that the linearization \widehat{f} of a Lipschitz map $f : M \rightarrow N$ is Dunford-Pettis if and only if it is Radon-Nikodým if and only if it does not fix any copy of L_1 . We also identify and study the corresponding metric property of f , which is a natural extension of the curve-flatness introduced in [5]. Further, we show that \widehat{f} is compact if and only if it does not fix any copy of ℓ_1 .

1. INTRODUCTION

The Lipschitz-free space $\mathcal{F}(M)$ over a pointed metric space M is a Banach space that is built around M in such a way that M is isometric to a subset $\delta(M)$ of $\mathcal{F}(M)$, and 0-preserving Lipschitz maps from $\delta(M)$ into any other Banach space X uniquely extend to bounded linear operators from $\mathcal{F}(M)$ into X (see Section 2.1 for details and references). It follows that, given two metric spaces M, N , for any 0-preserving Lipschitz map $f : M \rightarrow N$ there exists its canonical linearization $\widehat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ which is a bounded linear operator, called *Lipschitz operator* hereafter. The mapping $M \mapsto \mathcal{F}(M)$ and $f \mapsto \widehat{f}$ is a functor between the category of pointed metric spaces together with 0-preserving Lipschitz maps and the category of Banach spaces together with bounded linear maps. In order to gain a better understanding of either of these categories, it is intriguing to study both

- A) the relation between the metric properties of M and the linear properties of $\mathcal{F}(M)$, as well as
- B) the relation between properties of f and corresponding properties of \widehat{f} .

For example, in the direction A) it has been shown in [5, Theorem C]: *For a complete M it is equivalent*

- (i) M is purely-1-unrectifiable (p1u),
- (ii) $\mathcal{F}(M)$ has the Radon-Nikodým property (RNP),
- (iii) $\mathcal{F}(M)$ has the Schur property (SP),
- (iv) L_1 is not isomorphic to a subspace of $\mathcal{F}(M)$.

In this paper, we aim to proceed in the direction B), i.e. we want to study the part of the free functor acting on morphisms. Our main result is the following extension of the aforementioned Theorem C of [5]:

2020 *Mathematics Subject Classification*. Primary 47B01, 47B07, 47B33; Secondary 46B20, 54E35.

Key words and phrases. Lipschitz-free space, Lipschitz operator, Radon-Nikodym, Dunford-Pettis, p1u metric space, completely continuous, strictly singular.

Theorem 1.1. *For a 0-preserving Lipschitz map $f : M \rightarrow N$ between complete pointed metric spaces M and N , the following are equivalent:*

- (i) f is curve-flat (CF), that is, for every $K \subset \mathbb{R}$ compact and every Lipschitz map $\gamma : K \rightarrow M$,

$$\lim_{\substack{y \rightarrow x \\ y \in K}} \frac{d(f(\gamma(x)), f(\gamma(y)))}{|x - y|} = 0 \quad \text{for } \lambda\text{-almost every } x \in K.$$

- (ii) \widehat{f} is a Radon-Nikodým (RN) operator.
 (iii) \widehat{f} is a Dunford-Pettis (DP) operator.
 (iv) \widehat{f} does not fix any copy of L_1 .

We will prove Theorem 1.1 in Section 3 (the equivalence of (i), (iii) and (iv)) and in Section 4 (the equivalence of (i), (ii) and (iv)).

Theorem 1.1 is indeed an extension of [5, Theorem C] because of the functoriality mentioned above (in particular since $\widehat{Id}_M = Id_{\mathcal{F}(M)}$) and since we have

- (i) M is p1u if and only if $Id_M : M \rightarrow M$ is CF,
 (ii) X has the RNP if and only if $Id_X : X \rightarrow X$ is RN,
 (iii) X has the SP if and only if $Id_X : X \rightarrow X$ is DP,
 (iv) L_1 is isomorphic to a subspace of X if and only if $Id_X : X \rightarrow X$ fixes a copy of L_1 .

In the subsequent sections we explore some other operator properties which extend properties of Banach spaces through similar equivalences. Namely, notice that it is true that X has the RNP if and only if $Id_X : X \rightarrow X$ is a strong Radon-Nikodým operator (strong RN). In Section 5 we study the possibility of adding the item “ \widehat{f} is strong RN” to the equivalence of Theorem 1.1. While we show that this is not possible in general (i.e. there exists curve-flat f such that \widehat{f} is not strong RN; see Example 5.3), we show that for compact countably 1 rectifiable M such equivalence holds true (see Proposition 5.1).

Also notice that X has the SP if and only if $Id_X : X \rightarrow X$ factors through a Banach space Z with the Schur property. We do not know whether the item “ \widehat{f} factors through a Schur space” could be added to the equivalence of Theorem 1.1. In Section 6 we make several remarks in this respect. In general it is not possible to add an item claiming “ $f : M \rightarrow N$ factors through a p1u complete metric space”, i.e. there exists curve-flat f which does not factor through any complete p1u space (see Example 6.8). Notice that if each curve-flat function factored through a complete p1u metric space, Theorem 1.1 could be obtained by direct application of [5, Theorem C]. Thus, the above example shows that such a naive approach does not work in general. In fact, we characterize those compact metric spaces M such that every curve-flat $f : M \rightarrow N$ factors through a p1u complete metric space (see Theorem 6.5). It happens precisely when the second dual of the space of uniformly locally flat functions is the space of curve-flat functions. From a metric viewpoint, this happens exactly when collapsing all curve fragments in M leads to a p1u space.

In Section 7 we address another topic in the direction B) as described above. It has been proved in [2]: *For complete metric spaces M, N and a Lipschitz $f : M \rightarrow N$ the following are equivalent:*

- a) \widehat{f} is weakly compact,
 b) \widehat{f} is compact.

Here we prove that when \widehat{f} does not fix any copy of ℓ_1 then it is a compact operator. Thus we can extend the above characterization by adding the following equivalent items:

- c) \widehat{f} is strictly singular,
- d) \widehat{f} is Asplund,
- e) \widehat{f} does not fix any copy of ℓ_1 .

Finally, in an appendix, we discuss a short proof due to Sylvester Eriksson-Bique of Theorem A in [5] and its relation to our proof of Theorem 1.1.

2. PRELIMINARIES

Throughout the paper, we will use the letters M, N for metric spaces. If $p \in M$, $r \geq 0$ and S a non empty subset of M , we will use the notation:

$$\begin{aligned} B(p, r) &= \{x \in M \mid d(x, p) \leq r\} \\ d(p, S) &= \inf\{d(x, p) : x \in S\} \\ [S]_r &= \{x \in M : d(x, S) \leq r\} \\ \text{rad}(S) &= \sup\{d(0, x) : x \in S\}. \end{aligned}$$

The letters X, Y are reserved for real Banach spaces. As usual, X^* denotes the topological dual of X , and $\mathcal{L}(X, Y)$ is the space of all bounded operators from X to Y . Additionally, B_X and S_X denote the closed unit ball and the unit sphere of X , respectively. If $S \subset X$ then $\text{conv}(S)$ stands for the convex hull of S . Ordinarily, $L_1 = L_1([0, 1])$ represents the Banach space of Lebesgue integrable functions from $[0, 1]$ to \mathbb{R} . The Lebesgue measure is denoted by λ . Similarly, $\ell_1 = \ell_1(\mathbb{N})$ is the space of absolutely summable sequences in \mathbb{R} . To indicate that X is isometrically isomorphic to Y , we use the notation $X \equiv Y$.

2.1. Lipschitz-free spaces. By a *pointed metric space* we mean a metric space M containing a distinguished point, denoted 0 and called a *basepoint*. It is well known that while the basepoint is needed in order to elegantly formulate the following definitions and properties, the particular choice of this point is inconsequential.

If M and N are pointed metric spaces then $\text{Lip}_0(M, N)$ stands for the set of all Lipschitz functions $f : M \rightarrow N$ such that $f(0) = 0$. If $N = X$ is a real Banach space, then $\text{Lip}_0(M, X)$ naturally becomes a Banach space when equipped with the Lipschitz norm:

$$\forall f \in \text{Lip}_0(M, X), \quad \|f\|_L = \sup_{x \neq y \in M} \frac{\|f(x) - f(y)\|_X}{d(x, y)}.$$

In the case $X = \mathbb{R}$, it is customary to shorten the notation to $\text{Lip}_0(M) := \text{Lip}_0(M, \mathbb{R})$. Next, for $x \in M$, we let $\delta_M(x) \in \text{Lip}_0(M)^*$ be the evaluation functional defined by $\langle \delta_M(x), f \rangle = f(x)$, for all f in $\text{Lip}_0(M)$. It is readily seen that $\delta_M : x \in M \mapsto \delta_M(x) \in \text{Lip}_0(M)^*$ is an isometry. The *Lipschitz-free space over M* is then defined as the closed linear span of all such evaluation functionals:

$$\mathcal{F}(M) := \overline{\text{span}}^{\|\cdot\|} \{\delta(x) : x \in M\} \subset \text{Lip}_0(M)^*.$$

A classical reference on Lipschitz-free spaces is [42].

Let us first recall the universal extension property of Lipschitz-free spaces. Let X be a Banach space and $f \in \text{Lip}_0(M, X)$. Then there exists a unique $\widehat{f} \in$

$\mathcal{L}(\mathcal{F}(M), X)$ such that $f = \bar{f} \circ \delta_M$. That is, we have the following commutative diagram:

$$\begin{array}{ccc} M & \xrightarrow{f} & X \\ \delta_M \downarrow & \nearrow \bar{f} & \\ \mathcal{F}(M) & & \end{array}$$

Moreover $\|\bar{f}\|_{\mathcal{L}(\mathcal{F}(M), X)} = \|f\|_L$. In particular, $\text{Lip}_0(M, X) \equiv \mathcal{L}(\mathcal{F}(M), X)$ and $\mathcal{F}(M)$ is an isometric predual of $\text{Lip}_0(M)$, that is $\text{Lip}_0(M) \equiv \mathcal{F}(M)^*$.

From the previous factorization, one easily deduces another fundamental feature of Lipschitz-free spaces that we describe now. If M, N are pointed metric spaces and $f : M \rightarrow N$ is a basepoint preserving Lipschitz map, then there exists a unique bounded linear operator $\hat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ such that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \delta_M \downarrow & & \downarrow \delta_N \\ \mathcal{F}(M) & \xrightarrow{\hat{f}} & \mathcal{F}(N). \end{array}$$

Moreover, we have that $\|\hat{f}\| = \|f\|_L$. Operators of the form \hat{f} are commonly referred to as *Lipschitz operators* or, more pragmatically, as linearizations of Lipschitz maps. It is noteworthy that the adjoint of $\hat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ is the “pre-composition by f ” operator acting between the respective spaces of Lipschitz functions, that is $(\hat{f})^* = C_f : g \in \text{Lip}_0(N) \mapsto g \circ f \in \text{Lip}_0(M)$.

We also wish to recall some further important features of Lipschitz-free spaces. Firstly, note that, on bounded subsets of $\text{Lip}_0(M)$, the weak* topology coincides with the pointwise topology. Next, if $0 \in K \subset M$, then $\mathcal{F}(K) \equiv \mathcal{F}_M(K)$ where:

$$\mathcal{F}_M(K) := \overline{\text{span}}\{\delta_M(x) \mid x \in K\} \subset \mathcal{F}(M).$$

Under this identification, the *support* of $\gamma \in \mathcal{F}(M)$ is the smallest closed subset $K \subset M$ such that $\gamma \in \mathcal{F}_M(K)$. It is denoted by $\text{supp}(\gamma)$. For further details, we refer to [6, 8]. If $x \neq y \in M$, then the elementary molecule $m_{x,y} \in \mathcal{S}_{\mathcal{F}(M)}$ is defined by:

$$m_{x,y} := \frac{\delta(x) - \delta(y)}{d(x,y)}.$$

Finally, we will invoke McShane-Whitney’s extension theorem at various places. In essence, it allows one to extend a real valued Lipschitz map defined on a subset $S \subset M$ to the whole space M without increasing its Lipschitz constant; see e.g. [42, Theorem 1.33].

2.2. Preliminaries on operators and operator ideals. In this paper we will deal with the following well known operator ideals:

- **Dunford-Pettis operators.** An operator $T : X \rightarrow Y$ between Banach spaces is *Dunford-Pettis* (or *completely continuous*) whenever it is weak-to-norm sequentially continuous. Equivalently, $T(W)$ is a norm-compact subset of Y whenever W is a weakly compact subset of X . We will denote by DP the operator ideal in the sense of Pietsch [37] of Dunford-Pettis operators and by $\text{DP}(X, Y)$ the set of all DP operators from X to Y .

- **Radon-Nikodým operators.** An operator $T : X \rightarrow Y$ between Banach spaces is called *Radon-Nikodým operator* if T takes bounded X -valued martingales into martingales which converge pointwise almost everywhere (see also Section 2.4 below). For equivalent definitions see [20] and references therein. We will denote by \mathcal{RN} the operator ideal of Radon-Nikodým operators and by $\mathcal{RN}(X, Y)$ the set of all \mathcal{RN} operators from X to Y .
- **Operators factoring through a Banach space with the Schur property.** Partially following the notation of [37], the ideals of these operators will be called $Op(SP)$. While it is clear that $Op(SP) \subset DP$, the opposite inclusion does not hold. Indeed, the following can be extracted from the monograph of Pietsch [37, Proposition 3.1.8] (where the ideal of Dunford-Pettis operators is denoted by \mathfrak{D} or some similar letter): the canonical injection $T : C([0, 1]) \rightarrow L_2([0, 1])$ is Dunford-Pettis since it is weakly compact and since $C([0, 1])$ has the Dunford-Pettis property; but fails to factor through a Schur space. Indeed, if there existed a Schur space X and two operators $R : C([0, 1]) \rightarrow X$ and $S : X \rightarrow L_2([0, 1])$ such that $T = S \circ R$, then R would be weakly compact as $C([0, 1])$ has property (V) of Pelczynski [34] and S would be Dunford-Pettis. However, as a composition of a weakly compact map with a Dunford-Pettis map, T would be a compact operator, yielding a contradiction.
- **Operators factoring through a Banach space with the RNP.** Partially following the notation of [37], the ideals of these operators will be called $Op(RNP)$. Again we clearly have that $Op(RNP) \subset \mathcal{RN}$. And again the opposite inclusion does not hold: an example is constructed in [24] of a \mathcal{RN} operator which does not factor through any space with the RNP.
- **Operators factoring through an Asplund space, Asplund operators.** An operator $T : X \rightarrow Y$ is said to be an Asplund operator if its adjoint operator is Radon-Nikodým. Equivalently, T is Asplund if it factors through an Asplund space [20, 41].
- **Strictly singular operators.** An operator $T : X \rightarrow Y$ between Banach spaces is *strictly singular* if for every infinite-dimensional subspace $Z \subseteq X$ the restriction $T|_Z$ is not an isomorphic embedding. In other words: For any $\varepsilon > 0$ and any infinite dimensional subspace Z of X , there exists $u \in S_Z$ such that $\|Tu\| \leq \varepsilon$.
- **Operators not fixing any copy of ℓ_1 (respectively, L_1).** An operator $T : X \rightarrow Y$ does not fix any copy of ℓ_1 (respectively, L_1) if there is no subspace $Z \subset X$ which is isomorphic to ℓ_1 (respectively, L_1) and such that $T|_Z$ is an isomorphic embedding.

We will also briefly consider the class of strong Radon-Nikodým operators (which does not form an ideal). An operator $T : X \rightarrow Y$ between Banach spaces is a *strong Radon-Nikodým operator* if $\overline{T(B_X)}$ is a RNP set. Recall that a closed bounded subset C of a Banach space X is called a *Radon-Nikodým set* (RNP set, in short) if for every probability space $(\Omega, \Sigma, \mathbb{P})$ and any vector measure $m : \Sigma \rightarrow X$ such that the set $\{m(B)/\mathbb{P}(B) : B \in \Sigma, \mathbb{P}(B) > 0\}$ is contained in C , there exists a Bochner-integrable Radon-Nikodým derivative $f : \Omega \rightarrow X$ such that $m(B) = \int_B f d\mathbb{P}$ for every $B \in \Sigma$. It is known that if T is strong Radon-Nikodým, then it is Radon-Nikodým (see [20]) while the converse is not true in general (considering any quotient map from ℓ_1 onto c_0). Notice that a Banach space X has the RNP if

and only if the identity operator on X is strong Radon-Nikodým, equivalently, the closed unit ball of X is a RNP set.

Finally, for a finite measure space (Ω, Σ, μ) , a bounded operator $T : L_1(\mu) \rightarrow Y$ is said to be *representable* if there exists $g \in L^\infty(\mu, Y)$ such that:

$$\forall h \in L_1(\mu), \quad T(h) = \int_{\Omega} h g d\mu.$$

2.3. Preliminaries on curve-flat Lipschitz functions. A *Lipschitz curve* in a metric space M is a Lipschitz map $\gamma : K \rightarrow M$ where K is a compact subset of the real line. We warn the reader that our use of the term “Lipschitz curve” differs from a more traditional meaning where the domain of γ would be required to be an interval. The next concept has been introduced in [5, Definition 5.21] for functions with values in \mathbb{R} . We say that a Lipschitz map $f : M \rightarrow N$ is *curve-flat* if for every Lipschitz curve $\gamma : K \rightarrow M$,

$$\lim_{\substack{y \rightarrow x \\ y \in K}} \frac{d(f(\gamma(x)), f(\gamma(y)))}{|x - y|} = 0$$

for λ -almost every $x \in K$, where λ denotes the Lebesgue measure on \mathbb{R} . We will use the following notation

$$\text{CF}_0(M, N) := \{f \in \text{Lip}_0(M, N) : f \text{ is curve-flat}\}.$$

If $N = \mathbb{R}$ we abbreviate the notation by letting $\text{CF}_0(M) := \text{CF}_0(M, \mathbb{R})$. Let us mention that a routine argument shows that $\text{CF}_0(M, Y)$ is a closed subspace of $\text{Lip}_0(M, Y)$ when Y is a Banach space. We will see in Proposition 2.2 below that $\text{CF}_0(M)$ is even weak*-closed.

The curve-flat functions are close relatives of (but should not be confused with) *uniformly locally flat* functions, that is functions f such that

$$\lim_{\substack{d(x,y) \rightarrow 0 \\ x \neq y}} \frac{d(f(x), f(y))}{d(x, y)} = 0.$$

We will use the following notation

$$\text{lip}_0(M, N) := \{f \in \text{Lip}_0(M, N) : f \text{ is uniformly locally flat}\}.$$

If $N = \mathbb{R}$ we let $\text{lip}_0(M) := \text{lip}_0(M, \mathbb{R})$.

We always have $\text{lip}_0(M, N) \subset \text{CF}_0(M, N)$ but equality might not hold. Indeed, consider $M = \{0\} \cup \{\frac{1}{n} : n \in \mathbb{N}\}$ together with the metric d inherited from \mathbb{R} , then $d(\cdot, 0) \in \text{CF}_0(M) \setminus \text{lip}_0(M)$.

In this paper, we will handle curve-flat functions through equivalent descriptions based on classical results from metric geometry, namely Kirchheim’s theorems [31]. For the convenience of the reader, let us state the relevant results to be used in the sequel. From now on, \mathcal{H}^1 stands for the Hausdorff 1-measure.

Theorem (Area Formula, [31, Corollary 8]). *Let A be a Borel subset of \mathbb{R} and $\gamma : A \rightarrow M$ be a Lipschitz map. Let the metric differential of γ at $x \in A$ be the nonnegative number:*

$$MD(\gamma, x) := \lim_{\substack{y \rightarrow x \\ y \in A}} \frac{d(\gamma(x), \gamma(y))}{|x - y|}, \text{ whenever this limit exists.}$$

For each $y \in \gamma(A)$, $|\gamma^{-1}(y)| \in [1, \infty]$ denotes the cardinality of the preimage $\gamma^{-1}(y)$. Then

$$\int_A MD(\gamma, x) d\lambda(x) = \int_{\gamma(A)} |\gamma^{-1}(y)| d\mathcal{H}^1(y).$$

Consequently, $MD(\gamma, x) = 0$ for λ -almost every $x \in A$ if and only if $\mathcal{H}^1(\gamma(A)) = 0$.

Kirchheim's lemma. For every metric space (M, d) and every Lipschitz map $\gamma : A \rightarrow M$ with $A \subset \mathbb{R}$ Borel and $\mathcal{H}^1(\gamma(A)) > 0$, there exists a compact subset $S \subset A$ with $\lambda(S) > 0$ such that $\gamma|_S$ is bi-Lipschitz.

The previous lemma does not appear explicitly in but can be easily derived from [31]; for further details, see [5, Lemma 1.11]. We can now present the desired equivalent formulations of the curve-flat property.

Proposition 2.1. Let $f : M \rightarrow N$ be a Lipschitz map between metric spaces. The following assertions are equivalent :

- (i) f is curve-flat;
- (ii) for every Lipschitz curve $\gamma : K \rightarrow M$, $\mathcal{H}^1(f \circ \gamma(K)) = 0$.
- (iii) for every bi-Lipschitz curve $\gamma : K \rightarrow M$, $\mathcal{H}^1(f \circ \gamma(K)) = 0$.
- (iv) for every Lipschitz curve $\gamma : K \rightarrow M$ and every Lipschitz map $g : N \rightarrow \mathbb{R}$, $\mathcal{H}^1(g \circ f \circ \gamma(K)) = 0$.

Proof. The equivalence (i) \iff (ii) follows from the Area formula. The implications (ii) \implies (iii) and (ii) \implies (iv) are obvious. Let us prove the contrapositive of (iii) \implies (ii). Assume that there exists a Lipschitz curve $\gamma : K \rightarrow M$ such that $\mathcal{H}^1(f \circ \gamma(K)) > 0$. By Kirchheim's lemma, there exists a compact subset A of K such that $\mathcal{H}^1(A) = \lambda(A) > 0$ and $f \circ \gamma|_A$ is bi-Lipschitz. Obviously $(\gamma|_A)^{-1} = (f \circ \gamma|_A)^{-1} \circ f|_{\gamma(A)}$ is Lipschitz. So $\gamma : A \rightarrow M$ is the desired bi-Lipschitz curve. The proof of (iv) \implies (ii) is similar: Let $\gamma : K \rightarrow M$ a Lipschitz curve such that $\mathcal{H}^1(f \circ \gamma(K)) > 0$. Apply Kirchheim's lemma to obtain $A \subset K$ such that $\mathcal{H}^1(A) > 0$ and $f \circ \gamma|_A$ is bi-Lipschitz. Then define $g := (f \circ \gamma|_A)^{-1} : f \circ \gamma(A) \rightarrow A$ and extend g to N using McShane-Whitney's extension theorem. We denote this extension by \tilde{g} . Now observe that \tilde{g} is Lipschitz on N and such that $\mathcal{H}^1(\tilde{g} \circ f \circ \gamma(A)) = \mathcal{H}^1(A) > 0$, which is precisely the negation of (iv). \square

Proposition 2.2. For any metric space M , the space $CF_0(M)$ is weak*-closed in $Lip_0(M)$.

Proof. By Krein-Smulian theorem, it is enough to show that $B_{CF_0(M)}$ is w^* -closed. So let $f \in \overline{B_{CF_0(M)}}^{w^*}$. Clearly, $\|f\|_L \leq 1$. Let us assume that $f \notin CF_0(M)$. Then, by Proposition 2.1 there exists a Lipschitz curve $\gamma : K \rightarrow M$ such that $\mathcal{H}^1(f \circ \gamma(K)) > 0$. Thus by Kirchheim's lemma we may as well assume that $\lambda(K) > 0$ and $f \circ \gamma$ is bi-Lipschitz. Let $C > 0$ be the co-Lipschitz constant of $f \circ \gamma$, i.e., $C|x - y| \leq |f \circ \gamma(x) - f \circ \gamma(y)|$ for every $x, y \in K$. We apply Lebesgue's density theorem to obtain a point $a \in K \setminus \{\max K\}$ where the density of K is 1. Now, given any $g \in B_{CF_0(M)}$ and $t \in K \cap (a, \max K]$, note that if h is a McShane-Whitney's extension of $g \circ \gamma$ to $[a, \max K]$ then :

$$\frac{|g \circ \gamma(t) - g \circ \gamma(a)|}{|t - a|} \leq \frac{1}{|t - a|} \int_a^t |h'(s)| d\lambda(s) \leq \|\gamma\|_L \frac{\lambda([a, t] \setminus K)}{|t - a|}.$$

Since the density of K at a is 1, this last quantity goes to 0 as $t \rightarrow a+$. In particular,

$$\left\{ \varphi \in \text{Lip}_0(M) : \frac{|\langle \delta(\gamma(t)) - \delta(\gamma(a)), \varphi \rangle|}{|t - a|} > \frac{C}{2} \right\}$$

is a w^* -open neighborhood of f which does not intersect with $B_{\text{CF}_0(M)}$ if t is chosen close enough to a . Thus $f \notin \overline{B_{\text{CF}_0(M)}}^{w^*}$. This contradiction shows that $B_{\text{CF}_0(M)}$ is w^* -closed. \square

We conclude this section with the next central definition.

Definition 2.3. A metric space M is said to be *purely 1-unrectifiable* (p1u for short) if $\mathcal{H}^1(\gamma(A)) = 0$ for every $A \subset \mathbb{R}$ and every Lipschitz map $\gamma : A \rightarrow M$.

It is well known that, thanks to Kirchheim's lemma, M is purely 1-unrectifiable if and only if M contains no *curve fragments*, that is, no image of a bi-Lipschitz embedding $\gamma : K \rightarrow M$, where $K \subset \mathbb{R}$ is compact with $\lambda(K) > 0$. This, together with Proposition 2.1, shows that M is p1u if and only if $\text{Id}_M \in \text{CF}_0(M, M)$. We even get the following curious corollary.

Corollary 2.4. *If M is any metric space such that $\text{CF}_0(M)$ is w^* -dense in $\text{Lip}_0(M)$, then M is purely 1-unrectifiable.*

Proof. We get by Proposition 2.2 and by the hypothesis that $\text{CF}_0(M) = \text{Lip}_0(M)$. Thus, applying Proposition 2.1 (iv), we get $\text{Id}_M \in \text{CF}_0(M, M)$, and so M is p1u. \square

2.4. Preliminaries on martingales. Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space and let X be a Banach space. We denote by $L_1(\Omega, \mathcal{A}, \mathbb{P}; X)$ the space of (equivalence classes of) Bochner measurable functions $f : \Omega \rightarrow X$ such that $\int_{\Omega} \|f\|_X d\mathbb{P} < \infty$ equipped as usual with the norm

$$\|f\|_{L_1(\Omega, \mathcal{A}, \mathbb{P}; X)} = \mathbb{E}[\|f\|_X] = \int_{\Omega} \|f\|_X d\mathbb{P}.$$

In the sequel, we will suppress notation and write simply $L_1(X)$, or $L_1(\mathcal{A}; X)$ when the σ -algebra \mathcal{A} needs to be emphasized.

We recall that a sequence $(M_n)_{n=0}^{\infty}$ in $L_1(\mathcal{A}; X)$ is called a *martingale* if there exists an increasing sequence $(\mathcal{A}_n)_{n=0}^{\infty}$ of σ -subalgebras of \mathcal{A} (called a *filtration*) such that for each $n \geq 0$, M_n is \mathcal{A}_n -measurable and satisfies

$$M_n = \mathbb{E}^{\mathcal{A}_n}(M_{n+1}),$$

where $\mathbb{E}^{\mathcal{A}_n}$ denotes the X -valued conditional expectation relative to \mathcal{A}_n (see e.g. [38, Section 1.2]). We say moreover that $(M_n)_{n=0}^{\infty}$ is *uniformly integrable* if the sequence of non-negative random variables $(\|M_n\|_X)_{n=0}^{\infty}$ is uniformly integrable. More precisely, this means that $(\|M_n\|_X)_{n=0}^{\infty}$ is bounded in $L_1(\mathbb{R})$ and that, for any $\varepsilon > 0$, there is a $\delta > 0$ such that

$$\forall A \in \mathcal{A}, \quad \mathbb{P}(A) < \delta \implies \sup_{n \geq 0} \int_A \|M_n\|_X d\mathbb{P} < \varepsilon.$$

We will need the following characterization of Radon-Nikodým operators.

Lemma 2.5. *Let X and Y be Banach spaces. A bounded operator $T : X \rightarrow Y$ is Radon-Nikodým if and only if for every uniformly integrable X -valued martingale $(M_n)_n$, $(T(M_n))_n$ converges in $L_1(Y)$.*

The proof of the “if” implication follows the exact same lines as the proof of [38, Proposition 1.33]. We include it below for convenience.

Proof. When $(M_n) \subset L_1(X)$ is uniformly integrable the hypothesis and Vitali Convergence Theroem (see [40, Exercise 10 (b) on p. 133]) imply that the sequence $(T(M_n))_n$ converges in $L_1(Y)$.

It remains to prove the “if” direction. To this end, we fix a bounded X -valued martingale $(M_n)_n$ and claim that $(T(M_n))_n$ converges almost everywhere. For a fixed $t > 0$, define the following stopping time:

$$S_t = \begin{cases} \inf\{n \geq 0 : \|M_n\| > t\} & \text{if } \sup_n \|M_n\| > t, \\ \infty & \text{otherwise.} \end{cases}$$

By [38, Lemma 1.32], $(M_{n \wedge S_t})_n$ is a uniformly integrable martingale. Hence, by assumption, $(T(M_{n \wedge S_t}))_n$ converges in $L_1(Y)$ and, thanks to [38, Theorem 1.30], converges almost everywhere. Notice that if $\omega \in \Omega$ is such that $S_t(\omega) = \infty$, then $M_{n \wedge S_t} = M_n$ for every $n \in \mathbb{N}$. We will prove that $\mathbb{P}(\{S_t = \infty\})$ goes to 1 as t goes to infinity, which readily implies that $(T(M_n))_n$ itself converges almost everywhere, and therefore concludes the proof. By Doob’s maximal inequality (see [38, Corollary 1.29]), one has:

$$\mathbb{P}(\{S_t < \infty\}) = \mathbb{P}(\{\sup_n \|M_n\| > t\}) \leq \frac{\sup_n \|M_n\|_{L_1(X)}}{t}.$$

This probability can be made arbitrarily small by choosing t large enough, which implies that $\mathbb{P}(\{S_t < \infty\})$ goes to 0 as t goes to infinity, as desired. \square

2.5. General preliminaries. Here we aim to state a general version of a well known principle that weakly null normalized sequences stay away from any finite dimensional subspace.

Lemma 2.6. *Let X be a Banach space. Let σ be a Hausdorff locally convex topology on X such that $\|\cdot\|$ is lower semi-continuous and let $(D_n) \subset X$ be a bounded sequence such that $D_n \rightarrow 0$ in σ . Then for every finite dimensional $Y \subseteq X$, we have that*

$$\liminf_n \text{dist}(D_n, Y) \geq \frac{\inf_n \|D_n\|}{2}.$$

Proof. Let $(y_{n_k}) \subset Y$ be such that $\|D_{n_k} - y_{n_k}\| \rightarrow \liminf_n \text{dist}(D_n, Y)$. Moreover, by finite dimension of Y and the fact that σ is locally convex and Hausdorff we can assume that $y_{n_k} \rightarrow y$ in norm. It follows that

$$\|D_{n_k} - y\| \rightarrow \liminf_n \text{dist}(D_n, Y).$$

And so

$$(1) \quad \|y\| \leq \liminf_n \text{dist}(D_n, Y)$$

by the lower semi-continuity of $\|\cdot\|$ since $D_{n_k} - y \rightarrow -y$ in σ by the hypothesis. Also

$$(2) \quad \inf_n \|D_n\| - \|y\| \leq \liminf_k (\|D_{n_k}\| - \|y\|) \leq \lim_k \|D_{n_k} - y\| = \liminf_n \text{dist}(D_n, Y).$$

Thus summing (1) and (2) gives

$$\frac{\inf_n \|D_n\|}{2} \leq \liminf_n \text{dist}(D_n, Y).$$

□

It is well known that topologies which satisfy the hypothesis of the lemma are for instance the weak topology $w = \sigma(X, X^*)$ or, more generally, any topology $\sigma(X, S)$ where S is subspace of X^* which is norming for X , meaning that for all $x \in X$, $\|x\|_X = \sup\{x^*(x), x^* \in B_{X^*} \cap S\}$.

In particular, for a Banach space X , denote $S \subseteq L_\infty(\Omega, \mathcal{A}, \mathbb{P}, X^*)$ the normed linear space of X^* -valued simple functions, i.e. measurable functions with only finitely many distinct values. This space is easily seen to embed isometrically into $(L_1(\mathcal{A}, X))^*$ (see [17, p. 97]).

Corollary 2.7. *Let X be a Banach space and let $(D_n) \subset L_1(X)$ be a bounded sequence such that $D_n \rightarrow 0$ in $\sigma(L_1(X), S)$. Then for every finite dimensional $Y \subseteq L_1(X)$, we have that*

$$\liminf_n \text{dist}(D_n, Y) \geq \frac{\inf_n \|D_n\|}{2}.$$

Proof. It follows from the definition of the Bochner-Lebesgue space that the space S of simple functions is norming for $L_1(X)$. As we already mentioned it can be easily deduced that $\|\cdot\|$ is $\sigma(L_1(X), S)$ -lower semi-continuous. We can thus apply Lemma 2.6 to get the desired conclusion. □

3. DUNFORD-PETTIS LIPSCHITZ OPERATORS

In this section we prove the first “half” of Theorem 1.1.

Theorem 3.1. *Let M, N be complete pointed metric spaces and let $f \in \text{Lip}_0(M, N)$. Then the following properties are equivalent:*

- (1) f is curve-flat,
- (2) \hat{f} is Dunford-Pettis,
- (3) \hat{f} does not fix any copy of L_1 .

The majority of this section is dedicated to showing that (1) implies (2) in Theorem 3.1. Indeed, as we will later see in its proof, (2) implies (3) is direct since L_1 contains a normalized weakly null sequence, while (3) implies (1) follows from Kirchem’s Lemma and Godard’s work on Lipschitz-free spaces over subsets of the real line.

This section is divided into three subsections. In the first one we address (1) implies (2) of Theorem 3.1 in the case where M is any compact metric space, leveraging a profound result established by D. Bate [10]. In the second, we deal with the general case thanks to a compact reduction principle along the lines of [9]. In the last subsection we use both previous results to finally prove Theorem 3.1.

3.1. f is curve-flat implies \hat{f} is Dunford-Pettis: The compact case. As previously mentioned, a key ingredient in this setting is a result due to D. Bate [10]. To ensure clarity and ease of reference, we will present this result in its entirety. First, let us introduce some notation.

Definition 3.2. Given $\varepsilon > 0$ and $\theta > 0$, we say that a Lipschitz map $f: M \rightarrow N$ is ε -flat at radius θ if $d(f(x), f(y)) \leq \varepsilon d(x, y)$ for all $x, y \in M$ with $d(x, y) \leq \theta$. It is ε -flat if it is ε -flat at radius θ for some $\theta > 0$. Denote by $\text{lip}_0^\varepsilon(M)$ the set of ε -flat Lipschitz functions in $\text{Lip}_0(M)$. Note that if $\varepsilon_1 < \varepsilon_2$, then $\text{lip}_0^{\varepsilon_1}(M) \subset \text{lip}_0^{\varepsilon_2}(M)$.

In the next lemma, if $t \in K \subset \mathbb{R}$ and $\gamma : K \rightarrow M$ is Lipschitz, then $\text{Lip}(\gamma, t)$ stands for the *pointwise Lipschitz constant* of γ at t , defined as:

$$\text{Lip}(\gamma, t) := \limsup_{x \rightarrow t} \frac{d(\gamma(t), \gamma(x))}{|t - x|}.$$

Lemma 3.3 ([10, Lemma 3.4]). *Let M be a compact metric space, let X be a Banach space which contains M isometrically. Let $\delta > 0$ and let $x^* \in X^*$. Suppose that every bi-Lipschitz curve $\gamma : K \rightarrow X$ satisfying*

$$(x^* \circ \gamma)'(t) \geq \delta \|x^*\| \text{Lip}(\gamma, t), \quad \mathcal{H}^1\text{-a.e. in } t \in K$$

also satisfies $\mathcal{H}^1(\gamma(K) \cap M) = 0$.

Then, for every $\varepsilon > 0$, there exists a Lipschitz function $g : X \rightarrow \mathbb{R}$ such that:

- $\frac{|g(x) - g(y)|}{\|x - y\|} \leq \frac{|x^*(x) - x^*(y)|}{\|x - y\|} + 3\|x^*\|\delta$ for every $x \neq y \in X$. In particular, $\|g\|_L \leq \|x^*\|(1 + 3\delta)$,
- $|x^*(x) - g(x)| < \varepsilon$ for all $x \in X$, and
- there exists $\theta > 0$ such that $g|_M$ is 3δ -flat at radius θ .

Clearly, if $0 \in M \subset X$, the function g from the above lemma can be also assumed to satisfy $g(0) = 0$. This is how we will use it below. The next lemma and its proof are essentially contained in Section 3 of [10]. We present a direct proof here for clarity and convenience.

Lemma 3.4. *If M is compact then $B_{\text{CF}_0(M)} \subset \overline{B_{\text{lip}_0^\tau(M)}}^{w^*}$ for every $\tau > 0$.*

Proof. Let $f \in \text{CF}_0(M)$ with $\|f\|_L \leq 1$, and let $\tau > 0$. Since the weak* topology coincides with the topology of pointwise convergence on norm-bounded subsets of $\text{Lip}_0(M)$, it suffices to show that for every $\eta > 0$, there exists a function $g \in \text{lip}_0^\tau(M)$ with $\|g\|_{\text{Lip}} \leq 1$ such that $|f(x) - g(x)| < 2\eta$ for all $x \in M$.

We start by showing that the linear extension of f , i.e. the map $\bar{f} : \mathcal{F}(M) \rightarrow \mathbb{R}$, satisfies the hypothesis of Lemma 3.3 for every $\delta > 0$. Suppose that $\gamma : K \rightarrow \mathcal{F}(M)$ is a bi-Lipschitz curve satisfying

$$(\bar{f} \circ \gamma)'(t) \geq \delta \|\bar{f}\| \text{Lip}(\gamma, t), \quad \mathcal{H}^1\text{-a.e. in } t \in K.$$

We may of course assume that $f \neq 0$, so that the right-hand side is everywhere strictly positive. Now, arguing by contradiction, suppose that $\mathcal{H}^1(\gamma(K) \cap M) > 0$. Clearly $A := \gamma^{-1}(\gamma(K) \cap M) \subset K$ has positive measure, and the curve $\gamma|_A : A \rightarrow M$ is bi-Lipschitz with values in M . Since f is curve-flat, we obtain that $(f \circ \gamma|_A)'(t) = 0$ almost everywhere in A , which is a contradiction.

Now, we can apply Lemma 3.3 to \bar{f} , with $\varepsilon = \eta$ and $\delta = \delta(\tau, \eta) > 0$ to be determined later, to obtain $\tilde{g} \in \text{lip}_0^{3\delta}(M)$ with $\|\tilde{g}\|_L \leq (1 + 3\delta)$ and such that $|\tilde{g}(x) - f(x)| < \eta$ for all $x \in M$. We only need to normalize \tilde{g} to get the result. Indeed, consider $g = \frac{1}{1+3\delta}\tilde{g}$. Then g also belongs to $\text{lip}_0^{3\delta}(M)$ and now $\|g\|_L \leq 1$. Finally, we have for any $x \in M$:

$$\begin{aligned} |g(x) - f(x)| &\leq |\tilde{g}(x) - f(x)| + \left(1 - \frac{1}{1+3\delta}\right)|\tilde{g}(x)| \\ &\leq \eta + \left(1 - \frac{1}{1+3\delta}\right)(1+3\delta)\text{rad}(M) \\ &= \eta + 3\delta\text{rad}(M). \end{aligned}$$

Consequently, for $\delta = \min \left\{ \frac{\tau}{3}, \frac{\eta}{3\text{rad}(M)} \right\}$, we obtained the desired conclusion. \square

We can now prove the main result of this subsection. We will employ a gliding hump type argument inspired by Kalton's proof that the Lipschitz-free space over a snowflaked metric space has the Schur property ([29, Theorem 4.6]). For the next proof, recall the notation $[S]_\delta := \{x \in M : d(x, S) \leq \delta\}$ for S a non empty subset of M and $\delta > 0$.

Proposition 3.5. *Let M and N be metric spaces with M compact, and let $f \in \text{Lip}_0(M, N)$. If f is curve-flat, then $\widehat{f}: \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ is Dunford-Pettis.*

Proof. Suppose that $f: M \rightarrow N$ is a curve-flat Lipschitz map with $\|f\|_{\text{Lip}} = 1$. We will prove by contradiction that $\widehat{f}: \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ is Dunford-Pettis. Suppose there exists a weakly null sequence $(\gamma_k)_k \subset \mathcal{F}(M)$ such that $\|\widehat{f}(\gamma_k)\| = 1$ for all $k \in \mathbb{N}$. We may assume without loss of generality that the support of γ_k is finite for all $k \in \mathbb{N}$. Choose a decreasing sequence of positive numbers $(\varepsilon_n)_n$ such that $\sum_n \varepsilon_n < \frac{1}{4}$. We will construct, by induction, a bounded sequence of functions $(\varphi_n)_n \subset \text{Lip}_0(M)$ and a subsequence $(\gamma_{k_n})_n$ such that:

- (1) φ_n is $(\sum_{i=1}^n \varepsilon_i)$ -flat at some positive radius $0 < \theta_n < 1$,
- (2) $\|\varphi_n\|_L \leq 1 + \sum_{i=1}^n \varepsilon_i$, and
- (3) $\langle \varphi_n, \gamma_{k_j} \rangle \geq \frac{1}{2} - \sum_{i=1}^n \varepsilon_i$ for all $j = 1, \dots, n$.

For the initial step, let $\gamma_{k_1} = \gamma_1$. Since $\|\widehat{f}(\gamma_1)\| = 1$, there exists a Lipschitz function $g_1 \in \text{Lip}_0(N)$ such that $\langle g_1, \widehat{f}(\gamma_1) \rangle = 1 = \langle g_1 \circ f, \gamma_1 \rangle$. The function $g_1 \circ f$ is curve-flat, so by Lemma 3.4 there exists $\varphi_1 \in B_{\text{Lip}_0(M)}$ such that $\langle \varphi_1, \gamma_1 \rangle \geq 1 - \varepsilon_1$ and such that φ_1 is ε_1 -flat at some radius $\theta_1 \in (0, 1)$.

Let us fix $n \in \mathbb{N}$ and suppose we have constructed γ_{k_j} and φ_j for every $j \leq n$. Fix $\tau > 0$ to be determined later. By compactness of M , there exists a finite set E_{n+1} , containing the support of γ_{k_j} for all $j = 1 \leq n$, and such that $M = [E_{n+1}]_\tau$.

Since $f(E_{n+1})$ is finite as well, $\mathcal{F}(f(E_{n+1}))$ is finite-dimensional. Given that $(\widehat{f}(\gamma_k))_k$ is a normalized weakly null sequence, Lemma 2.6 gives that

$$\liminf_k \text{dist}(\widehat{f}(\gamma_k), \mathcal{F}(f(E_{n+1}))) \geq \frac{1}{2}.$$

Therefore, combining with the Hahn-Banach separation theorem, we can find $k_{n+1} > k_n$ and $g_{n+1} \in \text{Lip}_0(N)$ with $\|g_{n+1}\|_L \leq 1$, $g_{n+1}(f(E_{n+1})) = \{0\}$ and $\langle g_{n+1}, \widehat{f}(\gamma_{k_{n+1}}) \rangle > \frac{1}{2} - \tau$. Since $(\gamma_k)_k$ is weakly null, we may assume as well that $\langle \varphi_n, \gamma_{k_{n+1}} \rangle < \tau$.

The map $g_{n+1} \circ f \in \text{Lip}_0(M)$ is curve-flat, and satisfies

$$(g_{n+1} \circ f)(u) = 0, \quad \text{for all } u \in E_{n+1}, \quad \text{and } \langle g_{n+1} \circ f, \gamma_{k_{n+1}} \rangle > \frac{1}{2} - \tau.$$

Therefore, thanks to Lemma 3.4, it can be approximated by a function $h_{n+1} \in \text{Lip}_0(M)$ with $\|h_{n+1}\|_L \leq 1$ and such that:

- $|h_{n+1}(u)| < 2\tau$, for all $u \in M$,
- $\langle h_{n+1}, \gamma_{k_{n+1}} \rangle > \frac{1}{2} - 2\tau$,
- $\langle h_{n+1}, \gamma_{k_j} \rangle < \tau$ for all $j = 1, \dots, n$, and
- h_{n+1} is ε_{n+1} -flat at some radius θ_{n+1} , which can be chosen so that $0 < \theta_{n+1} \leq \theta_n$.

Now, define $\varphi_{n+1} = \varphi_n + h_{n+1}$. Let us check that φ_{n+1} satisfies conditions (1), (2) and (3). First, condition (1) is clear since φ_n is $(\sum_{i=1}^n \varepsilon_i)$ -flat and h_{n+1} is ε_{n+1} -flat. Next we estimate its Lipschitz constant. Let $x, y \in M$. If $d(x, y) < \theta_n$,

then, since φ_n is $(\sum_{i=1}^n \varepsilon_i)$ -flat at radius θ_n :

$$\begin{aligned} |\varphi_{n+1}(x) - \varphi_{n+1}(y)| &\leq |\varphi_n(x) - \varphi_n(y)| + |h_{n+1}(x) - h_{n+1}(y)| \\ &\leq \left(\sum_{i=1}^n \varepsilon_i + 1 \right) d(x, y) \end{aligned}$$

Otherwise, if $d(x, y) \geq \theta_n$ then:

$$\begin{aligned} |\varphi_{n+1}(x) - \varphi_{n+1}(y)| &\leq |\varphi_n(x) - \varphi_n(y)| + |h_{n+1}(x)| + |h_{n+1}(y)| \\ &\leq \left(1 + \sum_{i=1}^n \varepsilon_i \right) d(x, y) + 4 \frac{\tau}{\theta_n} d(x, y) \\ &= \left(1 + \sum_{i=1}^n \varepsilon_i + 4 \frac{\tau}{\theta_n} \right) d(x, y). \end{aligned}$$

Therefore, we conclude that $\|\varphi_{n+1}\|_L \leq \left(1 + \sum_{i=1}^n \varepsilon_i + 4 \frac{\tau}{\theta_n} \right)$.

Finally, let us examine the estimates in (3). If $j \leq n$:

$$\langle \varphi_{n+1}, \gamma_{k_j} \rangle \geq \langle \varphi_n, \gamma_{k_j} \rangle - |\langle h_{n+1}, \gamma_{k_j} \rangle| > \frac{1}{2} - \sum_{i=1}^n \varepsilon_i - \tau.$$

Simultaneously, we have:

$$\langle \varphi_{n+1}, \gamma_{k_{n+1}} \rangle \geq \langle h_{n+1}, \gamma_{k_{n+1}} \rangle - |\langle \varphi_n, \gamma_{k_{n+1}} \rangle| > \frac{1}{2} - 3\tau.$$

Choosing τ so that $4 \frac{\tau}{\theta_n} < \varepsilon_{n+1}$ yields condition (2). In particular, this implies $3\tau < \varepsilon_{n+1}$, thereby proving (3) as well. The induction process is now complete.

To conclude, passing to a subsequence if necessary, we may assume that $(\varphi_n)_n$ converges to a Lipschitz map $\tilde{\varphi} \in \text{Lip}_0(M)$ in the weak* topology. For a fixed $j \in \mathbb{N}$, we have that $\langle \varphi_n, \gamma_{k_j} \rangle > \frac{1}{2} - \sum_{k=1}^{\infty} \varepsilon_k > \frac{1}{4}$ whenever $n \geq j$. Therefore, $\langle \tilde{\varphi}, \gamma_{k_j} \rangle > \frac{1}{4}$ for all $j \in \mathbb{N}$. This is a contradiction with the fact that $(\gamma_{k_j})_j$ is weakly null. \square

3.2. Compact determination of Dunford-Pettis operators. Once we have established that (1) implies (2) in Theorem 3.1 for compact metric spaces, we proceed to extend this result to general metric spaces thanks to the following compact reduction result:

Proposition 3.6. *Let M, N be complete metric spaces, and let $f \in \text{Lip}_0(M, N)$. The property “ \widehat{f} is Dunford-Pettis” is compactly determined, that is, \widehat{f} is Dunford-Pettis if and only if for every compact $K \subset M$ such that $0 \in K$, $\widehat{f}|_K$ is Dunford-Pettis.*

Proof. One direction is clear: If \widehat{f} is Dunford-Pettis, then so is $\widehat{f}|_K$ for every $K \subset M$ compact containing 0. The converse will follow from the compact reduction principle described in [9]. Suppose that $f : M \rightarrow N$ is a Lipschitz map such that $\widehat{f}|_K$ is Dunford-Pettis for every $K \subset M$ compact containing 0. Let $(\gamma_n)_n \subset \mathcal{F}(M)$ be weakly null. We will prove that $(\widehat{f}(\gamma_n))_n$ is norm-null. Let $W = \{\gamma_n : n \in \mathbb{N}\}$ and fix $\varepsilon > 0$. By [9, Theorem 2.3], there exist a compact $K \subset M$ containing 0 and a linear mapping $T : \text{span}(W) \rightarrow \mathcal{F}(K)$ such that

- $\|\mu - T\mu\| \leq \varepsilon$ for all $\mu \in W$, and

- there is a sequence of bounded linear operators $T_k : \mathcal{F}(M) \rightarrow \mathcal{F}(M)$ such that $T_k \rightarrow T$ uniformly on W .

A classical exchange of limits argument (see [9, Corollary 2.4] for more details) yields that the sequence $(T(\gamma_n))_n$ is weakly null in $\mathcal{F}(K)$. Since $\widehat{f}|_K$ is Dunford-Pettis, $\|\widehat{f}(T\gamma_n)\| \rightarrow 0$. So there exists $n_0 \in \mathbb{N}$ such that for every $n \geq n_0$, $\|\widehat{f}(T\gamma_n)\| \leq \varepsilon$. By the triangle inequality we obtain:

$$\forall n \geq n_0, \quad \|\widehat{f}(\gamma_n)\| \leq \|\widehat{f}\| \|\gamma_n - T\gamma_n\| + \|\widehat{f}(T\gamma_n)\| \leq (\|f\|_L + 1)\varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, this proves that $(\widehat{f}(\gamma_n))_n$ is norm-null, and therefore \widehat{f} is Dunford-Pettis. \square

We now have all the ingredients to prove the main result of the section.

3.3. Proof of Theorem 3.1.

Proof of Theorem 3.1. The implication (1) \implies (2) follows from Propositions 3.5 and 3.6, while (2) \implies (3) is direct since the unit sphere of L_1 contains a weakly null sequence.

To show that (3) \implies (1), we proceed by contrapositive. Thus, suppose that f is not curve-flat. By Proposition 2.1 there exists $K \subset \mathbb{R}$ compact, $\gamma : K \rightarrow M$ bi-Lipschitz such that $\mathcal{H}^1(f \circ \gamma(K)) > 0$. By Kirchheim's lemma, there exists A compact, $A \subset K$ such that $\lambda(A) > 0$ and $f \circ \gamma|_A$ is bi-Lipschitz. Now, thanks to Godard's work [25], $\mathcal{F}(A)$ is isomorphic to L_1 . As $\gamma|_A$ and $f \circ \gamma|_A$ are bi-Lipschitz, $\widehat{\gamma|_A}$ and $\widehat{f \circ \gamma|_A}$ are isomorphisms from $\mathcal{F}(A)$ onto $\mathcal{F}(\gamma(A))$ and $\mathcal{F}(f \circ \gamma(A))$, respectively. Since $\widehat{f \circ \gamma} = \widehat{f} \circ \widehat{\gamma}$, it follows that $\widehat{f}|_{\mathcal{F}(\gamma(A))}$ is an isomorphism from $\mathcal{F}(\gamma(A))$ onto $\mathcal{F}(f \circ \gamma(A))$. This proves that \widehat{f} fixes a copy of L_1 . \square

Remark 3.7. It has been proved in [12] that if $T : L_1 \rightarrow L_1$ is not Dunford-Pettis, then it fixes a copy of $\ell_1(\ell_2)$. The equivalence of (2) and (3) in Theorem 3.1 shows that linearizations of Lipschitz maps which are not Dunford-Pettis satisfy a stronger property.

4. RADON-NIKODÝM LIPSCHITZ OPERATORS

In this section, we prove the second ‘‘half’’ of Theorem 1.1.

Theorem 4.1. *Let M, N be complete pointed metric spaces and let $f \in \text{Lip}_0(M, N)$. Then the following properties are equivalent:*

- (1) f is curve-flat,
- (2) \widehat{f} is Radon-Nikodým,
- (3) \widehat{f} does not fix any copy of L_1 .

Note that the proof of (2) \implies (3) is trivial as there exists a L_1 -bounded martingale in $L_1(L_1)$ which does not converge almost everywhere. On the other hand, (3) \implies (1) is shown in Theorem 3.1 already. Hence, it only remains to deal with the implication (1) \implies (2), which will be the focus of the present section.

The strategy to show (1) \implies (2) is the same as the one we followed in Section 3: first we show that it holds in compact metric spaces, and then we prove that the property ‘‘ \widehat{f} is Radon-Nikodým’’ is compactly determined. In both results we will be dealing with Banach space valued martingales and quasi-martingales.

Recall that a stochastic process $(S_n)_n$ in $L_1(\mathcal{A}, X)$ adapted to a filtration (\mathcal{A}_n) is said to be a *quasi-martingale* if

$$\sum_{n=1}^{\infty} \|\mathbb{E}^{\mathcal{A}_{n-1}}(S_n - S_{n-1})\| < \infty.$$

We refer to [38, Remark 2.16] for more details about quasi-martingales.

Lemma 4.2. *Let X be a Banach space, (M_n) be a bounded and uniformly integrable quasi-martingale in $L_1(X)$ and denote $D_n := M_n - M_{n-1}$ (with $M_0 := 0$). Then $D_n \rightarrow 0$ in $\sigma(L_1(X), S)$ where S denotes the space of simple functions in $L_\infty(\Omega, X^*) \subseteq L_1(\Omega, X)^*$.*

Proof. Let $g \in S$ be fixed. Note that the set $\{g(\omega) : \omega \in \Omega\}$ is a finite set; thus generates a finite dimensional subspace E of X^* . Let us fix a basis $\{x_1^*, \dots, x_k^*\}$ of E . Due to finite dimensionality, there exists $C > 0$ such that $\max_i |\lambda_i| \leq C \|x^*\|$ for every $x^* = \sum_{i=1}^k \lambda_i x_i^* \in E$. If $\omega \in \Omega$ then we can write $g(\omega) = \sum_{i=1}^k \lambda_i(\omega) x_i^*$ for some $\lambda_i(\omega) \in \mathbb{R}$, $i = 1, \dots, k$. Notice that $(x_i^* \circ M_n)_n$ is a scalar valued, bounded, and uniformly integrable quasi-martingale for each $i = 1, \dots, k$. Therefore it converges in L_1 (see [38, Remark 2.16]). This readily implies that each $(x_i^* \circ D_n)_n$ goes to 0 in L_1 as $n \rightarrow \infty$. We conclude with the following estimates:

$$\begin{aligned} |\langle g, D_n \rangle| &\leq \int_{\Omega} |\langle g(\omega), D_n(\omega) \rangle| d\mathbb{P}(\omega) \\ &\leq \int_{\Omega} \sum_{i=1}^k |\lambda_i(\omega)| \cdot |x_i^* \circ D_n(\omega)| d\mathbb{P}(\omega) \\ &\leq C \|g\|_{\infty} \sum_{i=1}^k \int_{\Omega} |x_i^* \circ D_n(\omega)| d\mathbb{P}(\omega) \rightarrow 0. \quad \square \end{aligned}$$

4.1. Proof of the compact case of Theorem 4.1. Here we will deal with the compact case. Up to the use of martingales, the proof will largely resemble that of Proposition 3.5 which established the implication “ $f \in \text{CF}_0$ ” \implies “ $\hat{f} \in \text{DP}$ ” (see Remark 4.5 for an in-depth comparison).

Proposition 4.3. *Let M and N be metric spaces with M compact, and let $f \in \text{Lip}_0(M, N)$. If f is curve-flat, then $\hat{f}: \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ is Radon-Nikodým.*

Proof. We assume that there is a bounded uniformly integrable martingale $(M_n) \subset L_1(\mathcal{F}(M))$ such that $(\hat{f} \circ M_n)_n$ does not converge in $L_1(\mathcal{F}(N))$.

Let $D_n = M_{n+1} - M_n$. Up to passing to a subsequence and multiplying by a positive constant, we may assume that $\|\hat{f} \circ D_n\| \geq 1$ for every $n \in \mathbb{N}$. We will employ a gliding hump argument to show that (M_n) is not bounded.

Let us fix $\varepsilon > 0$. First, using a standard approximation technique, we may assume that $(M_n)_{n=0}^{\infty}$ is adapted to a filtration $(\mathcal{A}_n)_{n=0}^{\infty}$ where each \mathcal{A}_n is finite (see the proof of [5, Proposition 4.3] for more details). Next, by density of the finitely supported elements in $\mathcal{F}(M)$, we may replace $(M_n)_n$ with a stochastic process $(\overline{M}_n)_n$ adapted to $(\mathcal{A}_n)_{n=0}^{\infty}$ satisfying

- \overline{M}_n is constant on each atom of \mathcal{A}_n and not just essentially constant,
- $\overline{M}_n(\omega)$ is finitely supported for every $n \in \mathbb{N}$ and $\omega \in \Omega$,
- $(\overline{M}_n)_n$ is $L_1(\mathcal{F}(M))$ -bounded,

- $\|\widehat{f} \circ \overline{D}_n\| \geq \frac{1}{2}$, where $\overline{D}_n := \overline{M}_{n+1} - \overline{M}_n$ for every $n \in \mathbb{N}$,
- for every $n \leq i$, $\|\mathbb{E}^{\mathcal{A}_n}(\overline{M}_i) - \overline{M}_n\|_{L_1} \leq \min(2^{-n}, \varepsilon)$.

Notice that $(\overline{M}_n)_n$ is still uniformly integrable and the last condition implies that $(\overline{M}_n)_{n=0}^\infty$ is a quasi-martingale. By Lemma 4.2, we have that the sequence $(\overline{D}_n)_n$ is $\sigma(L_1(\mathcal{F}(M)), S)$ -null, where S denotes the space of simple functions in $L_\infty(\Omega, \mathcal{F}(M))^* \subseteq L_1(\Omega, \mathcal{F}(M))^*$.

Now choose a decreasing sequence of positive numbers $(\varepsilon_n)_n$ in such a way that $\sum_n \varepsilon_n < \frac{1}{8}$. We will construct, by induction, a bounded sequence $(\varphi_n)_n$ and a subsequence $(\overline{D}_{k_n})_n$ such that:

- (1) $\varphi_n \in L_1(\mathcal{A}_{k_{n+1}}, \mathcal{F}(M))^* = L_\infty(\mathcal{A}_{k_{n+1}}, \text{Lip}_0(M))$,
- (2) For every $\omega \in \Omega$, $\varphi_n(\omega)$ is $(\sum_{i=1}^n \varepsilon_i)$ -flat at some positive radius $0 < \theta_n < 1$,
- (3) $\|\varphi_n\| \leq 1 + \sum_{i=1}^n \varepsilon_i$, and
- (4) $\langle \varphi_n, \overline{D}_{k_i} \rangle \geq \frac{1}{4} - \sum_{i=1}^n \varepsilon_i$ for all $i = 1, \dots, n$.

For the initial step, let $\overline{D}_{k_1} = \overline{D}_1$. Since $\|\widehat{f} \circ \overline{D}_1\| \geq \frac{1}{2}$, there exists a map $g_1 \in L_\infty(\mathcal{A}_2, \text{Lip}_0(N))$ with $\|g_1\| = 1$ such that $\langle g_1, \widehat{f} \circ \overline{D}_1 \rangle \geq \frac{1}{2}$. Notice that

$$\langle g_1, \widehat{f} \circ \overline{D}_1 \rangle = \int_{\Omega} \langle g_1(\omega), \widehat{f}(\overline{D}_1(\omega)) \rangle d\mathbb{P} = \int_{\Omega} \langle g_1(\omega) \circ f, \overline{D}_1(\omega) \rangle d\mathbb{P}.$$

For each $\omega \in \Omega$, the function $g_1(\omega) \circ f$ is curve-flat. So by Lemma 3.4, there exists a simple function $\varphi_1 \in L_\infty(\mathcal{A}_2, \text{Lip}_0(M))$ such that $\|\varphi_1\| \leq 1$, $\langle \varphi_1, \overline{D}_1 \rangle \geq \frac{1}{2} - \varepsilon_1$ and finally $\varphi_1(\omega)$ is ε_1 -flat at some radius $\theta_1 \in (0, 1)$ for every $\omega \in \Omega$.

Let us fix $n \in \mathbb{N}$ and suppose we have constructed \overline{D}_{k_i} and φ_i for every $i \leq n$. Fix $0 < \tau < \frac{1}{16}$ to be determined later. By compactness, there exists a finite set E_{n+1} containing

$$\bigcup_{i=1}^n \bigcup_{\omega \in \Omega} (\text{supp}(\overline{M}_{k_i}(\omega)) \cup \text{supp}(\overline{M}_{k_{i+1}}(\omega)))$$

and such that $M = [E_{n+1}]_\tau$.

Since $f(E_{n+1})$ is finite as well, $\mathcal{F}(f(E_{n+1}))$ is finite-dimensional. Notice that for each $g \in L_1(\mathcal{A}_{k_{n+1}}, \mathcal{F}(N))^* = L_\infty(\mathcal{A}_{k_{n+1}}, \text{Lip}_0(N))$, we have

$$\langle g, \widehat{f} \circ \overline{D}_k \rangle = \int_{\Omega} \langle g(\omega) \circ f, \overline{D}_k(\omega) \rangle d\mathbb{P} \rightarrow 0,$$

where the convergence is guaranteed thanks to Lemma 4.2. Thus, applying Corollary 2.7 to $(\widehat{f} \circ \overline{D}_k)_k$, $\mathcal{A}_{k_{n+1}}$ and $\mathcal{F}(f(E_{n+1}))$, we observe that

$$\liminf_k \text{dist}(\widehat{f} \circ \overline{D}_k, L_1(\mathcal{A}_{k_{n+1}}, \mathcal{F}(f(E_{n+1})))) \geq \frac{\inf_k \|\widehat{f} \circ \overline{D}_k\|}{2} \geq \frac{1}{4}.$$

So there exists $k_{n+1} > k_n$ such that $\text{dist}(\widehat{f} \circ \overline{D}_{k_{n+1}}, L_1(\mathcal{A}_{k_{n+1}}, \mathcal{F}(f(E_{n+1})))) > \frac{1}{4} - \tau$.

Since $\widehat{f} \circ \overline{D}_{k_{n+1}} \in L_1(\mathcal{A}_{k_{n+1}+1}, \mathcal{F}(N))$, by Hahn Banach separation theorem, there exists $g_{n+1} \in L_\infty(\mathcal{A}_{k_{n+1}+1}, \text{Lip}_0(N))$ with $\|g_{n+1}\| \leq 1$ such that

- $\langle g_{n+1}, L_1(\mathcal{A}_{k_{n+1}}, \mathcal{F}(f(E_{n+1}))) \rangle = \{0\}$, and
- $\langle g_{n+1}, \widehat{f} \circ \overline{D}_{k_{n+1}} \rangle > \frac{1}{4} - \tau$.

Knowing $(\overline{D}_k)_k$ is $\sigma(L_1(\mathcal{F}(M)), S)$ -null, we may assume that $|\langle \varphi_n, \overline{D}_{k_{n+1}} \rangle| < \tau$.

For every $\omega \in \Omega$, the map $g_{n+1}(\omega) \circ f \in \text{Lip}_0(M)$ is curve-flat, and satisfies that

$$(g_{n+1}(\omega) \circ f)(u) = 0, \quad \text{for all } u \in E_{n+1},$$

because g_{n+1} vanishes on $L_1(\mathcal{A}_{k_{n+1}}, \mathcal{F}(f(E_{n+1})))$. In particular, keeping in mind the construction of E_{n+1} , we also observe that

$$\langle g_{n+1}, \widehat{f} \circ \overline{D}_{k_i} \rangle = \int_{\Omega} \langle g_{n+1}(\omega) \circ f, \overline{D}_{k_i}(\omega) \rangle d\mathbb{P} = 0$$

for all $i = 1, \dots, n$. Moreover:

$$\langle g_{n+1}, \widehat{f} \circ \overline{D}_{k_{n+1}} \rangle = \int_{\Omega} \langle g_{n+1}(\omega) \circ f, \overline{D}_{k_{n+1}}(\omega) \rangle d\mathbb{P} > \frac{1}{4} - \tau.$$

Therefore, thanks to Lemma 3.4 and because g_{n+1} is finitely valued, there is $h_{n+1} \in L_{\infty}(\mathcal{A}_{k_{n+1}+1}, \text{Lip}_0(M))$ with $\|h_{n+1}\| \leq 1$ and such that:

- $|\langle h_{n+1}(\omega), \delta(u) \rangle| < 2\tau$, for all $u \in M$ and all $\omega \in \Omega$,
- $|\langle h_{n+1}, \overline{D}_{k_i} \rangle| < \tau$ for all $i = 1, \dots, n$,
- $\langle h_{n+1}, \overline{D}_{k_{n+1}} \rangle > \frac{1}{4} - 2\tau > \frac{1}{8}$, and
- $h_{n+1}(\omega)$ is ε_{n+1} -flat at some radius θ_{n+1} , which can be chosen so that $0 < \theta_{n+1} \leq \theta_n$, for every $\omega \in \Omega$.

Now, define $\varphi_{n+1} = \varphi_n + h_{n+1}$. Let us check that φ_{n+1} satisfies conditions (2) to (4). First, condition (2) is clear since $\varphi_n(\omega)$ is $(\sum_{i=1}^n \varepsilon_i)$ -flat at radius θ_{n+1} and $h_{n+1}(\omega)$ is ε_{n+1} -flat at radius θ_{n+1} .

Next, the estimation used in the proof of Proposition 3.5 shows that for every $\omega \in \Omega$, $\|\varphi_{n+1}(\omega)\|_L \leq \left(1 + \sum_{i=1}^n \varepsilon_i + 4\frac{\tau}{\theta_n}\right)$. Taking the maximum over $\omega \in \Omega$, we obtain (3) if τ is chosen small enough.

Finally, let us examine the estimates in (4). If $i \leq n$:

$$\langle \varphi_{n+1}, \overline{D}_{k_i} \rangle \geq \langle \varphi_n, \overline{D}_{k_i} \rangle - |\langle h_{n+1}, \overline{D}_{k_i} \rangle| > \frac{1}{4} - \sum_{i=1}^n \varepsilon_i - \tau.$$

Simultaneously, we also have:

$$\langle \varphi_{n+1}, \overline{D}_{k_{n+1}} \rangle \geq \langle h_{n+1}, \overline{D}_{k_{n+1}} \rangle - |\langle \varphi_n, \overline{D}_{k_{n+1}} \rangle| > \frac{1}{4} - 3\tau.$$

Choosing τ so that $4\frac{\tau}{\theta_n} < \varepsilon_{n+1}$ yields conditions (3) and (4). The induction process is now complete.

If we denote $h_1 = \varphi_1$ then it is clear that $\varphi_i = \sum_{n=1}^i h_n$ for every $i \in \mathbb{N}$. By the point (3) above we have that $\|\varphi_i\| \leq \frac{9}{8} =: C$ for every i . Observe that

$$\begin{aligned}
C \|\overline{D}_{k_i}\| &\geq \mathbb{E} \left(\left\langle \sum_{n=1}^i h_n, \overline{D}_{k_i} \right\rangle \right) \\
&= \mathbb{E} \left(\sum_{n=1}^i \langle h_n, \overline{M}_{k_{i+1}} \rangle \right) - \mathbb{E} \left(\sum_{n=1}^i \langle h_n, \overline{M}_{k_i} \rangle \right) \\
&= \mathbb{E} \left(\sum_{n=1}^i \langle h_n, \mathbb{E}^{\mathcal{A}_{k_{i+1}}}(\overline{M}_{k_{i+1}}) \rangle \right) - \mathbb{E} \left(\sum_{n=1}^i \langle h_n, \mathbb{E}^{\mathcal{A}_{k_n}}(\overline{M}_{k_i}) \rangle \right) \\
&= \mathbb{E} \left(\sum_{n=1}^i \langle h_n, \overline{D}_{k_n} \rangle \right) - \mathbb{E} \left(\sum_{n=1}^i \langle h_n, \overline{M}_{k_{n+1}} - \mathbb{E}^{\mathcal{A}_{k_{n+1}}}(\overline{M}_{k_{i+1}}) \rangle \right) \\
&\quad + \mathbb{E} \left(\sum_{n=1}^i \langle h_n, \overline{M}_{k_n} - \mathbb{E}^{\mathcal{A}_{k_n}}(\overline{M}_{k_i}) \rangle \right) \\
&\geq \frac{i}{8} - C \left(\sum_{n=1}^i \|\overline{M}_{k_{n+1}} - \mathbb{E}^{\mathcal{A}_{k_{n+1}}}(\overline{M}_{k_{i+1}})\| + \sum_{n=1}^i \|\overline{M}_{k_n} - \mathbb{E}^{\mathcal{A}_{k_n}}(\overline{M}_{k_i})\| \right) \\
&\geq \frac{i}{8} - C \left(\sum_{n=1}^i \frac{1}{2^{k_{n+1}}} + \frac{1}{2^{k_n}} \right) > \frac{i}{8} - 2C.
\end{aligned}$$

Since i was arbitrary, this contradicts the assumption that (\overline{M}_i) is bounded in $L_1(\mathcal{F}(M))$. \square

4.2. Compact determination of RN operators. Just like Dunford-Pettis operators, we claim that the property “ \widehat{f} is Radon-Nikodým” is compactly determined. The method of the proof follows the lines of the compact determination of the RNP for Lipschitz-free spaces (see [5, Corollary 4.5]).

Proposition 4.4. *Let M, N be complete metric spaces, and let $f \in \text{Lip}_0(M, N)$. The property “ \widehat{f} is Radon-Nikodým” is compactly determined, that is, $\widehat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ is Radon-Nikodým if and only if for every compact subset K of M containing 0, $f|_K$ is Radon-Nikodým.*

Proof. The “only if” implication is clear. We will prove the contrapositive of the “if” direction. Suppose that $f : M \rightarrow N$ is a Lipschitz map such that $\widehat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ is not Radon-Nikodým. By Lemma 2.5, there exists a bounded uniformly integrable martingale $(M_n)_n$ in $L_1(\mathcal{F}(M))$ such that $(\widehat{f}(M_n))_n$ does not converge in $L_1(\mathcal{F}(N))$. So there exists some $\delta > 0$ such that

$$(3) \quad \limsup_{n, m \rightarrow \infty} \|\widehat{f}(M_n) - \widehat{f}(M_m)\|_{L_1(\mathcal{F}(N))} > \delta.$$

By [5, Proposition 4.2] (with $\varepsilon = \frac{\delta}{4\|f\|_L}$, and using the fact that (M_n) has the *mean Kalton property* thanks to [5, Proposition 4.3 and Remark 4.4]), there exist a compact subset K of M containing 0, bounded linear operators $T_k : \mathcal{F}(M) \rightarrow \mathcal{F}(M)$ and a map $T : (M_n)_n \rightarrow L_1(\mathcal{F}(K))$ such that

$$(4) \quad \sup_{n \in \mathbb{N}} \|M_n - T(M_n)\|_{L_1(\mathcal{F}(M))} \leq \frac{\delta}{4\|f\|_L}$$

and

$$(5) \quad \limsup_{k \rightarrow \infty} \sup_{n \in \mathbb{N}} \|T_k(M_n) - T(M_n)\|_{L_1(\mathcal{F}(M))} = 0.$$

As demonstrated in the proof of [5, Corollary 4.5], we can observe that the sequence $(T(M_n))_n$ is a $L_1(\mathcal{F}(K))$ -bounded martingale using (4).

Moreover, the sequence $(T_k(M_n))_n$ is uniformly integrable for each k since each T_k is bounded operator. Also, condition (5) then implies that $(T(M_n))_n$ is an $L_1(\mathcal{F}(K))$ -uniformly integrable martingale. But we have

$$\limsup_{n, m \rightarrow \infty} \|\widehat{f} \circ T(M_n) - \widehat{f} \circ T(M_m)\|_{L_1(\mathcal{F}(N))} > \frac{\delta}{2}$$

by (3) and (4), so $(\widehat{f} \upharpoonright_K(T(M_n)))_n$ cannot converge in $L_1(\mathcal{F}(N))$. According to Lemma 2.5 again, this shows that $\widehat{f} \upharpoonright_K : \mathcal{F}(K) \rightarrow \mathcal{F}(N)$ is not Radon-Nikodým, completing the proof. \square

Finally we prove the main theorem of the section.

4.3. Proof of Theorem 4.1.

Proof of Theorem 4.1. By Theorem 3.1, we only need to show (1) \implies (2) and (2) \implies (3). The first implication follows by Propositions 4.3 and 4.4. The contrapositive of the second implication is clear, since L_1 fails the RNP, and thus there exists an L_1 -valued bounded martingale which does not converge almost surely. \square

Before closing this section, let us comment on the similarities in the proofs of Propositions 3.5 and 4.3.

Remark 4.5. In the proofs of Propositions 3.5 and 4.3 we use a gliding hump type argument in order to derive a contradiction with the fact that a particular sequence (a weakly null sequence in Proposition 3.5 and a uniformly integrable martingale in Proposition 4.3) is not convergent in norm. In fact, the inductive process of both proofs can be unified and generalized for sequences in $L_1(X)$ that are scalarly convergent in L_1 but not convergent in $L_1(X)$. Let us expand on this statement.

Following the terminology of [18], a sequence $(X_n)_n$ of X -valued random variables in $L_1(X)$ is *scalarly convergent in L_1* if for every $x^* \in X^*$, the sequence $(x^* \circ X_n)$ of scalar random variables in L_1 is convergent in norm. Note that we may identify any weakly convergent sequence in a Banach space X with a sequence of random variables in $L_1(X)$ scalarly convergent in L_1 in a natural way. Indeed, given such a sequence $(x_n)_n$ in X , let Ω be any nonempty set, put $\mathcal{A} = \{\emptyset, \Omega\}$, and consider \mathbb{P} as the trivial (and unique) probability measure on \mathcal{A} . Then, $L_1(\Omega, X)$ is isometrically isomorphic to X with the isometry $\varphi(x) = x1_\Omega$ for every $x \in X$. Moreover, the sequence given by $S_n = x_n1_\Omega \in L_1(\Omega, X)$ is scalarly convergent in L_1 if and only if $(x_n)_n$ is weakly convergent. Notice as well that every bounded and uniformly integrable martingale is also scalarly convergent in L_1 (see e.g. [38, Corollary 1.16]).

With this in mind, we could first generalize Lemma 4.2 to sequences of random variables that scalarly converge in L_1 . Then, following the proof of Proposition 4.3 with minimal changes, we could prove the following statement:

Fact 4.6. *Let M be a compact metric space and let $f: M \rightarrow N$ be a curve-flat Lipschitz function. Suppose $(X_n)_n$ is a sequence of $\mathcal{F}(M)$ -valued random variables*

in $L_1(\mathcal{F}(M))$ adapted to a filtration $(\mathcal{A}_n)_n$ of finite sub- σ -algebras, such that $(X_n)_n$ is scalarly convergent in L_1 but $\|\widehat{f} \circ D_n\| \geq 1$ for all $n \in \mathbb{N}$ (where $D_n = X_{n+1} - X_n$ for all $n \in \mathbb{N}$). Then, given a decreasing sequence of positive numbers $(\varepsilon_n)_n$ such that $\sum_n \varepsilon_n < \frac{1}{8}$, there exists a bounded sequence $(\varphi_n)_n$ and a subsequence $(D_{k_n})_n$ such that:

- (1) $\varphi_n \in L_1(\mathcal{A}_{k_n+1}, \mathcal{F}(M))^* = L_\infty(\mathcal{A}_{k_n+1}, \text{Lip}_0(M))$,
- (2) For every $\omega \in \Omega$, $\varphi_n(\omega)$ is $(\sum_{i=1}^n \varepsilon_i)$ -flat at some positive radius $0 < \theta_n < 1$,
- (3) $\|\varphi_n\| \leq 1 + \sum_{i=1}^n \varepsilon_i$, and
- (4) $\langle \varphi_n, D_{k_i} \rangle \geq \frac{1}{4} - \sum_{i=1}^n \varepsilon_i$ for all $i = 1, \dots, n$.

Since, as commented above, both weakly convergent sequences and uniformly integrable martingales are modelled by random variables that scalarly converge in L_1 , this fact could be used as a Lemma for both Propositions 3.5 and 4.3. Although this would shorten this article and allow us to avoid using similar arguments in both proofs, we chose not to follow this structure. This is mainly due to the fact that Proposition 3.5 can be shown without mention of random variables, and using the approach explained in this remark would arguably obscure the main idea of its proof.

Moreover, note that the extra properties of weakly convergent sequences in Proposition 3.5 and martingales in Proposition 4.3 are needed at the end of the proofs of the respective results. Certainly, it is not true in general that if $f: M \rightarrow N$ is a curve-flat Lipschitz function between compact metric spaces and $(X_n)_n$ is a sequence of random variables in $L_1(\mathcal{F}(M))$ that scalarly converges in L_1 , then $(\widehat{f} \circ X_n)_n$ converges in $L_1(\mathcal{F}(N))$. Indeed, let M be an infinite compact purely 1-unrectifiable metric space, and let $f: M \rightarrow M$ be the identity function. Then f is curve-flat, but since $\mathcal{F}(M)$ is infinite-dimensional, by [18, Corollary 4.4] there exists a sequence $(X_n)_n$ in $L_1(\mathcal{F}(M))$ that is scalarly convergent in L_1 but that admits no a.e. convergent subsequence, which in particular means that $(X_n)_n$ cannot converge in the norm of $L_1(\mathcal{F}(M))$.

5. REMARKS ON STRONG RADON-NIKODÝM AND REPRESENTABLE LIPSCHITZ OPERATORS

In this section, we discuss two classes of Lipschitz operators (strong Radon-Nikodým and representable Lipschitz operators, see Section 2.2) which, in some cases, can be added to the list of equivalences of Theorem 1.1.

5.1. Strong Radon-Nikodým Lipschitz operators. We start by showing that strong Radon-Nikodým and Radon-Nikodým Lipschitz operators are indeed the same provided the underlying metric space in the domain is compact and countably 1-rectifiable.

Following [21, Definition 3.2.14, Page 251] (see also [22, Section 3.2]), a metric space M is *countably 1-rectifiable* if there exist a countable collection of subsets $A_i \subset \mathbb{R}$ and Lipschitz maps $\gamma_i: A_i \rightarrow M$ such that $\mathcal{H}^1(M \setminus (\bigcup_i \gamma_i(A_i))) = 0$. Clearly, any subset of \mathbb{R} is countably 1-rectifiable. Notice also that M is purely 1-unrectifiable if and only if every countably 1-rectifiable subset of M is \mathcal{H}^1 -negligible.

Proposition 5.1. *Let M be a compact metric space which is countably 1-rectifiable, let N be any metric space and let $f \in \text{Lip}_0(M, N)$. The following are equivalent.*

- (1) $f(M)$ is \mathcal{H}^1 -negligible.

- (2) \widehat{f} is strong Radon-Nikodým.
(3) \widehat{f} does not fix a copy of L_1 .

Proof. (1) \implies (2). Since M is compact and $\mathcal{H}^1(f(M)) = 0$, $f(M)$ is complete and purely 1-unrectifiable. Thus, [5, Theorem C] implies that $\mathcal{F}(f(M))$ has the Radon-Nikodým property. Since $\widehat{f}(B_{\mathcal{F}(M)})$ is contained in $\mathcal{F}(f(M))$, this yields that \widehat{f} is strong Radon-Nikodým.

(2) \implies (3). This implication is true in general, since every strong Radon-Nikodým operator is Radon-Nikodým, and, as we discussed in the proof of Theorem 4.1, such an operator cannot fix a copy of L_1 .

(3) \implies (1). We do not use the compactness assumption on M for this implication. Suppose that $\mathcal{H}^1(f(M)) > 0$. Since M is countably 1-rectifiable, there exist a countable collection $(A_i)_{i \in I}$ of subsets of \mathbb{R} and Lipschitz maps $\gamma_i : A_i \rightarrow M$ such that $\mathcal{H}^1(M \setminus (\bigcup_{i \in I} \gamma_i(A_i))) = 0$. Since f is Lipschitz,

$$\begin{aligned} \mathcal{H}^1\left(f(M) \setminus f\left(\bigcup_{i \in I} \gamma_i(A_i)\right)\right) &\leq \mathcal{H}^1\left(f\left(M \setminus \left(\bigcup_{i \in I} \gamma_i(A_i)\right)\right)\right) \\ &\leq \|f\|_L \mathcal{H}^1\left(M \setminus \left(\bigcup_{i \in I} \gamma_i(A_i)\right)\right) = 0. \end{aligned}$$

Therefore, by σ -additivity of \mathcal{H}^1 , there exists $i \in I$ such that $\mathcal{H}^1(f \circ \gamma_i(A_i)) > 0$. By Kirchheim's lemma, there exists a compact subset B of A_i such that $\lambda(B) > 0$ and $(f \circ \gamma_i)|_B$ is bi-Lipschitz. Thus $\gamma_i|_B : B \rightarrow \gamma_i(B)$ and $f|_{\gamma_i(B)} : \gamma_i(B) \rightarrow f \circ \gamma_i(B)$ are both bi-Lipschitz as well, and in particular $\widehat{\gamma_i|_B}$ and $\widehat{f|_{\gamma_i(B)}}$ are isomorphisms onto their respective range. On the other hand, since $\lambda(B) > 0$, Godard's work [25] implies that $\mathcal{F}(B)$ contains an isomorphic copy E of L_1 . Thus $\widehat{\gamma_i|_B}(E)$ is also an isomorphic copy of L_1 . Restricting \widehat{f} to this copy of L_1 yields the desired result. \square

Remark 5.2. It is worth noting that the equivalence between (1) and (3) in Proposition 5.1 holds when M is only assumed to be complete and not necessarily compact. Indeed, we already pointed out that our proof of (3) \implies (1) does not need the compactness of M . Conversely, let M be a complete countably 1-rectifiable set, and suppose that $f(M)$ is \mathcal{H}^1 -negligible for a given $f \in \text{Lip}_0(M, N)$. It is clear that $f(K)$ is also \mathcal{H}^1 -negligible for any compact $K \subset M$. Then, (if moreover $0 \in K$) by Proposition 5.1 and Theorem 1.1, we see that $\widehat{f|_K}$ is Dunford-Pettis. As K can be any compact subset of M such that $0 \in K$, we can apply Proposition 3.6 to conclude that \widehat{f} is Dunford-Pettis, and thus cannot fix a copy of L_1 .

Notice however that completeness of M is necessary. For instance, consider $M = \mathbb{Q} \cap [0, 1]$, $N = [0, 1]$ and $f = Id_M$. Then \widehat{f} is conjugate to the identity on L_1 ; so it fixes a copy of L_1 while the set $f(M) = M$ is countable and thus \mathcal{H}^1 -negligible.

Thanks to Theorem 1.1, Proposition 5.1 shows in particular that, if M is a compact countably 1-rectifiable metric space, then $\widehat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ is Radon-Nikodým if and only if it is strong Radon-Nikodým. However, this is not true if M is only assumed to be complete, as shown by the following example.

Example 5.3. Let N be a complete, separable and non-purely 1-unrectifiable metric space, with diameter not greater than 1 (e.g. the unit interval in \mathbb{R}). Let $(x_k)_k$ be a countable dense subset of N , and let $\varphi : \mathbb{N} \rightarrow \tilde{\mathbb{N}} = \mathbb{N}^2 \setminus \{(n, n) : n \in \mathbb{N}\}$

be an enumeration. Write $\varphi(n) = (\varphi(n)_1, \varphi(n)_2)$ for each $n \in \mathbb{N}$. Consider T the unique \mathbb{R} -tree with only one branching point $0 \in T$ and with branches $B_n := [0, 1 + d(x_{\varphi(n)_1}, x_{\varphi(n)_2})]$ for every $n \in \mathbb{N}$. Denote by $e_{(n,1)}$ the point in B_n at distance 1 from 0, and by $e_{(n,2)}$ the point in B_n at distance $1 + d(x_{\varphi(n)_1}, x_{\varphi(n)_2})$ from 0 and thus at distance $d(x_{\varphi(n)_1}, x_{\varphi(n)_2})$ from $e_{(n,1)}$. Let $M = \bigcup_{n \in \mathbb{N}} \{e_{(n,1)}, e_{(n,2)}\} \cup \{0\} \subset T$, with the \mathbb{R} -tree distance inherited from T . It is clear that M is a complete metric space. Then, for every pair $k_1 \neq k_2 \in \mathbb{N}$, if we choose $n = \varphi^{-1}(k_1, k_2)$, it holds that $d_N(x_{k_1}, x_{k_2}) = d_M(e_{(n,1)}, e_{(n,2)})$. Then we define $f: M \rightarrow N$ as $f(0) = 0$, $f(e_{(n,1)}) = x_{\varphi(n)_1}$, and $f(e_{(n,2)}) = x_{\varphi(n)_2}$. Since the diameter of N is not greater than 1, it follows from the tree structure of M that f is 1-Lipschitz. Then, using that $(x_k)_k$ is dense in N , it easily follows that $C_f = (\widehat{f})^*$ is a linear isometry. Hence, \widehat{f} is a linear quotient from $\mathcal{F}(M)$ onto $\mathcal{F}(N)$. This implies that \widehat{f} is not strong Radon-Nikodým, since $\mathcal{F}(N)$ fails the RNP. However, \widehat{f} is Radon-Nikodým because M is purely 1-unrectifiable (so, $\mathcal{F}(M)$ has the RNP), and thus \widehat{f} is trivially curve-flat. Finally, since M is countable, we have $\mathcal{H}^1(M) = 0$ and so M is countably 1-rectifiable.

5.2. Representable Lipschitz operators. In this subsection we work with compact subsets of the real line, since it is in this context where we can properly deal with representable operators between Lipschitz-free spaces. Indeed, given a compact subset A of \mathbb{R} , thanks to Godard's work [25], there exists a finite measure μ_A on A such that $\mathcal{F}(A)$ is isometrically isomorphic to $L_1(\mu_A)$. Thus, any Lipschitz operator $\widehat{f}: \mathcal{F}(A) \rightarrow \mathcal{F}(N)$ is conjugate to a bounded operator $T_f: L_1(\mu_A) \rightarrow \mathcal{F}(N)$. More precisely, if $\Phi: L_1(\mu_A) \rightarrow \mathcal{F}(A)$ denotes Godard's isometry, then $T_f := \widehat{f} \circ \Phi$. We can thus study when the operator T_f is representable, based on the properties of f .

Our main result in this subsection is based on the following general fact. Although it is not commonly found explicitly stated in the literature, it is well-known among specialists. For the convenience of the reader, we provide some detailed explanations.

Lemma 5.4. *Let (Ω, Σ, μ) be a finite measure space and Y be a Banach space. Then a bounded operator $T: L_1(\mu) \rightarrow Y$ is representable if and only if it is Radon-Nikodým.*

Proof. Assume that $T: L_1(\mu) \rightarrow Y$ is Radon-Nikodým. Let us define the vector measure $G: \Sigma \rightarrow L_1(\mu)$ by $G(E) := \mathbf{1}_E$. Then G is of bounded variation; indeed its variation $|G|$ is nothing but μ . By the characterization of RN operators (see [20, p. 461]), $T \circ G: \Sigma \rightarrow Y$ admits a Bochner integrable density, i.e., there exists $g \in L_1(\mu, Y)$ such that $T(\mathbf{1}_E) = \int_E g d\mu$. This proves that T is representable thanks to [17, Lemma III.1.4]. Conversely, suppose that T is representable. By Lewis-Stegall theorem (see e.g. the proof of [17, Theorem III.1.8]), T factors through ℓ_1 : There exist $T_1: L_1(\mu) \rightarrow \ell_1$ and $T_2: \ell_1 \rightarrow Y$ such that the following diagram commutes

$$\begin{array}{ccc} L_1(\mu) & \xrightarrow{T} & Y \\ T_1 \downarrow & \nearrow T_2 & \\ \ell_1 & & \end{array}$$

Since ℓ_1 has the RNP, we have that T_1 is (strong) Radon-Nikodým. Since the class of Radon-Nikodým operators is an operator ideal [33, Theorem 5], we conclude that T is Radon-Nikodým. \square

We now obtain the following corollary. Note that several more items could be added to the equivalence therein according to Theorem 1.1.

Corollary 5.5. *Let A be a compact subset of \mathbb{R} containing 0, let N be a complete metric space and let $f \in \text{Lip}_0(A, N)$ be given. Recall that Φ denotes Godard's isomorphism from $L_1(\mu_A)$ onto $\mathcal{F}(A)$. Then the following assertions are equivalent.*

- (i) $T_f = \widehat{f} \circ \Phi : L_1(\mu_A) \rightarrow \mathcal{F}(N)$ is representable.
- (ii) \widehat{f} is Radon-Nikodým.

Proof. Applying Lemma 5.4 we obtain that T_f is representable if and only if T_f is Radon-Nikodým. Since $\widehat{f} = T_f \circ \Phi^{-1}$ and Radon-Nikodým operators form an operator ideal, we obtain the desired result. \square

5.3. Remarks on bounded operators from $\mathcal{F}(M)$ to X . Here we are going to show that there is no analog of Theorem 1.1 for the linearization $\bar{f} : \mathcal{F}(M) \rightarrow X$ of a Lipschitz map $f : M \rightarrow X$, where M is a metric space and X is a Banach space. This should be no surprise as every bounded linear operator $T : \mathcal{F}(M) \rightarrow X$ is of the form \bar{f} for some $f \in \text{Lip}_0(M, X)$.

Proposition 5.6. *Let M be a complete metric space, X be a Banach space and $f \in \text{Lip}_0(M, X)$. Then*

$$\begin{array}{ccc}
 f \in \text{CF} & \xRightarrow{\quad} & \bar{f} \in \text{DP} \\
 \Downarrow & \searrow & \Downarrow \\
 \bar{f} \in \text{RN} & \xRightarrow{\quad} & \bar{f} \text{ does not fix copy of } L_1
 \end{array}$$

and all the implications not shown on the diagram are false in general.

For the proof we will need the following proposition of independent interest.

Proposition 5.7. *Let \mathcal{I} be an operator ideal and let $\text{Space}(\mathcal{I})$ be the class of Banach spaces X such that $\text{Id}_X \in \mathcal{I}$. Then $X \in \text{Space}(\mathcal{I})$ implies $\beta_X \in \mathcal{I}$. If, in addition, $\text{Space}(\mathcal{I})$ is separably determined, then $\beta_X \in \mathcal{I}$ implies $X \in \text{Space}(\mathcal{I})$. In particular, X has the SP if and only if β_X is DP, and X has the RNP if and only if β_X is RN.*

Proof. The first implication follows since $\beta_X = \text{Id}_X \circ \beta_X$ and \mathcal{I} is an operator ideal.

To see the second implication, it is enough to show that every separable subspace Y of X belongs to $\text{Space}(\mathcal{I})$. Fix a separable subspace Y of X and observe the following commutative diagram

$$\begin{array}{ccc}
 \mathcal{F}(Y) & \xrightarrow{\widehat{\iota}} & \mathcal{F}(X) \\
 \beta_Y \downarrow & & \downarrow \beta_X \\
 Y & \xrightarrow{\iota} & X,
 \end{array}$$

where $\iota : Y \hookrightarrow X$ is the canonical linear embedding, and β_X and β_Y are linear quotient maps which are the left inverses of δ_X and δ_Y , respectively. That is, $\iota \circ \beta_Y = \beta_X \circ \widehat{\iota}$. Write $E = \iota(Y)$ and notice from [27, Theorem 3.1] that there exists a Godefroy-Kalton isometric linear lifting $T : Y \rightarrow \mathcal{F}(Y)$ of the map β_Y , i.e.

$$Id_Y = \beta_Y \circ T.$$

Noting $\iota^{-1} : E \rightarrow Y$ is bounded as well, we conclude

$$Id_Y = \iota^{-1} \circ \iota \circ \beta_Y \circ T = \iota^{-1} \circ \beta_X \circ \widehat{\iota} \circ T.$$

Since $\beta_X \in \mathcal{I}$, we have that $Id_Y \in \mathcal{I}$, equivalently, $Y \in Space(\mathcal{I})$.

The last part of the proposition holds as $SP = Space(DP)$, $RNP = Space(RN)$ and both classes SP and RNP are separably determined. \square

Proof of Proposition 5.6. Taking into account that $\bar{f} = \beta_X \circ \widehat{f}$, it follows from Theorem 1.1 that if f is curve-flat, then \bar{f} is DP, RN and does not fix any copy of L_1 . The converses of the three implications are easily seen to be false. Indeed, if $f : \mathbb{R} \rightarrow \mathbb{R}$ is the identity, then it is not curve-flat, while $\bar{f} : \mathcal{F}(\mathbb{R}) \rightarrow \mathbb{R}$ is of rank one and therefore DP, RN and not fixing copy of L_1 .

It is a general fact, that neither DP nor RN operators fix copies of L_1 .

Next, consider $Id_{\ell_2} : \ell_2 \rightarrow \ell_2$ the identity map. Since ℓ_2 has the RNP, the linearization $\overline{Id_{\ell_2}} : \mathcal{F}(\ell_2) \rightarrow \ell_2$ is RN (and fixes no copy of L_1). However, $\overline{Id_{\ell_2}}$ is nothing else than the quotient map β_{ℓ_2} ; so it is not DP by Proposition 5.7.

Similarly, let X be the Bourgain-Rosenthal space which has the SP but lacks the RNP [13]. Then β_X is DP (and therefore fixes no copy of L_1) but is not RN by Proposition 5.7. \square

Remark 5.8. The last part of the above proof can be achieved by a much more elementary example. Let $(A_n)_{n=1}^{\infty}$ be an enumeration of the dyadic intervals in $[0, 1]$. Note that $\lim_n \lambda(A_n) = 0$ and define $T : L_1 = L_1([0, 1]) \rightarrow c_0$ by $Tf = (\int_{A_n} f d\lambda)_{n=1}^{\infty}$. The operator T is easily seen to be DP. Indeed if $(f_k)_k$ is weakly null in L_1 , then $\{f_k, k \in \mathbb{N}\}$ is uniformly integrable. It readily follows that $(Tf_k)_k$ is norm null in c_0 . On the other hand, T is not RN. Indeed, For a Borel subset E of $[0, 1]$, define again $G(E)$ to be the indicator function of E . Clearly, G is an L_1 -valued measure that is absolutely continuous with respect to λ and $T \circ G(E) = (\lambda(E \cap A_n))_{n=1}^{\infty}$. Assume that T is representable, so that there exists $g = (g_n)_{n=1}^{\infty} \in L_1([0, 1]; c_0)$ such that for all E Borel, $T \circ G(E) = \int_E g d\lambda$. Then, for each $n \in \mathbb{N}$, g_n is the indicator function of A_n . But, $\limsup_n g_n(t) = 1$ for any $t \in [0, 1]$. This is a contradiction showing that T is not representable and therefore not RN. Note finally that, identifying $L_1([0, 1])$ with $\mathcal{F}([0, 1])$ via the linear isometry sending δ_t to the indicator function of $[0, t]$, we can view T as \bar{f} , where $f : [0, 1] \rightarrow c_0$ is the Lipschitz map defined by $f(t) = (\lambda([0, t] \cap A_n))_{n=1}^{\infty}$. Thus we have that \bar{f} is DP but not RN.

6. QUOTIENT METRIC SPACES AND FACTORIZATIONS

In general, given an operator $T : X \rightarrow Y$ between Banach spaces, it is clear that if T factors through a Schur (respectively RNP) space, then T is Dunford-Pettis (respectively Radon-Nikodým). However, the converse is not necessarily true, as we explained in Section 2.2.

In our context, it is still a natural question whether the conditions $\widehat{f} \in Op(SP)$ (resp. $\widehat{f} \in Op(RNP)$), i.e. “ \widehat{f} factors through a space with the SP (resp. the RNP)”, can be added to the list of equivalences in Theorem 1.1. Taking into account the characterization of Lipschitz-free spaces with the SP (resp. the RNP), we can also ask whether the stronger condition “ f factors through a complete purely 1-unrectifiable metric space” can also be included in Theorem 1.1. In Theorem 6.5, we will characterize the compact metric spaces M for which this stronger factorization condition holds, for every range space N . Then, we will present a concrete example showing that the answer is negative in general (Example 6.8). Although the “factorization through Schur spaces” question remains open, we will provide a few related observations in the last two subsections.

6.1. Factoring through p1u spaces: the compact case. In what follows, $Fact_{p1u}(M, N)$ stands for the set of those $f \in Lip_0(M, N)$ which factor through a complete p1u space. When M is compact, we can omit to stress the completeness of the p1u factor as that is automatic. We always have $Fact_{p1u}(M, N) \subset CF_0(M, N)$ as $Id_P : P \rightarrow P$ is curve-flat for any p1u metric space P . For maps $f \in Lip_0(M, N)$ and $g \in Lip_0(M, P)$ we will say that f factors through g if there exists $h \in Lip_0(P, N)$ such that $f = h \circ g$. We will denote $Fact_g(M, N)$ the set of such maps. Given a metric space M , a particular role is going to be played by the metric space M_{lip} defined as the canonical quotient of M where we identify the points $x, y \in M$ such that $d_{lip}(x, y) = 0$ where

$$d_{lip}(x, y) := \sup\{|f(x) - f(y)| : f \in B_{lip_0(M)}\}.$$

We equip M_{lip} with the metric $d_{lip}([x], [y]) = d_{lip}(x, y)$. Let us denote $\pi : M \rightarrow M_{lip}$ the canonical quotient map defined by $\pi(x) = [x]$ for every $x \in M$. Clearly $\|\pi\|_L \leq 1$. It is well known and easy to see that, given any metric space N , every $f \in lip_0(M, N)$ factors through $\pi : M \rightarrow M_{lip}$. In other words, we always have $lip_0(M, N) \subset Fact_{\pi: M \rightarrow M_{lip}}(M, N)$. Since M is compact, [42, Corollary 8.13] combined with [5, Theorem A] imply that M_{lip} is p1u. Thus, we arrive at the following observation.

Proposition 6.1. *Let M be a compact metric space and let N be any metric space. If $f \in lip_0(M, N)$, then f factors through M_{lip} which is compact and p1u.*

On the other hand, it follows from the results in [5] that, for compact M , given any p1u metric space P , every $f \in Lip_0(M, P)$ factors through $\pi : M \rightarrow M_{lip}$.

Lemma 6.2. *Let M be compact and P be p1u. Let $f \in Lip_0(M, P)$. Then f factors through $\pi : M \rightarrow M_{lip}$. More precisely, there exists $f_{lip} \in Lip_0(M_{lip}, P)$ such that $\|f_{lip}\|_L \leq \|f\|_L$ and $f = f_{lip} \circ \pi$.*

Proof. This follows from [5, Proposition 5.8] and [5, Theorem 5.9]. Here we provide a more direct proof: We define $f_{lip} : M_{lip} \rightarrow P$ by $f_{lip}([x]) = f(x)$. Let $x, y \in M$. We have

$$\begin{aligned} d(f_{lip}([x]), f_{lip}([y])) &= d(f(x), f(y)) \\ &= \sup\{\varphi(f(x)) - \varphi(f(y)) : \varphi \in B_{lip_0(P)}\} \\ &\leq \|f\|_L d_{lip}([x], [y]) \end{aligned}$$

where we have used [5, Theorem A] and the fact that $\varphi \circ f \in lip_0(M)$. This inequality shows at once that f_{lip} is well defined and that $\|f_{lip}\|_L \leq \|f\|_L$ \square

Proposition 6.3. *Let M be a compact metric space and let N be any metric space. Then*

$$\text{Fact}_{p1u}(M, N) = \text{Fact}_{\pi: M \rightarrow M_{\text{lip}}}(M, N).$$

Moreover, there exists a canonical bijective correspondence between this set and $\text{Lip}_0(M_{\text{lip}}, N)$. Namely, the pre-composition operator $C_\pi : \text{Lip}_0(M_{\text{lip}}, N) \rightarrow \text{Fact}_{p1u}(M, N)$, given by $u \mapsto u \circ \pi$, establishes this correspondence. It moreover satisfies $\|C_\pi(u)\|_L \leq \|u\|_L$ for every $u \in \text{Lip}_0(M_{\text{lip}}, N)$ and

$$\forall f \in \text{Fact}_{p1u}(M, N), \quad \|C_\pi^{-1}(f)\|_L = \inf \|h\|_L \|g\|_L,$$

where the infimum is taken over all factorizations $f = h \circ g$ through a complete p1u metric space.

Proof. Assume first that $f \in \text{Lip}_0(M, N)$ factors through $\pi : M \rightarrow M_{\text{lip}}$. As mentioned before Proposition 6.1, the metric space M_{lip} is p1u; so $f \in \text{Fact}_{p1u}(M, N)$.

Assume now $f \in \text{Fact}_{p1u}(M, N)$ and that P is a complete p1u metric space such that $f = h \circ g$, where $g \in \text{Lip}_0(M, P)$ and $h \in \text{Lip}_0(P, N)$. Since M is compact, we can write $f = h \circ g_{\text{lip}} \circ \pi$ where $\|g_{\text{lip}}\|_L \leq \|g\|_L$ by Lemma 6.2. Thus $f \in \text{Fact}_{\pi: M \rightarrow M_{\text{lip}}}(M, N)$.

Now observe that, since $\pi : M \rightarrow M_{\text{lip}}$ is surjective, the map $C_\pi : u \in \text{Lip}_0(M_{\text{lip}}, N) \mapsto u \circ \pi \in \text{Fact}_{p1u}(M, N)$ is injective and $\|C_\pi(u)\|_L \leq \|u\|_L$, since $\|\pi\|_L \leq 1$. The argument above shows that C_π is also surjective with $C_\pi^{-1}(f) = h \circ g_{\text{lip}}$ with $\|C_\pi^{-1}(f)\|_L \leq \|h\|_L \|g_{\text{lip}}\|_L$. The opposite inequality is clear as $f = C_\pi^{-1}(f) \circ \pi$ and $\|\pi\|_L \leq 1$. \square

As we mentioned in Section 2.3, the space $\text{CF}_0(M, Y)$ is a Banach space. Thus, the following is obtained as a consequence of Proposition 6.3 and the open mapping theorem.

Corollary 6.4. *Let M be a compact metric space and Y be a Banach space. Then $C_\pi : \text{Lip}_0(M_{\text{lip}}, Y) \rightarrow \text{Fact}_{p1u}(M, Y)$ is linear, bounded and pointwise-to-pointwise continuous.*

If moreover $\text{CF}_0(M, Y) \subset \text{Fact}_{p1u}(M, Y)$, then C_π is an isomorphism from $\text{Lip}_0(M_{\text{lip}}, Y)$ into $\text{CF}_0(M, Y)$.

Proof. The first part of the corollary follows immediately from Proposition 6.3. If $\text{CF}_0(M, Y) \subset \text{Fact}_{p1u}(M, Y)$ then $\text{CF}_0(M, Y) = \text{Fact}_{p1u}(M, Y)$, since the reverse inclusion is always true. We have that $\|C_\pi\| \leq \|\pi\|_L$ and Proposition 6.3 claims that this map is a bijection from $\text{Lip}_0(M_{\text{lip}}, Y)$ to $\text{Fact}_{p1u}(M, Y)$. Thus, by the open mapping theorem, C_π is an isomorphism. \square

Let us turn to conditions on M guaranteeing $\text{CF}_0(M, N) \subset \text{Fact}_{p1u}(M, N)$. The next theorem shows that when looking for curve-flat functions which do not factor through a complete p1u space, it is enough to consider functions with real values. Also, this theorem relates the inclusion $\text{CF}_0(M, N) \subset \text{Fact}_{p1u}(M, N)$ to the condition that the following natural quotient metric space is p1u. Let (M, d) be a compact metric space and define

$$\forall x, y \in M, \quad d_{\text{CF}}(x, y) := \sup\{|f(x) - f(y)| : f \in B_{\text{CF}_0(M)}\}.$$

We denote M_{CF} the canonical quotient space of M where we identify the points $x, y \in M$ such that $d_{\text{CF}}(x, y) = 0$, which we equip with the metric $d_{\text{CF}}([x], [y]) = d_{\text{CF}}(x, y)$. While, on the first sight, the metric space M_{CF} seems to be defined using

behavior of Lipschitz functions on M , in fact it admits a fully metric definition. This has been proved in [5, Proposition 5.22]. In particular, the metric spaces M such that M_{CF} is p1u are those where the iterative process of “removing curve fragments” stabilizes after at most 1 step.

Theorem 6.5. *Let M be a compact metric space. Then it is equivalent*

- (1) $CF_0(M) \subset Fact_{p1u}(M, \mathbb{R})$,
- (2) $CF_0(M, N) \subset Fact_{p1u}(M, N)$ for every metric space N ,
- (3) The metric space M_{CF} is compact p1u.
- (4) The metric spaces M_{CF} and M_{lip} are Lipschitz equivalent.
- (5) $B_{CF_0(M)} \subset \alpha \overline{B_{lip_0(M)}}^{w^*}$ for some $\alpha > 0$.
- (6) $CF_0(M) \subset \overline{lip_0(M)}^{w^*}$.

Moreover, under any of these conditions, the constant α in (5) can be taken equal 1 and the Lipschitz equivalence in (4) is an isometry. Also the inclusions in items (1), (2), (5) with $\alpha = 1$ and (6) are equalities.

Proof. (5) \implies (4). Recalling the definition of d_{lip} and d_{CF} , observe that, under assumption (5), we have that for $x \neq y \in M$

$$d_{lip}(x, y) \leq d_{CF}(x, y) \leq \alpha d_{lip}(x, y).$$

This implies that M_{CF} and M_{lip} are bi-Lipschitz equivalent.

(4) \implies (3). Indeed, by Proposition 6.1, M_{CF} is compact and p1u.

(3) \implies (2) is clear, as for any metric space N , any $f \in CF_0(M, N)$ factors through M_{CF} in a canonical way.

As (2) \implies (1) is trivial, we claim that (1) \implies (5). Let $f \in B_{CF_0(M)}$ be given. By Corollary 6.4 there is $u \in Lip_0(M_{lip})$ such that $\|u\|_L \leq \|C_\pi^{-1}\|$ and $C_\pi(u) = f$. Let $\alpha := \|C_\pi^{-1}\|$. Since M_{lip} is compact p1u, we have $lip_0(M_{lip})^{**} = Lip_0(M_{lip})$. So there is a sequence $(u_n) \subset \alpha B_{lip_0(M_{lip})}$ which converges pointwise to u . Thus, by Proposition 6.3, the sequence $u_n \circ \pi$ converges pointwise (and, being bounded, also weakly*) to $u \circ \pi = f$ which concludes the proof as $u_n \circ \pi \in lip_0(M)$.

Further (5) clearly implies (6). In order to prove (6) \implies (5), observe that Petunin-Pličko theorem [36] (see also [26]) yields the following fact: Let X be a separable Banach space and $Z \subseteq X^*$ be a separable normed-closed subspace contained in the set of norm-attaining functionals. Then $Z^{**} = \overline{Z}^{w^*}$ isometrically. Indeed, denote $Y = X/Z_\perp$. Then $Y^* = \overline{Z}^{w^*}$ isometrically. Moreover Z clearly separates points of Y . Also Z is a subset of norm-attaining functionals on Y . This last claim follows from the facts that $\|z\|_{Y^*} = \|z\|_{X^*}$ and $Z \subset NA(X)$. It then follows from Petunin-Pličko theorem that Y is isometric to Z^* , and therefore Z^{**} is isometric to \overline{Z}^{w^*} . Thus applying the above fact to $Z = lip_0(M)$ – notice that $lip_0(M)$ is contained in the set of norm attaining functionals on $\mathcal{F}(M)$ thanks to [15, Lemma 2.3] – we get $CF_0(M) \subseteq lip_0(M)^{**}$ and (5) follows with $\alpha = 1$ by Goldstine’s theorem.

The very last claim follows from the facts $Fact_{p1u}(M, N) \subseteq CF_0(M, N)$ in general and the weak*-closedness of $CF_0(M)$ due to Proposition 2.2. \square

We continue with some more concrete conditions on M that are sufficient for the inclusion $CF_0(M, N) \subset Fact_{p1u}(M, N)$ to hold for every metric space N .

Proposition 6.6. *Let M be compact and countably 1-rectifiable, and N be any metric space. If $f \in \text{CF}_0(M, N)$ then $f(M)$ is $p1u$. In particular, f factors through a compact $p1u$ space.*

Proof. Theorem 1.1 implies that \widehat{f} does not fix any copy of L_1 and so $\mathcal{H}^1(f(M)) = 0$ by Proposition 5.1. Thus $f(M)$ is compact and $p1u$ as claimed. \square

Remark 6.7. When M is compact and countably 1-rectifiable we have that $B_{\text{Lip}_0(M)}$ is weak*-dense in $B_{\text{CF}_0(M)}$. This can be of course seen from Theorem 6.5 combined with Proposition 6.6. But we can also get a bit stronger approximation: since $\widehat{f}(A) = f(A)$ by compactness, [23, Lemma 4.8] guarantees that there exists a sequence $(\varphi_n)_n$ of locally constant functions in $B_{\text{Lip}_0(A)}$ such that $\varphi_n \rightarrow f$ in the weak* topology.

6.2. A curve-flat function which does not factor through a $p1u$ space.

The following provides a negative answer to the question of whether the condition “ $f \in \text{Fact}_{p1u}$ ” can be added to the list of equivalences in Theorem 1.1.

Example 6.8. *There exists a compact M such that $\text{CF}_0(M) \not\subset \text{Fact}_{p1u}(M, \mathbb{R})$. In fact we have that each non-constant $f \in \text{CF}_0(M)$ does not factor through any $p1u$ space.*

Proof. The example of such M comes from [5, Example 5.25]. Consider the standard middle thirds Cantor set $\mathcal{C} \subset [0, 1]$ and let $\beta := \log_3(2)$ be the Hausdorff dimension of \mathcal{C} . The Cantor function $f : \mathcal{C}^\beta \rightarrow [0, 1]$ is monotone, surjective, and 1-Lipschitz ([19, Proposition 10.1]), where \mathcal{C}^β denotes the snowflake space. Let (M, d) be a metric space obtained by “filling in the gaps” of \mathcal{C}^β with geodesics. In fact there are many ways to do the above but we equip M by the metric which is the largest possible among all such metrics. We will not use this particularity until Remark 6.10. The Cantor function f extends to a 1-Lipschitz map $f : M \rightarrow [0, 1]$ that is constant on each geodesic. Observe that f is curve-flat because \mathcal{C}^β is purely 1-unrectifiable and f is constant on each of the countably many geodesics. On the other hand, note that every locally flat function is constant. Hence, $M_{\text{lip}} = \{0\}$ and so $f \notin \text{Fact}_{\pi: M \rightarrow M_{\text{lip}}}(M, \mathbb{R})$. \square

Remark 6.9. It might be an interesting research project to study for which (non- $p1u$) compacts K there exists compact M such that $M_{\text{CF}} = K$. For instance, it can be shown that M_{CF} corresponding to the space from Example 6.8 is isometric to the unit interval $[0, 1]$ with the standard metric. Indeed, if $f : M \rightarrow [0, 1]$ is the extended Cantor function described in Example 6.8, the function $f_{\text{CF}_0} : M_{\text{CF}} \rightarrow [0, 1]$, where $f_{\text{CF}_0}([x]) = f(x)$ for all $[x] \in M_{\text{CF}}$, is the desired isometry. To see this, note first that, since f is curve-flat and 1-Lipschitz, we have that $d([x_1], [x_2]) = d_{\text{CF}}(x_1, x_2) \geq |f(x_1) - f(x_2)|$ for all $x_1, x_2 \in M$. It is then enough to show that for any given curve-flat 1-Lipschitz function $g \in \text{Lip}_0(M)$, one has $|g(x_1) - g(x_2)| \leq |f(x_1) - f(x_2)|$ for all $x_1, x_2 \in M$. Hence, fix such a function g and two points $x_1, x_2 \in M$. Since g and f are curve-flat, they are constant on all geodesics of M . Therefore, there exist $\tilde{x}_1, \tilde{x}_2 \in \mathcal{C}^\beta \subset M$ such that $g(\tilde{x}_i) = g(x_i)$ and $f(\tilde{x}_i) = f(x_i)$ for $i = 1, 2$. We may assume then without loss of generality that x_1 and x_2 belong to \mathcal{C}^β with $x_1 < x_2$ (in the real line ordering). Now, using again that $g(M)$ is contained in $g(\mathcal{C}^\beta)$, we get the estimate

$$|g(x_1) - g(x_2)| \leq \lambda([g(x_1), g(x_2)]) \leq \mathcal{H}^1([x_1, x_2] \cap \mathcal{C}^\beta) = \mathcal{H}^\beta([x_1, x_2] \cap \mathcal{C}),$$

where $([x_1, x_2] \cap \mathcal{C})^\beta$ denotes the snowflake of $[x_1, x_2] \cap \mathcal{C}$, isometrically contained in M . Finally, by [19, Proposition 5.5], we have that

$$|f(x_1) - f(x_2)| = \lambda(f([x_1, x_2] \cap \mathcal{C})) = \mathcal{H}^\beta([x_1, x_2] \cap \mathcal{C}),$$

and the conclusion follows.

Remark 6.10. Let M be the compact metric space considered in Example 6.8. We claim that $\widehat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(\mathbb{R})$ factors through ℓ_1 for every $f \in \text{CF}_0(M)$. In fact, the following much stronger property is true: there exists a compact plu $P \subset M$ (such that $\mathcal{F}(P) \simeq \ell_1$) and a weak* continuous extension operator $E : \text{Lip}_0(P) \rightarrow \text{Lip}_0(M)$ such that $E \circ R(f) = f$ for every $f \in \text{CF}_0(M)$ where $R : \text{Lip}_0(M) \rightarrow \text{Lip}_0(P)$ is the restriction operator. It then follows that $\widehat{f} = \widehat{f} \circ R_* \circ E_*$ where $R_* : \mathcal{F}(P) \rightarrow \mathcal{F}(M)$ and $E_* : \mathcal{F}(M) \rightarrow \mathcal{F}(P)$ are the corresponding pre-adjoints. Indeed, let $P = \mathcal{C}^\beta$ and define E as the ‘‘affine extension’’ operator. In order to check, that this operator is bounded, we observe that $E = \bar{g}^*$ where $g : M \rightarrow \mathcal{F}(\mathcal{C}^\beta)$ is defined as follows: For each $x \in \mathcal{C}^\beta \subset M$, we let $g(x) := \delta(x)$. Next, if $x \in M \setminus \mathcal{C}^\beta$, then x belongs to a unique geodesic which fills a gap of \mathcal{C}^β , say between the points $x_\ell, x_r \in \mathcal{C}^\beta$. In this case, we put $\lambda_x = \frac{d(x, x_\ell)}{d(x_\ell, x_r)}$ and define $g(x) := (1 - \lambda_x)\delta(x_\ell) + \lambda_x\delta(x_r)$. It is not hard to prove that g is 1-Lipschitz. For instance, let $x \in M \setminus \mathcal{C}^\beta$ and $y \in \mathcal{C}^\beta$. Suppose that the point x lies in the geodesic $[x_\ell, x_r]$ with $d(x, y) = d(x, x_\ell) + d(x_\ell, y)$. Note that

$$\|g(x) - g(x_\ell)\| = \|(1 - \lambda_x)\delta(x_\ell) + \lambda_x\delta(x_r) - \delta(x_\ell)\| = \lambda_x d(x_\ell, x_r) = d(x, x_\ell)$$

and $\|g(x_\ell) - g(y)\| = d(x_\ell, y)$ since $x_\ell, y \in \mathcal{C}^\beta$. Thus, observe that

$$\|g(x) - g(y)\| \leq \|g(x) - g(x_\ell)\| + \|g(x_\ell) - g(y)\| = d(x, x_\ell) + d(x_\ell, y) = d(x, y).$$

Where the very last equality follows from our choice of the metric on M . The other cases being similar, this proves that g is 1-Lipschitz; thus its linearization $\bar{g} : \mathcal{F}(M) \rightarrow \mathcal{F}(\mathcal{C}^\beta)$ is a norm-one bounded linear operator. Finally, $\mathcal{F}(\mathcal{C}^\beta)$ is isomorphic to ℓ_1 thanks to [42, Theorem 8.49].

6.3. An excursion to the non compact setting. When M is not necessarily compact, we still have the next elementary but interesting fact.

Lemma 6.11. *Let M, N be metric spaces and $f : M \rightarrow N$ a Lipschitz function which factors through a compact purely 1-unrectifiable metric space P , i.e. there are Lipschitz functions g, h such that the following diagram commutes:*

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ & \searrow g & \nearrow h \\ & & P \end{array}$$

Then for all $x, y \in M$ such that $f(x) \neq f(y)$ and for all $\varepsilon > 0$, there is $\psi \in \text{lip}_0(M)$ with $\|\psi\|_L \leq 1$ such that

$$|\psi(x) - \psi(y)| \geq \frac{d(f(x), f(y))}{\|h\|_L \|g\|_L} - \varepsilon.$$

Proof. We first observe that

$$d(f(x), f(y)) = d(h(g(x)), h(g(y))) \leq \|h\|_L d(g(x), g(y)).$$

Note that $f \neq 0$, so $\|h\|_L > 0$ and $\|g\|_L > 0$. Since P is purely 1-unrectifiable and compact, by [5, Theorem A], there exists $\varphi \in \text{lip}_0(P)$ with $\|\varphi\|_L \leq 1$ such that

$$|\varphi(g(x)) - \varphi(g(y))| \geq d(g(x), g(y)) - \varepsilon \|g\|_L.$$

We set $\psi = \frac{1}{\|g\|_L} \varphi \circ g$ and the conclusion follows. \square

Finally, we deal with the case when M is a complete subset of \mathbb{R} below.

Proposition 6.12. *Let $M \subset \mathbb{R}$ and let N be a metric space, both M and N complete. If $f \in \text{CF}_0(M, N)$, then f factors through a complete purely 1-unrectifiable metric space.*

Proof. If M is bounded, hence compact, Proposition 6.6 yields the conclusion. Next we will deal with the case when $\min M = 0 =: x_0$ and $\sup M = +\infty$. Let $(x_n) \subset M$ be a strictly increasing sequence with $\lim x_n = \infty$. We denote $M_n = M \cap [x_{n-1}, x_n]$




FIGURE 1. The construction of the factorization is best understood on a picture.

for each $n \in \mathbb{N}$. Now by Proposition 5.1 the space $f(M_n)$ is a compact \mathcal{H}^1 -negligible space. We define a sequence (P_n) of almost disjoint copies of $f(M_n)$. More precisely, for every n , let $i_n : P_n \rightarrow f(M_n)$ be an isometry such that $i_n^{-1}(f(x_n)) = i_{n+1}^{-1}(f(x_n))$ and $P_n \cap P_{n+1} = \{i_n^{-1}(f(x_n))\}$.

We have $f \upharpoonright_{M_n} = i_n \circ (i_n^{-1} \circ f \upharpoonright_{M_n})$. Now we endow the set $P = \bigcup P_n$ with the following distance d_P : for $x \in P_n$ and $y \in P_l$ with $n, l \in \mathbb{N}$

$$d_P(x, y) := \begin{cases} d(i_n(x), i_n(y)), & \text{if } l = n \\ d(i_n(x), f(x_n)) + d(f(x_n), i_{n+1}(y)), & \text{if } l = n + 1 \\ d(i_n(x), f(x_n)) + \sum_{j=n}^{l-2} d(f(x_j), f(x_{j+1})) + d(f(x_{l-1}), i_l(y)) & \text{if } l \geq n + 2. \end{cases}$$

We define $g : M \rightarrow P$ and $h : P \rightarrow N$ by

$$g(x) = i_n^{-1} \circ f(x) \text{ if } x \in M_n \text{ and } h(x) = i_n(x) \text{ if } x \in P_n.$$

Notice that g and h are well-defined, and $h \circ g = f$. It is not difficult to check that g is $\|f\|_L$ -Lipschitz and h is 1-Lipschitz.

Consider the completion Q of P . We claim that $Q \setminus P$ is empty or a singleton. Indeed, let $(z_n) \subset P$ such that $z_n \rightarrow x$. If $(z_n) \subset \bigcup_{i \in F} P_i$ for a finite set F , then $x \in \bigcup_{i \in F} P_i$ by compactness of $\bigcup_{i \in F} P_i$. Thus $x \in P$. It remains to address the

case where no such finite set F exists. In this case, we will show that $g(x_n) \rightarrow x$ which will finish the proof of the claim. Let us denote by $m_n \in \mathbb{N}$ one of the indices such that $z_n \in P_{m_n}$. We may assume, by passing to a subsequence of (z_n) , that (m_n) is strictly increasing. Since (z_n) is Cauchy, passing to a further subsequence, we may assume that $\sum d_P(z_n, z_{n+1}) < \infty$. Taking into account that $z_n \in P_{m_n}$ and $z_{n+1} \in P_{m_{n+1}}$, observe from the definition of d_P that

$$\begin{aligned} d_P(g(x_{m_n}), g(x_{m_n} + 1) + \dots + d_P(g(x_{m_{n+1}-1}), g(x_{m_{n+1}}))) \\ \leq d_P(z_n, z_{n+2}) \leq d_p(z_n, z_{n+1}) + d_P(z_{n+1}, z_{n+2}) \end{aligned}$$

for every $n \in \mathbb{N}$. This implies that $\sum d_P(g(x_n), g(x_{n+1})) < \infty$. In particular, the sequence $(g(x_n))$ converges in Q . On the other hand, since $d_P(z_n, g(x_{m_n})) \leq d_P(z_n, z_{n+1})$ by definition of d_P , we conclude that $g(x_n) \rightarrow x$ and our claim is proved.

Therefore Q is complete and p1u. Since $\|h\|_L \leq 1$ and since N is complete, h extends to Q with the same Lipschitz constant.

In the case when M is unbounded below as well as above we will get by a similar argument that $Q \setminus P$ has at most two elements. The details are left to the reader. \square

6.4. Factoring through Schur spaces and RNP spaces. We now turn to the “factorization through Schur or RNP spaces” problem:

Question 1. Let $f: M \rightarrow N$ be a curve-flat Lipschitz function.

- (1) Does it follow that \widehat{f} factors through a Schur space?
- (2) Does it follow that \widehat{f} factors through a space with the RNP?

This question remains largely open, we only have a few observations. Of course, as we already mentioned, if f factors through a complete p1u metric space then \widehat{f} factors through a Schur space with RNP. Still in the way of positive results towards Question 1, we have the following “reduction to bounded sets” principle which we will apply to several cases.

Proposition 6.13. *Let (P) be any property stable under ℓ_1 sums (such as the Schur property or the RNP). Let $f \in \text{Lip}_0(M, N)$ such that there exists $C > 0$ for which, for every bounded set $B \subset M$, $\widehat{f}|_{\mathcal{F}_M(B)}$ factors through a space Z that has the property (P) with $\widehat{f}|_{\mathcal{F}_M(B)} = \phi \circ q$ where $\phi: Z \rightarrow \mathcal{F}(N)$ and $q: \mathcal{F}_M(B) \rightarrow Z$ satisfy $\|\phi\| \|q\| \leq C$. Then \widehat{f} factors through a space with property (P) .*

The proof is based on Kalton’s decomposition [29, Section 4]. In fact, we will use the similar construction presented in [7, Section 2.2]. For every $n \in \mathbb{Z}$ and $x \in M$, let

$$\Lambda_n(x) = \begin{cases} 0 & \text{if } d(x, 0) \leq 2^{n-1} \\ 2^{-n+1}d(x, 0) - 1 & \text{if } 2^{n-1} \leq d(x, 0) \leq 2^n \\ 2 - 2^{-n}d(x, 0) & \text{if } 2^n \leq d(x, 0) \leq 2^{n+1} \\ 0 & \text{if } d(x, 0) \geq 2^{n+1}. \end{cases}$$

Note that $\|\Lambda_n\|_L = \frac{1}{2^{n-1}}$. Let us denote $M_n = B(0, 2^{n+1}) \setminus B(0, 2^{n-1})$ so that $\text{supp}(\Lambda_n) \subset M_n$. Next, for each $n \in \mathbb{Z}$, we introduce the pointwise multiplication operator $T_n: \text{Lip}_0(M) \rightarrow \text{Lip}_0(M)$ given by $T_n(f)(x) = f(x)\Lambda_n(x)$. As explained in [7, Section 2.2], T_n defines a weak*-to-weak* continuous linear operator from

$\text{Lip}_0(M)$ into $\text{Lip}_0(M)$ with $\|T_n\| \leq 5$. We will denote by $W_n : \mathcal{F}(M) \rightarrow \mathcal{F}(M)$ its pre-adjoint operator. Notice that:

$$\forall x \in M, \quad W_n(\delta(x)) = \Lambda_n(x)\delta(x).$$

Since $\text{supp}(\Lambda_n) \subset M_n$, it is readily seen that $\text{supp}(W_n(\gamma)) \subset M_n$ for every $\gamma \in \mathcal{F}(M)$. Therefore we can view W_n as an operator from $\mathcal{F}(M)$ to $\mathcal{F}_M(M_n) \equiv \mathcal{F}(M_n \cup \{0\})$. Now for every $\gamma \in \mathcal{F}(M)$, one has

$$\sum_{n \in \mathbb{Z}} \|W_n \gamma\| \leq 45 \|\gamma\| \quad \text{and moreover} \quad \gamma = \sum_{n \in \mathbb{Z}} W_n \gamma$$

where the convergence of the series is in norm. So $T : \mathcal{F}(M) \rightarrow \left(\sum_{n \in \mathbb{Z}} \mathcal{F}_M(M_n)\right)_{\ell_1}$ given by

$$\forall \gamma \in \mathcal{F}(M), \quad T(\gamma) = (W_n(\gamma))_{n \in \mathbb{Z}},$$

defines a bounded operator with norm $\|T\| \leq 45$.

Proof. Using the notation introduced above, we define $S : \left(\sum_{n \in \mathbb{Z}} \mathcal{F}_M(M_n)\right)_{\ell_1} \rightarrow \ell_1(\mathcal{F}(N))$ by:

$$\forall (\gamma_n)_n \in \left(\sum_{n \in \mathbb{Z}} \mathcal{F}_M(M_n)\right)_{\ell_1}, \quad S((\gamma_n)_n) := \left(\widehat{f}|_{\mathcal{F}_M(M_n)}(\gamma_n)\right)_n.$$

By assumption, each map $\widehat{f}|_{\mathcal{F}_M(M_n)}$ factors through a space Z_n with property (P):

$$\begin{array}{ccc} \mathcal{F}_M(M_n) & \xrightarrow{\widehat{f}|_{\mathcal{F}_M(M_n)}} & \mathcal{F}(N) \\ & \searrow q_n & \nearrow \phi_n \\ & & Z_n \end{array}$$

with $\|\phi_n\| \|q_n\| \leq C$. Of course, after rescaling, we may as well assume that $\|q_n\| \leq 1$ and $\|\phi_n\| \leq C$. We readily deduce the next factorization for S :

$$\begin{array}{ccc} \left(\sum_{n \in \mathbb{Z}} \mathcal{F}_M(M_n)\right)_{\ell_1} & \xrightarrow{S} & \ell_1(\mathcal{F}(N)) \\ & \searrow (q_n)_n & \nearrow (\phi_n)_n \\ & & \left(\sum_{n \in \mathbb{Z}} Z_n\right)_{\ell_1} \end{array}$$

where the operators $(q_n)_n$ and $(\phi_n)_n$ are well-defined bounded linear operators, with $\|(q_n)_n\| \leq 1$ and $\|(\phi_n)_n\| \leq C$. Notice that $\left(\sum_{n \in \mathbb{Z}} Z_n\right)_{\ell_1}$ has the property (P), since this property is stable under ℓ_1 -sums. To conclude, we introduce $R : \ell_1(\mathcal{F}(N)) \rightarrow \mathcal{F}(N)$ given by $R((\mu_n)_n) = \sum_n \mu_n$. It is an easy exercise to check that $\widehat{f} = R \circ S \circ T = R \circ (q_n)_n \circ (\phi_n)_n \circ T$. \square

Practically speaking, in the Propositions 6.1 and 6.6 concerning Fact_{p1u} , we haven't explicitly mentioned it, but the factorizations are actually uniformly bounded: Let M be a proper metric space and N any metric space. If $f \in \text{lip}_0(M, N)$, then \widehat{f} factors through a Schur space with RNP. Indeed, let $B \subseteq M$ be

a bounded set. Then B is compact and $f|_B \in \text{lip}_0(B, N)$. Thanks to Proposition 6.1, we have that the following diagram commutes:

$$\begin{array}{ccc} B & \xrightarrow{f|_B} & N \\ & \searrow \pi_B & \nearrow (f|_B)_{\text{lip}} \\ & & B_{\text{lip}} \end{array}$$

Notice that $\|\pi_B\| \leq 1$, $\|(f|_B)_{\text{lip}}\|_L \leq \|f\|_L$ (the upper bounds are not depending on the choice of B), and B_{lip} is p1u. Now, Proposition 6.13 concludes that \widehat{f} factors through a Schur space with RNP.

On the other hand, suppose that M is a proper metric space which is countably 1-rectifiable and $f \in \text{CF}_0(M, N)$. Then \widehat{f} does not fix a copy of L_1 ; so by Remark 5.2 we obtain that $f(M)$ is \mathcal{H}^1 -negligible. It follows that for any bounded subset B of M , the set $f(B)$ is compact and \mathcal{H}^1 -negligible. Therefore, f satisfies the assumption in Proposition 6.13. Summarizing, we observe the following corollary.

Corollary 6.14. *Let M be a proper metric space and N any metric space.*

- (1) *If $f \in \text{lip}_0(M, N)$, then $\widehat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$ factors through a Schur space with RNP.*
- (2) *Assuming M is countably 1-rectifiable, if $f \in \text{CF}_0(M, N)$, then \widehat{f} factors through a Schur space with RNP.*

7. LIPSCHITZ OPERATORS THAT DO NOT FIX A COPY OF ℓ_1

We conclude this paper with the discussion of Lipschitz operators that do not fix a copy of ℓ_1 . In fact, we will deal with a more general setting by considering the weighted version of Lipschitz operators. Let $w : M \rightarrow \mathbb{R}$ be any map (referred to as “the weight”). Let $f \in \text{Lip}_0(M, N)$. Inspired by the definition of \widehat{f} , one can define a linear map $w\widehat{f} : \text{span}(\delta(M)) \rightarrow \mathcal{F}(N)$ by setting

$$w\widehat{f}\left(\sum_{i=1}^n a_i \delta(x_i)\right) = \sum_{i=1}^n a_i w(x_i) \delta(f(x_i)).$$

When this map is bounded, it has a unique extension still denoted by $w\widehat{f} : \mathcal{F}(M) \rightarrow \mathcal{F}(N)$. Such bounded operators are named *weighted Lipschitz operators*, and they are characterized in [3, 14]. It is proved in [3] that $w\widehat{f}$ is compact if and only if it is weakly compact. The goal of this section is to prove that this is also equivalent to $w\widehat{f}$ not fixing a copy of ℓ_1 . In turn, this will let us easily show that it is also equivalent to $w\widehat{f}$ being strictly singular and to $w\widehat{f}$ being Asplund.

Theorem 7.1. *Let M, N be infinite pointed metric spaces, $f : M \rightarrow N$ and $w : M \rightarrow \mathbb{K}$ in such a way that $w\widehat{f}$ defines a bounded operator. Then the following assertions are equivalent*

- (i) *$w\widehat{f}$ is compact.*
- (ii) *$w\widehat{f}$ is strictly singular.*
- (iii) *$w\widehat{f}$ is Asplund.*
- (iv) *$w\widehat{f}$ does not fix any copy of ℓ_1 .*

Proof. The implications (i) \implies (ii) and (ii) \implies (iv) are clear. On the other hand, (i) \implies (iii) follows from the fact that every weakly compact operator

(in particular every compact operator) factors through a reflexive space (see [16]). The implication (iii) \implies (iv) follows from the fact that every Asplund operator factors through an Asplund space.

We will prove the contrapositive of (iv) \implies (i) using [3, Proposition 4.1], which states that $w\widehat{f}$ is compact if and only if

$$S := \left\{ \frac{w(x)\delta(f(x)) - w(y)\delta(f(y))}{d(x,y)} : x \neq y \in M \right\}$$

is relatively compact.

Assuming that $w\widehat{f}$ is not compact, there exists a sequence $(\gamma_n)_n \subset S$ which does not have any convergent subsequence. Let us express this sequence as:

$$\forall n \in \mathbb{N}, \quad \gamma_n = \frac{w(x_n)\delta(f(x_n)) - w(y_n)\delta(f(y_n))}{d(x_n, y_n)}.$$

Since the support of each γ_n contains at most 2 elements, it follows from [2, Theorem C] that $(\gamma_n)_n$ does not admit any weakly Cauchy subsequence. Thus, by Rosenthal's ℓ_1 -theorem and by passing to a subsequence, we may assume that $(\gamma_n)_n$ is equivalent to the ℓ_1 canonical basis.

At the same time, $(m_{x_n, y_n})_n$ cannot have any weakly Cauchy subsequence since $w\widehat{f}$ is weakly continuous and $w\widehat{f}(m_{x_n, y_n}) = \gamma_n$. So, by Rosenthal's ℓ_1 -theorem and by passing to a further subsequence if necessary, we may also assume that $(m_{x_n, y_n})_n$ is equivalent to the ℓ_1 canonical basis. Now, $X := \overline{\text{span}}\{m_{x_n, y_n} : n \in \mathbb{N}\}$ is isomorphic to ℓ_1 , and $w\widehat{f}|_X$ is an isomorphism onto its range. \square

It is clear that if \widehat{f} is strictly singular, then for any infinite subset $E \subset M$, $f|_E$ is not bi-Lipschitz. The following example shows that the converse is not true.

Example 7.2. Let $M = \{0\} \cup (x_n)_n \cup (y_n)_n$ be a countable metric space, with the distance defined by:

$$d(p, q) = \begin{cases} \frac{1}{n^2}, & \text{if } p = x_n, q = y_n, n \in \mathbb{N} \\ \frac{1}{n}, & \text{if } p \in \{x_n, y_n\}, q = 0 \\ \frac{1}{n} + \frac{1}{m} & \text{if } p \in \{x_n, y_n\}, q \in \{x_m, y_m\}, n \neq m \in \mathbb{N}. \end{cases}$$

Essentially, it is a pair of sequences $(x_n)_n, (y_n)_n$ converging to 0 at a slower rate than the distance $d(x_n, y_n)$ converges to 0. Now, consider the function $f: M \rightarrow [0, 1]$ defined by $f(0) = f(y_n) = 0$ and $f(x_n) = \frac{1}{n^2}$ for all $n \in \mathbb{N}$. Note that f is 1-Lipschitz. Moreover f attains its Lipschitz constant at each pair (x_n, y_n) , which means that f is not uniformly locally flat. Particularly, thanks to [2, Theorem A], this indicates that \widehat{f} is not compact and so not strictly singular, as shown in Theorem 7.1.

However, for every $\rho > 0$ there exists a $n_0 \in \mathbb{N}$ such that $\frac{1}{n_0} < \rho$. Straightforward computations show that if $n \neq m \geq n_0$, then

$$\forall p \in \{x_n, y_n\} \text{ and } \forall q \in \{x_m, y_m\} \implies d(f(p), f(q)) < \rho d(p, q).$$

Therefore f is not bi-Lipschitz when restricted to any infinite subset of M .

APPENDIX: SHORT PROOF OF THEOREM A IN [5]

Theorem A in [5] is used repeatedly along this article. Its original proof in [5] is fairly complex, and it uses some underlying principles previously used to show Lemma 3.4 in [10]. In this appendix, we include a short proof of Theorem A in [5]

based directly on Lemma 3.4 in [10]. This proof was communicated to the authors of [5] by Sylvester Eriksson-Bique. We are grateful to him for allowing us to include it in this paper.

Proposition 7.3. *Let M be a compact purely 1-unrectifiable metric space. Then $B_{\text{lip}_0}(M)$ is w^* -dense in $B_{\text{Lip}_0}(M)$.*

Proof. Let f in $B_{\text{Lip}_0}(M)$ and $\varepsilon > 0$ be arbitrary. Set $f_0 := f$ and $\theta_0 := \text{diam}(M)$. Since M is purely 1-unrectifiable, f_0 is curve-flat. Applying Lemma 3.3, as we did in the proof of Lemma 3.4, we obtain a function $f_1 \in B_{\text{lip}_{\frac{1}{2}}}(M)$ such that $\|f_0 - f_1\|_\infty < \frac{\varepsilon}{2}$ and $\frac{|f_1(x) - f_1(y)|}{d(x,y)} \leq \frac{|f_0(x) - f_0(y)|}{d(x,y)} + \frac{1}{2}$ for every $x \neq y \in M$. Denote by θ_1 the radius at which f_1 is $\frac{1}{2}$ -flat. Importantly, f_1 is again curve-flat since M is purely 1-unrectifiable.

Continuing inductively in this way, we obtain sequences $(f_n)_n \subset B_{\text{Lip}_0}(M)$ and $(\theta_n)_n$ such that for all $n \in \mathbb{N}$ and all $x \neq y \in M$:

- (1) f_n is $\frac{1}{2^n}$ -flat at radius θ_n ,
- (2) $\|f_n - f_{n-1}\|_\infty < \frac{\varepsilon}{2^n}$, and
- (3) $\frac{|f_n(x) - f_n(y)|}{d(x,y)} \leq \frac{|f_{n-1}(x) - f_{n-1}(y)|}{d(x,y)} + \frac{1}{2^n}$ for every $x \neq y \in M$.

Let f_∞ be the limit of $(f_n)_n$ in the topology of uniform convergence. In particular, $(f_n)_n$ converges to f_∞ in the weak*-topology and $\|f - f_\infty\|_\infty < \varepsilon$.

It only remains to show that f_∞ is uniformly locally flat. Let $x \neq y \in M$ and fix an arbitrary $n \in \mathbb{N}$. For any $m \geq n$, we have by induction

$$\frac{|f_m(y) - f_m(x)|}{d(x,y)} \leq \frac{|f_n(y) - f_n(x)|}{d(x,y)} + \frac{1}{2^n}.$$

Taking $m \rightarrow \infty$ yields $\frac{|f_\infty(y) - f_\infty(x)|}{d(x,y)} \leq \frac{|f_n(y) - f_n(x)|}{d(x,y)} + \frac{1}{2^n}$. Notice that this implies that $\|f_\infty\|_L \leq 1$. Moreover, since f_n is $\frac{1}{2^n}$ -flat at radius θ_n , we obtain that f_∞ is $\frac{1}{2^{n-1}}$ -flat at radius θ_n . As it holds for arbitrary $n \in \mathbb{N}$, the map f_∞ is uniformly locally flat. \square

Let us note that this approach does not work to obtain Theorems 3.1 and 4.1. Particularly, one may attempt to use a similar argument in Propositions 3.5 and 4.3, approximating a curve-flat function by flatter and flatter functions, and hoping to take the uniform limit of a sequence of such approximations to obtain a uniformly locally flat function, whose linearization can easily be seen to be DP and RN respectively (indeed, when M is compact it is a compact operator as shown in [2, 28, 30]). However, we crucially need each approximation to be again curve-flat in order to continue the induction, which need not be true outside of the purely 1-unrectifiable domain case. In fact, this approach simply cannot work, since by Example 5.25 in [5] (also used in Example 6.8) there exists a compact metric space that admits a non-trivial curve-flat function, but where only constant functions are uniformly locally flat. Therefore, such a curve-flat function cannot be approximated at all by a uniformly locally flat function.

ACKNOWLEDGMENTS

We are grateful to Sylvester Eriksson-Bique for letting us include his short proof of Proposition 7.3. G. Flores was supported by Universidad de O'Higgins (112011002-PD-17706171-9). M. Jung was supported by June E Huh Center for

Mathematical Challenges (HP086601) at Korea Institute for Advanced Study. G. Lancien, C. Petitjean, A. Procházka and A. Quilis were partially supported by the French ANR project No. ANR-20-CE40-0006. A. Quilis was also supported by MCIN/AEI/10.13039/501100011033 (PID2021-122126NB-C33) and CNRS project PEPS JCJC 2024.

REFERENCES

- [1] A. Abbar, C. Coine and C. Petitjean, *On the dynamics of Lipschitz operators*, Integral Equations Operator Theory 93 (2021), no. 4, Paper No. 45, 27 pp.
- [2] A. Abbar, C. Coine and C. Petitjean, *Compact and weakly compact Lipschitz operators*, Proceedings of the Royal Society of Edinburgh: Section A Mathematics, 2023, vol. 153, no 3, p. 1002–1020. [1](#), [7](#), [7.2](#), [7](#)
- [3] A. Abbar, C. Coine and C. Petitjean, *A Pre-adjoint Approach on Weighted Composition Operators Between Spaces of Lipschitz Functions*, Results Math 79, 85 (2024). [7](#), [7](#)
- [4] D. Achour, P. Rueda, E.A. Sánchez-Pérez, R. Yahi, *Lipschitz operator ideals and the approximation property*, J. Math. Anal. Appl. 436 (2016), 217–236.
- [5] R. J. Aliaga, C. Gartland, C. Petitjean and A. Procházka, *Purely 1-unrectifiable spaces and locally flat Lipschitz functions*, Trans. Amer. Math. Soc. 375 (2022), 3529–3567. ([document](#)), [1](#), [1](#), [2.3](#), [4.1](#), [4.2](#), [4.2](#), [4.2](#), [5.1](#), [6.1](#), [6.1](#), [6.1](#), [6.1](#), [6.2](#), [6.3](#), [7](#), [7](#)
- [6] R. J. Aliaga and E. Pernecká, *Supports and extreme points in Lipschitz-free spaces*, Rev. Mat. Iberoam. 36 (2020), no. 7, 2073–2089. [2.1](#)
- [7] R. J. Aliaga, and E. Pernecká, *Integral representation and supports of functionals on Lipschitz spaces*, Int. Math. Res. Not. IMRN (2023), no. 4, 3004–3072. [6.4](#)
- [8] R. J. Aliaga, E. Pernecká, C. Petitjean and A. Procházka, *Supports in Lipschitz-free spaces and applications to extremal structure*, J. Math. Anal. Appl. 489 (2020), no. 1, 124128. [2.1](#)
- [9] R. J. Aliaga, C. Petitjean and A. Procházka, *Compact reduction in Lipschitz free spaces*, Studia Math. 260 (2021), no. 3, 341–359. [3](#), [3.2](#)
- [10] D. Bate, *Purely unrectifiable metric spaces and perturbations of Lipschitz functions*, Acta Math. 224 (2020), no.1, 1–65. [3](#), [3.1](#), [3.3](#), [3.1](#), [7](#)
- [11] J. Bourgain, *Dunford-Pettis operators on L^1 and the Radon-Nikodým property*, Israel J. Math. 37 (1980), no.1-2, 34–47.
- [12] J. Bourgain, *A characterization of non-Dunford-Pettis operators on L_1* , Israel J. Math. 37, 48–53 (1980). [3.7](#)
- [13] J. Bourgain, and H. P. Rosenthal, *Martingales valued in certain subspaces of L^1* , Israel J. Math. **37**, no. 1-2, 54–75. [5.3](#)
- [14] Sh. Behrouzi, A. Golbaharan and H. Mahyar, *Weighted composition operators between pointed Lipschitz spaces*, Results Math. 77 (2022), no. 4, Paper No. 157, 16 pp. [7](#)
- [15] A. Dalet, *Free spaces over countable compact metric spaces*, Proc. Amer. Math. Soc. (2015), 143 (8) : 3537–3546. [6.1](#)
- [16] W. J. Davis, T. Figiel, W. B. Johnson, A. Pełczyński, *Factoring weakly compact operators*, J. Functional Analysis 17(1974), 311–327. [7](#)
- [17] J. Diestel and J. J. Uhl, *Vector measures*, American Mathematical Society, Providence, R.I., 1977. [2.5](#), [5.2](#)
- [18] S. J. Dilworth and M. Girardi, *On Various Modes of Scalar Convergence in $L_0(X)$* , J. Math. Anal. Appl. **259** (2001), no. 2, 660–684 [4.5](#), [4.5](#)
- [19] O. Dovgoshey, O. Martio, V. Ryazanov and M. Vuorinen, *The Cantor function*, Expo. Math. **24** (2006), no. 1, 1–37. [6.2](#), [6.9](#)
- [20] G. A. Edgar, *Asplund operators and a.e. convergence*, J. Multivariate Anal. 10 (1980), no.3, 460–466. [2.2](#), [5.2](#)
- [21] H. Federer, *Geometric measure theory*, Die Grundlehren der mathematischen Wissenschaften, Band 153, Springer-Verlag New York, Inc., New York, 1969. [5.1](#)
- [22] D. Freeman and C. Gartland, *Lipschitz functions on quasiconformal trees*, Fund. Math. 262 (2023), no.2, 153–203. [5.1](#)
- [23] L. García-Lirola, C. Petitjean and A. Procházka, *Injectivity of Lipschitz operators*, Bull. Malays. Math. Sci. Soc. 46, 68 (2023). [6.7](#)

- [24] N. Ghoussoub, W. B. Johnson, *Counterexamples to several problems on the factorization of bounded linear operators*, Proc. Amer. Math. Soc. 92 (1984), 233–238. [2.2](#)
- [25] A. Godard, *Tree metrics and their Lipschitz-free spaces*, Proc. Amer. Math. Soc. 138 (2010), 4311–4320. [3.3](#), [5.1](#), [5.2](#)
- [26] G. Godefroy, *Boundaries of a convex set and interpolation sets*, Math. Ann., 277 (2) : 173–184, 1987. [6.1](#)
- [27] G. Godefroy and N. J. Kalton, *Lipschitz-free Banach spaces*, Studia Math. 159 (2003), 121–141. [5.3](#)
- [28] A. Jiménez-Vargas and M. Villegas-Vallecillos, *Compact composition operators on noncompact Lipschitz spaces*, J. Math. Anal. Appl. 398 (2013), no. 1, 221–229. [7](#)
- [29] N. J. Kalton, *Spaces of Lipschitz and Hölder functions and their applications*, Collect. Math. 55 (2004), no. 2, 171–217. [3.1](#), [6.4](#)
- [30] H. Kamowitz, S. Scheinberg, *Some properties of endomorphisms of Lipschitz Algebras*, Studia Math. 96 (1990), 61–67. [7](#)
- [31] B. Kirchheim, *Rectifiable metric spaces: local structure and regularity of the Hausdorff measure*, Proc. Amer. Math. Soc. 121 (1994), 113–123. [2.3](#)
- [32] P. Lefèvre *When Strict Singularity Of Operators Coincides With Weak Compactness*, Journal of Operator Theory, Vol. 67, No. 2 (2012), 369–378.
- [33] W. Linde, *An operator ideal in connection with the Radon-Nikodym property of Banach spaces*, Math. Nachr. 71 (1976), 65–73. [5.2](#)
- [34] A. Pełczyński, *Banach spaces on which every unconditionally converging operator is weakly compact*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.10 (1962), 641–648. [2.2](#)
- [35] A. Pełczyński, *On Strictly Singular and Strictly Cosingular Operators*, Vols. I and II, Bull. Acad. pol. sei. III (1965), Sér. Sci. math. astr. phys. 13, No 1.
- [36] Ju. Ī Petuniĭn and A. N. Pliĭčko, *Some properties of the set of functionals that attain a supremum on the unit sphere*, Ukrain. Mat. Ž., 26 : 102–106, 143, 1974. [6.1](#)
- [37] A. Pietsch, *Operator Ideals*, North-Holland Publishing Company, Mathematical Studies, 1980. [2.2](#)
- [38] G. Pisier, *Martingales in Banach spaces*, Cambridge University Press, Cambridge, 2016. [2.4](#), [2.4](#), [4](#), [4](#), [4.5](#)
- [39] O. I. Reĭnov, *The Radon-Nikodým property, and integral representations of linear operators*, Funkcional. Anal. i Priložen, Vol. 9 (1975), no. 4, 87–88.
- [40] W. Rudin, *Real and complex analysis*, 3rd. ed., McGraw-Hill, Singapore, 1987. [2.4](#)
- [41] C. Stegall, *The Radon-Nikodym property in conjugate Banach spaces. II*, Trans. Amer. Math. Soc. 264 (1981), no. 2, 507–519. [2.2](#)
- [42] N. Weaver, *Lipschitz algebras*, 2nd ed., World Scientific Publishing Co., River Edge, NJ, 2018. [2.1](#), [6.1](#), [6.10](#)

(G. Flores) UNIVERSIDAD DE O’HIGGINS, INSTITUTO DE CIENCIAS DE LA INGENIERÍA, AV. LIB. GRAL. BERNARDO O’HIGGINS 611, RANCAGUA, CHILE
Email address: gonzalo.flores@uoh.cl

(M. Jung) JUNE E HUH CENTER FOR MATHEMATICAL CHALLENGES, KOREA INSTITUTE FOR ADVANCED STUDY, 02455 SEOUL, REPUBLIC OF KOREA
Email address: jmingoo@kias.re.kr

(G. Lancien) UNIVERSITÉ DE FRANCHE-COMTÉ, CNRS, LMB (UMR 6623), F-25000 BESANÇON, FRANCE.

(C. Petitjean) UNIV GUSTAVE EIFFEL, UNIV PARIS EST CRETEIL, CNRS, LAMA UMR8050, F-77447 MARNE-LA-VALLÉE, FRANCE
Email address: colin.petitjean@univ-eiffel.fr

(A. Prochazka) UNIVERSITÉ DE FRANCHE-COMTÉ, CNRS, LMB (UMR 6623), F-25000 BESANÇON, FRANCE.
Email address: antonin.prochazka@univ-fcomte.fr

(A. Quilis) UNIVERSITÉ DE FRANCHE-COMTÉ, CNRS, LMB (UMR 6623), F-25000 BESANÇON, FRANCE.
Email address: andresqsa@gmail.com