



**HAL**  
open science

## **Radio signatures of star–planet interactions, exoplanets and space weather**

J. Callingham, B. Pope, R. Kavanagh, S. Bellotti, S. Daley-Yates, M. Damasso, J.-M. Grießmeier, M. Güdel, M. Günther, M. Kao, et al.

### ► **To cite this version:**

J. Callingham, B. Pope, R. Kavanagh, S. Bellotti, S. Daley-Yates, et al.. Radio signatures of star–planet interactions, exoplanets and space weather. *Nature Astronomy*, 2024, 8, pp.1359-1372. <10.1038/s41550-024-02405-6>. <hal-04785169>

**HAL Id: hal-04785169**

**<https://hal.science/hal-04785169v1>**

Submitted on 15 Jul 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

# Radio Signatures of Star-Planet Interactions, Exoplanets, and Space Weather

J. R. Callingham<sup>1,2\*</sup>, B. J. S. Pope<sup>3,4</sup>, R. D. Kavanagh<sup>1</sup>, S. Bellotti<sup>2</sup>, S. Daley-Yates<sup>5</sup>, M. Damasso<sup>6</sup>, J.-M. Grießmeier<sup>7,8</sup>, M. Güdel<sup>9</sup>, M. Günther<sup>10</sup>, M. M. Kao<sup>11,12</sup>, B. Klein<sup>13</sup>, S. Mahadevan<sup>14,15</sup>, J. Morin<sup>16</sup>, J. D. Nichols<sup>17</sup>, R. A. Osten<sup>18,19</sup>, M. Pérez-Torres<sup>20</sup>, J. S. Pineda<sup>21</sup>, J. Rigney<sup>22,23,24</sup>, J. Saur<sup>25</sup>, G. Stefánsson<sup>26,27</sup>, J. D. Turner<sup>28,29</sup>, H. Vedantham<sup>1,29</sup>, A. A. Vidotto<sup>2</sup>, J. Villadsen<sup>30</sup> and P. Zarka<sup>31,8</sup>

<sup>1</sup>ASTRON, Netherlands Institute for Radio Astronomy, Oude Hoogeveensedijk 4, Dwingeloo, 7991 PD, The Netherlands.

<sup>2</sup>Leiden Observatory, Leiden University, PO Box 9513, 2300 RA, Leiden, The Netherlands.

<sup>3</sup>School of Mathematics and Physics, University of Queensland, St Lucia, QLD 4072, Australia.

<sup>4</sup>Centre for Astrophysics, University of Southern Queensland, West Street, Toowoomba, QLD 4350, Australia.

<sup>5</sup>School of Physics and Astronomy, University of St Andrews, North Haugh, St Andrews, Fife KY16 YSS, UK.

<sup>6</sup>INAF - Osservatorio Astrofisico di Torino, Via Osservatorio 20, I10025 Pino Torinese, Italy.

<sup>7</sup>LPC2E, OSUC, Univ Orleans, CNRS, CNES, Observatoire de Paris, F-45071 Orleans, France.

<sup>8</sup>Observatoire Radioastronomique de Nançay (ORN), Observatoire de Paris, CNRS, PSL, Université d'Orléans, OSUC, route de Souesmes, Nançay, 18330, France.

<sup>9</sup>Department of Astrophysics, University of Vienna, Türkenschanzstr. 17, 1180, Vienna, Austria.

<sup>10</sup>European Space Agency (ESA), European Space Research and Technology Centre (ESTEC), Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands.

<sup>11</sup>Department of Astronomy & Astrophysics, University of California, Santa Cruz, CA, USA.

<sup>12</sup>Lowell Observatory, Flagstaff, AZ, USA.

<sup>13</sup>Department of Physics, University of Oxford, OX13RH, Oxford, UK.

<sup>14</sup>Department of Astronomy & Astrophysics, Pennsylvania State University, University Park, PA, 16802, USA.

<sup>15</sup>Center for Exoplanets and Habitable Worlds, Pennsylvania State University, University Park, PA, 16802, USA.

<sup>16</sup>LUPM, Université de Montpellier, CNRS, Place Eugène Bataillon, F-34095 Montpellier, France.

<sup>17</sup>Department of Physics and Astronomy, University of Leicester, Leicester, UK.

<sup>18</sup>Space Telescope Science Institute, Baltimore, MD 21218, USA.

<sup>19</sup>Center for Astrophysical Sciences, Department of Physics and Astronomy, Johns Hopkins University, 3400 North Charles Street, Baltimore, MD 21218, USA.

<sup>20</sup>Instituto de Astrofísica de Andalucía (IAA-CSIC), Consejo Superior de Investigaciones Científicas (CSIC), Glorieta de la Astronomía, s/n, Granada, E-18008, Spain.

<sup>21</sup>Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, Boulder, CO, USA.

<sup>22</sup>Astronomy & Astrophysics Section, DIAS Dunsink Observatory, Dublin Institute for Advanced Studies, Dublin, D15 XR2R, Ireland.

<sup>23</sup>Armagh Observatory and Planetarium, College Hill, Armagh BT61 9DG, N. Ireland.

<sup>24</sup>School of Mathematics and Physics, Queen's University Belfast, University Road, Belfast BT7 1NN, N. Ireland.

<sup>25</sup>Institute of Geophysics and Meteorology, University of Cologne, Albertus-Magnus-Platz, 50923 Cologne, Germany.

<sup>26</sup>Department of Astrophysical Sciences, Princeton University, 4 Ivy Lane, Princeton, NJ 08540, USA.

<sup>27</sup>NASA Sagan Fellow.

<sup>28</sup>Department of Astronomy and Carl Sagan Institute, Cornell University, Ithaca, NY, USA.

<sup>29</sup>Kapteyn Astronomical Institute, University of Groningen, PO Box 72, 97200 AB, Groningen, The Netherlands.

<sup>30</sup>Department of Physics & Astronomy, Bucknell University, Lewisburg, PA, USA.

<sup>31</sup>LESIA, Observatoire de Paris, Université PSL, CNRS,  
Sorbonne Université, Université de Paris, 5 place Jules Janssen,  
Meudon, 92195, France.

\*Corresponding author(s). E-mail(s): [callingham@astron.nl](mailto:callingham@astron.nl);

### Abstract

Radio detections of stellar systems provide a window onto stellar magnetic activity and the space weather conditions of extrasolar planets, information that is difficult to attain at other wavelengths. There have been recent advances observing auroral emissions from radio-bright low-mass stars and exoplanets largely due to the maturation of low-frequency radio instruments and the plethora of wide-field radio surveys. To guide us in placing these recent results in context, we introduce the foremost local analogues for the field: Solar bursts and the aurorae found on Jupiter. We detail how radio bursts associated with stellar flares are foundational to the study of stellar coronae, and time-resolved radio dynamic spectra offers one of the best prospects of detecting and characterising coronal mass ejections from other stars. We highlight the prospects of directly detecting coherent radio emission from exoplanetary magnetospheres, and early tentative results. We bridge this discussion to the field of brown dwarf radio emission, in which their larger and stronger magnetospheres are amenable to detailed study with current instruments. Bright, coherent radio emission is also predicted from magnetic interactions between stars and close-in planets. We discuss the underlying physics of these interactions and implications of recent provisional detections for exoplanet characterisation. We conclude with an overview of outstanding questions in theory of stellar, star-planet interaction, and exoplanet radio emission, and the prospects of future facilities in answering them.

We know that the vast majority of stars host planets [1] – now the focus of exoplanet science has shifted to understanding their detailed physics. Determining the composition of exoplanetary atmospheres, geospheres, and associated space weather conditions is considered paramount for assessing the potential habitability of a planet [2]. While transmission spectroscopy is beginning to reveal the composition of exoplanets [3, 4, 5, 6], the space weather and radiation environment that shape their atmospheric evolution and habitability remain relatively unknown.


The largest contributors to space weather in the Solar System are coronal mass ejections (CMEs) [7, 8], which launch a significant amount of hot, dense plasma from the Sun into the Solar System. The persistent impact of CMEs on a terrestrial planet has the potential to erode its atmosphere [9, 10], as may

the quiescent solar wind, particularly for planets without intrinsic magnetic fields [11, 12]. Despite the importance of understanding the plasma physics of CMEs, there has yet to be an unambiguous detection of a CME from a star other than our Sun. The winds of low-mass main sequence stars are also generally too tenuous to detect via current technology [13, 14] for all but a handful of stars [15]. Furthermore, there has not yet been a direct measurement of an exoplanet's magnetic field, which is crucial information for understanding its atmospheric evolution, as a planet's magnetic field could protect the atmosphere from the impact of stellar plasma [16, 17].

Radio observations, particularly at low frequencies ( $\lesssim 300$  MHz), are a unique probe of stellar and planetary plasma environments [18, 19, 20]. As observed on our Sun, CMEs are often accompanied by bursty, low-frequency radio emission that encodes the kinematics of the plasma as it is ejected into interplanetary space [18, 21]. Furthermore, the incident solar wind on the magnetised planets drives auroral emission [22], particularly at radio wavelengths [23]. While optical emission from exoplanet aurorae will likely be difficult to observe even with extremely large telescopes [24], direct detection of auroral radio emission from an exoplanet has been a long-standing goal of the radio astronomy community. This would allow us to infer the presence, topology, and strength of exoplanetary magnetic fields for the first time [25, 26, 23]. At other wavelength regimes, such information about the magnetic field of an exoplanet is either model dependent [27] or untraceable [20].

While the goals of detecting CMEs, stellar winds, and planetary magnetic fields in exoplanetary systems have been pursued for decades, recent observational progress has been made due to both the maturation of low-frequency radio arrays and the increased sensitivity at gigahertz-frequencies. For example, observations from the LOW-Frequency ARray (LOFAR) [28], Giant Metrewave Radio Telescope (GMRT), Karl G. Jansky Very Large Array (JVLA), Five-hundred-meter Aperture Spherical Telescope (FAST), and Australian Square Kilometre Array Pathfinder (ASKAP) [29] have begun to uncover new and diverse signals from radio stellar systems [30], some potentially consistent with interactions between stars and planets analogous to some phenomena observed in our Solar System [31, 32, 33, 34, 35, 36, 37]. This advancement in observational radio astronomy has also been paired with a revolution in optical and near-infrared facilities, in particular the Transiting Exoplanet Survey Satellite (TESS) [38], near-infrared radial velocity [39, 40] and Zeeman-Doppler imaging (ZDI) [41, 42] surveys. The combination of these radio and optical/near-infrared facilities provides an unparalleled opportunity to trace key stellar activity indicators across the electromagnetic spectrum, informing us about whether emission is driven by coronal or magnetospheric processes, and providing probes of extrasolar space weather environments.

This manuscript is focused on communicating the recent observational advances that have been made on coherent radio signatures of star-planet interactions, exoplanets, ultracool dwarfs, and space weather, and the associated basic foundational theories – with an aim to be understandable and a primer for new doctoral students and scholars to the field.




**Fig. 1** Schematic representation of a CME and the radio dynamic spectrum of the event that shows Type II and III bursts. The Type II burst is produced in the coronal shock front, as represented by the blue region emanating from the red star in the left panel. The magnetic reconnection event that is allowing mass to escape the magnetosphere of the star is shown as the yellow region. Type III bursts are produced on open field lines surrounding the magnetic reconnection event. Some structure and higher frequency harmonic emission is evident in and around the Type II burst in the right panel. The dynamic spectrum is in total intensity.

## 1 Radio emission in the Solar System

The properties of radio emission observed from the Sun and Solar System planets are often used as templates for interpreting radio emission from stellar systems. Solar radio emission is often incoherent, implying the electrons producing the radio emission do not act in phase, and is observed over a broad range of frequencies up to a few gigahertz [18, 43]. Such incoherent radio emission is often associated with magnetic reconnection events in the Solar corona, or active regions on the Sun, where mildly-relativistic electrons produce gyrosynchrotron emission by their motion along magnetic field lines.

Coherent solar radio emission is predominantly generated by plasma emission via plasma waves. These plasma waves are excited by beams of energetic electrons that have been accelerated by magnetic reconnection events or shocks [18, 43]. Phenomenologically, the two most important types of solar bursts for tracing space weather are classed as Type II and Type III bursts. Type II bursts, lasting from several minutes to hours, are usually produced by the acceleration of electrons by shocks at the leading edges of outward-moving CMEs. Type II bursts tend to slowly decrease in frequency over several minutes, providing a measure of how the plasma density decreases as the wave propagates out of the solar corona. Type III bursts are short ( $\sim$  few seconds) events associated with electron beams accelerated by reconnection events, and often accompany Type II bursts. A schematic of the set up of a CME, and the dynamic spectrum of Type II and III bursts, is provided in Figure 1. In Figure 2 we provide a practical, heuristic guide of differentiating between the different radio emission mechanisms operating in stellar systems.

In contrast, radio emission from the magnetised Solar System planets is dominated by auroral processes. What this loosely refers to is emission that is



**Fig. 2** Schematic for distinguishing the emission mechanism operating when a radio stellar system is detected. The emission characteristics listed for differentiating between the different mechanisms should be treated as an approximate guideline. For a more physical differentiation based on the conditions of the plasma, the reader is referred to [18, 43].  $\delta\nu/\Delta\nu$  represents the amount of bandwidth  $\delta\nu$  the emission occupies relative to the available bandwidth  $\Delta\nu$ , assuming a relative large fractional bandwidth. Differentiating between plasma emission and emission from the electron cyclotron maser (ECM) instability can be difficult if the time-frequency structure of the radio emission can not be resolved. In that case, arguments can be made in favour of one emission mechanism based on the coronal scale height of the radio star, as derived from the stellar X-ray luminosity [44, 45]. Fundamental and harmonic plasma emission have different circular polarisation fractions and maximum brightness temperatures, with harmonic plasma emission able to reach the highest brightness temperatures but limited to  $\lesssim 60\%$  circular polarisation fraction [45]. Note that the polarisation fractions reported do not take into account propagation effects, which often suppresses the fractional amount [46].

generated close to the magnetic poles of the planet. A key ingredient to driving auroral emission is electron acceleration. The injection of a high velocity electron population into a quiescent magnetosphere powers bright, coherent radio emission via the electron cyclotron maser (ECM) instability [47, 48]. ECM emission is highly circularly polarised and occurs at the local cyclotron frequency. The maximum frequency of ECM emission is directly proportional to the ambient magnetic field strength at the emitting point. Therefore, ECM emission is a direct probe of the magnetic field strength of the emitting body [18]. Another characteristic of ECM emission is that it is beamed, in that the radio waves propagate outwards on the surface of a cone that is near-perpendicular to the local magnetic field. A result of this beaming is that the emission is only generally visible for brief windows in time and can exhibit complicated periodicity [49].

Several distinct mechanisms can produce the population of accelerated electrons in planetary magnetospheres. The first is due to the interaction of the solar wind directly with a planet's magnetosphere. In this case, the incident solar wind is magnetised and compresses the planet's intrinsic magnetosphere, resulting in magnetic reconnection on the nightside of the planet. This process is often referred to as the Dungey cycle [22], and interaction of the high energy electrons with the atmosphere produces the aurora australis and borealis on

Earth. Interestingly, the observed auroral radio power from the magnetised Solar System planets is seen to directly scale with the power of the incident solar wind, both kinetically and magnetically [23, 50, 51, 52] – with the kinetic relation often referred to as the radiometric Bode’s law, and the magnetic relation as the radio-magnetic scaling law.

Despite being the brightest low-frequency radio emitter in the Solar System, Jupiter’s auroral emission deviates from this narrative as two different processes dominate the auroral radio emission: the breakdown of co-rotation and the Jupiter-satellite interaction [53, 54]. For the breakdown of co-rotation model, Jupiter’s inner magnetosphere is continuously supplied with plasma from Io’s volcanic outgassing. The inner plasma is forced to co-rotate with the magnetic field of Jupiter as it is centrifugally expelled outward. At a certain distance in Jupiter’s magnetosphere, the magnetic field is not strong enough to force co-rotation of the plasma, producing a shear against the outer plasma environment. This shearing generates a current system in which electrons are accelerated from the equator to the poles, producing the auroral ring [55, 56, 57].

The brighter and more localised radio emission from Jupiter is associated with the magnetic field lines of Jupiter connecting it to its Galilean moons, particularly to Io [58, 59, 54, 60]. The driving mechanism of this emission is thought to either be due to Alfvén waves<sup>1</sup> generated by the perturbation of Jupiter’s magnetic field by the inner moons [59], or magnetic reconnection occurring between the magnetic fields of Jupiter and the moons [61, 62, 63]. In either case, electrons are accelerated, subsequently producing bright circularly-polarised radio emission via the ECM mechanism. A constraint on this form of emission is that it can only be produced if the perturbing body orbits inside the Alfvén surface<sup>2</sup> of the host body for star-planet interactions, a region where the host body’s magnetic field dominates (see Section 5.1 for further details).


The observed radio emission processes from Jupiter serve as archetypes for what is expected from extrasolar planetary systems, where it is possible to produce emission that is orders of magnitude brighter than that observed in Solar System by scaling the interaction by mass, obstacle size, magnetic field strength of the stellar wind and of the planetary obstacle, and orbital distance (see Section 5.1 for quantitative expressions of this statement) [23, 52, 64]. These interactions are often referred to as magnetic star-planet interactions in the literature, which will refer to generally as star-planet interactions (SPI) in this manuscript.

As illustrated in Figure 3, there are two sub-types of SPI that are relevant in producing ECM radio emission from radio stellar systems that contain a low-mass host star:

---

<sup>1</sup>Alfvén waves are transverse plasma waves that travel along magnetic field lines. They are produced when ions oscillate in response to the restoring force provided by the tension of magnetic field lines.

<sup>2</sup>The Alfvén surface is defined as the three-dimensional boundary that separates a star’s corona from its stellar wind – the boundary at which information in the stellar wind can not propagate back to the surface of the star. This is the locus where the Alfvénic Mach number is unity.



**Fig. 3** Sketch illustrating the two putative sources of electron cyclotron maser (ECM) emission in exoplanetary systems. *Left Panel:* Emission induced on a star by a close-in planet. If the planet orbits inside the Alfvén surface of the star, it can perturb the star’s magnetic field, producing Alfvén waves which propagate back towards the star. These waves interact with electrons, accelerating them towards the stellar surface. Electrons with sufficiently large pitch angles undergo a magnetic mirroring effect and are reflected, producing ECM emission which propagates in a hollow cone. *Right Panel:* Auroral emission induced on a planet by the interaction of its magnetosphere with the incident wind of its host star. The magnetic field carried by the stellar wind causes the planet’s magnetosphere to open up on the dayside (left). These field lines are pushed towards the nightside (right), where they subsequently reconnect. The energy released in the magnetic reconnection accelerates electrons back towards the poles along the field line highlighted in red, where they reflect and power ECM emission in a similar manner to that described for the left panel. For clarity, the emission cone is only shown for the Northern hemisphere in both cases.

### 1. Sub-Alfvénic interactions:

Analogous to the Jupiter-Io interaction. Radio emission occurs along the stellar magnetic field line connecting the star and planet, driven by interactions between the field line and the planet. It requires the planet to be orbiting inside the Alfvén surface, with either an intrinsic or induced magnetic field. It is sometimes referred to as magnetic star-planet interactions. The power produced by this interaction appears to only follow the radio-magnetic scaling law [65].

### 2. Wind-magnetosphere interactions:

Auroral radio emission is produced in a ring-like configuration near the planet’s magnetic poles, driven by the interaction of the incident stellar wind and planetary magnetosphere. The power produced by this interaction appears to follow both the radiometric Bode’s law and the radio-magnetic scaling law for most auroral sources in the Solar System.

Naturally, the SPI terminology is no longer accurate once either the star or planet are replaced with a brown dwarf or moon. However, the underlying physical processes are still relevant in those scenarios. For instance, an

exomoon could produce detectable radio emission on an exoplanet via a sub-Alfvénic interaction. Stochastic flaring and breakdown of co-rotation [66] are also pathways to generate coherent radio emission in these systems, however since these processes do not explicitly rely on the presence of a planet, they do not fall under the umbrella of SPI. We also note that the term SPI often more broadly also refers to the irradiation of the planetary atmospheres by the host star’s light and its subsequent interactions with the wind of the host star [67] and tidal interactions between the two bodies [68]. However, neither of these processes are expected to directly produce bright signatures in the radio, therefore we do not discuss them in this manuscript.

## 2 Stellar Flares and Coronal Mass Ejections

Radio observations of the Sun and other solar-like and low-mass stars have shown that stellar flare production involves all layers of the stellar atmosphere, from the photosphere to the corona. The large extent of physical scales involved in producing flares implies that a range of processes are responsible, such as coronal heating and particle acceleration [69]. A consensus picture is emerging that connects the radio, optical, ultraviolet, and X-ray flare luminosities of stars to coronal heating. However, one remaining challenge is using time-resolved, simultaneous multiwavelength observations [70] to trace coronal mass ejections, both to understand stellar mass loss and to directly probe the space weather environments faced by exoplanets.

There is a well-established model for the non-thermal incoherent radio emission from solar and stellar flares. The model is motivated by a remarkable correlation between the quasi-steady gigahertz-frequency radio  $L_{\nu,\text{rad}}$  and soft X-rays  $L_X$  luminosities:  $L_X \propto L_{\nu,\text{rad}}^{0.73}$  [71, 72]. This correlation is often referred to as the Güdel-Benz relation. Such a relation is canonically explained via the fact magnetic reconnection events produce streams of non-thermal, accelerated coronal electrons. These streams impact the chromosphere and cause evaporation of million-degree plasma, depositing their kinetic energy as thermal energy in the corona [73]. The free electrons produced by the magnetic reconnection event are observationally traced by the gigahertz-frequency gyrosynchrotron emission as they radiate in the stellar magnetic field, while the heated plasma is identified by its thermal soft (0.2-2 keV) X-ray emission. It is important to note that while the explanation presented here is qualitatively plausible, it has difficulties in quantitatively describing the Güdel-Benz relation [74]. The Güdel-Benz relation equally applies over many orders of magnitude to most solar and stellar flares, suggesting that very hot, active stellar coronae are heated by the integrated energy of flares – including those flares that may be too low energy to detect individually [75, 76].

Small scale, stochastic flares could indeed be responsible for the quasi-steady soft X-ray and low circularly polarised, steady gigahertz-frequency radio emission from stellar systems. If the energy distribution  $E$  of stellar radio flares follows a power law, low energy flares will be largely responsible for

heating the corona if the power-law index  $\alpha$  is  $> 2$  in  $dN/dE \propto E^{-\alpha}$ . However, this has been challenging to test in the radio: observational flare distributions are limited at low  $E$  by the instrumental detection limit, depends on stellar distance and flare peak luminosity, and long time series are required.

At short UV, optical, and the non-thermal hard X-ray ranges, Solar and stellar flare statistics do indeed reveal power-law distributions down to the light curve noise, mostly with  $\alpha \gtrsim 2$  for active stars [77, 78, 79, 80, 81, 82, 83, 84, 85, 86]. These optical stellar flare statistical studies have been revolutionised since the *Kepler* [87] and TESS [88] space telescopes have produced hundreds of thousands of high-cadence, high-precision broadband optical light curves. Homogeneous statistical catalogs of flares [89, 90, 91, 92, 93, 94, 95, 96] have been used to investigate the population level statistics [97, 98], flares characteristics of planet-hosts [99, 100, 101], and have revealed Carrington-analog superflares on solar-like stars [102, 103, 104, 105, 106, 107].

New radio facilities are for the first time capable of probing whether large optical flares are accompanied by coronal mass ejections. While this is true for the largest of the Sun's flares [108], the commonly flaring M dwarfs have different magnetic geometries and stronger global fields [42, 109]. Recent modelling efforts [110] have shown that large overlying magnetic fields can prevent mass breakout, resulting in the occurrence of confined flares, which do not have an accompanying eruption. The ongoing, all-sky TESS mission uniquely complements modern radio telescopes for multiwavelength flare studies: simultaneous ASKAP and TESS observations have traced a radio burst from Proxima Centauri [32], and also radio emission with no optical counterpart [111]. The TESS flare rates of LOFAR-detected stars are correlated with their X-ray luminosities, consistent with the Güdel-Benz mechanism, but several show high ECM radio luminosity and few or no flares or X-ray emission. Such a deviation has been taken as evidence in support of the radio emission being generated by star-planet magnetic interaction since radio emission generated by plasma or gyro-synchrotron mechanisms are expected to correlate with magnetic events that often have an optical/X-ray flare counterpart [112].

While Type II bursts are the only unambiguous radio proof of CME material escaping from the stellar magnetosphere, there are no firm detections of Type II bursts from other stars despite significant observational effort [113, 114, 115, 116]. The non-detections of extrasolar Type II bursts may be due to sensitivity limitations, magnetic confinement of CMEs on radio-bright stars [117], or an Alfvén speed that prevents shock formation [118]. Instead, radio observations of active M dwarfs have found other types of coherent radio bursts, including hours-long events attributed to ECM emission [115, 119, 116, 120] that do not have a direct Solar analogue. Other promising evidence of stellar CMEs [121] includes blueshifts of chromospheric lines and EUV/X-ray coronal dimming [122], but such measurements do not inform us if the material was retained in the stellar magnetosphere – implying the plasma never impacts a putative planet. Low-frequency, wide-field surveys with  $< 10$  mJy sensitivity on timescales of minutes are the most likely avenue to

detect an extrasolar Type II burst and confirm that mass has been ejected since such bursts are stochastic and the emission frequency is likely lower in weaker magnetic field strengths [18, 123]. Such a detection would allow a measurement of the plasma density as the radio emission occurs at the plasma frequency  $\nu_p \propto n_e^{1/2}$ , where  $n_e$  is the electron density, allowing us to trace the particle flux at the point of impact of an exoplanet [124].

Finally, the Sun exhibits steady mass loss through the solar wind. Measuring stellar winds have been notoriously difficult. Radio observations of thermal bremsstrahlung emission have so far only placed upper limits on stellar winds from young solar analogs [13, 14], and indirect ultraviolet wind measurements of M dwarfs suggest a fairly large spread of inferred mass loss rates with surface X-ray flux [15]. There is promise with the Next Generation Very Large Array (ngVLA; [125]) to be able to detect mass loss from the steady stellar wind of nearby M dwarfs [126].

### 3 Radio Emission Directly from Exoplanets

Exoplanetary systems are observed in the radio primarily to search for auroral emission powered by breakdown of co-rotation, wind-magnetospheric interactions, or emission induced on the host star by an exoplanet via sub-Alfvénic interactions. However, the energetics of these different models imply any radio detection of an exoplanet will likely be near the sensitivity limits of current radio facilities [23, 53, 127, 128]. Therefore, the direct detection of exoplanetary auroral radio emission has been elusive despite decades of searching [129, 130, 128]. Following several foundational works [?, 23, 51, 52, 131], a large collection of theoretical work has been published (e.g. [132, 133, 134, 135, 53, 136, 137, 138, 139, 140, 141, 142, 143, 144]) predicting the intensity and frequency of the radio emission of exoplanets observed from Earth.

As stated in Section 1, radio emission directly from an exoplanet can be powered by the energy deposited on the planetary magnetosphere from the incident stellar wind [51], or via a sub-Alfvénic interaction with a moon [145, 146, 147]. The radiometric Bode’s law [51, 132] and the radio-magnetic scaling law [23, 52, 148] have been used to predict the radio intensity of exoplanets, directly relating the incident stellar wind power (kinetic or magnetic) to the emitted radio power. However, the uncertainty on the predicted flux density can be greater than an order of magnitude, and the uncertainty on the predicted maximum emission frequency can be off by a factor of 2-3 [148]. Therefore, these predictions should be used with caution.

A large number of observational campaigns have been performed to search for radio emission from exoplanets [20, 149], resulting in clear non-detections [25, 150, 151, 152, 153, 154, 155, 156, 127, 157, 141, 158, 159, 160, 161, 162] and a few tentative but contested detections [163, 164, 165, 35, 166]. No claim of detection has yet been confirmed by follow-up observations. The causes for radio non-detections directly from exoplanets are degenerate [129, 130, 128, 167]. Namely, it is possible that we have no clear detections because:

1. Observations have not been sufficiently sensitive;
2. The planetary magnetic field is too weak, implying ECM emission is not produced at accessible observing frequencies;
3. Earth was outside the beaming pattern of the radio emission at the time of the observation [135, 168];
4. The ECM conditions are varying in response to variations in the stellar wind conditions and/or the electron velocity distribution, causing the radio emission to drop below the detection threshold of the observation [169, 170];
5. Inhibition of the ECM process from, for example, inflated ionospheres (e.g. [171, 172, 173, 140, 174]);
6. The wind of the host star absorbs or prevents the escape of the emission [167];
7. The presence of high plasma densities in the planetary magnetosphere or surrounding stellar wind suppresses wave propagation [18, 172], or;
8. Difficulties disentangling whether the detected coherent emission is stellar or exoplanetary in origin since stars also can intrinsically produce coherent emission (e.g. [116, 162]).

It is possible all of these effects are in play for each individual system, though their relative contributions have not yet been quantified – implying it is difficult to infer physically meaningful information about a single system from a non-detection of ECM emission.

Progress in detecting radio emission directly from an exoplanet will largely come from complete phase coverage of the orbit of the exoplanet, preferably several times, and increased sensitivity at low frequencies. Further development of sensitive instruments that observe at  $\lesssim 300$  MHz is vital if we want to probe systems at Jupiter-mass or below since ECM emission scales linearly with the magnetic field strength of the body, and from dynamo theory it is unlikely planets with mass less than 10 times that of Jupiter will possess magnetic field strengths significantly exceeding  $\sim 100$  G [175]. For example, Jovian radio emission cuts off above  $\approx 40$  MHz since its magnetic field does not exceed  $\approx 14$  G [176]).

## 4 Auroral Radio Emission from Ultracool Dwarfs

Due to the difficulty in conclusively detecting radio emission directly from exoplanets, we can leverage the considerable effort that has gone into observing ultracool dwarfs (UCDs; spectral types  $>M7$ ), to examine magnetic processes across the substellar regime, which is expected to have the same underlying physics as direct emission from exoplanets and SPIs. Some UCDs have exhibited bright radio bursts [177] generated by ECM emission [178], allowing us to trace auroral processes in magnetospheres more similar to Jupiter than the Sun [179, 180, 181]. The Jupiter-like analogy for UCDs has recently gained further support with the detection of synchrotron radiation belts around a

UCD [182, 183], a key radio morphological characteristic possessed by Jupiter's magnetosphere.

The first serendipitous discovery of bursting radio emission from a brown dwarf [177] and ensuing UCD radio detections, demonstrated that their radio activity can strongly depart from well-established stellar coronal/flaring relationships [184, 185]. In light of the subsequent discovery that UCD radio bursts can exhibit periodic timing [46], their strong circular polarization and high brightness temperatures implies the ECM process is at least partly responsible for the radio emission from some UCDs [186, 178]. Despite thorough gigahertz-frequency radio searches of UCDs, detections rates have remained persistently low  $\sim 10\%$  detection rates [187, 188, 189, 190, 191, 192, 193, 194, 195].

Periodically bursting radio emission traces extrasolar aurorae in UCDs. Simultaneous radio and optical observations of the M8.5 dwarf LSR J1835+3259 showed that the impacting electron current traced by its bursting radio emission creates a surface feature that is spectrally distinct from typical magnetic spots seen on stars [179]. UCDs with optical-infrared periodic variability or H $\alpha$  emission can have an 80% radio detection rate [180]. Magnetospheric current systems modeled off of Jupiter can explain the generally correlated radio and optical/atmospheric behaviors across the UCD population [181, 196]. In this picture, brown dwarfs are magnetic analogues to gas giants and laboratories for studying planetary dynamos [180, 197, 198]. Furthermore, the discovery of hitherto unknown brown dwarfs via their radio emission [196, 199, 198] demonstrates that large-field radio surveys can access a new discovery space for brown dwarfs or planets that other observational means may overlook. In particular, it is expected the next generation of wide-field radio surveys will likely identify the lowest mass or most distant UCDs, since infrared surveys are reasonably complete within 25 pc for UCDs of spectral types earlier than T4 [200].

The broad success of the UCD auroral paradigm naturally leads to a substantive open question: what is the source of the magnetospheric plasma in such systems? This magnetospheric condition may determine the overall low radio detection rate ( $< 10\%$ ) [194, 193]. A potentially viable solution is exoplanet companions [181, 201]. In the same way that Io seeds the Jovian magnetosphere with plasma, these satellites could provide the plasma for their UCD hosts. Although challenging, searches for planets around UCDs are ongoing [202, 203, 204], with TRAPPIST-1 [205] providing a prototypical example. The radio emission itself is also enabling companion searches, and astrometric radio monitoring via very long baseline interferometry has already yielded evidence for a companion around an auroral UCD [206].

A statistically significant coincidence between UCD planetary systems and bright radio emission remains unproven. Nevertheless, because a satellite population likely exists, the strong magnetic fields of the lowest-mass stars and brown dwarfs [197] make them exceptional candidates for searching for magnetic host-planet interactions [207]. Alternatively, the radio emission could be similar to the breakdown of co-rotation seen in the Jupiter system, with

the plasma supplied by magnetic reconnection events in the UCD's magnetosphere. Long term radio monitoring of the systems, and determination of the UCDs rotation period, is required to determine which model is correct since the periodicity will either correlate with the rotation of the UCD or the orbit of a putative satellite.

## 5 Radio Emission from Star-Planet Interactions

As well as radio emission directly from a planetary magnetosphere, it is thought to be possible to drive radio emission via the magnetic connection between a close-in planet and its star – à la a scaled-up Jupiter-Io interaction [23, 52, 148, 208, 209], with a star taking the place of Jupiter and a planet taking the place of Io.

Radio emission from SPI via ECM emission can potentially reach higher frequencies than planetary magnetospheric radio emission (up to gigahertz frequencies for some M dwarfs, massive, or young stars due to their relatively strong magnetic fields) since the auroral emission is occurring in the magnetosphere of the star. This higher frequency of emission makes radio from SPI easier to detect with ground-based radio observatories than that expected directly from exoplanetary magnetospheres. Note that the radio emission from SPI only gives a direct measure of the magnetic field of the star, not the planet. However, potentially it is possible to model a planet's magnetic field in such an interaction (see, e.g. [130, 210, 211]), similar as is done for Ganymede [212]. Emission from SPI may also manifest as variation in X-rays [213] and ultraviolet/optical/near-infrared activity indicators [214, 215, 27, 216] – but all claims of detection of SPI at these wavelengths are disputed [217, 218, 219, 220].

Similar to radio emission from exoplanets, there has been no confirmed detection of radio emission from SPI. However, some recent publications show encouraging signals that merit follow-up. LOFAR observations have shown low-frequency radio emission from the quiescent M dwarf GJ 1151, and several others, at 144 MHz [31, 34], which has been attributed to the interaction of the star with a close-in Earth-size planet due to the emission and stellar properties. An intensive campaign of precise radial velocity observations with HARPS-N [221], HPF [222] and CARMENES [223] has detected a  $> 10.6 M_{\oplus}$  companion to GJ 1151 in a 390 d orbit at a separation of 0.57 au, but only a  $1.2 M_{\oplus}$  upper limit on the mass and orbit of any companion close enough to be in the sub-Alfvénic region of the star [224].

Several other recent unconfirmed SPI detections also merit followup with intensive observing campaigns, particularly in synergy with spectropolarimetric monitoring aimed at modelling the large-scale field of the host stars. At gigahertz frequencies there have been suggested detections of ECM emission possibly modulated by planetary orbits from Proxima Centauri [33] and YZ Ceti [166, 225]. Modulation of optical activity tracers close to the period of AU Microscopii b has been interpreted as SPI [216], and suggests AU Mic as a

promising target for SPI. However, only evidence of stellar rotation modulation in the radio has been detected from AU Mic [162].

## 5.1 Theory of magnetic star-planet interactions

Considering the likelihood of detection of radio emission from SPI with the current generation of instruments, in this section we outline the theory of SPI and how it can be used to derive physical characteristics of SPI systems.

SPI radio emission is produced by the ECM process near the local electron cyclotron frequency  $\nu_c$ , which peaks at  $\nu_c = 2.8B_*$  MHz, where  $B_*$  is the magnetic field strength of the star in Gauss. Such emission is produced in a rarefied ( $\nu_c \gg \nu_p$ ) plasma by an unstable electron population with characteristic energy of  $\sim 1\text{--}20$  keV [48, 226, 227, 228, 229, 230].

Furthermore, SPI radio emission is predominately emitted via the extraordinary (x) magneto-ionic mode, implying the polarisation of the emission is highly circular or weakly elliptical. The emission is beamed at a large angle from the magnetic field, typically  $60^\circ\text{--}90^\circ$ , and in a thin conical sheet of  $1^\circ\text{--}2^\circ$  thickness [226]. We schematically show such emission geometry in Figure 3. Modelling the ECM emission often involves assumptions about the stellar magnetic field topology, making it possible to produce a ‘visibility curve’ for radio emission for a SPI system [167, 231]. In the limit of a dipolar stellar magnetic field, radio emission from SPI can have a very low duty cycle ( $< 10\%$ ) and a sensitive dependence on the obliquity between the axes of the stellar dipole, rotation, and planetary orbit [49].

The radio power produced by variable ECM emission is hard to predict from first principles: ECM growth rates and the wave path along which it operates depend on the details of the distribution of the unstable  $\sim 1\text{--}20$  keV electrons in the auroral regions, and on the magnetic field topology and ambient plasma distribution in the radio sources. However, observations collected in the Solar System reveal that the emitted auroral radio power  $P_r$  averaged over time, frequency, and solid angle is approximately proportional to both the incident kinetic  $P_{\text{kin}}$  and Poynting  $P_B$  fluxes [23, 149]:

$$P_r = \alpha P_{\text{kin}} = \alpha \rho v^3 \pi R_{\text{obs}}^2 \quad (1)$$

$$P_r = \beta P_B = \beta \frac{B_\perp^2}{\mu} v \pi R_{\text{obs}}^2, \quad (2)$$

where  $\rho$  is the density of the wind incident on the planet’s magnetosphere,  $R_{\text{obs}}$  is the effective size of the obstacle,  $v$  the incident flow velocity in the obstacle’s frame,  $B_\perp$  the magnetic field in the flow perpendicular to  $v$ , and  $\mu$  is the permeability of space. The proportionality factors  $\alpha$  and  $\beta$  are the overall efficiencies of the conversion of the kinetic and Poynting flux into the emitted radio power, which are empirically estimated to be of order of  $10^{-5}$  and  $10^{-3}$  respectively from Solar System observations [52, 65]. When auroral

and satellite-Jupiter radio emissions are considered together, only equation Equation 2 holds. Its high efficiency ( $\beta \sim 10^{-3}$ ) explains why the Jupiter-Io interaction is the brightest object in the sky below 40 MHz, outshining even the quiet Sun.

The detailed magnetohydrodynamics of SPI models lead to a wide range of estimates of the Poynting flux. For sub-Alfvénic interactions we can have 1) force-free models, where the electric current and magnetic field vectors are parallel [232] or; 2) Alfvén wing models [64, 233] with pairs of standing waves in the magnetised wind, which separately may connect the planet either to the star or interplanetary space.

For the Alfvén wing model, the power transmitted in one Alfvén wing was first estimated as  $P_{AW} = \epsilon P_B$ , where  $\epsilon = (1 + M_A^{-2})^{-1/2}$ .  $M_A$  is the Alfvén Mach number, which is defined as the plasma flow speed divided by the Alfvén velocity. For small  $M_A$ ,  $\epsilon \approx M_A$  [23, 52]. For reference,  $M_A = 0.3 \pm 0.1$  for Io's interaction [234]. Based on a nonlinear solution of Alfvén wings [59], the total power can be expressed in closed form for small  $M_A$  as [64]

$$P_{AW} = 2\pi R^2 \bar{\alpha}^2 M_A \frac{B_{\perp}^2}{\mu_0} v_0. \quad (3)$$

Here  $\bar{\alpha}$  is the interaction strength ( $0 \leq \bar{\alpha} \leq 1$ ) and  $R$  is the radius of the central Alfvén tube carrying the energy away along the background field. It is often approximated by the effective radius of the obstacle [64].  $\bar{\alpha}$  is expected to be close to one if the planet possesses a dense atmosphere/ionosphere. This Alfvén wing expression is widely used [63, 169, 235], and is a more exact solution than the early estimate  $\approx M_A P_B$  presented by [23]. The energy flux in Equation 3 is a factor of two larger as than that presented by [23] as it considers the energy fluxes inside and outside the central Alfvén tube defined by  $\pi R_{obs}^2$  [64]. The dependence on  $M_A$  also highlights that low density stellar winds will have a lower amount of power emitted compared to high density stellar winds.

In force-free models of SPI powered by reconnection, the energy fluxes have been estimated as  $P_{FF} = \gamma P_B$  with the efficiency factor  $0 \leq \gamma \leq 1$  [232]. These fluxes are roughly the same order as those in the Alfvén wing models. In force-free models of the Jupiter-Io interaction, such as the unipolar inductor, the energy fluxes are controlled by the conductance of Jupiter's ionosphere and not by the wave impedance of the Alfvén waves [236, 207].

One commonly used form of a force-free model is called the stretch-and-break model [237, 238], where the planetary polar magnetic field  $B_P$  is stretched and energy is released by reconnection. In this case the power produced is

$$P_{SB} = 2\pi R_P^2 \frac{B_P^2}{\mu_0} v_0 f_{AP}, \quad (4)$$

where  $R_P$  is the radius of the planet and  $f_{AP}$  is the area fraction of the planet where magnetic field lines connect to the stellar wind.  $f_{AP}$  might have a typical value of order 0.1 depending on the strength and configuration of the planetary and wind magnetic fields. The direct occurrence of the planet's internal magnetic field  $B_P$  is the primary difference between the Alfvén and stretch and break model. It would mean that  $P_{SB}/P_{AW}$  can be 100-1000 [238]. However, the Poynting flux in Equation 4 is primarily perpendicular and not directed along the flux tube towards the star, and the flow velocities  $v_0$  over the poles will be reduced as well. The fluxes in Equation 4 have neither been seen in numerical simulations [235], nor take place at Jupiter's magnetized moon Ganymede. We suggest further work on the applicability of Equation 4 is necessary if the stretch and break model is to be used for predictions.

A general prediction from all of these models is that the brightest radio emission from SPI will be produced by the largest planet as close as possible to a star with the strongest magnetic field. This heuristic can inform future SPI searches but needs to be used with caution as such a heuristic neglects potentially important second order effects, such as ECM inhibition due to inflated ionospheres [140, 171, 172, 173, 174].

## 5.2 Stellar wind environments around planet-hosting stars

Whether or not radio emission from SPI occurs, as well its morphology and strength, depends on the plasma environment that the planet is embedded. We require detailed knowledge of this plasma if we are to predict and interpret signatures of SPI. For low-mass stars, the primary source of the interplanetary plasma is its stellar wind, which expands outward from the stellar surface, likely via a combination of magnetic and thermal pressure forces [239]. Additional sources may further add to the interplanetary environment, such as atmospheric mass loss from the planet itself [240] and CMEs [241].

Unlike our Sun, we lack detailed knowledge of the stellar wind environments around other low-mass stars [242]. This is primarily because their winds are very tenuous, implying they do not produce strong observational signatures. However, sophisticated magnetohydrodynamic (MHD) models can be deployed to obtain three-dimensional snapshots of the winds of low-mass stars. Measurements of the mass-loss rates of the winds of low-mass stars are currently limited to a handful of cases, and rely on indirect methods (see, [13, 243, 244]). Nevertheless, this has not stopped trends relating mass-loss rates to more readily available information such as surface X-ray fluxes being established [124]. In turn, these trends are used to inform MHD models.

Another key constraint that can be implemented into MHD models of the winds of low-mass stars are reconstructed surface magnetic field maps. Since the inception of the ZDI method [41], such data have become readily available for a relatively large population of low-mass stars [42, 216]. The magnetic field is embedded in the stellar wind and strongly influence the dynamics of the flow [167]. Additionally, the wind magnetic field is essential in predicting

and interpreting signatures of star-planet interaction for at least two reasons. Firstly, it is necessary to estimate the location of the Alfvén surface, and thus predict whether the planet has a sub- or super-Alfvénic orbit [245]. Secondly, the local magnetic field is also required to estimate the Poynting flux reaching the planet, which dictates the strength of the interaction [246]. By implementing such maps as boundary conditions in MHD simulations of stellar winds, we can construct models of the plasma environment around the star. These in turn can be used to estimate quantities relevant to SPI.

## 6 Outlook

We expect the emerging trend of claimed radio emission from stellar CMEs, SPI and planetary aurora from LOFAR, FAST, GMRT, JVLA, and ASKAP will become a flood in the next decade, as the expected construction and commissioning of the much more sensitive Square Kilometre Array (SKA; [247]) and ngVLA are completed. Scaling from the sample of LOFAR detected stars would suggest that SKA1-Low ( $\sim 50 - 350$  MHz) could detect  $\sim 10^3$  M dwarf systems [34, 248], and SKA1-Mid ( $\sim 350$  MHz – 15.4 GHz) and ngVLA (1.2 – 116 GHz) will be sensitive even to quiescent radio emission from the nearest stars [249]. These relatively higher frequencies of SKA1-Mid/ngVLA ( $\sim 1$ –2 GHz) will likely only probe stellar magnetic fields via SPI, while the lower frequencies of LOFAR/SKA-Low could directly detect auroral emission from Jupiter-mass and larger planets. For tracing aurora from lower mass planets, such as an Earth twin, observations will need to be conducted from space since the Earth’s ionosphere reflects emission  $\lesssim 10$  MHz back into space. In the long-term, low-frequency exoplanet science will require radio interferometers on the far side of the Moon [250, 251], the best location in the Solar System for avoiding terrestrial radio frequency interference – provided that lunar satellites and missions do not pollute that pristine location by the time such telescopes can be established.

The new groundbased radio facilities will need to be matched with both improved theoretical models of stellar plasma environments, and multiwavelength observations that independently constrain stellar activity and exoplanet companions. One of the most important questions that MHD modelling could begin to answer is the extent to which sub-Alfvénic SPI affects planetary atmospheric mass loss and habitability.

Furthermore, more statistical robustness is needed in the field before claiming the detection of radio periodicity from SPI considering the emission mechanism is inherently variable and often dependent on unknown stellar magnetic field properties that can vary in time [252]. The expected periodicity can also bear a complicated relationship to the orbital and rotational periods [49]. We suggest the following rule of thumb should be adopted in the field: at least three detections of a stellar system at the same phase, preferably distinct from the rotational phase of the star, should be required before a reliable claim of detection of SPI can be made. Such a standard would prevent premature

declaration of the detection of radio emission from SPI, with off-phase observations also vital in ensuring the alignment of detections at a set phase is due to a biased sampling window. Currently, no claim of SPI in the literature has met this standard.

Detecting these variable and periodic signals may benefit from treatment of methods with greater statistical sophistication. For example, using Gaussian Processes [253] or generalized periodograms [254] could potentially aid in a reliable determination of periodicity from heavily undersampled datasets. A firm attribution of observed radio emission to SPI also depends critically on knowledge of the stellar magnetic field, implying it will be essential to complement SKA facilities with dedicated facilities for contemporaneous spectropolarimetric monitoring of interesting targets.

As radio emission from SPI may even be biased *against* transiting or edge-on systems [49], it will be necessary to continue precise radial velocity (PRV) surveys to determine the existence and orbits of exoplanets inferred from the radio. Red-sensitive and NIR instruments (such as SPIRou [255], the Habitable Planet Finder [40], and CARMENES [39]) will be especially important for probing the population around M dwarfs where the Alfvén surface encompasses a large fraction of planetary systems and reaches into habitable zone orbits. As well as stabilized spectrographs for ZDI and PRVs, short-cadence time series photometry is essential for detecting stellar flares. Photometry by itself can determine a stellar flare rate and can help distinguish between SPI and coronal emission [112], but simultaneous photometry is ideal since have been observed correlation between some radio bursts and optical flares [32].

Considering the significant commitment of observatory resources required to determine radio detections of SPI, it is important the field invests wisely on the most likely candidates. To first order, it is possible to identify the current most likely systems to produce SPI radio emission based on the size of the exoplanet, its proximity to its host star, and the distance of the stellar system from Earth [128, 256]. In this case, the top five candidate systems are 51 Pegasi, HIP 65 A, Tau Boötis A, 55 Cancri A, and WASP-18 – many of which have been searched for radio signatures already.

However, many of these stars likely do not have strong enough magnetic fields to produce ECM emission at  $\gtrsim 150$  MHz, where our most sensitive telescopes currently operate. Such stars also likely have complicated magnetic field topologies, implying that radio emission may not be consistently beamed to Earth. Therefore, planets close to M dwarfs may be the best systems to followup considering M dwarfs can possess kilogauss-strong dipolar magnetic fields. GJ 367, GJ 436, GJ 1252, GJ 3253, GJ 625, YZ Ceti, AU Microscopii, and Proxima Centauri could be ideal candidates in such a case – of which many have been searched [33, 162]. The main issue with focusing on M dwarfs for SPI radio signatures is that the star itself is also known to produce radio emission. As discussed above, this implies that a robust detection of SPI will likely involve  $\gtrsim 200$  hr on radio telescopes since the planets will have a  $\sim 1$  to 10 day orbital period.

Regardless of the difficulties faced in determining radio emission from SPI, exoplanets, and stars, the potential scientific return is invaluable. Radio observations of stellar systems can provide information about the planet, star, and space weather that are not directly attainable at any other wavelength. Determining the potential habitability of exoplanets will be a focus of astronomy in the coming decades – and radio astronomy is poised to provide key pieces to this puzzle.

**Acknowledgments.** This project was initiated at the Lorentz Center workshop *Life Around a Radio Star*, held 27 June - 1 July 2022 in Leiden, the Netherlands.

JRC thanks the following graduate students and postdoctoral scholars for providing comments on the manuscript from the perspective of scientists new to the field: Sanne Bloot (ASTRON), Cristina Cordun (ASTRON), Evan Fitzmaurice (Penn. State), David Konijn (ASTRON), Kristo Ment (Penn. State), and Timothy Yiu (ASTRON).

This research has made use of the NASA Exoplanet Archive, which is operated by the California Institute of Technology, under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program.

BJSP acknowledges and pays respect to the traditional owners of the land on which the University of Queensland is situated, and to their Ancestors and descendants, who continue cultural and spiritual connections to Country. He acknowledges funding from the ARC DECRA DE21 scheme and the Big Questions Institute.

RDK acknowledges funding from the Dutch Research Council (NWO) for the e-MAPS (exploring magnetism on the planetary scale) project (project number VI.Vidi.203.093) under the NWO talent scheme Vidi.

SB acknowledges funding from the Dutch Research Council (NWO) for the ‘Exo-space weather and contemporaneous signatures of star-planet interactions’ of the research programme ‘Open Competition Domain Science- M’ (number OCENW.M.22.215).

MD acknowledges support from the INAF funding scheme Fundamental Research in Astrophysics 2022 (mini grant “A pilot study to explore the potential of SRT in detecting nearby radio-emitting stars with confirmed or candidate exoplanets, supported by a radial velocity follow-up”).

PZ acknowledges funding from the ERC N°101020459–Exoradio.

SM acknowledges funding from NSF AST-2108512 for a precision NIR M dwarf radial velocity survey with HPF, from NASA XRP investigating radio detected M dwarfs.

JM acknowledges funding from the French National Research Agency (ANR) under contract number ANR-18-CE31-0019 (SPLaSH).

AAV acknowledges funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 817540, ASTROFLOW).

GS acknowledges support provided by NASA through the NASA Hubble Fellowship grant HST-HF2-51519.001-A awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS5-26555.

BK acknowledge funding from the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement No 865624, GPRV).

JDT was supported for this work by NASA through the NASA Hubble Fellowship grant #HST-HF2-51495.001-A awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555.

JS received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 884711).

JMG acknowledges support by the "Programme National de Planétologie" (PNP) of CNRS/INSU co-funded by CNES and by the "Programme National de Physique Stellaire" (PNPS) of CNRS/INSU co-funded by CEA and CNES.

MMK acknowledges support from the Heising-Simons Foundation through the 51 Pegasi b Fellowship grant 2021-2943.

## Declarations

- This project was partly funded by the Lorentz Centre at Leiden University.
- Conflict of interest: The authors do not have any conflicts of interest to report.
- Authors' contributions: Callingham organized overall structure, acted as chief editor, led the replies to the referees, and edited all contributions into a cohesive text with Pope and Kavanagh. Kavanagh produced Figure 3. Callingham, Nichols, Rigney, Saur, Turner, and Zarka were the principal contributors to the section on Jupiter and the Sun; Daley-Yates, Güdel, Günther, Osten, Pope, and Villadsen, were the principal contributors to the stellar flares and CMEs section; Callingham, Kavanagh, Perez-Torres, Saur, Vedantham, Vidotto, and Zarka were the principal contributors to the SPI section; Griebmeier and Turner about radio emission from planet aurorae. All authors reviewed the final text.

## References

- [1] Gaudi, B. S., Meyer, M. & Christiansen, J. in *The Demographics of Exoplanets* (ed. Madhusudhan, N.) *ExoFrontiers; Big Questions in Exoplanetary Science* 2–1 (2021).
- [2] Madhusudhan, N. Exoplanetary Atmospheres: Key Insights, Challenges, and Prospects. *Annu. Rev. Astron. Astrophys.* **57**, 617–663 (2019).
- [3] JWST Transiting Exoplanet Community Early Release Science Team *et al.* Identification of carbon dioxide in an exoplanet atmosphere. *Nature* **614**, 649–652 (2023).

- [4] Kempton, E. M. R. *et al.* A reflective, metal-rich atmosphere for GJ 1214b from its JWST phase curve. *Nature* **620**, 67–71 (2023).
- [5] Greene, T. P. *et al.* Thermal emission from the Earth-sized exoplanet TRAPPIST-1 b using JWST. *Nature* **618**, 39–42 (2023).
- [6] Zieba, S. *et al.* No thick carbon dioxide atmosphere on the rocky exoplanet TRAPPIST-1 c. *Nature* **620**, 746–749 (2023).
- [7] Kouloumvakos, A. *et al.* Properties of solar energetic particle events inferred from their associated radio emission. *Astron. Astrophys.* **580**, A80 (2015).
- [8] Badruddin, A. & Falak, Z. Study of the geoeffectiveness of coronal mass ejections, corotating interaction regions and their associated structures observed during Solar Cycle 23. *Astrophys. Space Sci.* **361**, 253 (2016).
- [9] Lammer, H. *Origin and Evolution of Planetary Atmospheres* (2013).
- [10] Scalo, J. *et al.* M Stars as Targets for Terrestrial Exoplanet Searches And Biosignature Detection. *Astrobiology* **7**, 85–166 (2007).
- [11] Kulikov, Y. N. *et al.* A Comparative Study of the Influence of the Active Young Sun on the Early Atmospheres of Earth, Venus, and Mars. *Space Sci. Rev.* **129**, 207–243 (2007).
- [12] Curry, S. M. *et al.* Response of Mars O<sup>+</sup> pickup ions to the 8 March 2015 ICME: Inferences from MAVEN data-based models. *Geophys. Res. Lett.* **42**, 9095–9102 (2015).
- [13] Fichtinger, B. *et al.* Radio emission and mass loss rate limits of four young solar-type stars. *Astron. Astrophys.* **599**, A127 (2017).
- [14] Ó Fionnagáin, D. *et al.* The solar wind in time - II. 3D stellar wind structure and radio emission. *Mon. Not. R. Astron. Soc.* **483**, 873–886 (2019).
- [15] Wood, B. E. *et al.* New Observational Constraints on the Winds of M dwarf Stars. *Astrophys. J.* **915**, 37 (2021).
- [16] Owen, J. E. & Adams, F. C. Magnetically controlled mass-loss from extrasolar planets in close orbits. *Mon. Not. R. Astron. Soc.* **444**, 3761–3779 (2014).
- [17] Vidotto, A. A. & Cleary, A. Stellar wind effects on the atmospheres of close-in giants: a possible reduction in escape instead of increased erosion. *Mon. Not. R. Astron. Soc.* **494**, 2417–2428 (2020).
- [18] Dulk, G. A. Radio emission from the sun and stars. *Annu. Rev. Astron. Astrophys.* **23**, 169–224 (1985).
- [19] Villadsen, J. & Hallinan, G. Ultra-wideband Detection of 22 Coherent Radio Bursts on M Dwarfs. *Astrophys. J.* **871**, 214 (2019).
- [20] Lazio, T. J. W. Radio Observations as an Extrasolar Planet Discovery and Characterization: Interior Structure and Habitability. *arXiv e-prints* arXiv:2404.12348 (2024).
- [21] McLean, D. J. & Labrum, N. R. *Solar radiophysics : studies of emission from the sun at metre wavelengths* (1985).
- [22] Dungey, J. W. The Steady State of the Chapman-Ferraro Problem in Two Dimensions. *J. Geophys. Res.* **66**, 1043–1047 (1961).

- [23] Zarka, P. Plasma interactions of exoplanets with their parent star and associated radio emissions. *Planet. Space Sci.* **55**, 598–617 (2007).
- [24] Luger, R. *et al.* The Pale Green Dot: A Method to Characterize Proxima Centauri b Using Exo-Aurorae. *Astrophys. J.* **837**, 63 (2017).
- [25] Yantis, W. F., Sullivan, I., W. T. & Erickson, W. C. A Search for Extra-Solar Jovian Planets by Radio Techniques. *Bulletin of the American Astronomical Society* **9**, 453 (1977).
- [26] Lazio, W., T. Joseph *et al.* The Radiometric Bode’s Law and Extrasolar Planets. *Astrophys. J.* **612**, 511–518 (2004).
- [27] Cauley, P. W., Shkolnik, E. L., Llama, J. & Lanza, A. F. Magnetic field strengths of hot Jupiters from signals of star-planet interactions. *Nature Astronomy* 408 (2019).
- [28] van Haarlem, M. P. *et al.* LOFAR: The LOw-Frequency ARray. *Astron. Astrophys.* **556**, A2 (2013).
- [29] Johnston, S. *et al.* Science with ASKAP. The Australian square-kilometre-array pathfinder. *Experimental Astronomy* **22**, 151–273 (2008).
- [30] Driessen, L. N. *et al.* The Sydney Radio Star Catalogue: properties of radio stars at megahertz to gigahertz frequencies. *arXiv e-prints arXiv:2404.07418* (2024).
- [31] Vedantham, H. K. *et al.* Coherent radio emission from a quiescent red dwarf indicative of star-planet interaction. *Nature Astronomy* (2020).
- [32] Zic, A. *et al.* A Flare-type IV Burst Event from Proxima Centauri and Implications for Space Weather. *Astrophys. J.* **905**, 23 (2020).
- [33] Pérez-Torres, M. *et al.* Monitoring the radio emission of Proxima Centauri. *Astron. Astrophys.* **645**, A77 (2021).
- [34] Callingham, J. R. *et al.* The population of M dwarfs observed at low radio frequencies. *Nature Astronomy* **5**, 1233–1239 (2021).
- [35] Turner, J. D. *et al.* The search for radio emission from the exoplanetary systems 55 Cancri,  $\nu$  Andromedae, and  $\tau$  Boötis using LOFAR beam-formed observations. *Astron. Astrophys.* **645**, A59 (2021).
- [36] Pritchard, J. *et al.* A circular polarization survey for radio stars with the Australian SKA Pathfinder. *Mon. Not. R. Astron. Soc.* **502**, 5438–5454 (2021).
- [37] Zhang, J. *et al.* Fine Structures of Radio Bursts from Flare Star AD Leo with FAST Observations. *Astrophys. J.* **953**, 65 (2023).
- [38] Ricker, G. R. *et al.* Transiting Exoplanet Survey Satellite (TESS). *Journal of Astronomical Telescopes, Instruments, and Systems* **1**, 014003 (2015).
- [39] Quirrenbach, A. *et al.* CARMENES: Calar Alto high-resolution search for M dwarfs with exo-earths with a near-infrared Echelle spectrograph. *Ground-based and Airborne Instrumentation for Astronomy III* **7735**, 773513 (2010).
- [40] Mahadevan, S. *et al.* The habitable-zone planet finder: a stabilized fiber-fed NIR spectrograph for the Hobby-Eberly Telescope. *Ground-based*

- and Airborne Instrumentation for Astronomy IV **8446**, 84461S (2012).
- [41] Semel, M. Zeeman-Doppler imaging of active stars. I - Basic principles. *Astron. Astrophys.* **225**, 456–466 (1989).
- [42] Morin, J. *et al.* Large-scale magnetic topologies of late M dwarfs\*. *Mon. Not. R. Astron. Soc.* **407**, 2269–2286 (2010).
- [43] Bastian, T. S., Benz, A. O. & Gary, D. E. Radio Emission from Solar Flares. *Annu. Rev. Astron. Astrophys.* **36**, 131–188 (1998).
- [44] Johnstone, C. P. & Güdel, M. The coronal temperatures of low-mass main-sequence stars. *Astron. Astrophys.* **578**, A129 (2015).
- [45] Vedantham, H. K. Prospects for radio detection of stellar plasma beams. *Astron. Astrophys.* **639**, L7 (2020).
- [46] Hallinan, G. *et al.* Rotational Modulation of the Radio Emission from the M9 Dwarf TVLM 513-46546: Broadband Coherent Emission at the Substellar Boundary? *Astrophys. J.* **653**, 690–699 (2006).
- [47] Wu, C. S. & Lee, L. C. A theory of the terrestrial kilometric radiation. *Astrophys. J.* **230**, 621–626 (1979).
- [48] Treumann, R. A. The electron-cyclotron maser for astrophysical application. *Astron. Astrophys. Rev.* **13**, 229–315 (2006).
- [49] Kavanagh, R. D. & Vedantham, H. K. Hunting for exoplanets via magnetic star-planet interactions: Geometrical considerations for radio emission. *Mon. Not. R. Astron. Soc.* (2023).
- [50] Desch, M. D. & Kaiser, M. L. Predictions for Uranus from a radiometric Bode’s law. *Nature* **310**, 755–757 (1984).
- [51] Farrell, W. M., Desch, M. D. & Zarka, P. On the possibility of coherent cyclotron emission from extrasolar planets. *J. Geophys. Res.* **104**, 14025–14032 (1999).
- [52] Zarka, P., Treumann, R. A., Ryabov, B. P. & Ryabov, V. B. Magnetically-Driven Planetary Radio Emissions and Application to Extrasolar Planets. *Astrophys. Space. Sci.* **277**, 293–300 (2001).
- [53] Nichols, J. D. Magnetosphere-ionosphere coupling at Jupiter-like exoplanets with internal plasma sources: implications for detectability of auroral radio emissions. *Mon. Not. R. Astron. Soc.* **414**, 2125–2138 (2011).
- [54] Saur, J., Neubauer, F. M., Connerney, J. E. P., Zarka, P. & Kivelson, M. G. in *Plasma interaction of Io with its plasma torus* (eds Bagenal, F., Dowling, T. E. & McKinnon, W. B.) *Jupiter. The Planet, Satellites and Magnetosphere*, Vol. 1 537–560 (2004).
- [55] Terasawa, T., Maezawa, K. & Machida, S. Solar wind effect on Jupiter’s non-Io-related radio emission. *Nature* **273**, 131–132 (1978).
- [56] Zarka, P. & Genova, F. Low-frequency Jovian emission and solar wind magnetic sector structure. *Nature* **306**, 767–768 (1983).
- [57] Genova, F., Zarka, P. & Barrow, C. H. Voyager and Nancay observations of the Jovian radio-emission at different frequencies - Solar wind effect and source extent. *Astron. Astrophys.* **182**, 159–162 (1987).
- [58] Bigg, E. K. Influence of the Satellite Io on Jupiter’s Decametric Emission.

- Nature* **203**, 1008–1010 (1964).
- [59] Neubauer, F. M. Nonlinear standing Alfvén wave current system at Io: Theory **85**, 1171–1178 (1980).
- [60] Marques, M. S. *et al.* Statistical analysis of 26 yr of observations of decametric radio emissions from Jupiter. *Astron. Astrophys.* **604**, A17 (2017).
- [61] Ip, W.-H., Kopp, A. & Hu, J.-H. On the Star-Magnetosphere Interaction of Close-in Exoplanets. *Astrophys. J. Lett.* **602**, L53–L56 (2004).
- [62] Lanza, A. F. Star-planet magnetic interaction and activity in late-type stars with close-in planets. *Astron. Astrophys.* **544**, A23 (2012).
- [63] Turnpenney, S., Nichols, J. D., Wynn, G. A. & Burleigh, M. R. Exoplanet-induced Radio Emission from M Dwarfs. *Astrophys. J.* **854**, 72 (2018).
- [64] Saur, J., Grambusch, T., Duling, S., Neubauer, F. M. & Simon, S. Magnetic energy fluxes in sub-Alfvénic planet star and moon planet interactions. *Astron. Astrophys.* **552**, A119 (2013).
- [65] Zarka, P. *et al.* Jupiter radio emission induced by Ganymede and consequences for the radio detection of exoplanets. *A&A* **618**, A84 (2018).
- [66] Nichols, J. D. *et al.* Origin of Electron Cyclotron Maser Induced Radio Emissions at Ultracool Dwarfs: Magnetosphere-Ionosphere Coupling Currents. *Astrophys. J.* **760**, 59 (2012).
- [67] Lecavelier des Etangs, A. *et al.* Temporal variations in the evaporating atmosphere of the exoplanet HD 189733b. *Astron. Astrophys.* **543**, L4 (2012).
- [68] Cuntz, M., Saar, S. H. & Musielak, Z. E. On Stellar Activity Enhancement Due to Interactions with Extrasolar Giant Planets. *Astrophys. J. Lett.* **533**, L151–L154 (2000).
- [69] Benz, A. O. & Güdel, M. Physical Processes in Magnetically Driven Flares on the Sun, Stars, and Young Stellar Objects. *Annu. Rev. Astron. Astrophys.* **48**, 241–287 (2010).
- [70] Howard, W. S. *et al.* The Mouse That Squeaked: A Small Flare from Proxima Cen Observed in the Millimeter, Optical, and Soft X-Ray with Chandra and ALMA. *Astrophys. J.* **938**, 103 (2022).
- [71] Güdel, M. & Benz, A. O. X-Ray/Microwave Relation of Different Types of Active Stars. *Astrophys. J. Lett.* **405**, L63 (1993).
- [72] Benz, A. O. & Güdel, M. X-ray/microwave ratio of flares and coronae. *Astron. Astrophys.* **285**, 621–630 (1994).
- [73] Antonucci, E., Gabriel, A. H. & Dennis, B. R. The energetics of chromospheric evaporation in solar flares. *Astrophys. J.* **287**, 917–925 (1984).
- [74] Airapetian, V. S. & Holman, G. D. Atmospheric Heating and Quiescent Radio Emission in Active Stars. *Astrophys. J.* **501**, 805–812 (1998).
- [75] Airapetian, V. S. *et al.* Impact of space weather on climate and habitability of terrestrial-type exoplanets. *International Journal of Astrobiology*

- 19**, 136–194 (2020).
- [76] Ó Fionnagáin, D. *et al.* Coronal Mass Ejections and Type II Radio Emission Variability during a Magnetic Cycle on the Solar-type Star Eridani. *Astrophys. J.* **924**, 115 (2022).
- [77] Crosby, N. B., Aschwanden, M. J. & Dennis, B. R. Frequency distributions and correlations of solar X-ray flare parameters. *Sol. Phys.* **143**, 275–299 (1993).
- [78] Audard, M., Güdel, M. & Guinan, E. F. Implications from Extreme-Ultraviolet Observations for Coronal Heating of Active Stars. *Astrophys. J. Lett.* **513**, L53–L56 (1999).
- [79] Audard, M., Güdel, M., Drake, J. J. & Kashyap, V. L. Extreme-Ultraviolet Flare Activity in Late-Type Stars. *Astrophys. J.* **541**, 396–409 (2000).
- [80] Kashyap, V. L., Drake, J. J., Güdel, M. & Audard, M. Flare Heating in Stellar Coronae. *Astrophys. J.* **580**, 1118–1132 (2002).
- [81] Güdel, M., Audard, M., Kashyap, V. L., Drake, J. J. & Guinan, E. F. Are Coronae of Magnetically Active Stars Heated by Flares? II. Extreme Ultraviolet and X-Ray Flare Statistics and the Differential Emission Measure Distribution. *Astrophys. J.* **582**, 423–442 (2003).
- [82] Arzner, K. & Güdel, M. Are Coronae of Magnetically Active Stars Heated by Flares? III. Analytical Distribution of Superposed Flares. *Astrophys. J.* **602**, 363–376 (2004).
- [83] Stelzer, B. *et al.* A statistical analysis of X-ray variability in pre-main sequence objects of the Taurus molecular cloud. *Astron. Astrophys.* **468**, 463–475 (2007).
- [84] Maehara, H. *et al.* Superflares on solar-type stars. *Nature* **485**, 478–481 (2012).
- [85] Aschwanden, M. J. Thresholded Power law Size Distributions of Instabilities in Astrophysics. *Astrophys. J.* **814**, 19 (2015).
- [86] Yang, H. & Liu, J. The Flare Catalog and the Flare Activity in the Kepler Mission. *Astrophys. J. Suppl. Ser.* **241**, 29 (2019).
- [87] Borucki, W. J. *et al.* Kepler Planet-Detection Mission: Introduction and First Results. *Science* **327**, 977 (2010).
- [88] Ricker, G. R. *et al.* Transiting Exoplanet Survey Satellite (TESS). *Journal of Astronomical Telescopes, Instruments, and Systems* **1**, 014003 (2015).
- [89] Walkowicz, L. M. *et al.* White-light Flares on Cool Stars in the Kepler Quarter 1 Data. *Astron. J.* **141**, 50 (2011).
- [90] Hawley, S. L. *et al.* Kepler Flares. I. Active and Inactive M Dwarfs. *Astrophys. J.* **797**, 121 (2014).
- [91] Davenport, J. R. A. The Kepler Catalog of Stellar Flares. *Astrophys. J.* **829**, 23 (2016).
- [92] Yang, H. & Liu, J. The Flare Catalog and the Flare Activity in the Kepler Mission. *Astrophys. J. Suppl. Ser.* **241**, 29 (2019).
- [93] Günther, M. N. *et al.* Stellar Flares from the First TESS Data Release:

- Exploring a New Sample of M Dwarfs. *Astron. J.* **159**, 60 (2020).
- [94] Feinstein, A. D. *et al.* Flare Statistics for Young Stars from a Convolutional Neural Network Analysis of TESS Data. *Astron. J.* **160**, 219 (2020).
- [95] Gao, D.-Y., Liu, H.-G., Yang, M. & Zhou, J.-L. Correcting Stellar Flare Frequency Distributions Detected by TESS and Kepler. *Astron. J.* **164**, 213 (2022).
- [96] Pietras, M., Falewicz, R., Siarkowski, M., Bicz, K. & Preś, P. Statistical Analysis of Stellar Flares from the First Three Years of TESS Observations. *Astrophys. J.* **935**, 143 (2022).
- [97] Davenport, J. R. A. *et al.* The Evolution of Flare Activity with Stellar Age. *Astrophys. J.* **871**, 241 (2019).
- [98] Feinstein, A. D., Seligman, D. Z., Günther, M. N. & Adams, F. C. Testing Self-organized Criticality across the Main Sequence Using Stellar Flares from TESS. *Astrophys. J. Lett.* **925**, L9 (2022).
- [99] Howard, W. S. The flaring TESS Objects of Interest: flare rates for all two-minute cadence TESS planet candidates. *Mon. Not. R. Astron. Soc.* **512**, L60–L65 (2022).
- [100] Gilbert, E. A. *et al.* Flares, Rotation, and Planets of the AU Mic System from TESS Observations. *Astron. J.* **163**, 147 (2022).
- [101] Feinstein, A. D. *et al.* AU Microscopii in the Far-UV: Observations in Quiescence, during Flares, and Implications for AU Mic b and c. *Astron. J.* **164**, 110 (2022).
- [102] Shibayama, T. *et al.* Superflares on Solar-type Stars Observed with Kepler. I. Statistical Properties of Superflares. *Astrophys. J. Suppl. Ser.* **209**, 5 (2013).
- [103] Shibata, K. *et al.* Can Superflares Occur on Our Sun? *Publ. Astron. Soc. Jpn* **65**, 49 (2013).
- [104] Maehara, H. *et al.* Statistical properties of superflares on solar-type stars based on 1-min cadence data. *Earth, Planets and Space* **67**, 59 (2015).
- [105] Notsu, Y. *et al.* Superflares on Solar-type Stars Observed with Kepler II. Photometric Variability of Superflare-generating Stars: A Signature of Stellar Rotation and Starspots. *Astrophys. J.* **771**, 127 (2013).
- [106] Notsu, Y. *et al.* Do Kepler Superflare Stars Really Include Slowly Rotating Sun-like Stars?—Results Using APO 3.5 m Telescope Spectroscopic Observations and Gaia-DR2 Data. *Astrophys. J.* **876**, 58 (2019).
- [107] Cliver, E. W., Schrijver, C. J., Shibata, K. & Usoskin, I. G. Extreme solar events. *Living Reviews in Solar Physics* **19**, 2 (2022).
- [108] Yashiro, S., Akiyama, S., Gopalswamy, N. & Howard, R. A. Different Power-Law Indices in the Frequency Distributions of Flares with and without Coronal Mass Ejections. *Astrophys. J. Lett.* **650**, L143–L146 (2006).
- [109] Donati, J. F. & Landstreet, J. D. Magnetic Fields of Nondegenerate Stars. *Annu. Rev. Astron. Astrophys.* **47**, 333–370 (2009).

- [110] Alvarado-Gómez, J. D. *et al.* Coronal Response to Magnetically Suppressed CME Events in M-dwarf Stars. *Astrophys. J. Lett.* **884**, L13 (2019).
- [111] Rigney, J. *et al.* Searching for stellar flares from low-mass stars using ASKAP and TESS. *Mon. Not. R. Astron. Soc.* **516**, 540–549 (2022).
- [112] Pope, B. J. S. *et al.* The TESS View of LOFAR Radio-emitting Stars. *Astrophys. J. Lett.* **919**, L10 (2021).
- [113] Crosley, M. K. *et al.* The Search for Signatures of Transient Mass Loss in Active Stars. *Astrophys. J.* **830**, 24 (2016).
- [114] Crosley, M. K. & Osten, R. A. Low-frequency Radio Transients on the Active M-dwarf EQ Peg and the Search for Coronal Mass Ejections. *Astrophys. J.* **862**, 113 (2018).
- [115] Villadsen, J. & Hallinan, G. Ultra-wideband Detection of 22 Coherent Radio Bursts on M Dwarfs. *Astrophys. J.* **871**, 214 (2019).
- [116] Callingham, J. R. *et al.* Low-frequency monitoring of flare star binary CR Draconis: long-term electron-cyclotron maser emission. *Astron. Astrophys.* **648**, A13 (2021).
- [117] Alvarado-Gómez, J. D., Drake, J. J., Cohen, O., Moschou, S. P. & Garraffo, C. Suppression of Coronal Mass Ejections in Active Stars by an Overlying Large-scale Magnetic Field: A Numerical Study. *Astrophys. J.* **862**, 93 (2018).
- [118] Alvarado-Gómez, J. D. *et al.* Tuning the Exospace Weather Radio for Stellar Coronal Mass Ejections. *Astrophys. J.* **895**, 47 (2020).
- [119] Zic, A. *et al.* ASKAP detection of periodic and elliptically polarized radio pulses from UV Ceti. *Mon. Not. R. Astron. Soc.* **488**, 559–571 (2019).
- [120] Bastian, T. S., Cotton, W. D. & Hallinan, G. Radio Emission from UV Cet: Auroral Emission from a Stellar Magnetosphere. *Astrophys. J.* **935**, 99 (2022).
- [121] Osten, R. A., Vidotto, A. A., Fossati, L. & Vink, J. (eds) *Observations of Winds and CMEs of Low-Mass Stars*. (eds Vidotto, A. A., Fossati, L. & Vink, J.) *Winds of Stars and Exoplanets*, Vol. Proceedings of IAU Symposium 370, submitted (2023).
- [122] Veronig, A. M. *et al.* Indications of stellar coronal mass ejections through coronal dimmings. *Nature Astronomy* **5**, 697–706 (2021).
- [123] Güdel, M. Stellar Radio Astronomy: Probing Stellar Atmospheres from Protostars to Giants. *Annu. Rev. Astron. Astrophys.* **40**, 217–261 (2002).
- [124] Vidotto, A. A. *et al.* Characterization of the HD 219134 multi-planet system II. Stellar-wind sputtered exospheres in rocky planets b & c. *Mon. Not. R. Astron. Soc.* **481**, 5296–5306 (2018).
- [125] Di Francesco, J. *et al.* The Next Generation Very Large Array. *Canadian Long Range Plan for Astronomy and Astrophysics White Papers* **2020**, 32 (2019).
- [126] Osten, R. A. & Crosley, M. K. Quantifying the ngVLA’s Contribution to Exo-Space Weather: Results of a Community Studies Report Next Generation VLA Memo #31. *arXiv e-prints* arXiv:1711.05113 (2017).

- [127] Lynch, C. R., Murphy, T., Kaplan, D. L., Ireland, M. & Bell, M. E. A search for circularly polarized emission from young exoplanets. *Mon. Not. R. Astron. Soc.* **467**, 3447–3453 (2017).
- [128] Grießmeier, J.-M. Fischer, G., Mann, G., Panchenko, M. & Zarka, P. (eds) *The search for radio emission from giant exoplanets*. (eds Fischer, G., Mann, G., Panchenko, M. & Zarka, P.) *Planetary Radio Emissions VIII*, 285–300 (Austrian Academy of Sciences Press, Vienna, 2017).
- [129] Zarka, P., Lazio, J. & Hallinan, G. Magnetospheric Radio Emissions from Exoplanets with the SKA. *Advancing Astrophysics with the Square Kilometre Array (AASKA14)* 120 (2015).
- [130] Grießmeier, J.-M. Lammer, H. & Khodachenko, M. (eds) *Detection Methods and Relevance of Exoplanetary Magnetic Fields*. (eds Lammer, H. & Khodachenko, M.) *Astrophysics and Space Science Library*, Vol. 411 of *Astrophysics and Space Science Library*, 213 (2015).
- [131] Zarka, P. *et al.* Rucker, H. O., Bauer, S. J. & Lecacheux, A. (eds) *Ground-Based High Sensitivity Radio Astronomy at Decameter Wavelengths*. (eds Rucker, H. O., Bauer, S. J. & Lecacheux, A.) *Planetary Radio Emission IV*, 101 (1997).
- [132] Lazio, T. J., W. *et al.* The Radiometric Bode’s Law and Extrasolar Planets. *Astrophys. J.* **612**, 511–518 (2004).
- [133] Stevens, I. R. Magnetospheric radio emission from extrasolar giant planets: the role of the host stars. *Mon. Not. R. Astron. Soc.* **356**, 1053–1063 (2005).
- [134] Jardine, M. & Collier Cameron, A. Radio emission from exoplanets: the role of the stellar coronal density and magnetic field strength. *Astron. Astrophys.* **490**, 843–851 (2008).
- [135] Hess, S. L. G. & Zarka, P. Modeling the radio signature of the orbital parameters, rotation, and magnetic field of exoplanets. *Astron. Astrophys.* **531**, A29 (2011).
- [136] Vidotto, A. A. *et al.* The stellar wind cycles and planetary radio emission of the  $\tau$  Boo system. *Mon. Not. R. Astron. Soc.* **423**, 3285–3298 (2012).
- [137] See, V., Jardine, M., Fares, R., Donati, J.-F. & Moutou, C. Time-scales of close-in exoplanet radio emission variability. *Mon. Not. R. Astron. Soc.* **450**, 4323–4332 (2015).
- [138] Vidotto, A. A., Fares, R., Jardine, M., Moutou, C. & Donati, J.-F. On the environment surrounding close-in exoplanets. *Mon. Not. R. Astron. Soc.* **449**, 4117–4130 (2015).
- [139] Nichols, J. D. & Milan, S. E. Stellar wind-magnetosphere interaction at exoplanets: computations of auroral radio powers. *Mon. Not. R. Astron. Soc.* **461**, 2353–2366 (2016).
- [140] Weber, C. *et al.* Supermassive hot Jupiters provide more favourable conditions for the generation of radio emission via the cyclotron maser instability - a case study based on Tau Bootis b. *Mon. Not. R. Astron. Soc.* **480**, 3680–3688 (2018).
- [141] Lynch, C. R., Murphy, T., Lenc, E. & Kaplan, D. L. The detectability

- of radio emission from exoplanets. *Mon. Not. R. Astron. Soc.* **478**, 1763–1775 (2018).
- [142] Wang, X. & Loeb, A. Nonthermal Emission from the Interaction of Magnetized Exoplanets with the Wind of Their Host Star. *Astrophys. J. Lett.* **874**, L23 (2019).
- [143] Kavanagh, R. D. *et al.* MOVES - II. Tuning in to the radio environment of HD189733b. *Mon. Not. R. Astron. Soc.* **485**, 4529–4538 (2019).
- [144] Turnpenney, S., Nichols, J. D., Wynn, G. A. & Jia, X. Magnetohydrodynamic modelling of star-planet interaction and associated auroral radio emission. *Mon. Not. R. Astron. Soc.* **494**, 5044–5055 (2020).
- [145] Noyola, J. P., Satyal, S. & Musielak, Z. E. On the Radio Detection of Multiple-exomoon Systems due to Plasma Torus Sharing. *Astrophys. J.* **821**, 97 (2016).
- [146] Narang, M. *et al.* Radio-loud Exoplanet-exomoon Survey: GMRT Search for Electron Cyclotron Maser Emission. *Astron. J.* **165**, 1 (2023).
- [147] Narang, M. *et al.* uGMRT observations of the hot-Saturn WASP-69b: Radio-Loud Exoplanet-Exomoon Survey II (RLEES II). *Mon. Not. R. Astron. Soc.* **522**, 1662–1668 (2023).
- [148] Grießmeier, J.-M., Zarka, P. & Spreeuw, H. Predicting low-frequency radio fluxes of known extrasolar planets. *A&A* **475**, 359–368 (2007).
- [149] Zarka, P. in *Star-Planet Interactions in the Radio Domain: Prospect for Their Detection* (eds Deeg, H. J. & Belmonte, J. A.) *Handbook of Exoplanets* 22 (2018).
- [150] Winglee, R. M., Dulk, G. A. & Bastian, T. S. A search for cyclotron maser radiation from substellar and planet-like companions of nearby stars. *ApJl* **309**, L59–L62 (1986).
- [151] Bastian, T. S., Dulk, G. A. & Leblanc, Y. A Search for Radio Emission from Extrasolar Planets. *ApJ* **545**, 1058–1063 (2000).
- [152] Lazio, T. J. W. & Farrell, W. M. Magnetospheric Emissions from the Planet Orbiting  $\tau$  Bootis: A Multiepoch Search. *ApJ* **668**, 1182–1188 (2007).
- [153] Smith, A. M. S. *et al.* Secondary radio eclipse of the transiting planet HD 189733 b: an upper limit at 307–347 MHz. *MNRAS* **395**, 335–341 (2009).
- [154] Lazio, T. J. W. *et al.* A Blind Search for Magnetospheric Emissions from Planetary Companions to Nearby Solar-Type Stars. *AJ* **139**, 96–101 (2010).
- [155] Hallinan, G. *et al.* Looking for a Pulse: A Search for Rotationally Modulated Radio Emission from the Hot Jupiter,  $\tau$  Boötis b. *ApJ* **762**, 34 (2013).
- [156] Murphy, T. *et al.* Limits on low-frequency radio emission from southern exoplanets with the Murchison Widefield Array. *Mon. Not. R. Astron. Soc.* **446**, 2560–2565 (2015).
- [157] Lenc, E., Murphy, T., Lynch, C. R., Kaplan, D. L. & Zhang, S. N. An all-sky survey of circular polarization at 200 MHz. *Mon. Not. R. Astron.*

- Soc.* **478**, 2835–2849 (2018).
- [158] O’Gorman, E. *et al.* A search for radio emission from exoplanets around evolved stars. *Astron. Astrophys.* **612**, A52 (2018).
- [159] de Gasperin, F., Lazio, T. J. W. & Knapp, M. Radio Observations of HD80606 Near Planetary Periastron: II. LOFAR Low Band Antenna Observations at 30-78 MHz. *arXiv e-prints* arXiv:2011.05696 (2020).
- [160] Cendes, Y., Williams, P. K. G. & Berger, E. A Pilot Radio Search for Magnetic Activity in Directly Imaged Exoplanets. *Astron. J.* **163**, 15 (2022).
- [161] Route, M. & Wolszczan, A. ROME. III. The Arecibo Search for Star-Planet Interactions at 5 GHz. *Astrophys. J.* **952**, 118 (2023).
- [162] Bloot, S. *et al.* Phenomenology and periodicity of radio emission from the stellar system AU Microscopii. *Astron. Astrophys.* **682**, A170 (2024).
- [163] Lecavelier des Etangs, A., Sirothia, S. K., Gopal-Krishna & Zarka, P. Hint of 150 MHz radio emission from the Neptune-mass extrasolar transiting planet HAT-P-11b. *A&A* **552**, A65 (2013).
- [164] Sirothia, S. K., Lecavelier des Etangs, A., Gopal-Krishna, Kantharia, N. G. & Ishwar-Chandra, C. H. Search for 150 MHz radio emission from extrasolar planets in the TIFR GMRT Sky Survey. *A&A* **562**, A108 (2014).
- [165] Vasylieva, I. *Pulsars and transients survey, and exoplanet search at low-frequencies with the UTR-2 radio telescope: methods and first results.* Theses, Paris Observatory (2015). URL <https://tel.archives-ouvertes.fr/tel-01246634>.
- [166] Pineda, J. S. & Villadsen, J. Coherent radio bursts from known M-dwarf planet-host YZ Ceti. *Nature Astronomy* **7**, 569–578 (2023).
- [167] Kavanagh, R. D. *et al.* MOVES - II. Tuning in to the radio environment of HD189733b. *Mon. Not. R. Astron. Soc.* **485**, 4529–4538 (2019).
- [168] Ashtari, R., Sciola, A., Turner, J. D. & Stevenson, K. Detecting Magnetospheric Radio Emission from Giant Exoplanets. *Astrophys. J.* **939**, 24 (2022).
- [169] Fischer, C. & Saur, J. Time-variable Electromagnetic Star-Planet Interaction: The TRAPPIST-1 System as an Exemplary Case. *apj* **872**, 113 (2019).
- [170] Elekes, F. & Saur, J. Space environment and magnetospheric Poynting fluxes of the exoplanet  $\tau$  Boötis b. *Astron. Astrophys.* **671**, A133 (2023).
- [171] Weber, C. *et al.* How expanded ionospheres of Hot Jupiters can prevent escape of radio emission generated by the cyclotron maser instability. *Mon. Not. R. Astron. Soc.* **469**, 3505–3517 (2017).
- [172] Weber, C. *et al.* Fischer, G., Mann, G., Panchenko, M. & Zarka, P. (eds) *On the Cyclotron Maser Instability in Magnetospheres of Hot Jupiters - Influence of ionosphere models.* (eds Fischer, G., Mann, G., Panchenko, M. & Zarka, P.) *Planetary Radio Emissions VIII*, 317–329 (Austrian Academy of Sciences Press, Vienna, 2017).
- [173] Daley-Yates, S. & Stevens, I. R. Inhibition of the electron cyclotron

- maser instability in the dense magnetosphere of a hot Jupiter. *Mon. Not. R. Astron. Soc.* **479**, 1194–1209 (2018).
- [174] Erkaev, N. V. *et al.* Can radio emission escape from the magnetosphere of *v* Andromedae b - a new method to constrain the minimum mass of Hot Jupiters. *Mon. Not. R. Astron. Soc.* **512**, 4869–4876 (2022).
- [175] Brain, D. A., Kao, M. M. & O'Rourke, J. G. Exoplanet Magnetic Fields. *arXiv e-prints* arXiv:2404.15429 (2024).
- [176] Connerney, J. E. P. *et al.* A New Model of Jupiter's Magnetic Field at the Completion of Juno's Prime Mission. *Journal of Geophysical Research (Planets)* **127**, e07055 (2022).
- [177] Berger, E. *et al.* Discovery of radio emission from the brown dwarf LP944-20. *Nature* **410**, 338–340 (2001).
- [178] Hallinan, G. *et al.* Periodic Bursts of Coherent Radio Emission from an Ultracool Dwarf. *Astrophys. J. Lett.* **663**, L25–L28 (2007).
- [179] Hallinan, G. *et al.* Magnetospherically driven optical and radio aurorae at the end of the stellar main sequence. *Nature* **523**, 568–571 (2015).
- [180] Kao, M. M. *et al.* Auroral Radio Emission from Late L and T Dwarfs: A New Constraint on Dynamo Theory in the Substellar Regime. *Astrophys. J.* **818**, 24 (2016).
- [181] Pineda, J. S., Hallinan, G. & Kao, M. M. A Panchromatic View of Brown Dwarf Aurorae. *Astrophys. J.* **846**, 75 (2017).
- [182] Kao, M. M., Mioduszewski, A. J., Villadsen, J. & Shkolnik, E. L. Resolved imaging of an extrasolar radiation belt around an ultracool dwarf. *arXiv e-prints* arXiv:2302.12841 (2023).
- [183] Climent, J. B., Guirado, J. C., Pérez-Torres, M., Marcaide, J. M. & Peña-Moñino, L. Evidence of a radiation belt around a brown dwarf. *arXiv e-prints* arXiv:2303.06453 (2023).
- [184] Berger, E. *et al.* The Magnetic Properties of an L Dwarf Derived from Simultaneous Radio, X-Ray, and H $\alpha$  Observations. *Astrophys. J.* **627**, 960–973 (2005).
- [185] Williams, P. K. G., Cook, B. A. & Berger, E. Trends in Ultracool Dwarf Magnetism. I. X-Ray Suppression and Radio Enhancement. *Astrophys. J.* **785**, 9 (2014).
- [186] Osten, R. A., Hawley, S. L., Allred, J. C., Johns-Krull, C. M. & Roark, C. From Radio to X-Ray: Flares on the dMe Flare Star EV Lacertae. *Astrophys. J.* **621**, 398–416 (2005).
- [187] Osten, R. A. & Jayawardhana, R. Radio Constraints on Activity in Young Brown Dwarfs. *Astrophys. J. Lett.* **644**, L67–L70 (2006).
- [188] Berger, E. *et al.* Simultaneous Multi-Wavelength Observations of Magnetic Activity in Ultracool Dwarfs. III. X-ray, Radio, and H $\alpha$  Activity Trends in M and L dwarfs. *Astrophys. J.* **709**, 332–341 (2010).
- [189] Antonova, A. *et al.* Volume-limited radio survey of ultracool dwarfs. *Astron. Astrophys.* **549**, A131 (2013).
- [190] Burgasser, A. J., Melis, C., Zauderer, B. A. & Berger, E. Detection

- of Radio Emission from the Hyperactive L Dwarf 2MASS J13153094-2649513AB. *Astrophys. J. Lett.* **762**, L3 (2013).
- [191] Route, M. & Wolszczan, A. The Arecibo Detection of the Coolest Radio-flaring Brown Dwarf. *Astrophys. J. Lett.* **747**, L22 (2012).
- [192] Route, M. & Wolszczan, A. The 5 GHz Arecibo Search for Radio Flares from Ultracool Dwarfs. *Astrophys. J.* **773**, 18 (2013).
- [193] Route, M. & Wolszczan, A. The Second Arecibo Search for 5 GHz Radio Flares from Ultracool Dwarfs. *Astrophys. J.* **830**, 85 (2016).
- [194] Lynch, C. *et al.* Radio detections of southern ultracool dwarfs. *Mon. Not. R. Astron. Soc.* **457**, 1224–1232 (2016).
- [195] Kao, M. M. & Shkolnik, E. L. The Occurrence Rate of Quiescent Radio Emission for Ultracool Dwarfs using a Generalized Semi-Analytical Bayesian Framework. *arXiv e-prints* arXiv:2306.16460 (2023).
- [196] Vedantham, H. K. *et al.* Direct Radio Discovery of a Cold Brown Dwarf. *Astrophys. J. Lett.* **903**, L33 (2020).
- [197] Kao, M. M., Hallinan, G., Pineda, J. S., Stevenson, D. & Burgasser, A. The Strongest Magnetic Fields on the Coolest Brown Dwarfs. *Astrophys. J. Suppl. Ser.* **237**, 25 (2018).
- [198] Vedantham, H. K. *et al.* Polarised radio pulsations from a new T-dwarf binary. *Astron. Astrophys.* **675**, L6 (2023).
- [199] Rose, K. *et al.* Periodic Radio Emission from the T8 Dwarf WISE J062309.94-045624.6. *Astrophys. J. Lett.* **951**, L43 (2023).
- [200] Best, W. M. J., Sanghi, A., Liu, M. C., Magnier, E. A. & Dupuy, T. J. A Volume-Limited Sample of Ultracool Dwarfs. II. The Substellar Age and Mass Functions in the Solar Neighborhood. *arXiv e-prints* arXiv:2401.09535 (2024).
- [201] Kao, M. M., Hallinan, G. & Pineda, J. S. Constraints on magnetospheric radio emission from Y dwarfs. *Mon. Not. R. Astron. Soc.* **487**, 1994–2004 (2019).
- [202] Delrez, L. *et al.* Marshall, H. K. & Spyromilio, J. (eds) *SPECULOOS: a network of robotic telescopes to hunt for terrestrial planets around the nearest ultracool dwarfs*. (eds Marshall, H. K. & Spyromilio, J.) *Ground-based and Airborne Telescopes VII*, Vol. 10700 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 107001I (2018).
- [203] Tamburo, P. *et al.* The Perkins Infrared Exosatellite Survey (PINES) I. Survey Overview, Reduction Pipeline, and Early Results. *Astron. J.* **163**, 253 (2022).
- [204] Limbach, M. A. *et al.* On the Detection of Exomoons Transiting Isolated Planetary-mass Objects. *Astrophys. J. Lett.* **918**, L25 (2021).
- [205] Gillon, M. *et al.* Seven temperate terrestrial planets around the nearby ultracool dwarf star TRAPPIST-1. *Nature* **542**, 456–460 (2017).
- [206] Curiel, S., Ortiz-León, G. N., Mioduszewski, A. J. & Torres, R. M. An Astrometric Planetary Companion Candidate to the M9 Dwarf TVLM 513-46546. *Astron. J.* **160**, 97 (2020).
- [207] Saur, J. *et al.* Brown dwarfs as ideal candidates for detecting UV

- aurora outside the Solar System: Hubble Space Telescope observations of 2MASS J1237+6526 **655**, A75 (2021).
- [208] Zarka, P. *Star-Planet Interactions in the Radio Domain: Prospect for Their Detection*, 22 (2018).
- [209] Griessmeier, J. M. *Future Exoplanet Research: Radio Detection and Characterization*, 159 (2018).
- [210] Preusse, S., Kopp, A., Büchner, J. & Motschmann, U. A magnetic communication scenario for hot Jupiters **460**, 317–322 (2006).
- [211] Kopp, A., Schilp, S. & Preusse, S. Magnetohydrodynamic Simulations of the Magnetic Interaction of Hot Jupiters with Their Host Stars: A Numerical Experiment. *Astrophys. J.* **729**, 116 (2011).
- [212] Louis, C. K., Louarn, P., Allegrini, F., Kurth, W. S. & Szalay, J. R. Ganymede-Induced Decametric Radio Emission: In Situ Observations and Measurements by Juno. *Geophys. Res. Lett.* **47**, e90021 (2020).
- [213] Scharf, C. A. Possible Constraints on Exoplanet Magnetic Field Strengths from Planet-star Interaction. *Astrophys. J.* **722**, 1547–1555 (2010).
- [214] Shkolnik, E., Walker, G. A. H., Bohlender, D. A., Gu, P.-G. & Kürster, M. Hot Jupiters and Hot Spots: The Short- and Long-Term Chromospheric Activity on Stars with Giant Planets. *ApJ* **622**, 1075–1090 (2005).
- [215] Shkolnik, E., Bohlender, D. A., Walker, G. A. H. & Collier Cameron, A. The On/Off Nature of Star-Planet Interactions. *ApJ* **676**, 628–638 (2008).
- [216] Klein, B. *et al.* One year of AU Mic with HARPS - II. Stellar activity and star-planet interaction. *Mon. Not. R. Astron. Soc.* **512**, 5067–5084 (2022).
- [217] Poppenhaeger, K. & Schmitt, J. H. M. M. A Correlation Between Host Star Activity and Planet Mass for Close-in Extrasolar Planets? *Astrophys. J.* **735**, 59 (2011).
- [218] Scandariato, G. *et al.* A coordinated optical and X-ray spectroscopic campaign on HD 179949: searching for planet-induced chromospheric and coronal activity. *Astron. Astrophys.* **552**, A7 (2013).
- [219] Miller, B. P., Gallo, E., Wright, J. T. & Pearson, E. G. A Comprehensive Statistical Assessment of Star-Planet Interaction. *Astrophys. J.* **799**, 163 (2015).
- [220] Viswanath, G., Narang, M., Manoj, P., Mathew, B. & Kartha, S. S. A Statistical Search for Star-Planet Interaction in the Ultraviolet Using GALEX. *Astron. J.* **159**, 194 (2020).
- [221] Pope, B. J. S. *et al.* No Massive Companion to the Coherent Radio-emitting M Dwarf GJ 1151. *Astrophys. J. Lett.* **890**, L19 (2020).
- [222] Mahadevan, S. *et al.* The Habitable-zone Planet Finder Detects a Terrestrial-mass Planet Candidate Closely Orbiting Gliese 1151: The Likely Source of Coherent Low-frequency Radio Emission from an Inactive Star. *Astrophys. J. Lett.* **919**, L9 (2021).

- [223] Perger, M. *et al.* The CARMENES search for exoplanets around M dwarfs. No evidence for a super-Earth in a 2-day orbit around GJ 1151. *Astron. Astrophys.* **649**, L12 (2021).
- [224] Blanco-Pozo, J. *et al.* The CARMENES search for exoplanets around M dwarfs. A long-period planet around GJ 1151 measured with CARMENES and HARPS-N data. *Astron. Astrophys.* **671**, A50 (2023).
- [225] Trigilio, C. *et al.* Star-Planet Interaction at radio wavelengths in YZ Ceti: Inferring planetary magnetic field. *arXiv e-prints* arXiv:2305.00809 (2023).
- [226] Zarka, P. Auroral radio emissions at the outer planets: Observations and theories. *J. Geophys. Res.* **103**, 20159–20194 (1998).
- [227] Lamy, L. *et al.* The low-frequency source of Saturn’s kilometric radiation. *Science* **362**, aat2027 (2018).
- [228] Louis, C. K. *et al.* ExPRES: an Exoplanetary and Planetary Radio Emissions Simulator. *Astron. Astrophys.* **627**, A30 (2019).
- [229] Sulaiman, A. H. *et al.* Jupiter’s Low-Altitude Auroral Zones: Fields, Particles, Plasma Waves, and Density Depletions. *Journal of Geophysical Research (Space Physics)* **127**, e30334 (2022).
- [230] Lamy, L. *et al.* Determining the Beaming of Io Decametric Emissions: A Remote Diagnostic to Probe the Io-Jupiter Interaction. *Journal of Geophysical Research (Space Physics)* **127**, e30160 (2022).
- [231] Fares, R. *et al.* Searching for star-planet interactions within the magnetosphere of HD189733. *Mon. Not. R. Astron. Soc.* **406**, 409–419 (2010).
- [232] Lanza, A. F. Stellar coronal magnetic fields and star-planet interaction. *Astron. Astrophys.* **505**, 339–350 (2009).
- [233] Strugarek, A. Physics of star-planet magnetic interactions. *arXiv e-prints* arXiv:2104.05968 (2021).
- [234] Kivelson, M. G. *et al.* in *Magnetospheric interactions with satellites* (eds Bagenal, F., Dowling, T. E. & McKinnon, W. B.) *Jupiter. The Planet, Satellites and Magnetosphere*, Vol. 1 513–536 (2004).
- [235] Strugarek, A., Brun, A. S., Matt, S. P. & Réville, V. Magnetic Games between a Planet and Its Host Star: The Key Role of Topology **815**, 111 (2015).
- [236] Goldreich, P. & Lynden-Bell, D. Io, a jovian unipolar inductor. *Astrophys. J.* **156**, 59–78 (1969).
- [237] Lanza, A. F. Star-planet magnetic interaction and evaporation of planetary atmospheres. *aap* **557**, A31 (2013).
- [238] Strugarek, A. *et al.* MOVES - V. Modelling star-planet magnetic interactions of HD 189733. *Mon. Not. R. Astron. Soc.* **512**, 4556–4572 (2022).
- [239] Shoda, M. *et al.* Alfvén-wave-driven Magnetic Rotator Winds from Low-mass Stars. I. Rotation Dependences of Magnetic Braking and Mass-loss Rate. *Astrophys. J.* **896**, 123 (2020).
- [240] Fossati, L. *et al.* Absorbing Gas around the WASP-12 Planetary System.

- Astrophys. J. Lett.* **766**, L20 (2013).
- [241] Osten, R. A. & Wolk, S. J. Connecting Flares and Transient Mass-loss Events in Magnetically Active Stars. *Astrophys. J.* **809**, 79 (2015).
- [242] Vidotto, A. A. The evolution of the solar wind. *Living Reviews in Solar Physics* **18**, 3 (2021).
- [243] Wood, B. E. Astrospheres and Solar-like Stellar Winds. *Living Reviews in Solar Physics* **1**, 2 (2004).
- [244] Jardine, M. & Collier Cameron, A. Slingshot prominences: nature’s wind gauges. *Mon. Not. R. Astron. Soc.* **482**, 2853–2860 (2019).
- [245] Kavanagh, R. D. *et al.* Planet-induced radio emission from the coronae of M dwarfs: the case of Prox Cen and AU Mic. *Mon. Not. R. Astron. Soc.* **504**, 1511–1518 (2021).
- [246] Fischer, C. & Saur, J. Time-variable Electromagnetic Star-Planet Interaction: The TRAPPIST-1 System as an Exemplary Case. *Astrophys. J.* **872**, 113 (2019).
- [247] Dewdney, P. E., Hall, P. J., Schilizzi, R. T. & Lazio, T. J. L. W. The Square Kilometre Array. *IEEE Proceedings* **97**, 1482–1496 (2009).
- [248] Callingham, J. R. *et al.* V-LoTSS: The circularly polarised LOFAR Two-metre Sky Survey. *Astron. Astrophys.* **670**, A124 (2023).
- [249] Pope, B. J. S., Withers, P., Callingham, J. R. & Vogt, M. F. Exoplanet transits with next-generation radio telescopes. *Mon. Not. R. Astron. Soc.* **484**, 648–658 (2019).
- [250] Burns, J. O. Transformative science from the lunar farside: observations of the dark ages and exoplanetary systems at low radio frequencies. *Philosophical Transactions of the Royal Society of London Series A* **379**, 20190564 (2021).
- [251] Burns, J. O. *et al.* Low Radio Frequency Observations from the Moon Enabled by NASA Landed Payload Missions. *Planet. Sci. J.* **2**, 44 (2021).
- [252] Bellotti, S. *et al.* Monitoring the large-scale magnetic field of AD Leo with SPIRou, ESPaDOnS, and Narval. Towards a magnetic polarity reversal? *Astron. Astrophys.* **676**, A56 (2023).
- [253] Aigrain, S. & Foreman-Mackey, D. Gaussian Process regression for astronomical time-series. *arXiv e-prints* arXiv:2209.08940 (2022).
- [254] VanderPlas, J. T. Understanding the Lomb-Scargle Periodogram. *Astrophys. J. Suppl. Ser.* **236**, 16 (2018).
- [255] Artigau, É. *et al.* SPIRou: the near-infrared spectropolarimeter/high-precision velocimeter for the Canada-France-Hawaii telescope. *Ground-based and Airborne Instrumentation for Astronomy V* **9147**, 914715 (2014).
- [256] Vidotto, A. A., Feeney, N. & Groh, J. H. Can we detect aurora in exoplanets orbiting M dwarfs? *Mon. Not. R. Astron. Soc.* **488**, 633–644 (2019).