



HAL
open science

Exploring the variations in ambient BTEX in urban Europe and its environmental health implications

Xiansheng Liu, Xun Zhang, Marvin Dufresne, Tao Wang, Lijie Wu, Rosa Lara, Roger Seco, Marta Monge, Ana Maria Yáñez-Serrano, Marie Gohy, et al.

► **To cite this version:**

Xiansheng Liu, Xun Zhang, Marvin Dufresne, Tao Wang, Lijie Wu, et al.. Exploring the variations in ambient BTEX in urban Europe and its environmental health implications. *Atmospheric Chemistry and Physics*, 2025, 25 (1), pp.625-638. <10.5194/acp-25-625-2025>. <hal-04786203>

HAL Id: hal-04786203

<https://hal.science/hal-04786203v1>

Submitted on 22 May 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



Measurement report: Exploring the variations in ambient BTEX in urban Europe and their environmental health implications

Xiansheng Liu¹, Xun Zhang², Marvin Dufresne³, Tao Wang⁴, Lijie Wu², Rosa Lara¹, Roger Seco¹, Marta Monge¹, Ana Maria Yáñez-Serrano^{1,5,6}, Marie Gohy⁷, Paul Petit⁷, Audrey Chevalier⁸, Marie-Pierre Vagnot⁹, Yann Fortier⁹, Alexia Baudic¹⁰, Véronique Gherzi¹⁰, Grégory Gille¹¹, Ludovic Lanzi¹¹, Valérie Gros¹², Leïla Simon¹², Heidi Hellen¹³, Stefan Reimann¹⁴, Zoé Le Bras¹⁴, Michelle Jessy Müller¹⁴, David Beddows¹⁵, Siqi Hou¹⁵, Zongbo Shi¹⁵, Roy M. Harrison¹⁵, William Bloss¹⁵, James Darnie¹⁶, Stéphane Sauvage³, Philip K. Hopke^{17,18}, Xiaoli Duan¹⁹, Taicheng An²⁰, Alastair C. Lewis²¹, James R. Hopkins²¹, Eleni Liakakou²², Nikolaos Mihalopoulos^{22,23}, Xiaohu Zhang²⁴, Andrés Alastuey¹, Xavier Querol¹, and Thérèse Salameh³

¹Institute of Environmental Assessment and Water Research (IDAEA-CSIC), 08034 Barcelona, Spain

²Beijing Key Laboratory of Big Data Technology for Food Safety, School of Computer and Artificial Intelligence, Beijing Technology and Business University, Beijing 100048, China

³IMT Nord Europe, Institut Mines-Télécom, Univ. Lille, Centre for Energy and Environment, 59000 Lille, France

⁴Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention, Department of Environmental Science and Engineering, Fudan University, Shanghai 200433, China

⁵CREAF, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain

⁶CSIC, Global Ecology Unit, CREAM-CSIC-UAB, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain

⁷Institut Scientifique de Service Public (ISSeP), 4000 Liège, Belgium

⁸Atmo Grand-Est (AtmoGE), 67300 Schiltigheim, France

⁹Atmo Auvergne-Rhône-Alpes (AtmoAURA), 69500 Bron, France

¹⁰Airparif, Air Quality Monitoring Network for the Greater Paris Area, 7 rue Crillon, 75004 Paris, France

¹¹AtmoSud, 13006 Marseille, France

¹²Laboratoire des Sciences du Climat et de l'Environnement (CEA-CNRS-UVSQ, IPSL), CAE/Orme des Merisiers, 91191 Gif-sur-Yvette, France

¹³Atmospheric Composition Research, Finish Meteorological Institute (FMI), 00560 Helsinki, Finland

¹⁴Empa, Laboratory for Air Pollution / Environmental Technology, 8600 Dübendorf, Switzerland

¹⁵School of Geography Earth and Environmental Sciences, University of Birmingham, B15 2TT Birmingham, United Kingdom

¹⁶Ricardo Energy & Environment, Gemini Building, Fermi Avenue, Harwell, Oxfordshire, OX11 0QR, UK

¹⁷Institute for a Sustainable Environment, Clarkson University, Potsdam, NY 13699, USA

¹⁸Departments of Public Health Sciences and Environmental Medicine, University of Rochester School of Medicine and Dentistry, Rochester, NY 14642, USA

¹⁹School of Energy and Environmental Engineering, University of Science and Technology Beijing, Beijing 100083, China

²⁰School of Environmental Science and Engineering, Guangdong University of Technology, Guangzhou 510006, China

²¹National Centre for Atmospheric Science (NCAS), Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK

²²Institute for Environmental Research and Sustainable Development, National Observatory of Athens, 15236 Athens, Greece

²³Environmental Chemical Processes Laboratory, Department of Chemistry,
University of Crete, 70013 Crete, Greece

²⁴National Engineering and Technology Center for Information Agriculture,
Nanjing Agricultural University, Nanjing 210095, China

Correspondence: Xun Zhang (zhangxun@btbu.edu.cn) and Thérèse Salameh
(therese.salameh@imt-nord-europe.fr)

Received: 23 July 2024 – Discussion started: 31 July 2024

Revised: 12 November 2024 – Accepted: 21 November 2024 – Published: 17 January 2025

Abstract. BTEX (benzene, toluene, ethylbenzene, and *m*-xylene, *p*-xylene, and *o*-xylene) are significant urban air pollutants. This study examines BTEX variability across 7 European countries using data from 22 monitoring sites in different urban settings (urban background, traffic, industry, and suburban background). Results indicate that the relative abundance of BTEX in urban areas follows the order toluene > benzene > *m*, *p*-xylene > *o*-xylene > ethylbenzene, with median mixing ratios of 266 ± 152 , 163 ± 74 , 129 ± 88 , 53 ± 35 , and 45 ± 27 ppt during the years 2017–2022, respectively. Seasonal trends show benzene had similar median concentrations across urban background, traffic, and industrial sites, indicating mixed sources. Toluene levels were highest in traffic and industrial areas, highlighting road traffic and industrial emissions. Ethylbenzene and xylenes showed equivalent levels in traffic and industrial areas but were lower in urban backgrounds. Peak BTEX levels occurred during morning and evening rush hours, linked to traffic, heating, and atmospheric stagnation. B/T ratios ranged from 0.29 ± 0.11 to 1.35 ± 0.95 , and X/E ratios ranged from 1.75 ± 0.91 to 3.68 ± 0.30 , indicating primary pollution from local traffic, followed by solvents, coatings, and biomass burning. Lifetime cancer risk from BTEX exposure was below the definite risk threshold (10^{-4}) but above the permissible risk level (10^{-6}), suggesting moderate risk from benzene and ethylbenzene, particularly in traffic and industrial areas. Additionally, the health index of BTEX at monitoring sites was generally lower than the threshold limit value, suggesting a low non-carcinogenic risk overall. This study offers essential insights into BTEX pollution in urban European environments.

1 Introduction

Atmospheric volatile organic compounds (VOCs) are significant precursors of tropospheric ozone (O₃) (Grosjean and Seinfeld, 1989) and secondary organic aerosols (SOAs) (Derwent et al., 2010). Among the most prevalent VOCs, BTEX – benzene, toluene, ethylbenzene, and xylenes – are a typical component of air pollution (Miri et al., 2016). Due to their reactivity with hydroxyl radicals (•OH) during daylight and nitrate radicals (NO₃•) both day and night, they can generate additional radicals, such as peroxyalkyl and hydroperoxy radicals, which further oxidize nitric oxide (NO) to nitrogen dioxide (NO₂) (Garg and Gupta, 2019; Ghaffari et al., 2021). This process contributes to increased concentrations of tropospheric O₃ and SOA formation (Ng et al., 2007). Additionally, exposure to BTEX is heightened by solvent evaporation, vehicular traffic, and emissions from fossil fuel extraction, which are increasingly occurring near densely populated areas (Bolden et al., 2015; Salameh et al., 2019; AQEG-EU, 2020; Liu et al., 2023). In addition, the removal of toxic octane enhancers like lead and methyl tertiary-butyl ether (MTBE) from gasoline allowed refiners to increase the volume of aromatics in gasoline to meet anti-knock requirements and enhance octane ratings (Yang et al., 2019). Thus,

due to its utility, ubiquity, and economic importance, BTEX are likely to remain persistent environmental pollutants in the air for the foreseeable future (Bolden et al., 2015; Davidson et al., 2021).

Furthermore, there is a body of evidence that BTEX have the potential to irritate various organs within the human body, including the respiratory tract, lungs, bronchi, skin, and heart, and that exposure to elevated concentrations can result in acute effects such as dizziness and vomiting (Davidson et al., 2021; Li et al., 2021; Ogbodo et al., 2022). According to the International Agency for Research on Cancer (IARC), benzene is recognized as a significant public health threat substance and is classified as a carcinogen (WHO, 2016). Additionally, the US EPA's carcinogenicity classification system categorizes benzene as a human carcinogen (Group A), supported by ample evidence of its carcinogenic effects in humans. On the other hand, toluene, ethylbenzene, and xylenes are not classified to human carcinogenicity (Group D) with inadequate (or no) evidence, while they have important non-carcinogenic toxic potentials (Durmusoglu et al., 2010). In 1990, the US Clean Air Act Amendments classified seven categories of VOCs (<https://www.epa.gov/haps>, 21 April 2024), including BTEX and three other VOCs (*n*-hexane, 1,3-butadiene, and styrene) as hazardous pollu-

tants, highlighting their potential to cause human cancer and other serious health issues. Simultaneously, the Environmental Protection Directory issued by the Ministry of Ecology and Environment of the People's Republic of China designated BTEX and styrene as high-pollution and high-risk environmental pollutants (Song et al., 2018). Furthermore, the European Air Quality Index (DIRECTIVE(EU), 2024) establishes the latest limit value of $3.4 \mu\text{g m}^{-3}$ for the annual mean of benzene in ambient air, with all member countries required to monitor and report this carcinogenic compound to the European Environment Agency (EEA) database (<https://www.eea.europa.eu/en>, last access: 21 April 2024). Since BTEX members are small molecules with lipophilic characteristics, they easily penetrate the human body (Zahed et al., 2024). However, studies focusing on BTEX within the European Union (EU) have primarily centered on member states. For instance, Borbon et al. (2018) conducted long-term observations of hydrocarbons, including TEX, in traffic and urban background locations in London, Paris, and Strasbourg. They estimated the relative importance of traffic emissions for TEX in each city and found that traffic emissions would no longer dominate TEX concentrations in urban areas of Europe. Despite this, studies addressing the overall BTEX pollution status and health risk assessment at the EU level remain relatively insufficient.

RI-URBANS (Research Infrastructures Services Reinforcing Air Quality Monitoring Capacities in European Urban and Industrial Areas, funded by the EU's Horizon 2020 research and innovation program, 101036245) is a European research project that demonstrates the application of advanced air quality service tools in urban Europe to improve the assessment of air quality policies, including a more accurate evaluation of health effects. In this context, this study focused on collecting and evaluating both online and offline BTEX data from 19 cities (total 22 monitoring sites) across 7 European countries (Belgium, Finland, France, Greece, Switzerland, Spain, and the United Kingdom). The study's objectives included (i) comparing BTEX concentrations across Europe, (ii) identifying sources of BTEX, and (iii) assessing the health impacts of BTEX. By comprehensively assessing BTEX levels in major European cities and their harmful effects on human health, this study aims to provide data support for the coordinated management of BTEX monitoring across Europe.

2 Methodology

2.1 Instrumentation

The instrumentation used for measuring BTEX at different stations, as described in Table 1, encompassed 3 industrial (IND) sites, 2 traffic (TR) sites, 16 urban background (UB) sites, and 1 suburban background (SUB) site. Briefly, in this study the BTEX were measured by thermal desorption gas chromatography with flame ionization detectors (TD-GC-

FID/2FID), thermal desorption gas chromatography–mass spectrometry (TD-GC-MS), proton-transfer-reaction time-of-flight mass spectrometry (PTR-TOF-MS), PTR-Quad-MS, and passive samplers. It should be noted that comparing such heterogeneous datasets, without an intercomparison exercise, is a limitation of the study.

In addition, as shown in Table 1, the monitoring periods differed for each site. We considered only BTEX data obtained from urban sites between the years 2017 and 2022, with the exception of HEL_UB (Helsinki) (only February 2016) and ATH_UB (Athens) (only 2016–2017), which provided shorter time series, to reduce the uncertainties when comparing BTEX mixing ratios and health risk assessment between sites due to long-term trends at each site (Table 1, Fig. 1). Also, BCN_UB (Barcelona) and PAR_SUB (Paris – SIRTIA) monitored the combined levels of ethylbenzene and xylenes. However, because the PTR-MS technique used at these sites does not distinguish between isomers (Simon et al., 2023), we excluded these data from our analysis when comparing BTEX mixing ratios. Additionally, BAQS_UB (Birmingham) did not detect *m*-xylene or *p*-xylene.

2.2 Risk assessment

Health risk assessments were conducted to evaluate the impacts on human health from exposure to atmospheric pollutants through inhalation, ingestion, and dermal penetration (Li et al., 2014; Yao et al., 2019; Nie et al., 2020; Zahed et al., 2024). Among the aforementioned pathways, inhalation is regarded as the main route. Therefore, in this study, the health risks associated with inhalation exposure were assessed. The lifetime cancer risk (LCR) was calculated based on the method proposed by the US Environmental Protection Agency (EPA) in 2009 (EPA-540-R-070-002) (EPA, 2009). This method was employed to evaluate carcinogenic risks of inhaled benzene and ethylbenzene to the population. The formula for calculation is as follows: $\text{LCR} = \text{CA} \times \text{IUR}$ (inhalation unit risk), where CA represents the concentration of pollutants in the atmosphere ($\mu\text{g m}^{-3}$). The mixing ratio was converted into concentrations for benzene and ethylbenzene. The IUR is the unit risk value for inhalation exposure ($\mu\text{g m}^{-3}$)⁻¹, indicating the maximum probability of developing cancer when exposed to a certain dose of a pollutant through inhalation. The IUR values for benzene and ethylbenzene are 7.8×10^{-6} and 2.5×10^{-6} ($\mu\text{g m}^{-3}$)⁻¹, respectively, as derived from the US EPA IRIS (EPA, 2020).

According to US EPA standards, LCR represents the probability of cancer occurrence in the exposed population, usually expressed as the ratio of the number of cancer cases per unit population. If the LCR falls below 10^{-6} (indicating an increase of 1 cancer case per 1 million people in a lifetime), it is widely considered negligible. Some jurisdictions consider risks as high as 10^{-4} as tolerable and those higher than this as requiring urgent action. Specially for this study, we adopted US EPA values as they provide a standardized framework

Table 1. Details on measuring sites and instrumentation used for offline and online BTEX datasets in this study (IND, industry; TR, traffic; UB, urban background; SUB, suburban background).

City (country)	Abbreviation	Instrument type/model	Start date	End date	Types
Athens (GR)	ATH_UB	TD-GC-FID	1 Mar 2016	28 Feb 2017	Online
Angleur (BE)	ANG_UB	Passive samplers	1 Jan 2018	31 Dec 2021	Offline
Birmingham (UK)	BAQS_UB	TD-GC-FID	14 Apr 2021	6 Jun 2022	Online
Barcelona (ES)	BCN_UB	PTR-TOF-MS	7 Oct 2021	31 Dec 2022	Online
Charleroi (BE)	CHM_UB	Passive samplers	2 Jan 2010	30 Dec 2021	Offline
Grenoble (FR)	GRE_UB	TD-GC-2FID PerkinElmer	16 Jan 2015	29 Dec 2022	Offline
Helsinki (FI)	HEL_UB	TD (Markes)-GC-MS (Agilent)	20 Jan 2011	18 Nov 2011	Online
		TD-GC-MS PerkinElmer	28 Jan 2016	24 Feb 2016	Online
Herstal (BE)	HET_UB	Passive samplers	1 Jan 2013	31 Dec 2021	Offline
Lodelinsart (BE)	LDS_UB	Passive samplers	2 Jan 2018	30 Dec 2021	Offline
Eltham (UK)	LND_UB	TD-GC-FID	1 Jan 2008	1 Jan 2022	Online
Marseille (FR)	MAR_UB	TD-GC-2FID PerkinElmer	31 Jan 2019	13 Aug 2020	Online
Mons (BE)	MON_UB	Passive samplers	2 Jan 2010	30 Dec 2021	Offline
Namur (BE)	NAM_UB	Passive samplers	1 Jan 2018	31 Dec 2021	Offline
Paris – Paris 1er Les Halles (FR)	PAR_UB	TD-GC-FID (30 min)	1 Jan 2020	1 Jan 2022	Online
Strasbourg Ouest (FR)	STB_UB	TD-GC-FID	8 Jul 2002	30 Aug 2021	Online
Zurich (CH)	ZUR_UB	TD-GC-2FID	1 Jan 2016	31 Dec 2017	Online
Helsinki (FI)	HEL_TR	TD-GC-MS PerkinElmer	21 Aug 2019	11 Sep 2019	Online
London (UK)	LND_TR	TD-GC-FID	1 Jan 2007	1 Jan 2022	Online
Lyon – Feyzin stade (FR)	LYO1_IND	TD-GC-2FID PerkinElmer	20 Oct 2004	10 Feb 2023	Online
Lyon – Vernaison (FR)	LYO2_IND	TD-GC-2FID PerkinElmer	30 Mar 2009	10 Feb 2023	Online
Mouscron (BE)	MSR_IND	Passive samplers	2 Jan 2010	30 Dec 2021	Offline
Paris – SIRTa (FR)	PAR_SUB	PTR-Quad-MS IONICON	18 Jan 2020	30 Dec 2021	Online

specifically designed for environmental exposures (Phillips and Moya, 2013). The EPA's health risk assessment values (LCR and RfC, reference concentration) are widely utilized due to their systematic approach and extensive data basis for chronic exposure scenarios. While the EU's occupational exposure limits (OELs) are established for workplace environments, they primarily address acute exposure risks in occupational settings (Högberg and Järnberg, 2023). Therefore, they are not entirely suitable for assessing chronic, low-level exposure risks among the general public.

Additionally, the non-cancer health risks of human exposure to all identified BTEX compounds at the monitoring sites were assessed. The results are presented as the hazard index (HI), defined as the ratio of long-term intake to the reference dose for respiratory exposure. According to US EPA standards, an HI value equal to or greater than 1 indicates potential adverse health effects for the exposed population under current environmental conditions. The HI is calculated using the following formula: $HI = CA \times 1/RfC$, where RfC represents the reference concentration of VOC species. The RfC values for BTEX compounds are 30, 5000, 1000, and 100 ($\mu\text{g m}^{-3}$), respectively, as determined by the US EPA (EPA, 2020).

2.3 Data treatment

The data quality was assessed and evaluated by two ACTRIS CiGas (<https://www.actris.eu/topical-centre/cigas>, last access: 21 April 2024) units at IMT Nord Europe (France) and EMPA (Switzerland), based on ACTRIS recommendations and guidelines (Laj et al., 2024). Some sites indicated that they follow the ACTRIS guidelines especially for GC measurements (https://www.actris.eu/sites/default/files/inline-files/WP3_D3.17_M42_0.pdf, last access: 21 April 2024). However, the data evaluation showed that some sites had outliers, which were deleted when analyzed. Meanwhile, the International Union of Pure and Applied Chemistry strongly recommends using mixing ratios (Schwartz and Warneck, 1995). Therefore, in this study, BTEX levels are presented as median mixing ratios (MED) \pm median absolute deviation (MAD) in parts per trillion by volume (ppt) to reflect the non-normal distribution of the data. Statistically significant differences in BTEX mixing ratios at various monitoring sites and during different seasons were assessed using the Mann–Whitney *U* test for pairwise comparisons and the Kruskal–Wallis test for overall differences in medians (Kruskal and Wallis, 1952). These statistical analyses were performed using the SPSS software (IBM SPSS Statistics 25, Chicago, IL, USA).

3 Results and discussion

3.1 Status of BTEX data availability and mixing ratios in urban Europe

The results showed that, in urban areas, the relative abundance of BTEX followed this order: toluene > benzene > *m*, *p*-xylene > *o*-xylene > ethylbenzene with median mixing ratios of 266 ± 152 , 163 ± 74 , 129 ± 88 , 53 ± 35 , and 45 ± 27 ppt during the years 2017–2022, respectively (Fig. 1). Comparing different types of locations, the average and median mixing ratios of BTEX were higher at industry (IND) and traffic (TR) sites, while lower levels were found at urban background (UB) sites (Fig. S1 in the Supplement). These observed differences in BTEX concentrations suggest potential contributions from transportation and industrial activities in the studied urban areas. Therefore, to strengthen our conclusions, we applied urban enhancement ratios (ERs) (Salameh et al., 2019), estimating the slopes of least-squares linear regressions between each TEX compound and benzene. By using ERs, we reduce the sensitivity of the analysis to background conditions, dilution, and air mass mixing compared to using absolute concentrations (Salameh et al., 2019). Our results show spatial differences in the ER values for TEX/B, with the highest ratios observed at TR sites, followed by UB sites, and the lowest at IND sites. Specifically, the slopes were 2.09 ± 0.05 for T/B, 0.37 ± 0.01 for E/B, 1.21 ± 0.03 for *m*, *p*-X/B, and 0.48 ± 0.01 for *o*-X/B at TR sites; 1.57 ± 0.02 for T/B, 0.23 ± 0.00 for E/B, 0.71 ± 0.01 for *m*, *p*-X/B, and 0.27 ± 0.00 for *o*-X/B at UB sites; and 0.37 ± 0.01 for T/B, 0.13 ± 0.00 for E/B, 0.29 ± 0.01 for *m*, *p*-X/B, and 0.13 ± 0.00 for *o*-X/B at IND sites. A similar trend was observed in the seasonal variations, with ER values generally following the order TR > UB > IND. Notably, for UB and TR sites, the ER for TEX/B was higher in summer, while for IND sites, the ER was lowest during summer (Table S1 in the Supplement). These findings suggest that the additional evaporative sources, potentially related to traffic or solvent usage, particularly at urban background sites, may contribute to the observed seasonal and spatial variations.

3.2 Spatial variations of BTEX mixing ratios

Spatial variation in BTEX mixing ratios across urban Europe is significant. For instance, ATH_UB (Athens) had the highest median BTEX levels (2768 ± 4117 ppt) among all monitoring sites, followed by BCN_UB (Barcelona) (622 ± 312 ppt). This is primarily because these two sites are located in the eastern part of the Mediterranean basin, where the combination of a temperate climate (mild and rainy winters versus hot and dry summers) favors the development of severe air pollution episodes (Monks et al., 2009; Im and Kanakidou, 2012). Nonetheless, compared to the period before 2000, the levels of benzene and other BTEX compounds

have shown a decreasing trend due to the successful implementation of air quality measures in Greece, such as the extension of metro lines and the use of catalytic converters in cars (Panopoulou et al., 2021). For instance, benzene levels at traffic sites have decreased significantly, dropping by as much as 8-fold, from approximately 12 520 ppt in 1994 to about 1565 ppt in 2016 (Panopoulou et al., 2021). Similarly, at urban monitoring stations, benzene concentrations fell sharply, from around 4695 ppt during the period of 1993–1996 to between 313 and 1565 ppt in 2016 (Panopoulou et al., 2021).

However, compared to other European cities, pollution in Athens remains relatively severe, largely due to traffic but especially to wood-burning impact for heating (Panopoulou et al., 2018). For more details on the spatiotemporal variation and source apportionment of BTEX and other VOCs at this site, refer to Panopoulou et al. (2021).

3.3 Temporal variations of BTEX mixing ratios

Figure S2 depicts the temporal variations of BTEX mixing ratios at the different monitoring stations. The mixing ratio curves of BTEX species exhibited irregular fluctuations over the whole sampling period, but the mixing ratios of each component showed highly similar temporal variations. These results indicated a significant correlation among their concentrations and consequently suggested similar or identical sources of the BTEX species (Table S2). The highest correlation was observed between *o*-xylene and *m*-xylene and *p*-xylene at all monitoring stations ($r^2 = 0.62$ – 0.99), suggesting that they had similar emission sources (e.g., paints, finishing, fuels, and solvents). Conversely, the correlation between benzene and xylene exhibited some variability across different sites. In the majority of locations, the correlation falls within the range of $r^2 = 0.30$ to 0.59 . However, lower correlations ($r^2 < 0.20$) were observed at BAQS_UB (Birmingham) and LYO1_IND (Lyon – Feyzin stade), while stronger and significant correlations ($r^2 > 0.70$) were noted at ATH_UB (Athens), HEL_TR, (Helsinki) and LDS_UB (Lodelinsart). This variability may be attributed to their respective emission sources, atmospheric lifetimes, and differences in their photo-reactivity within distinct environments (Monod et al., 2001). Indeed, benzene has a rate constant (kOH) of reaction with hydroxyl radicals ($\bullet\text{OH}$) of $1.2 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}$ and an atmospheric lifetime of about 9.5 d. In contrast, xylene has a significantly higher kOH, ranging from $14 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}$ for *o*-xylene to $23.1 \times 10^{-12} \text{ cm}^3 \text{ molec.}^{-1} \text{ s}^{-1}$ for *m*-xylene, with a much shorter atmospheric lifetime between 10 and 20 h (Atkinson and Arey, 2003; Liu et al., 2023). These differing properties suggest that both photo-reactivity and source emissions crucially influence atmospheric behavior and concentrations.




Figure 1. Average mixing ratios of BTEX at 22 European sites (left panel) and the contribution of BTEX at each monitoring site (right panel) during 2017–2022 (but HEL_UB only during February 2016 and ATH_UB only 2016–2017). Please note that the BTEX observations at BCN_UB and PAR_SUB were not able to distinguish the mixing ratios of ethylbenzene separate from xylenes, and BAQS_UB did not include *m*-xylene and *p*-xylene data. ©OpenStreetMap contributors 2024. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.

3.3.1 Seasonal variation

A comparative analysis of seasonal variations (spring, March–May; summer, June–August; autumn, September–November; winter, December–February) in BTEX mixing ratios across different types (urban background (UB), traffic (TR), and industry (IND)) for the 21 urban sites was also performed (Fig. 2). The results indicate that benzene reached very similar median mixing ratios at UB, TR, and IND in all seasons, pointing to a mix of source contributions. However, for toluene the average and median mixing ratios followed the trend $TR \geq IND > UB$ for all seasons, further indicating a relatively greater influence of traffic sources on this species (Baudic et al., 2016; Salameh et al., 2019). For ethylbenzene and xylene (EX), averaged mixing ratios measured seem to be equivalent for TR and IND (and significantly lower at UB sites).

Figure S3 shows the BTEX mixing ratios by season for each site. Overall, BTEX mixing ratios at all monitoring sites showed substantial seasonal differences using the Mann–Whitney U test and the Kruskal–Wallis test ($p = 0.05$). The extent to which there were statistically significant differences in each BTEX concentration between seasons varied slightly across different monitoring sites (Fig. S3). Specifically, except for sites BAQS_UB (Birmingham), HEL_TR (Helsinki), and LY01_IND (Lyon – Feyzin stade), the BTEX mixing ratios were lower during spring and summer and higher during autumn and winter. This finding can be attributed to seasonal differences in photochemical reactions, atmospheric conditions, and the intensity of different emission sources. When analyzing meteorological data from selected monitoring sites (Fig. S4), it becomes evident that winter exhibits lower temperatures and mixing layer height, un-

favorable for pollutant dispersion. Conversely, summer features more conducive diffusion conditions, likely resulting in lower pollutant concentrations. Meanwhile, the increase in BTEX mixing ratios in winter may be influenced by the emissions from residential heating and road traffic emissions (Boynard et al., 2014; Panopoulou et al., 2018), and the lower spring and summer concentrations might be also linked to high photochemical oxidation (Hui et al., 2019). At BAQS_UB (Birmingham), benzene was lower in summer and higher in winter, while TEX (toluene, ethylbenzene, and xylene) did not vary significantly, suggesting that in combination urban gasoline and solvent emission rates have been stabilized (AQEG-EU, 2020). Due to only summer and autumn data being available, HEL_TR (Helsinki) exhibited higher mixing ratios for BTEX in summer compared to autumn, which can be attributed to increased evaporative emissions during the summer months. For LY01_IND (Lyon – Feyzin stade), benzene levels were lower in summer and higher in winter, while TEX levels were lower in spring and higher in summer. This indicates that during the summer, TEX mixing ratios can mainly be influenced by additional pollution sources, such as the volatilization of paints, finishing products, solvents, and fuel evaporation.

Additionally, seasonal variations in gasoline formulations could also play a significant role. For example, increasing ethanol content in gasoline can reduce emissions of BTEX (Yao et al., 2011; Karavalakis et al., 2014). The most common formulation in Europe is E5 (containing 5% ethanol), with ethanol concentrations generally higher in the summer than in the winter (Dunmore et al., 2016).




Figure 2. Seasonal variations of BTEX (benzene, toluene, ethylbenzene, xylene) mixing ratios for the different types sites (UB, urban background; TR, traffic; IND, industry). The box represents the 5th–95th percentiles of ratios. The middle line and middle square represent the median and mean values of ratios, respectively.

3.3.2 Diel Variation

Only online monitoring sites with hourly resolution BTEX data were considered (ATH_UB (Athens), BAQS_UB (Birmingham), BCN_UB (Barcelona), HEL_UB (Helsinki), LND_UB (London), MAR_UB (Marseille), PAR_UB (Paris – Paris 1er Les Halles), STB_UB (Strasbourg Ouest), ZUR_UB (Zurich), HEL_TR (Helsinki), LND_TR (London), LYO1_IND (Lyon – Feyzin stade), LYO2_IND (Lyon – Vernaison), and PAR_SUB (Paris – SIRTA)). Diel variations of BTEX (Fig. 3) indicated that at some sites, such as ATH_UB (Athens), BCN_UB (Barcelona), LND_UB (London), MAR_UB (Marseille), HEL_TR (Helsinki), LND_TR (London), and LYO1_IND (Lyon – Feyzin stade), double peaks vary in intensity, amplitude, and time of the maxima. Generally, the concentration of aromatic compounds is higher at night than during the day. The chemical removal process of BTEX mainly involves photochemical reactions with OH and NO₃ radicals (Atkinson and Arey, 2003; Carter, 2007). •OH radicals are mainly present during the daytime, and their reaction rates with VOCs are faster than the NO₃

radicals present at night (Zou et al., 2015). Additionally, the higher mixing layer height during the day aids in diluting pollutants, while nighttime emissions from activities like firewood for leisure and winter heating result in higher BTEX concentrations at night compared to daytime levels (Wu et al., 2016; Liu et al., 2020).

The first of the two major diel BTEX peaks coincides with the morning road traffic rush hours. Particularly, BTEX mixing ratios in ATH_UB (Athens) and LYO1_IND (Lyon – Feyzin stade) are significantly ($p = 0.001$) higher than at the other sites, which may be closely related to industrial sources, off-road transport, types of vehicles, traffic flow, and other combustion-related sources. Similarly, influenced by traffic, the peaks of BTEX in LND_TR (London) were also more pronounced. After 17:00 LT, when people mainly leave from work, traffic increases, and the mixing layer height decreases. A similar phenomenon was observed in HEL_TR (Helsinki), although this site only provided data for 1 month in 2019 (Table 1). Consequently, an evening peak in BTEX levels is regularly observed between approximately 18:00–19:00 at these sites (Fig. 3). However, LYO1_IND (Lyon – Feyzin stade) is hardly affected by the evening rush hour. This is primarily due to the site’s exposure to industrial processes, as well as wind speed and direction, particularly when the monitoring site is downwind of the industrial area. The highest levels are observed at 23:00. These phenomena further indicate that BTEX are largely associated with anthropogenic activities and specific meteorological factors. Furthermore, in addition to traffic emissions, evaporative sources contribute significantly to the diel patterns observed in some of the BTEX. Given that evaporative emissions typically rise in the afternoon as temperatures increase (Nguyen et al., 2009), any elevation in TEX mixing ratios may be linked to their evaporative sources (Yurdakul et al., 2018). It is striking how similar the levels and variability of toluene are in BAQS_UB (Birmingham), HEL_UB (Helsinki), and PAR_SUB (Paris – SIRTA), likely reflecting similar traffic influences. In contrast, benzene levels show more variation, indicating regional influences. Focusing on Paris, similar to the findings of Simon et al. (2023), we observe comparable levels and variability of benzene between PAR_UB and PAR_SUB (Paris – SIRTA), suggesting regional influence. However, for toluene, levels are higher in PAR_UB (Paris – Paris 1er Les Halles) compared to PAR_SUB (Paris – SIRTA), indicating more local traffic influences in Paris.

Similarly, the analysis of the seasonal and weekly diel average variations of BTEX at 14 online monitoring sites (Figs. S5 and S6) showed patterns similar to the overall diel trend. Specifically, certain monitoring sites (i.e., ATH_UB (Athens), BCN_UB (Barcelona), LND_UB (London), MAR_UB (Marseille), PAR_UB (Paris – Paris 1er Les Halles), ZUR_UB (Zurich), HEL_TR (Helsinki), LND_TR (London), LYO1_IND (Lyon – Feyzin stade)) exhibited prominent morning and evening peaks during weekdays (Monday to Friday). This phenomenon is attributed to in-




Figure 3. Diel variations of hourly average BTEX mixing ratios for 14 of the 22 studied sites, presented in local time over the entire sampling period for each site. Please note that the two vertical axes of each compound span different ranges of mixing ratios.

creased emissions from vehicle exhaust, as suggested by measurements in tunnels (Ammoura et al., 2014). However, during weekends (Saturday and Sunday), particularly on Sunday mornings, there were either no peaks or much weaker ones due to lower traffic density. Compared to the corresponding peaks on weekdays, the peaks on Saturday and Sunday evenings ($\sim 22:00\text{--}24:00$) were delayed. This delay is linked to the fact that people tend to stay at home during weekends, engaging in leisure activities (thus leading to higher emissions from cooking and household heating), and/or go out socially in the evening. These phenomena are similar to our research on levels of lung-deposited surface area (LDSA) across various monitoring sites in Europe (Liu et al., 2023) and the diel variations of black carbon across different seasons in Augsburg (Liu et al., 2022).

3.4 Specific ratios of BTEX

3.4.1 Benzene/toluene (B/T)

Benzene and toluene have been found to be relatively more stable than xylenes, having lifetimes of 9.5 d, 2.1 d, and 7.8 h, respectively (Atkinson and Arey, 2003; Liu et al., 2008). Therefore, B/T remains relatively constant close to emission sources. However, this ratio can change over time and with the atmospheric processing of air masses (Seco et al.,




Figure 4. The ratios of benzene to toluene (B/T) and *meta*-xylene to ethylbenzene (X/E) at all sites (a1, b1) and different types of sites (UB, urban background; TR, traffic; IND, industry; a2, b2). The box represents the 5th–95th percentiles of ratios. The middle line and middle square represent the median and mean values of ratios, respectively. Blank means no available value.

2013). B/T values around 0.5 suggest significant influences from vehicle exhaust, while values significantly higher than 0.5 indicate that other sources such as industry, coal combustion, and biomass burning are major sources of BTEX species (Barletta et al., 2005; Baudic et al., 2016; Salameh et al., 2019). Additionally, higher B/T ratios can also be attributed to the fact that emissions and/or air masses further away from the source are more aged, leading to relatively more toluene disappearance due to its higher reactivity compared to benzene.

As shown in Fig. 4a-1, the B/T ratio ranges from 0.29 ± 0.11 to 1.35 ± 0.95 , with the majority of sites averaging around 0.4–0.6 (BCN_UB (Barcelona), CHM_UB (Charleroi), GRE_UB (Grenoble), HET_UB (Herstal), MAR_UB (Marseille), PAR_UB (Paris – Paris 1er Les Halles), ZUR_UB (Zurich), LND_TR (London), and MSR_IND (Mouscron)). Similarly, Salameh et al. (2019) found that at UB sites with weekly measurements, B/T was on the same order of magnitude as the TR sites with an average ratio of 0.34 in Paris. This finding was indirectly confirmed by the diel variation of BTEX mixing ratios (Fig. 3) and the B/T trends in different site typologies (TR: 0.41 ± 0.29 ; UB: 0.73 ± 0.68 ; IND: 1.03 ± 1.00 , Fig. 4a-1). Notably, the B/T ratio is higher at industrial (IND) sites, which can be attributed to the different atmospheric lifetimes of toluene and benzene. Although toluene is commonly emitted in greater quantities from industrial sources, its atmospheric lifetime is much shorter than that of benzene (toluene: 2.1 d, benzene: 9.5 d). As a result, even though toluene emissions may be significant, the shorter lifetime of toluene leads to its rapid degradation in the atmosphere compared to benzene. This allows benzene to accumulate relative to toluene, particularly near industrial sources, resulting in a




Figure 5. Regression of benzene to toluene (a) and *m*-xylene and *p*-xylene to ethylbenzene (b) ratios in different urban locations (UB, urban background; TR, traffic; IND, industry) across different seasons (green, spring; red, summer; gold, autumn; gray, winter).

higher B/T ratio despite toluene's greater initial emissions (Atkinson and Arey, 2003; Liu et al., 2008)

In addition, based on only summer and autumn data being available (Table 1), the B/T at HEL_TR (Helsinki), serving as a traffic site, was 0.29 ± 0.11 , suggesting vehicular exhaust emissions as the primary source of benzene and toluene in the study area's ambient air, primarily through fuel evaporative emissions. At HEL_UB (Helsinki), the B/T ratio is relatively high, which can be attributed to the limited monitoring data available only from February 2016, introducing a potential comparability bias. For PAR_SUB (Paris – SIRTa), the elevated B/T ratio can be attributed to seasonal factors (Simon et al., 2023). Benzene concentrations typically increase from September to April, driven by more active sources during the winter months, such as residential wood burning (Languille et al., 2020), and limited dispersion due to a lower boundary layer (Simon et al., 2023). This trend is further supported by the presence of wood-burning tracers like furfural and benzenediol, which exhibit similar seasonal patterns. Although toluene does not show as strong a seasonal variation as benzene, it also has higher levels during autumn and winter. The primary source of toluene, traffic, remains important throughout the year, and more stagnant conditions in these seasons contribute to the accumulation of pollutants.

Further, considering the seasonal B/T ratios in different urban environments (Fig. 5a), each monitoring site (Fig. S7), and temporal trends (Fig. S8) in B/T across monitoring sites, the seasonal fluctuations of benzene and toluene are consistent with B/T, showing seasonal differences to a certain extent. To analyze these variations more accurately, major axis regression (MA) was applied, accounting for uncertainties in both the *x* and *y* variables. Specifically, the mean and median

values of B/T across all monitoring sites exhibit lower values in summer and higher values in winter. This pattern is consistent across different types of environments. The higher B/T values in winter further indicate influences from biomass combustion sources.

3.4.2 *m*-xylene–*p*-xylene / ethylbenzene (X/E)

In the atmosphere, *m*-xylene and *p*-xylene and ethylbenzene are homologous, and the reaction rate of *m*-xylene and *p*-xylene with the OH radical is 3 times faster than that of ethylbenzene (Martins et al., 2008; Han et al., 2017). Hence, the X/E ratio can serve as an indicator reflecting the extent of photochemical reactions (McKeen and Liu, 1993), where higher and lower X/E ratios suggest local sources and external transport, respectively (Zalé and Broday, 2008).

Typical X/E ratios measured in the urban atmosphere are around 3 (Monod et al., 2001; Hsieh et al., 2011; Amodio et al., 2013). Wang et al. (2015) observed significant variations in X/E ratios (ranging from 1.2 to 2.8) in tunnels characterized by different types of vehicles in downtown Shanghai. In this study, the mean X/E ratio across monitoring sites ranged from 1.75 ± 0.91 to 3.68 ± 0.30 (Fig. 4b-1), with X/E values oscillating around 3 at all sites. Additionally, there were minimal differences in the average X/E values across seasons (Fig. 5b and S9) and different types of sites (Fig. 4b-2). This result suggests that pollutants at the monitoring sites are primarily emitted directly from local sources (with higher X/E values). Specifically, considering the temporal variations of X/E at each monitoring site, except for the monitoring period at HEL_UB (Helsinki) in 2016, this ratio was generally greater than 2 at the other sites (Fig. S7). This result further suggests that direct emissions from local sources dominate at each moni-

toring site. Notably, in general, the slightly lower X/E value observed at the urban background (UB) sites (i.e., ANG_UB (Angleur), CHM_UB (Charleroi), GRE_UB (Grenoble), HET_UB (Helsinki), MON_UB (Mons), NAM_UB (Nanur), LDS_UB (Lodelinsart), and ZUR_UB (Zurich)) in summer and the higher X/E value observed at the traffic (TR) site (i.e., LND_TR (London)) in winter can be attributed to seasonal differences in photochemical reactions and atmospheric conditions, the impact of changing gasoline formulations, and the significant increase in BTEX concentrations during winter cold starts (George et al., 2015; Zhang et al., 2022) (Fig. 5b). In summer, increased solar radiation and higher temperatures promote photochemical reactions (Rad et al., 2014), and since the reaction rate of *m*-xylene and *p*-xylene with the OH radical is 3 times faster than that of ethylbenzene, the X/E ratio decreases. Conversely, in winter, the formation of a stable atmospheric layer and lower temperatures may cause pollutants to remain in the atmosphere for longer periods, leading to an increased X/E ratio.

3.5 Health risk assessment of BTEX

Figure 6a presents LCR values associated with benzene and ethylbenzene exposure through inhalation at all monitoring sites. Across all sites, the average LCR values range from 2.6×10^{-6} to 1.9×10^{-5} , showing that all values are below the definite risk of cancer threshold (10^{-4}) but exceed the threshold established by the US EPA (2001) guidelines of 1×10^{-6} . This result indicates that the risk from benzene and ethylbenzene is at a moderate level and still warrants attention, especially in traffic (TR) and industry (IND) environments. It should be noted that the EPA values are based on the assumption of continuous exposure, which typically does not occur outdoors, where individuals spend less time compared to indoors. Thus, our findings apply primarily to outdoor exposure, and indoor air concentrations of BTEX compounds would need to be considered to draw comprehensive health-related conclusions.

The variability in LCR values across different sites highlights the importance of localized assessments and tailored interventions. Urban areas, for example, might need different pollution control strategies due to differences in pollutant sources and population density. Industrial and traffic-related areas often have higher concentrations of these pollutants due to emissions from vehicles and industrial activities. Therefore, populations in these areas might be at greater risk and require more stringent monitoring and regulatory measures to mitigate exposure.

His all fall below the threshold limit value (1) (Fig. 6b) set by the US Environmental Protection Agency in 2009 (EPA, 2009). This indicates a generally low non-carcinogenic risk from outdoor exposure to BTEX in the region, with levels mostly within safe thresholds. Therefore, there is no immediate risk of developing non-cancer diseases due to the inhalation of BTEX at the observed levels outdoors. How-




Figure 6. The lifetime cancer risk (LCR) values for the carcinogenic risk of benzene and ethylbenzene and the hazard index (HI) for the carcinogenic risk of benzene, toluene, ethylbenzene, *m*-xylene and *p*-xylene, and *o*-xylene through inhalation for individuals at all sites. Please note that the LCR values of BCN_UB and PAR_SUB have only been calculated with benzene data (marked gray), and HI values of BAQS_UB did not detect *m*-xylene or *p*-xylene.

ever, it is important to note that long-term exposure, even within these safe limits, can still adversely affect health. Notably, sites LYO1_IND (Lyon – Feyzin stade) and LND_TR (London) exhibit higher HI values compared to others, attributed to elevated concentrations of benzene. This observation aligns with the findings reported by Jia et al. (2021) regarding VOC HI values in the Chinese delta region and by Bretón et al. (2022) regarding BTEX HI values in southeastern Mexico.

Furthermore, the PTR-MS systems used at BCN_UB (Barcelona) and PAR_SUB (Paris – SIRTa) were not able to distinguish the mixing ratios of ethylbenzene separate from xylenes. This limitation affects the comparability of LCR values across different monitoring sites, as the absence of ethylbenzene data at these sites results in an incomplete assessment of the total carcinogenic risk. Despite these limitations, the comparisons still offer valuable insights into the spatial distribution of cancer risks and highlight areas where more comprehensive monitoring is needed.

To enhance the reliability and comprehensiveness of future assessments, it is crucial to establish a standardized observation platform. Such a platform would ensure consistent data collection methods, pollutant measurement techniques, and risk assessment criteria across all monitoring sites. This would enable better-informed decision-making for public health interventions specifically focused on outdoor exposures.

4 Conclusions

This study conducted a comprehensive assessment of long- and short-term mixing ratios of benzene, toluene, ethylbenzene, *m*-xylene and *p*-xylene, and *o*-xylene (BTEX)

across 22 monitoring sites in 7 European countries (Belgium, France, Finland, Greece, Spain, Switzerland, and the United Kingdom.). These sites included 3 industrial (IND) sites, 2 traffic (TR) sites, 16 urban background (UB) sites, and 1 suburban background (SUB) site. The median of BTEX mixing ratios was 772 ppt across all monitoring sites. Our findings demonstrate significant seasonal and diel variabilities in BTEX mixing ratios ($p = 0.01$) at each monitoring site, indicating the influence of changes in traffic volume, emission sources, photo-reactivity, and meteorological factors. The B/T ratio ranged from 0.29 ± 0.11 to 1.35 ± 0.95 , and the X/E ratio ranged from 1.75 ± 0.91 to 3.68 ± 0.30 , demonstrating spatial variations in BTEX emission sources across monitoring sites. These results highlight that local traffic emissions are the major source of BTEX pollution, with additional contributions from industrial processes, solvent use, coatings, biomass burning, and fuel evaporation. Furthermore, the health index (HI) values of BTEX at monitoring sites were generally lower than the threshold limit value, suggesting a low non-carcinogenic risk overall. However, our health risk assessment indicates that the lifetime cancer risk (LCR) from inhaling benzene and ethylbenzene, although relatively low ($< 10^{-4}$), still warrants attention due to the potential effects of long-term exposure.

In conclusion, this study provides new insights into the temporal and spatial variabilities of BTEX mixing ratios across different types of sites in multiple European countries. It emphasizes the significant impact of traffic-related emissions and seasonal factors on BTEX levels and underscores the need for continued monitoring and regulation to mitigate potential health risks. While we recognize that transportation and industrial activities are key contributors to BTEX pollution, our current data do not allow us to quantify their individual contributions. The findings also suggest that future studies should focus on the impact of changing gasoline formulations and vehicle technologies on BTEX emissions, as well as the effectiveness of current air quality regulations in reducing BTEX exposure. However, our primary aim is to provide a comprehensive assessment of the health risks associated with BTEX exposure, highlighting the urgent need for effective management strategies.

Data availability. The dataset for this paper can be accessed at <https://data.mendeley.com/drafts/n8hggjyjs2> (Liu, 2024). VOC data were assessed and evaluated by two ACTRIS CiGas (<https://www.actris.eu/topical-centre/cigas>, last access: 21 April 2024) units at IMT Nord Europe in France and EMPA in Switzerland based on ACTRIS recommendations and guidelines.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-25-625-2025-supplement>.

Author contributions. XL: writing (original draft and review and editing), conceptualization. XZ and TW: methodology, formal analysis. LW: software. MM: project administration. MD, RL, RS, AMY, MG, PP, AC, MV, YF, AB, VG, GG, LL, VG, LS, HH, SR, ZB, MJM, DB, SH, ZD, RMH, WB, JD, SS, AL, JH, EL, and NM: data curation, writing (review and editing). XD, TA, XZ, PKH, AA, and TS: writing (review and editing). XQ: writing (review and editing), supervision, project administration, funding acquisition.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Atmospheric Chemistry and Physics*. The peer-review process was guided by an independent editor. In addition, a referee published with an author during the past 3 years.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. The measurements at the SIRTa station have benefited from the support of the research infrastructure ACTRIS-FR, registered on the Roadmap of the French Ministry of Research. Nikolaos Mihalopoulos and Eleni Liakakou acknowledge Anastasia Panopoulou for measurements and data curation. We would also like to thank the Swiss Federal Office for the Environment (FOEN). Marvin Dufresne, Stéphane Sauvage, and Thérèse Salameh acknowledge the French Agency for Ecological Transition (ADEME) and the Mines-Telecom Institute Nord625 Europe (IMT Nord-Europe) for funding the PhD of Marvin Dufresne and acknowledge the INSU and the MISTRALS program for the financial support of the Marseille campaign.

Financial support. This research has been supported by the RI-URBANS project (Research Infrastructures Services Reinforcing Air Quality Monitoring Capacities in European Urban and Industrial Areas, European Union's Horizon 2020 research and innovation program, Green Deal, European Commission, contract 101036245). It has also received funding from the National Natural Science Foundation of China (grant nos. 42407566, 42205099), the Chunhui Project Foundation of the Education Department of China (grant no. HZKY20220053), and the Natural Science Foundation of Xinjiang Uygur Autonomous Region (grant no. 2023D01A57). Additional support was provided by the Agencia Estatal de Investigación of the Spanish Ministry of Science, including the Ramón y Cajal grant (grant nos. RYC2021-032519-I and RYC2020-029216-I, funded by MICIU/AEI/10.13039/501100011033 and "ESF Investing in your future"), the Consolidación Investigadora project (grant no. CNS2022-135757), the Severo Ochoa Centre of Research Excellence (MCIN/AEI, Project CEX2018-000794-S), and the H2020 ACTRIS-2 project (grant no. 654109).

Review statement. This paper was edited by Eva Y. Pfannerstill and reviewed by two anonymous referees.

References

- Actris: ACTRIS Centre for Reactive Trace Gases In Situ Measurements (CiGas), <https://www.actris.eu/topical-centre/cigas> (last access: 21 April 2024), 2024.
- Ammoura, L., Xueref-Remy, I., Gros, V., Baudic, A., Bonsang, B., Petit, J.-E., Perrussel, O., Bonnaire, N., Sciare, J., and Chevallier, F.: Atmospheric measurements of ratios between CO₂ and co-emitted species from traffic: a tunnel study in the Paris megacity, *Atmos. Chem. Phys.*, 14, 12871–12882, <https://doi.org/10.5194/acp-14-12871-2014>, 2014.
- Amodio, M., de Gennaro, G., Marzocca, A., Trizio, L., and Tutino, M.: Assessment of impacts produced by anthropogenic sources in a little city near an important industrial area (Modugno, Southern Italy), *The Sci. World J.*, 2013, 150397, <https://doi.org/10.1155/2013/150397>, 2013.
- AQEG-EU: Non-methane Volatile Organic Compounds in the UK. Department for Environment, Food and Rural Affairs; Scottish Government; Welsh Government; and Department of Agriculture, Environment and Rural Affairs in Northern Ireland, <https://uk-air.defra.gov.uk> (last access: 21 April 2024), 2020.
- Atkinson, R. and Arey, J.: Atmospheric Degradation of Volatile Organic Compounds, *Chem. Rev.*, 103, 4605–4638, <https://doi.org/10.1021/cr0206420>, 2003.
- Barletta, B., Meinardi, S., Rowland, F. S., Chan, C., Wang, X., Zou, S., Chan, L. Y., and Blake, D. R.: Volatile organic compounds in 43 Chinese cities, *Atmos. Environ.*, 39, 5979–5990, <https://doi.org/10.1016/j.atmosenv.2005.06.029>, 2005.
- Baudic, A., Gros, V., Sauvage, S., Locoge, N., Sanchez, O., Sarda-Estève, R., Kalogridis, C., Petit, J.-E., Bonnaire, N., Baisnée, D., Favez, O., Albinet, A., Sciare, J., and Bonsang, B.: Seasonal variability and source apportionment of volatile organic compounds (VOCs) in the Paris megacity (France), *Atmos. Chem. Phys.*, 16, 11961–11989, <https://doi.org/10.5194/acp-16-11961-2016>, 2016.
- Bolden, A. L., Kwiatkowski, C. F., and Colborn, T.: New look at BTEX: are ambient levels a problem?, *Environ. Sci. Technol.*, 49, 5261–5276, <https://pubs.acs.org/doi/10.1021/es505316f>, 2015.
- Borbon, A., Boynard, A., Salameh, T., Baudic, A., Gros, V., Gauduin, J., Perrussel, O., and Pallares, C.: Is traffic still an important emitter of monoaromatic organic compounds in European urban areas?, *Environ. Sci. Technol.*, 52, 513–521, <https://doi.org/10.1021/acs.est.7b01408>, 2018.
- Boynard, A., Borbon, A., Leonardis, T., Barletta, B., Meinardi, S., Blake, D. R., and Locoge, N.: Spatial and seasonal variability of measured anthropogenic non-methane hydrocarbons in urban atmospheres: Implication on emission ratios, *Atmos. Environ.* (1994), 82, 258–267, [10.1016/j.atmosenv.2013.09.039](https://doi.org/10.1016/j.atmosenv.2013.09.039), 2014.
- Bretón, R. M. C., Bretón, J. G. C., Kahl, J. W., Chi, M. P. U., Lozada, S. E. C., de la Luz Espinosa Fuentes, M., Martínez, R. G., and Del Carmen Lara Severino, R.: Seasonal and diurnal variations of btex in ambient air from a site impacted by the oil industry in southeast Mexico, *Bull. Environ. Contam. Toxicol.*, 108, 212–218, <https://doi.org/10.1007/s00128-021-03379-1>, 2022.
- Carter, W. P. L.: A detailed mechanism for the gas-phase atmospheric reactions of organic compounds, *Atmos. Environ.* (1994), 41, 80–117, <https://doi.org/10.1016/j.atmosenv.2007.10.061>, 2007.
- Davidson, C. J., Hannigan, J. H., and Bowen, S. E.: Effects of inhaled combined Benzene, Toluene, Ethylbenzene, and Xylenes (BTEX): Toward an environmental exposure model, *Environ. Toxicol. Pharmacol.*, 81, 103518, <https://doi.org/10.1016/j.etap.2020.103518>, 2021.
- Derwent, R. G., Jenkin, M. E., Utembe, S. R., Shallcross, D. E., Murrells, T. P., and Passant, N. R.: Secondary organic aerosol formation from a large number of reactive man-made organic compounds, *Sci. Total Environ.*, 408, 3374–3381, <https://doi.org/10.1016/j.scitotenv.2010.04.013>, 2010.
- DIRECTIVE(EU): DIRECTIVE(EU) 2024/2881 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 23 October 2024 on ambient air quality and cleaner air for Europe (recast), 2024, Official Journal of the European Union, https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=OJ:L_202402881#tit_1 (last access: 21 November 2024), 2024.
- Dunmore, R. E., Whalley, L. K., Sherwen, T., Evans, M. J., Heard, D. E., Hopkins, J. R., Lee, J. D., Lewis, A. C., Lidster, R. T., and Rickard, A. R.: Atmospheric ethanol in London and the potential impacts of future fuel formulations, *Faraday Discuss.*, 189, 105–120, <https://doi.org/10.1039/C5FD00190K>, 2016.
- Durmusoglu, E., Taspinar, F., and Karademir, A.: Health risk assessment of BTEX emissions in the landfill environment, *J. Hazard. Mater.*, 176, 870–877, <https://doi.org/10.1016/j.jhazmat.2009.11.117>, 2010.
- EPA, U. S.: Assessment, Probabilistic Risk, “Risk Assessment guidance for superfund: volume III-part a”, Process for Conducting Probabilistic Risk Assessment, United States Environmental Protection Agency Washington; Washington, D.C., United States, 2001.
- Garg, A. and Gupta, N. C.: A comprehensive study on spatio-temporal distribution, health risk assessment and ozone formation potential of BTEX emissions in ambient air of Delhi, India, *Sci. Total Environ.*, 659, 1090–1099, <https://doi.org/10.1016/j.scitotenv.2018.12.426>, 2019.
- George, I. J., Hays, M. D., Herrington, J. S., Preston, W., Snow, R., Faircloth, J., George, B. J., Long, T., and Baldauf, R. W.: Effects of cold temperature and ethanol content on VOC emissions from light-duty gasoline vehicles, *Environ. Sci. Technol.*, 49, 13067–13074, <https://doi.org/10.1021/acs.est.5b04102>, 2015.
- Ghaffari, H. R., Kamari, Z., Hassanvand, M. S., Fazlzadeh, M., and Heidari, M.: Level of air BTEX in urban, rural and industrial regions of Bandar Abbas, Iran; indoor-outdoor relationships and probabilistic health risk assessment, *Environ. Res.*, 200, 111745, <https://doi.org/10.1016/j.envres.2021.111745>, 2021.
- Grosjean, D. and Seinfeld, J. H.: Parameterization of the formation potential of secondary organic aerosols, *Atmos. Environ.* (1967), 23, 1733–1747, [https://doi.org/10.1016/0004-6981\(89\)90058-9](https://doi.org/10.1016/0004-6981(89)90058-9), 1989.
- Han, D., Wang, Z., Cheng, J., Wang, Q., Chen, X., and Wang, H.: Volatile organic compounds (VOCs) during non-haze and haze days in Shanghai: characterization and secondary organic aerosol (SOA) formation, *Environ. Sci. Pollut. Res.*, 24, 18619–18629, <https://doi.org/10.1007/s11356-017-9433-3>, 2017.

- Högberg, J. and Järnberg, J.: Approaches for the setting of occupational exposure limits (OELs) for carcinogens, *Crit. Rev. Toxicol.*, 53, 131–167, 2023.
- Hsieh, L., Wang, Y., Yang, H., and Mi, H.: Measurements and Correlations of MTBE and BTEX in Traffic Tunnels, *Aerosol Air Qual. Res.*, 11, 763–775, <https://doi.org/10.4209/aaqr.2011.03.0035>, 2011.
- Hui, L., Liu, X., Tan, Q., Feng, M., An, J., Qu, Y., Zhang, Y., and Cheng, N.: VOC characteristics, sources and contributions to SOA formation during haze events in Wuhan, Central China, *Sci. Total Environ.*, 650, 2624–2639, <https://doi.org/10.1016/j.scitotenv.2018.10.029>, 2019.
- Im, U. and Kanakidou, M.: Impacts of East Mediterranean megacity emissions on air quality, *Atmos. Chem. Phys.*, 12, 6335–6355, <https://doi.org/10.5194/acp-12-6335-2012>, 2012.
- IRIS (United States Environmental Protection Agency Integrated Risk Information System), 2020. IRIS assessments.: https://cfpub.epa.gov/ncea/iris_drafts/atoz.cfm?list_type=alpha, (last access: 21 April 2024), 2020.
- Jia, H., Gao, S., Duan, Y., Fu, Q., Che, X., Xu, H., Wang, Z., and Cheng, J.: Investigation of health risk assessment and odor pollution of volatile organic compounds from industrial activities in the Yangtze River Delta region, China, *Ecotox. Environ. Safe.*, 208, 111474, <https://doi.org/10.1016/j.ecoenv.2020.111474>, 2021.
- Karavalakis, G., Short, D., Russell, R. L., Jung, H., Johnson, K. C., Asa-Awuku, A., and Durbin, T. D.: Assessing the impacts of ethanol and isobutanol on gaseous and particulate emissions from flexible fuel vehicles, *Environ. Sci. Technol.*, 48, 14016–14024, <https://doi.org/10.1021/es5034316>, 2014.
- Kruskal, W. H. and Wallis, W. A.: Use of Ranks in One-Criterion Variance Analysis, *J. Am. Stat. Assoc.*, 47, 583–621, <https://doi.org/10.1080/01621459.1952.10483441>, 1952.
- Laj, P., Lund Myhre, C., Riffault, V., Amiridis, V., Fuchs, H., Eleftheriadis, K., Petäjä, T., Salameh, T., Kivekäs, N., and Juurola, E.: Aerosol, Clouds and Trace Gases Research Infrastructure–ACTRIS, the European research infrastructure supporting atmospheric science, *B. Am. Meteorol. Soc.*, 105, E1098–E1136, <https://doi.org/10.1175/BAMS-D-23-0064.1>, 2024.
- Languille, B., Gros, V., Petit, J., Honoré, C., Baudic, A., Perrussel, O., Foret, G., Michoud, V., Truong, F., and Bonnaire, N.: Wood burning: A major source of Volatile Organic Compounds during wintertime in the Paris region, *Sci. Total Environ.*, 711, 135055, <https://doi.org/10.1016/j.scitotenv.2019.135055>, 2020.
- Li, A. J., Pal, V. K., and Kannan, K.: A review of environmental occurrence, toxicity, biotransformation and biomonitoring of volatile organic compounds, *Environ. Chem. Ecotoxicol.*, 3, 91–116, <https://doi.org/10.1016/j.enceco.2021.01.001>, 2021.
- Li, L., Li, H., Zhang, X., Wang, L., Xu, L., Wang, X., Yu, Y., Zhang, Y., and Cao, G.: Pollution characteristics and health risk assessment of benzene homologues in ambient air in the northeastern urban area of Beijing, China, *J. Environ. Sci.*, 26, 214–223, [https://doi.org/10.1016/S1001-0742\(13\)60400-3](https://doi.org/10.1016/S1001-0742(13)60400-3), 2014.
- Liu, B., Yang, Y., Yang, T., Dai, Q., Zhang, Y., Feng, Y., and Hopke, P. K.: Effect of photochemical losses of ambient volatile organic compounds on their source apportionment, *Environ. Int.*, 172, 107766, [10.1016/j.envint.2023.107766](https://doi.org/10.1016/j.envint.2023.107766), 2023.
- Liu, P. G., Yao, Y., Tsai, J., Hsu, Y., Chang, L., and Chang, K.: Source impacts by volatile organic compounds in an industrial city of southern Taiwan, *Sci. Total Environ.*, 398, 154–163, <https://doi.org/10.1016/j.scitotenv.2008.02.053>, 2008.
- Liu, X.: Exploring the variations in ambient BTEX in urban Europe and its environmental health implications, *Mendeley Data*, V1 [data set], <https://doi.org/10.17632/n8hggjys2.1>, 2024.
- Liu, X., Schnelle-Kreis, J., Zhang, X., Bendl, J., Khedr, M., Jakobi, G., Schloter-Hai, B., Hovorka, J., and Zimmermann, R.: Integration of air pollution data collected by mobile measurement to derive a preliminary spatiotemporal air pollution profile from two neighboring German–Czech border villages, *Sci. Total Environ.*, 722, 137632, <https://doi.org/10.1016/j.scitotenv.2020.137632>, 2020.
- Liu, X., Hadiatullah, H., Schnelle-Kreis, J., Xu, Y., Yue, M., Zhang, X., Querol, X., Cao, X., Bendl, J., and Cyrus, J.: Levels and drivers of urban black carbon and health risk assessment during pre-and COVID19 lockdown in Augsburg, Germany, *Environ. Pollut.*, 120529, 316, <https://doi.org/10.1016/j.envpol.2022.120529>, 2022.
- Liu, X., Hadiatullah, H., Zhang, X., Trechera, P., Savadkoobi, M., Garcia-Marlès, M., Reche, C., Pérez, N., Beddows, D. C., and Salma, I.: Ambient air particulate total lung deposited surface area (LDSA) levels in urban Europe, *Sci. Total Environ.*, 165466, 898, <https://doi.org/10.1016/j.scitotenv.2023.165466>, 2023.
- Martins, L., Vieira, K. M., Rios, L. M., and Cardoso, D.: Basic catalyzed Knoevenagel condensation by FAU zeolites exchanged with alkylammonium cations, *Catal. Today*, 133, 706–710, <https://doi.org/10.1016/j.cattod.2007.12.043>, 2008.
- McKeen, S. A. and Liu, S. C.: Hydrocarbon ratios and photochemical history of air masses, *Geophys. Res. Lett.*, 20, 2363–2366, <https://doi.org/10.1029/93GL02527>, 1993.
- Miri, M., Shendi, M. R. A., Ghaffari, H. R., Aval, H. E., Ahmadi, E., Taban, E., Gholizadeh, A., Aval, M. Y., Mohammadi, A., and Azari, A.: Investigation of outdoor BTEX: Concentration, variations, sources, spatial distribution, and risk assessment, *Chemosphere*, 163, 601–609, <https://doi.org/10.1016/j.chemosphere.2016.07.088>, 2016.
- Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Akimoto, H., Amann, M., Baklanov, A., Baltensperger, U., and Bey, I.: Atmospheric composition change–global and regional air quality, *Atmos. Environ.*, 43, 5268–5350, <https://doi.org/10.1016/j.atmosenv.2009.08.021>, 2009.
- Monod, A., Sive, B. C., Avino, P., Chen, T., Blake, D. R., and Rowland, F. S.: Monoaromatic compounds in ambient air of various cities: a focus on correlations between the xylenes and ethylbenzene, *Atmos. Environ.*, 35, 135–149, [https://doi.org/10.1016/S1352-2310\(00\)00274-0](https://doi.org/10.1016/S1352-2310(00)00274-0), 2001.
- Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmos. Chem. Phys.*, 7, 3909–3922, <https://doi.org/10.5194/acp-7-3909-2007>, 2007.
- Nguyen, H. T., Kim, K., and Kim, M.: Volatile organic compounds at an urban monitoring station in Korea, *J. Hazard. Mater.*, 161, 163–174, <https://doi.org/10.1016/j.jhazmat.2008.03.066>, 2009.
- Nie, E., Zheng, G., and Ma, C.: Characterization of odorous pollution and health risk assessment of volatile organic compound emissions in swine facilities, *Atmos. Environ.* (1994), 223, 117233, [10.1016/j.atmosenv.2019.117233](https://doi.org/10.1016/j.atmosenv.2019.117233), 2020.
- Ogbodo, J. O., Arazu, A. V., Iguh, T. C., Onwodi, N. J., and Ezike, T. C.: Volatile organic compounds: A proinflammatory

- activator in autoimmune diseases, *Front. Immunol.*, 13, 928379, <https://doi.org/10.3389/fimmu.2022.928379>, 2022.
- Panopoulou, A., Liakakou, E., Gros, V., Sauvage, S., Locoge, N., Bonsang, B., Psiloglou, B. E., Gerasopoulos, E., and Mihalopoulos, N.: Non-methane hydrocarbon variability in Athens during wintertime: the role of traffic and heating, *Atmos. Chem. Phys.*, 18, 16139–16154, <https://doi.org/10.5194/acp-18-16139-2018>, 2018.
- Panopoulou, A., Liakakou, E., Sauvage, S., Gros, V., Locoge, N., Bonsang, B., Salameh, T., Gerasopoulos, E., and Mihalopoulos, N.: Variability and sources of non-methane hydrocarbons at a Mediterranean urban atmosphere: The role of biomass burning and traffic emissions, *The Sci. Total Environ.*, 800, 149389, <https://doi.org/10.1016/j.scitotenv.2021.149389>, 2021.
- Phillips, L. and Moya, J.: The evolution of EPA's Exposure Factors Handbook and its future as an exposure assessment resource, *J. Expos. Sci. Environ. Epidemiol.*, 23, 13–21, 2013.
- Rad, H. D., Babaei, A. A., Goudarzi, G., Angali, K. A., Ramezani, Z., and Mohammadi, M. M.: Levels and sources of BTEX in ambient air of Ahvaz metropolitan city, *Air Quality, Atmos. Health*, 7, 515–524, <https://doi.org/10.1007/s11869-014-0254-y>, 2014.
- Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part F, Supplemental Guidance for Inhalation Risk Assessment), United States Environmental Protection Agency, Washington, DC (2009): <https://www.epa.gov/risk/risk-assessment-guidance-superfund-rags-part-f>, (last access: 21 April 2024), 2009.
- Salameh, T., Sauvage, S., Locoge, N., Gauduin, J., Perrussel, O., and Borbon, A.: Spatial and temporal variability of BTEX in Paris megacity: Two-wheelers as a major driver, *Atmos. Environ. X*, 1, 100003, <https://doi.org/10.1016/j.aeoa.2018.100003>, 2019.
- Schwartz, S. E. and Warneck, P.: Units for use in atmospheric chemistry (IUPAC Recommendations 1995), *Pure Appl. Chem.*, 67, 1377–1406, <https://doi.org/10.1351/pac199567081377>, 1995.
- Seco, R., Peñuelas, J., Filella, I., Llusia, J., Schallhart, S., Metzger, A., Müller, M., and Hansel, A.: Volatile organic compounds in the western Mediterranean basin: urban and rural winter measurements during the DAURE campaign, *Atmos. Chem. Phys.*, 13, 4291–4306, <https://doi.org/10.5194/acp-13-4291-2013>, 2013.
- Simon, L., Gros, V., Petit, J.-E., Truong, F., Sarda-Estève, R., Kalalian, C., Baudic, A., Marchand, C., and Favez, O.: Two years of volatile organic compound online in situ measurements at the Site Instrumental de Recherche par Télédétection Atmosphérique (Paris region, France) using proton-transfer-reaction mass spectrometry, *Earth Syst. Sci. Data*, 15, 1947–1968, <https://doi.org/10.5194/essd-15-1947-2023>, 2023.
- Song, M., Tan, Q., Feng, M., Qu, Y., Liu, X., An, J., and Zhang, Y.: Source apportionment and secondary transformation of atmospheric nonmethane hydrocarbons in Chengdu, Southwest China, *J. Geophys. Res.-Atmos.*, 123, 9741–9763, <https://doi.org/10.1029/2018JD028479>, 2018.
- Wang, H., Wang, Q., Chen, J., Chen, C., Huang, C., Qiao, L., Lou, S., and Lu, J.: Do vehicular emissions dominate the source of C6–C8 aromatics in the megacity Shanghai of eastern China? *J. Environ. Sci.*, 27, 290–297, <https://doi.org/10.1016/j.jes.2014.05.033>, 2015.
- WHO: Agents classified by the IARC monographs, World Health Organization, International Agency for Research on Cancer, <http://monographs.iarc.fr/ENG/Classification> (last access: 21 April 2024), 2016.
- Wu, R., Li, J., Hao, Y., Li, Y., Zeng, L., and Xie, S.: Evolution process and sources of ambient volatile organic compounds during a severe haze event in Beijing, China, *Sci. Total Environ.*, 560, 62–72, <https://doi.org/10.1016/j.scitotenv.2016.04.030>, 2016.
- Yang, J., Roth, P., Durbin, T., and Karavalakis, G.: Impacts of gasoline aromatic and ethanol levels on the emissions from GDI vehicles: Part 1. Influence on regulated and gaseous toxic pollutants, *Fuel (Guildford)*, 252, 799–811, <https://doi.org/10.1016/j.fuel.2019.04.143>, 2019.
- Yao, X., Ma, R., Li, H., Wang, C., Zhang, C., Yin, S., Wu, D., He, X., Wang, J., and Zhan, L.: Assessment of the major odor contributors and health risks of volatile compounds in three disposal technologies for municipal solid waste, *Waste Manage.*, 91, 128–138, <https://doi.org/10.1016/j.wasman.2019.05.009>, 2019.
- Yao, Y., Tsai, J., and Chou, H.: Air pollutant emission abatement using application of various ethanol-gasoline blends in high-mileage vehicles, *Aerosol Air Qual. Res.*, 11, 547–559, <https://doi.org/10.4209/aaqr.2011.04.0044>, 2011.
- Yurdakul, S., Civan, M., Kuntasal, Ö., Doğan, G., Pekey, H., and Tuncel, G.: Temporal variations of VOC concentrations in Bursa atmosphere, *Atmos. Pollut. Res.*, 9, 189–206, <https://doi.org/10.1016/j.apr.2017.09.004>, 2018.
- Zahed, M. A., Salehi, S., Khoei, M. A., Esmaeili, P., and Mohajeri, L.: Risk assessment of Benzene, Toluene, Ethyl benzene, and Xylene (BTEX) in the atmospheric air around the world: A review, *Toxicol. Vitro*, 105825, 98, <https://doi.org/10.1016/j.tiv.2024.105825>, 2024.
- Zalel, A. and Broday, D. M.: Revealing source signatures in ambient BTEX concentrations, *Environ. Pollut.*, 156, 553–562, <https://doi.org/10.1016/j.envpol.2008.01.016>, 2008.
- Zhang, Z., Man, H., Zhao, J., Jiang, Y., Zeng, M., Cai, Z., Huang, C., Huang, W., Zhao, H., and Jing, S.: Primary organic gas emissions in vehicle cold start events: Rates, compositions and temperature effects, *J. Hazard. Mater.*, 435, 128979, <https://doi.org/10.1016/j.jhazmat.2022.128979>, 2022.
- Zou, Y., Deng, X. J., Zhu, D., Gong, D. C., Wang, H., Li, F., Tan, H. B., Deng, T., Mai, B. R., Liu, X. T., and Wang, B. G.: Characteristics of 1 year of observational data of VOCs, NO_x and O₃ at a suburban site in Guangzhou, China, *Atmos. Chem. Phys.*, 15, 6625–6636, <https://doi.org/10.5194/acp-15-6625-2015>, 2015.