



**HAL**  
open science

# AQUEOUS ALTERATION FINGERPRINTS IN THE MOLECULAR COMPLEXITY OF CARBONACEOUS CHONDRITES

Francois-Regis Orthous-Daunay, Junko Isa, L. Flandinet, Véronique Vuitton

► **To cite this version:**

Francois-Regis Orthous-Daunay, Junko Isa, L. Flandinet, Véronique Vuitton. AQUEOUS ALTERATION FINGERPRINTS IN THE MOLECULAR COMPLEXITY OF CARBONACEOUS CHONDRITES. 86th Annual Meeting of the Meteoritical Society, Jul 2024, Bruxelles, Belgium. hal-04788745

**HAL Id: hal-04788745**

**<https://hal.science/hal-04788745v1>**

Submitted on 18 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## AQUEOUS ALTERATION FINGERPRINTS IN THE MOLECULAR COMPLEXITY OF CARBONACEOUS CHONDRITES.

F.-R. Orthous-Daunay<sup>1</sup>, J. Isa<sup>2,3</sup>, L. Flandinet<sup>1</sup> and V. Vuitton<sup>1</sup> <sup>1</sup>Univ. Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France (CS 40700 38058 Grenoble Cedex 9, France (francois-regis.orthous-daunay@univ-grenoble-alpes.fr), <sup>2</sup>Cold Pine Observatory (Chigasaki-shi Kanagawa, Japan [jisa@cpo.org](mailto:jisa@cpo.org)), <sup>3</sup>ELSI, Tokyo Institute of Technology (2-12-1-IE-1 Ookayama, Meguro-ku Tokyo 152-8550, JAPAN, [jisa@elsi.jp](mailto:jisa@elsi.jp)).

**Introduction:** The use of High Resolution Mass Spectrometry has yield a remarkable insight on the organic composition of carbonaceous chondrites and extraterrestrial samples [1]–[4]. The capability of counting every single mass in a given range came with the description of the sample mixture as a whole. In this new data, all the planetary processes affecting the minerals and the organic matter may have left their fingerprints. As far as carbonaceous chondrites are concerned, the origin and the survivability of organic matter are always interpreted with the asteroidal processes taking place and modifying or creating molecules.

The finding of polymerization patterns in pristine meteorites [5] and laboratory proxies [6] helps understanding the molecular mixture because of chemical selection. As a synthesis pattern can be isolated, any other feature in the chemical complexity can be interpreted as a modification or an alternative synthetic pathway. This way, the effect of UV irradiation has been measured [5] and more recently, the chemical modification due to asteroidal aqueous alteration has been highlighted [7].

The canonical synthesis pattern in a bell shape mass distribution due to geometrical growth of the molecules. This pattern is seen in artificial polymers as well as in carbonaceous chondrites. It is described by fitting a model [7], [8] among the measured distributions. One parameter, the broadness of the distributions, is systematically higher in the carbonaceous chondrites. One exception to the typically extraterrestrial broadness has been found in altered part of Tagish Lake [7]. We present here an augmented series of data and its implication in the quantification of the aqueous alteration.

**Results:** Tagish Lake lithologies known as TL1 li and TL5b, respectively the most altered and the most pristine [9] were prepared for Orbitrap analysis following protocol in [7]. Sub-spectra were isolated and fitted with a Wesslau geometrical growth model. On top of these two samples, a large variety of artificial synthetic samples have been treated the same way. Mixtures like in [10] and [11] are standards to monitor aqueously synthesized complex mixtures.

The altered parts of Tagish Lake are systematically narrower than the pristine part in good agreement with the lower entropy described in [7]. The artificial samples seem to have the same narrow distributions. This can be interpreted as a common chemical selection process leading to a limitation of the diversity in mass. One remarkable fact is the discrepancy between Orgueil and the most altered part of Tagish Lake. Orgueil seems to be on the canonical pristine trend. This may be paradoxical since Orgueil is thought to have undergone extensive aqueous alteration. We will discuss the implication of Orgueil having « normal » organics.

**References:**[1] Schmitt-Kopplin, P. *et al.* Feb. 2010.*Proc. Natl. Acad. Sci.*vol. 107, no. 7.pp. 2763–2768[2] Schmitt-Kopplin, P. *et al.* 2023.*Nat. Commun.*vol. 14, no. 1.pp. 1–9[3] Oba, Y. *et al.* Dec. 2022.*Nat. Commun.*vol. 13, no. 1.p. 2008[4] Furukawa, Y. *et al.* 2023.*Sci. Rep.*vol. 13, no. 1.pp. 1–12[5] Orthous-Daunay, F. R. *et al.* 2019.*Geochem. J.*vol. 53, no. 1.pp. 21–32[6] Moran, S. E. *et al.* 2020.*Am. Astron. Soc. Meet. Abstr.* 235.vol. 235.pp. 204–248[7] Isa, J.Orthous-Daunay, F.-R.Beck, P.Herd, C. D. K.Vuitton, V.and Flandinet, L. Oct. 2021.*Astrophys. J. Lett.*vol. 920, no. 2.p. L39[8] Weßlau, V. H. 1956.*Die Makromol. Chemie.*vol. 20, no. 1.pp. 111–142[9] Herd, C. D. K. *et al.* Jun. 2011.*Science (80- )*.vol. 332, no. 6035.pp. 1304–1307[10] Vinogradoff, V.Bernard, S.Le Guillou, C.and Remusat, L. 2017.*Icarus.*vol. 305.pp. 358–370[11] Isono, Y.Tachibana, S.Naraoka, H.Orthous-Daunay, F. R.Piani, L.and Kebukawa, Y. 2019.*Geochem. J.*vol. 53, no. 1.pp. 41–51