

## Declining juvenile survival of Adélie penguins in Antarctica

Téo Barracho, Gaël Bardon, Rémi Choquet, Aymeric Houstin, Alexander Winterl, Michaël Beaulieu, Pierrick Blanchard, Cindy Cornet, Robin Cristofari, Thierry Raclot, et al.

### ▶ To cite this version:

Téo Barracho, Gaël Bardon, Rémi Choquet, Aymeric Houstin, Alexander Winterl, et al.. Declining juvenile survival of Adélie penguins in Antarctica. 2024. hal-04791136

## HAL Id: hal-04791136 https://hal.science/hal-04791136v1

Preprint submitted on 20 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# Declining juvenile survival of Adélie penguins in Antarctica

3

#### 4 Authors:

- 5 Téo Barracho <sup>1,2,3,4\*</sup>, Gaël Bardon <sup>1,4</sup>, Rémi Choquet <sup>3</sup>, Aymeric Houstin <sup>1,4,7</sup>, Alexander
- 6 Winterl <sup>5</sup>, Michaël Beaulieu <sup>8</sup>, Pierrick Blanchard <sup>9</sup>, Cindy C. Cornet <sup>4,10</sup>, Robin Cristofari <sup>6</sup>,
- 7 Thierry Raclot<sup>1</sup>, Sebastian Richter<sup>5</sup>, Nicolas Chatelain<sup>1</sup>, Julien Courtecuisse<sup>1</sup>, Matthieu
- 8 Brucker <sup>1</sup>, Daniel P. Zitterbart <sup>5,7</sup>, Nicolas Lecomte <sup>2\*†</sup> and Céline Le Bohec <sup>1,3,4\*†</sup>
- 9

#### 10 <sup>†</sup> Co-last authors

- 11 \* tbarracho@centrescientifique.mc
- 12 \* nicolas.lecomte@umoncton.ca
- 13 \* celine.le-bohec@cnrs.fr
- 14

#### **15** Affiliations:

- <sup>16</sup> <sup>1</sup> Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France
- <sup>17</sup> <sup>2</sup> Canada Research Chair in Polar and Boreal Ecology and Centre d'Études Nordiques,
- 18 Department of Biology, University of Moncton, Moncton, NB, Canada
- <sup>19</sup> <sup>3</sup> CEFE, Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France
- <sup>20</sup> <sup>4</sup> Centre Scientifique de Monaco, Département de Biologie Polaire, Monaco, Principality of
- 21 Monaco
- <sup>5</sup> Friedrich-Alexander-University Erlangen-Nürnberg, Department of Physics, Erlangen,
- 23 Germany
- <sup>6</sup> Institute of Biotechnology, University of Helsinki, Helsinki, Finland
- <sup>25</sup> <sup>7</sup>Woods Hole Oceanographic Institution, Applied Ocean Physics and Engineering Department,
- 26 Woods Hole, Massachusetts, USA
- <sup>8</sup> Wild Animal Initiative 5123 W 98th St, 1204 Minneapolis, MN 55437
- <sup>9</sup> Centre de Recherche sur la Biodiversité et l'Environnement (CRBE), Université de Toulouse,
- 29 CNRS, IRD, Toulouse INP, Université Toulouse 3 Paul Sabatier (UT3), Toulouse, France
- <sup>30</sup> <sup>10</sup>Centre for Blue Governance, Faculty of Business and Law, University of Portsmouth, UK
- 31
- 32

#### **33 Abstract:**

As summer sea ice around Antarctica reaches historical lows, quantifying the demographic 34 35 response of polar species to such environmental changes becomes critical. To achieve this, synthesizing results from across species' ranges and elucidating the environmental factors 36 driving population dynamics are key. Adélie penguins are considered reliable indicators of 37 changes in Antarctica but the pathways through which sea ice and other environmental factors 38 shape their population dynamics are still unclear, especially for the juvenile stage. Using a 17-39 year dataset of electronically tagged Adélie penguins from Adélie Land (Antarctica), we found 40 that juvenile survival probability was most impacted by sea ice concentration near their natal 41 colony right after fledging, with lower ice concentrations detrimental to survival. Importantly, 42 43 we found that juvenile survival declined by 32% from 2007 to 2020, mirroring trends at other distant colonies. The emergence of similar patterns at opposite ends of the continent may be an 44 45 early signal for shifts in population trends expected from climate change.

46

#### 47 Teaser:

48 Similar declines in the survival of juvenile Adélie penguins have emerged at distant locations
49 around Antarctica and receding sea ice could be at play.

#### 50 MAIN TEXT

51

#### 52 Introduction:

Antarctica is at the center of large and rapid environmental shifts, with both accelerating 53 climate changes (1, 2) and increasing exploitation pressure by fisheries (3). Quantifying the 54 response of animal populations to such environmental forcing is essential to determine their 55 potential dynamics and persistence (4, 5). Yet, our understanding of how Antarctic species 56 57 respond to such changes is still fragmented. Typically, the heterogeneous dynamics of climate 58 (6) and environmental parameters such as sea ice (7) makes identifying drivers of any demographic rate challenging (8–10). This hampers our ability to project species response to 59 environmental changes, especially over scales relevant to ecosystem management (11, 12). 60

61

The Adélie penguin (*Pygoscelis adeliae*), an iconic Antarctic species, exemplifies this challenge particularly well. Early observations of contrasting population trends between the Ross Sea and the Antarctic Peninsula in response to sea ice declines have fostered a population dynamics' model, where abundance is maximal at intermediate sea ice concentrations (*13*, *14*).

This model has been instrumental in projecting the response of Adélie penguins to climate 66 67 variability across large spatial and temporal scales (15-17). However, more localized and shorter-term predictions are still hindered by the inherent variability and noise in abundance 68 time series which most studies are constrained to rely on (17-19). Since abundance results from 69 the complex interplay of several demographic rates, each influenced by specific environmental 70 71 drivers, and some of which are poorly understood (20), focusing on single vital rates could enhance the accuracy of projecting species responses to environmental change (21). The need 72 73 for such an approach is twofold; it can provide short-term feedback for ecological management at the local scale and be integrated at larger spatial scales as more studies become available 74 75 across the species range (22-25).

76

In long-lived species such as the Adélie penguin, adult survival typically holds the greatest 77 potential to influence population dynamics (26). Yet, juvenile survival appears to play a 78 disproportionate role in driving Adélie penguin population fluctuations. For example, Adélie 79 penguin's abundance (as measured by the number of breeding pairs) correlates with sea ice 80 conditions with a 5-years lag (27); which is consistent with effects of sea ice on juvenile 81 survival during their first months at sea (28). In the Antarctic Peninsula (29, 30) and East 82 Antarctica (31), population size decreased when juvenile survival also did, further emphasizing 83 its central role in shaping overall population trends. Despite such findings, we still lack clear 84 evidence for a relationship between juvenile survival and environmental parameters known to 85 affect population dynamics, such as sea ice concentration (11). Although adult survival has 86 been convincingly linked to environmental variables (e.g., Southern Oscillation Index, winter 87 sea ice concentration) in several populations around Antarctica (29, 32-35), the drivers of 88 juvenile survival remain unclear (29, 31, 34). Given the importance of this demographic 89 parameter, evaluating its trends across the species' range and elucidating its environmental 90 91 drivers are required to adequately predict the species' response to future environmental changes 92 facing Antarctica (36).

93

Here, we quantify juvenile survival probabilities for the first time in the Western Pacific Ocean sector (90°E to 160°E, Fig. S1) using a 17-year-long dataset of known-age, electronically tagged Adélie penguins from Pointe Géologie archipelago, Adélie Land. By investigating trends in survival for this population located thousands of kilometers away from other studied colonies and synthesizing previously published studies on this vital rate, we provide a comprehensive picture of trends in Adélie penguin juvenile survival across Antarctica. To

uncover the mechanisms behind variations in juvenile survival, we also investigate the drivers
 of juvenile survival at Pointe Géologie by linking survival probabilities to intrinsic and
 environmental parameters such as body mass at fledging and sea ice concentrations.

103

#### **104 Results:**

We used a multi-state capture-recapture framework accounting for short-range dispersion 105 within a 10-km radius to model the survival probability of juvenile Adélie penguins (0-2 years 106 old). This model (see methods) enabled us to quantify juvenile survival probabilities between 107 108 2007 and 2020. Over this period, juvenile survival estimates averaged  $0.42 \pm 0.18$  and showed high inter-annual variability (CV = 44%, Fig. 2A). Notably, juvenile survival probabilities 109 110 exhibited a negative trend (Fig. 2A; Table 1, model 30, analysis of deviance: *p*-ANODEV = 0.016). The annual rate of change was of -2.5% ( $\pm 95\%$ CI: -3.2, -1.9), translating into a 32% 111 112 loss between 2007 and 2020. An exhaustive literature search showed that juvenile survival had previously been quantified at 5 other locations around Antarctica ((29, 31, 33, 37, 38), 113 summarized in Fig. 2). Temporal variation was investigated at only two of them, and juvenile 114 115 survival probabilities were found to decline at both sites (Admiralty Bay, King George Island, Western Antarctic Peninsula, -1.3% per year for 1982-2000 (29); Béchervaise Island, Mac. 116 Robertson Land, East Antarctica, -1.8% per year for 1992-2015 (31); Fig. 2). 117

118

To understand the drivers behind the decline observed at our study site, we used our previously 119 developed model to test the effect of various intrinsic (mass at fledging, cohort size) and 120 environmental covariates such as sea ice concentrations (SIC), large-scale climatic indices, and 121 windchill temperatures on juvenile survival probabilities. Among the original set of covariates, 122 we considered (Table S2), only autumn sea ice concentrations near the natal colony 123 (SIC.autumn) accounted for more than 20% of the temporal variance (DEV) in juvenile 124 survival rates (Table 1, model 22, %DEV = 28.2). After decomposing this variable into 125 monthly data (SIC.march and SIC.april), only SIC.april reached the 20% DEV threshold (Table 126 127 1, model 26, %DEV = 39.4). By decomposing SIC.april into the fraction of total ice area (15-100% SIC) covered by the marginal ice zone (15-80% SIC, MIZFRAC.april) and the fraction 128 of total ice area covered by dense sea ice (80-100% SIC, DENSEFRAC.april), we then revealed 129 that the amount of loose sea ice in the vicinity of the colony (MIZFRAC.april) was negatively 130 related to juvenile survival rates (slope  $\pm 95\%$  CI = -0.97 (-0.99, -0.94), Fig. 3B) and accounted 131 for 78.8% of the temporal variation (Table 1, model 27). Conversely, the amount of dense sea 132

ice (DENSEFRAC.april) was positively related to juvenile survival rates (slope  $\pm$  95%CI = 0.85 (0.83, 0.87), Fig. 3C) and accounted for 39.7% of the temporal variation (Table 1, model 28). This amount of April dense sea ice near the colony declined by 13% during the time of our study (linear model: F = 7.65, *p* = 0.016, R<sup>2</sup> = 0.37, Fig. 3D).

137

138 Linear models showed that chick body mass at tagging, an important driver of post-fledging survival in birds, decreased slightly over the study period (F = 57.87, p < 0.001, R<sup>2</sup> = 0.02, 139 2007-2022 average =  $3.60 \pm 0.61$  kg), and varied more within (average of annual mass SD = 140 0.54 kg) than across cohorts (SD of annual average mass = 0.33 kg, Fig. S8). Multi-state 141 142 capture-recapture modeling showed that chick body mass was positively related to juvenile survival probability, both across cohorts (mean annual fledging mass, MFM, Table 1, model 143 29, %DEV = 22.0, slope  $\pm$  95%CI = 0.04 (0.04, 0.04)), and also at the individual level, with a 144 model-predicted 5.4 % increase in survival probability for every 500 grams of chick body mass 145 (i.e. within cohorts, Fig. 4, LRT,  $\chi^2_1 = 48.014$ , p < 0.001). The slope of this relationship did not 146 differ among years, as indicated by the additive model having a lower AIC than the interactive 147 model (Table 1, models 33 and 34, respectively). 148

149

To ensure that our study colony was representative of the larger Pointe Géologie archipelago 150 population (see methods), we compared the breeding productivity of the study colony to that 151 of the entire population. Over the 2011-2024 period, breeding productivity for our study colony 152 averaged  $0.63 \pm 0.44$  chicks per breeding pair (Fig. S5) and was highly correlated to that of the 153 entire Pointe Géologie archipelago (2011-2017, Pearson correlation: r = 0.99, p < 0.001, Fig. 154 S6). The number of breeding pairs in the study colony averaged  $273 \pm 62$  between 2011 and 155 2024 and remained comparable with two undisturbed control colonies located on the same 156 island (Spearman rank correlations, both p < 0.022, Fig. S7). 157

158

#### **159 Discussion:**

Using 17 years of individual electronic monitoring, we observed a loss of 2.5% per year in Adélie penguin juvenile survival over 14 years (2007-2020) in an Adélie Land population, a rate likely stronger than the declines already observed in the only two other populations for which juvenile survival time series are available (in Mac. Robertson Land, East Antarctica, -1.8% per year (*31*), and in the Western Antarctic Peninsula, -1.3% per year (*29*)). By quantifying the survival of juvenile Adélie penguins for the first time in a region located thousands of kilometers (3400 km and 6300 km, respectively) away from other monitored colonies, our results point to emerging negative trends across the continent. We also highlight potential mechanisms at play, since the decline in juvenile survival probability recorded in Adélie Land was associated with decreasing amounts of landfast (dense) sea ice in the vicinity of the natal colony within two months after fledging. As sea ice declines are predicted to shrink Adélie penguin populations in the latter part of the century (*15*, *16*), steady declines in juvenile survival may already foreshadow such shifts.

- 173
- 174

#### • Range-wide declines in juvenile survival

175

Permanent dispersion of marked individuals away from study sites can bias the results of capture-recapture studies when incorrectly accounted for (*39*). Here, our deployment of a mobile RFID antenna network, radiating 10 kilometers around the main study colony (Fig. 1), has unveiled compelling evidence of high local philopatry, allowing a robust estimation of survival probability in spite of local dispersion. Long-distance permanent dispersion of juveniles, which could bias our survival estimates low, remains nonetheless largely unknown and should be addressed in the future.

183

Because of these difficulties and exceptional logistic constraints, temporal variations in the 184 survival of juvenile Adélie penguins had only been investigated at two other locations in 185 Antarctica before this study, and only at a single one (Béchervaise Island) without the negative 186 impact of flipper bands on penguin vital rates (40, 41). Hinke et al. (29) were the first to report 187 an annual decline in juvenile survival probability, with a decline of -1.3% per year (1982-2000) 188 for Admiralty Bay (King George Island, Western Antarctic Peninsula). Survival probability at 189 this site then seemed to stabilize up to 2011, although a change in marking methodology (from 190 191 aluminum to stainless steel flipper bands) and high inter-annual variability in survival 192 probabilities made it difficult to compare the two periods (29, 30). More recently, Emmerson and Southwell (31) extended the analysis of the capture-recapture dataset from Béchervaise 193 Island (Mac. Robertson Land, East Antarctica) to reveal that juvenile survival had also declined 194 (-1.8% per year, 1992-2015) in this population located halfway between the Antarctic 195 Peninsula and Adélie Land (Fig. 2). 196

197

With an annual decrease of -2.5% between 2007 and 2020 in Adélie Land, our study adds to
the picture of declining juvenile survival across the species range (Fig. 2). Juvenile survival

has now declined at all three locations where time trends have been investigated (Fig 2). The time series used here (14 years) is shorter than those for Admiralty Bay (18 years) and Béchervaise Island (24 years), but it is worth noting that the -2.5% annual decline at Pointe Géologie exceeds that of these two other colonies. In addition, such an annual decline in Pointe Géologie exceeds the average inter-annual variability in survival probabilities (2.50% vs. 1.79%) as in Béchervaise Island (1.80% vs. 1.57%) but unlike Admiralty Bay (1.30% vs. 2.90%).

- 207
- 208 209

#### Drivers of juvenile survival

The existence of a relationship between Adélie penguin population dynamics and sea ice is 210 well established (11, 16, 38, 42). However, in spite of the importance of juvenile survival in 211 driving population trends (31, 43, 44), previous studies have found relatively limited evidence 212 for a relationship between SIC and juvenile survival (29, 31, 34). Here, unlike for the two 213 populations previously studied (29, 31), we found that local sea ice concentration within 200 214 km of the natal colony in the timeframe of one to two months after fledging (April) accounted 215 for up to 40% of the temporal variation in survival probability (Fig. 3). Specifically, higher SIC 216 in April had a positive effect on juvenile survival. The timing of this relationship, together with 217 the fact that we did not detect effects of environmental variables later in the year, is highly 218 consistent with a survival bottleneck occurring in the weeks following fledging, as identified 219 for *Pygoscelis* penguins in the Antarctic Peninsula (45). A mortality peak immediately 220 following fledging is also consistent with previous studies in mammals (46, 47) and birds (48, 47)221 49). 222

223

Our finding that SIC in April, but not in March, was correlated to juvenile survival may be 224 225 particularly informative on the mechanisms linking sea ice to the survival of young Adélie penguins. Although the winter freeze-up of the Antarctic Ocean begins in March, sea ice 226 usually consolidates markedly and expands equatorward in April (Fig. S10, (50, 51)). Newly 227 fledged Adélie penguins therefore appear to be reliant on the rapid formation of sea ice at the 228 onset of winter, and especially dense landfast ice (Fig. 3, Fig. S11). By decomposing April SIC 229 into its loose (the marginal ice zone, between 15 and 80% SIC) and dense (80-100%) fractions, 230 we were able to show that juveniles survived better in years when sea ice was made of large, 231 fully frozen areas, rather than patchy ice (Fig. 3, Fig. S11). 232

233

To explain this relationship, we suggest that dense sea ice could provide more suitable resting 234 235 grounds for fledglings, in a context where they are at high risk of energy reserves depletion (45). Furthermore, the formation of large areas of solid ice allows the emergence of polynyas 236 and flaw leads (52), where prey may be more abundant (53) and/or more localized and thus 237 easier to detect and capture, especially for inexperienced individuals likely to display lower 238 foraging abilities compared to adults as shown in king penguins (Aptenodytes patagonicus) 239 (54). Finally, patchier sea ice may favor predation, especially by leopard seals (Hydrurga 240 *leptonyx*) (55), although predation pressure during fall and winter has never been investigated. 241 Biologging studies have shown that during the winter, adult Adélie penguins tend to be 242 associated with dense sea ice areas (56), with higher SIC promoting higher survival to the next 243 breeding season (35). For juveniles, however, knowledge of post-fledging behavior is 244 extremely limited. Only two studies so far have investigated their post-fledging movements 245 (45, 57), while their diving behavior and association with specific sea ice types remain 246 completely unexplored. 247

248

While SIC measured right after fledging accounted for most of the inter-annual variation in 249 juvenile survival, conditions during the breeding season may also play a significant role, as 250 juvenile survival was positively related to chick mass prior to fledgling (Fig. 4). The decline in 251 juvenile survival we observed over the study period may therefore be partly accounted for by 252 a concurrent but slight decrease in fledging mass (Fig. S8). Both the diet (58, 59) and the 253 foraging efficiency of breeders (60, 61) have indeed been linked to sea ice conditions during 254 chick rearing, and these parameters may directly affect chick fledging mass (62, 63). Land-255 based, food-independent, factors may also act, with higher precipitation and stronger winds 256 late in the breeding season negatively affecting chick fledging mass (64), likely because of 257 higher thermoregulatory costs for the chicks (63). Interestingly, the individual-level and 258 positive effect of juvenile body mass at fledging on survival probabilities was similar across 259 years, corroborating results from a previous study in the Ross Sea (62). These findings suggest 260 that heavier chicks may maintain their survival advantage over lighter ones across a range of 261 environmental conditions. 262

- 263
- 264

#### • Demographic consequences and perspectives

265

In long-lived species, the survival of young age classes can have major impacts on population
dynamics (65). For Adélie penguins, this is exemplified by declines in abundance of breeders

that followed declines in juvenile survival probabilities in two distinct populations from the 268 269 Western Antarctic Peninsula (29, 30) and East Antarctica (31). As in the present study, it was not possible to identify the onset of the decline in juvenile survival in these two populations, 270 since trends were observable right from the beginning of the monitoring (Fig. 2). Nonetheless, 271 evidence from East Antarctica suggests that declining juvenile survival may not cascade into 272 an abundance decline immediately, with a ~10 years lag between the onset of decline in 273 juvenile survival and the breakup in the abundance time series ((31), Fig. 2). Furthermore, 274 events of low adult survival and repeatedly low breeding productivity may have acted in 275 conjunction to generate the decline in abundance at Admiralty Bay (30) and Béchervaise Island 276 (31), respectively. This may help explain why abundance at Pointe Géologie has not yet shown 277 signs of decline ((66), Fig. 2). However, given the occurrence of two near-total breeding 278 failures in the past decade alone (67) and the stronger rate of decline in juvenile survival 279 observed at Pointe Géologie compared to other studies and to local inter-annual variability in 280 281 this vital rate, it is possible that the Pointe Géologie population will also decline in the coming 282 years.

283

While Adélie penguin population declines for the Western Antarctic Peninsula are well 284 documented (68–71) and consistent with the stronger rate of warming in the region (72), 285 declines in other regions and notably in East Antarctica were not expected before the later part 286 of the 21st century (16), with the Ross sea likely being the main refugia for the species (15). 287 However, our findings in the Western Pacific Ocean sector of Antarctica, combined with recent 288 evidence that an entire metapopulation elsewhere in East Antarctica has already declined by 289 almost half in the past decade alone (31) suggest that changes could happen earlier than 290 expected. 291

292

293 Such rapid changes are likely to challenge the adaptive abilities of the Adélie penguin. One of 294 the main mechanisms through which Adélie penguins may respond to these changes is dispersal towards areas where environmental conditions remain or become favorable as ice melts (16,295 73). Although mid-range (< 120 km) dispersion is possible in adult Adélie penguins (74), we 296 still lack estimations of long-distance dispersion over time scales relevant with the current rate 297 of environmental change. Similarly, we have highlighted that no clear universal environmental 298 drivers of juvenile survival stand out, although similar temporal trends have emerged across 299 the species range. 300

301

In this context, the next steps would first be to harmonize demographic databases across study 302 sites and analyze them collectively. Specifically, quantifying the temporal variation in juvenile 303 survival for the two Ross sea populations (33, 74, 75) is key. Completing the transition from 304 flipper bands to RFID-tags (40) could also facilitate comparisons of demographic rates among 305 populations. Second, conducting biologging studies of juvenile movement and diving behavior, 306 in conjunction with sea ice analyses (akin to (35)), would help identify range-wide 307 environmental drivers of survival. Third, rigorous efforts should be made to assess the dispersal 308 309 capabilities of juvenile Adélie penguins under climate change, such as deploying autonomous RFID-tag detection systems (Fig. S2) in populations adjacent to the main colonies where this 310 311 method is used. With the forthcoming fifth International Polar Year (2032-2033) and the likely large-scale decline in the survival of juvenile Adélie penguins, a significant window of 312 opportunity emerges for closely monitoring the demographic trends of vulnerable species and 313 unraveling the underlying factors driving these trends. 314

315

```
316 Methods:
```

- 317
- **318** *Study area and design*
- 319

Fieldwork was carried out at Pointe Géologie archipelago (Adélie Land, Antarctica; Fig. 1),
close to the Dumont d'Urville research station (66°40′S, 140°01′E). This archipelago has hosted
30,000-50,000 breeding pairs of Adélie penguins annually for the last two decades (66).

323

From the 2006-2007 breeding season onwards (breeding season will be referred to by the 324 fledging year, e.g., 2007 in this case), Radio Frequency IDentification tags (RFID-tags) were 325 implanted in chicks from a colony of about 270 breeding pairs located in a natural canyon 326 (hereafter study colony). Unlike flipper bands (40, 41, 76), RFID-tags are not known to 327 negatively affect penguin vital rates. All chicks still alive right before fledging were tagged (n 328 = 3,138). A few RFID-tagged chicks (n = 76) were found dead within the colony prior to 329 fledging and were consequently excluded from subsequent analyses, bringing the final dataset 330 to 3,062 individuals (range = 0-350/year, annual average = 192 from 2007 to 2022). 331

332

The study colony was fenced off in 2009 and instrumented with RFID gateways similar to those used in other penguin populations (Adélie penguins: (77–79); King penguins: (80, 81);

Southern rockhopper penguins: (82); Macaroni penguins: (83); Little penguins: (84)). This 335 336 setting made it impossible to miss tagged individuals visiting the colony. To account for dispersal from the study colony to neighboring colonies within a 10-km radius (Fig. 1), a grid 337 of mobile RFID detection units (2-8 units, Fig. S2) was deployed every year from November 338 to March, starting in 2013. Mobile RFID units were positioned at natural bottlenecks frequently 339 used by penguins traveling between colonies and the sea. Units were moved every one to two 340 weeks to maximize the number of detections. Among the 1061 individuals reobserved at least 341 once after their tagging year, only 19 (1.8%) were detected exclusively by mobile RFID 342 antennas, i.e. away from their natal colony. The mobile RFID-units resighting effort was more 343 344 intensive closer to the study colony (< 200 m) but was performed up to 10 km from it (Fig. 1).

345

#### 346

#### • Colony and individual-scale parameters

347

Colony-scale breeding productivity was defined as the number of fledglings, divided by the 348 annual maximum number of adults inside the colony divided by two (a proxy for the number 349 of breeding pairs). Individuals present in the colony were photo-counted weekly during the 350 whole breeding season, starting in 2011. Cohort size was taken as the total number of RFID-351 tagged chicks that fledged. We ensured that observations from the study colony were 352 representative of the larger-scale Pointe Géologie archipelago by comparing our local estimate 353 of breeding productivity with data available from the literature for Pointe Géologie (2011-2017, 354 (85)). Similarly, to ensure that our monitoring setup did not impact breeding abundance at the 355 colony, we compared the trend in number of breeding pairs in the study colony with that of two 356 other undisturbed colonies located on the same island for the 2011-2024 period (Fig. S7). 357

358

All RFID-tagged chicks were weighed during tagging to the nearest 0.05 kg. The timing of tagging was adjusted to annual chick departure phenology (i.e. starting when the first molting chick was observed in the colony), ensuring mass was comparable across years. Each year, tagging occurred 10-15 days prior to fledging, making chick mass at tagging a reliable and consistent proxy for fledging mass (62). In Pointe Géologie archipelago, fledging occurs from late February to early March.

- 365
- 366

Capture Mark Recapture (CMR) modeling and statistical analyses

367

To model apparent survival and recapture probabilities of juvenile Adélie penguins, we constructed capture histories for all individuals (n = 3,062) fledged between 2007 and 2022, and subsequently observed between the 2009 and 2023 breeding seasons. Out of these 3,062 fledglings, 1,061 were observed again in subsequent years.

372

Given the dispersal of some birds from the study colony to neighboring colonies, and the higher 373 resighting effort on and near the study colony, we adopted a multi-state framework to model 374 survival, recapture, and transition probabilities (86). We implemented this framework in the 375 program MARK v.9.0 (87). Different resigning probabilities among colonies were accounted 376 377 for using a two-state classification: a local state (1) for the study colony and adjacent colonies within 200 meters, and a distant state (2) for all colonies located more than 200 meters from 378 the study colony. This allowed closing the system and enhancing the robustness of survival 379 estimates despite short-range dispersal dynamics. As the detection probability of 1-year-old 380 individuals was near 0 (only a single bird was detected at age 1, Fig. S3), it was not possible to 381 compute survival probability between ages 0 and 1. The first recapture occasion following 382 tagging was thus deleted for all individuals (supplementary text). Consequently, juvenile 383 384 survival was estimated between 0 and 2 years old.

385

We used U-CARE software (88) to evaluate the goodness of fit (GOF) of the two-states time-386 dependent (JollyMoVe, JMV) model to our dataset (89). The JMV model did not fit the original 387 dataset because of transients (i.e. individuals never reencountered after tagging, Table S1a). 388 We thus tested the GOF of the JMV model to a dataset where the first capture was set to 0 for 389 each individual (i.e. effectively making capture history of each individual start at first 390 reencounter following tagging, see supplementary text), thereby allowing to test the fit of a 391 model with extended age-dependence on survival and recapture probabilities. The fit of the 392 393 JMV model to this dataset was satisfactory (Table S1), albeit with indications of further age 394 structure (i.e. > two age classes) in survival probabilities (significant test 3G.SM, Table S1b) and capture probabilities (almost significant Test M.ITEC, Table S1b). This validated the use 395 of an initial umbrella model incorporating  $\geq$  two age classes on recapture, survival, and 396 transition probabilities (model 1, Table S3). 397

398

In Adélie penguins, presence at the breeding colony (and thus availability for detection) is conditional on age, with reproductively mature individuals more likely to visit colonies than pre-breeders. Survival is also age-dependent, with lower survival probability in juveniles

compared to adults (33). Earlier studies of known-age Adélie penguins have therefore 402 403 accounted for age-structured recapture and survival probabilities, with either two (29), three (34), or five age classes (33). Here, we started from a model with five age classes on survival, 404 recapture, and transition probabilities, and explored all the possibilities in each submodel down 405 to only two age classes. Finally, our starting (umbrella) model was adjusted to account for the 406 low number of individuals detected in distant colonies. This model was framed with age 407 variation only (no time effects) for all transition probabilities, and recapture probabilities in the 408 distant colonies (state 2). The survival probabilities in both states (i.e. local and distant) were 409 also set to be equal in that model, which appears as a reasonable assumption considering that 410 all colonies belong to the same population of Pointe Géologie archipelago (Fig. 1). We 411 successively looked for the most appropriate age and time structure for recapture, transition, 412 and survival probabilities, using the previously retained best structure for each submodel. The 413 most parsimonious model was retained when two models were within two AIC points of each 414 other. This selection process yielded an optimal model (model 13, Table S3) including two age 415 classes (0-2 and 3+) and time effects within each age class for survival probabilities. Recapture 416 probabilities were best modeled using three age classes in both states. The best structure 417 included additional time effects in each of these three age classes for the local state only (Table 418 S3). Transition probabilities were best modeled with state and age effects (five age classes in 419 each state, Table S3). Because survival and recapture probabilities are not separable for the last 420 year (2021) in time-dependent CMR models, this model allowed to estimate juvenile (0-2) 421 survival from 2007 to 2020 only. Estimates of recapture probability in the local state were low 422 and displayed high inter-annual variability for the first age class (mean  $\pm$  SD: 0.33  $\pm$  0.24, Fig. 423 S9), but were high and stable in the last age class (mean  $\pm$  SD: 0.96  $\pm$  0.05, Fig. S9). Recapture 424 probabilities in the distant state were estimated for the last age class only and were particularly 425 low (0.02, SE = 0.009, Fig. S9). This model was then used as a basis for testing the effect of 426 selected covariates on juvenile survival (see below). 427

- 428
- 429

#### • Covariate selection and definition

430

Candidate environmental variables were considered based on relevance to the Adélie penguin's
ecology and previous studies of Antarctic seabirds (Table S2). We first considered two largescale indices, indicators of broad climatic variations in Antarctica and the surrounding Southern
Ocean: the Southern Oscillation Index (SOI) and the Southern Annular Mode (SAM).
Together, they affect wind patterns, temperatures, and sea ice dynamics, with complex

regional-specific variability (6, 90). Such variability has been linked to opposite responses both 436 437 among and within species. For instance, positive SOI values were previously linked to lower survival in adult Adélie penguins in Adélie Land (32) but no support for such an effect was 438 found in Mac. Robertson Land (34), the Western Ross Sea (33), or the Antarctic Peninsula 439 (29). High SAM values were also found to affect the survival of other Antarctic seabirds, either 440 positively (adult Cape petrel, *Daption capense*, (91); juvenile Emperor penguins, *Aptenodytes* 441 forsteri, (92) or negatively (juvenile Snow petrels, Pagodroma nivea, (93)). Monthly SOI data 442 were downloaded from the NOAA National Centers for Environmental Information 443 (https://www.cpc.ncep.noaa.gov/data/indices/soi) and daily SAM data from the NOAA 444 445 Climate Prediction Center (https://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily ao index/aao/aao.shtml). 446

447

Regional sea ice extent anomalies (SIEA) were also considered because of the quadratic relationship previously observed between SIEA and the survival of adult Adélie penguins in the Western Ross Sea (*33*). Monthly data for the Western Pacific sector (90°E to 150°E), which encompasses our study site, were downloaded from the National Snow and Ice Data Center (NSDIC, <u>https://nsidc.org/data/g02135/versions/3</u>). To capture the conditions experienced by juveniles after fledging and during their first winter at sea, each index (SOI, SAM, SIEA) was averaged from March to September (*34*), resulting in one value per year per index.

455

At the local scale, we considered both linear and quadratic effects of the average windchill temperatures (WCT) between April and September, based on the quadratic relationship previously observed between WCT and the survival of juvenile Adélie penguins (*31*). Ten-day average data of minimum temperature and average relative humidity from the Dumont D'Urville weather station were downloaded from Meteo France (<u>https://meteo.data.gouv.fr/</u>) and used to calculate WCT following the Australian Bureau of Meteorology calculation (as in (*31*)).

463

Finally, we considered sea ice concentrations (SIC, average sea ice concentration in a specific area) at three different spatio-temporal scales (Fig. S4). As tracking data for juveniles is currently not available in our focal region, we considered broadly similar migration routes as those of adults previously studied at Pointe Géologie and found to gradually move westward after reproduction following the Antarctic coastal current (94). This pattern is corroborated by the tracking of both juveniles and adults in another east Antarctica population (34, 57) and

therefore likely in our study area. To reflect post-fledging (autumn) conditions experienced by 470 birds near their natal colony, we considered March-April SIC between 135°E and 145°E 471 (SIC.autumn). We also considered winter (May-July, SIC.winter) and late winter (August-472 September, SIC.latewinter) conditions, both between 110°E and 135°E. Sea ice concentrations 473 were computed by averaging daily concentration values from 25×25 km grids downloaded 474 from the NSDIC ((95), https://nsidc.org/data/g02202/versions/4) over the spatio-temporal 475 windows defined above. Following initial analyses, we further adjusted our sea ice metrics by 476 first decomposing SIC.autumn into March and April SIC (SIC.march, SIC.april). Because 477 different types of ice may affect seabirds differently (96), we then calculated the April fraction 478 479 of the total ice area (15-100% SIC) covered by the marginal ice zone (loose sea ice between 15-80% SIC, MIZFRAC.april) and the April fraction of the total ice area covered by dense 480 pack ice (80-100% SIC, DENSEFRAC.april). 481

482

We also considered cohort size and average mass at tagging (as a proxy for fledging mass, (62)) as likely intrinsic drivers of juvenile survival in Adélie penguins. Cohort size was shown to correlate positively with fledgling survival in a previous study (31), while fledging mass is a common predictor for juvenile survival in birds (e.g., (97)), including penguins (62, 98). The effect of fledging mass on inter-annual differences in survival probabilities was investigated by using mean fledging mass (MFM) for each cohort. Inter-individual, within-season effects of fledging mass were also investigated, using chick mass as an individual covariate.

490

To avoid including covariates that are correlated with each other (multicollinearity, see (23)), 491 we checked for pair-wise correlations among our initial set of covariates (SOI, SAM, SIEA, 492 SIC.autumn, SIC.winter, SIC.latewinter, WCT, MFM, cohort size). Only SIC.winter was 493 correlated strongly enough (r > 0.7, (99)) with two other closely related sea ice metrics (SIEA 494 495 and SIC.latewinter) so that we decided to drop this covariate from analyses. Remaining covariates were scaled to improve model convergence and facilitate parameters comparisons. 496 Covariates were tested in isolation and considered influential when the 95% confidence interval 497 (CI) of the effect size did not overlap 0 and when accounting for  $\geq 20\%$  of the temporal 498 variation in survival rates (23). The support for a linear trend in juvenile survival was assessed 499 using ANODEV (23). The effect of individual mass on survival was investigated using 500 Likelihood Ratio Tests (LRT). 501

502

Homoscedasticity and normality assumptions were assessed where appropriate. Values are reported as means  $\pm$  SD unless mentioned otherwise. Statistical analyses other than the CMR modeling were conducted in R version 4.3.2 (*100*).

506

#### 507 **References:**

- A. Constable, J. Melbourne-Thomas, S. Corney, K. Arrigo, D. Barnes, N. Bindoff, P. Boyd, A.
   Brandt, D. Costa, A. Davidson, H. Ducklow, L. Emmerson, M. Fukuchi, J. Gutt, M. Hindell, E.
   Hofmann, G. Hosie, T. Iida, P. Ziegler, Climate change and Southern Ocean ecosystems I: How
   changes in physical habitats directly affect marine biota. *Glob. Change Biol.* 20 (2014).
- S. L. Chown, R. I. Leihy, T. R. Naish, C. M. Brooks, P. Convey, B. J. Henley, A. N. Mackintosh,
   L. M. Phillips, M. C. K. Ii, S. M. Grant, Antarctic Climate Change and the Environment.
- C. M. Brooks, D. G. Ainley, P. A. Abrams, P. K. Dayton, R. J. Hofman, J. Jacquet, D. B. Siniff,
   Antarctic fisheries: factor climate change into their management. *Nature* 558, 177–180 (2018).
- M. C. Urban, G. Bocedi, A. P. Hendry, J.-B. Mihoub, G. Pe'er, A. Singer, J. R. Bridle, L. G.
   Crozier, L. De Meester, W. Godsoe, A. Gonzalez, J. J. Hellmann, R. D. Holt, A. Huth, K. Johst,
   C. B. Krug, P. W. Leadley, S. C. F. Palmer, J. H. Pantel, A. Schmitz, P. A. Zollner, J. M. J.
   Travis, Improving the forecast for biodiversity under climate change. *Science* 353, aad8466
   (2016).

5. H. Lee, K. Calvin, D. Dasgupta, G. Krinner, A. Mukherji, P. W. Thorne, C. Trisos, J. Romero, P. 521 Aldunce, K. Barrett, G. Blanco, W. W. L. Cheung, S. Connors, F. Denton, A. Diongue-Niang, D. 522 Dodman, M. Garschagen, O. Geden, B. Hayward, C. Jones, F. Jotzo, T. Krug, R. Lasco, Y.-Y. 523 Lee, V. Masson-Delmotte, M. Meinshausen, K. Mintenbeck, A. Mokssit, F. E. L. Otto, M. 524 Pathak, A. Pirani, E. Poloczanska, H.-O. Pörtner, A. Revi, D. C. Roberts, J. Roy, A. C. Ruane, J. 525 Skea, P. R. Shukla, R. Slade, A. Slangen, Y. Sokona, A. A. Sörensson, M. Tignor, D. Van 526 Vuuren, Y.-M. Wei, H. Winkler, P. Zhai, Z. Zommers, J.-C. Hourcade, F. X. Johnson, S. 527 528 Pachauri, N. P. Simpson, C. Singh, A. Thomas, E. Totin, P. Arias, M. Bustamante, I. Elgizouli, 529 G. Flato, M. Howden, C. Méndez-Vallejo, J. J. Pereira, R. Pichs-Madruga, S. K. Rose, Y. Saheb, R. Sánchez Rodríguez, D. Ürge-Vorsatz, C. Xiao, N. Yassaa, A. Alegría, K. Armour, B. Bednar-530 Friedl, K. Blok, G. Cissé, F. Dentener, S. Eriksen, E. Fischer, G. Garner, C. Guivarch, M. 531 Haasnoot, G. Hansen, M. Hauser, E. Hawkins, T. Hermans, R. Kopp, N. Leprince-Ringuet, J. 532 Lewis, D. Ley, C. Ludden, L. Niamir, Z. Nicholls, S. Some, S. Szopa, B. Trewin, K.-I. Van Der 533 Wijst, G. Winter, M. Witting, A. Birt, M. Ha, J. Romero, J. Kim, E. F. Haites, Y. Jung, R. 534 535 Stavins, A. Birt, M. Ha, D. J. A. Orendain, L. Ignon, S. Park, Y. Park, A. Reisinger, D.

536 Cammaramo, A. Fischlin, J. S. Fuglestvedt, G. Hansen, C. Ludden, V. Masson-Delmotte, J. B. R.

- 537 Matthews, K. Mintenbeck, A. Pirani, E. Poloczanska, N. Leprince-Ringuet, C. Péan, "IPCC,
- 538 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to
- the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing]
- 540 Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland." (Intergovernmental Panel on
- 541 Climate Change (IPCC), 2023); https://doi.org/10.59327/IPCC/AR6-9789291691647.
- 6. R. L. Fogt, G. J. Marshall, The Southern Annular Mode: Variability, trends, and climate impacts
  across the Southern Hemisphere. *WIREs Clim. Change* 11, e652 (2020).
- 544 7. C. L. Parkinson, D. J. Cavalieri, Antarctic sea ice variability and trends, 1979–2010. *The*545 *Cryosphere* 6, 871–880 (2012).
- F. Olivier, J. A. van Franeker, J. C. S. Creuwels, E. J. Woehler, Variations of snow petrel
   breeding success in relation to sea-ice extent: detecting local response to large-scale processes?
   *Polar Biol.* 28, 687–699 (2005).
- 549 9. S. Descamps, A. Tarroux, S.-H. Lorentsen, O. P. Love, Ø. Varpe, N. G. Yoccoz, Large-scale
  550 oceanographic fluctuations drive Antarctic petrel survival and reproduction. *Ecography* 39, 496–
  551 505 (2016).
- 552 10. S. Labrousse, A. D. Fraser, M. Sumner, F. Le Manach, C. Sauser, I. Horstmann, E. Devane, K.
  553 Delord, S. Jenouvrier, C. Barbraud, Landfast ice: a major driver of reproductive success in a polar
  554 seabird. *Biol. Lett.* 17, 20210097 (2021).
- 555 11. D. T. Iles, H. Lynch, R. Ji, C. Barbraud, K. Delord, S. Jenouvrier, Sea ice predicts long-term
  556 trends in Adélie penguin population growth, but not annual fluctuations: Results from a range557 wide multiscale analysis. *Glob. Change Biol.* 26, 3788–3798 (2020).
- 12. B. Şen, C. Che-Castaldo, K. M. Krumhardt, L. Landrum, M. M. Holland, M. A. LaRue, M. C.
  Long, S. Jenouvrier, H. J. Lynch, Spatio-temporal transferability of environmentally-dependent
  population models: Insights from the intrinsic predictabilities of Adélie penguin abundance time
  series. *Ecol. Indic.* 150, 110239 (2023).
- 13. W. R. Fraser, W. Z. Trivelpiece, *Factors Controlling the Distribution of Seabirds: Winter- Summer Heterogeneity in the Distribution of Adélie Penguin Populations* (American Geophysical
  Union, Washington, D. C., 1996; http://doi.wiley.com/10.1029/AR070)vol. 70 of *Antarctic Research Series*.
- 14. R. C. Smith, D. Ainley, K. Baker, E. Domack, S. Emslie, B. Fraser, J. Kennett, A. Leventer, E.

567		Mosley-Thompson, S. Stammerjohn, M. Vernet, Marine Ecosystem Sensitivity to Climate
568		Change: Historical observations and paleoecological records reveal ecological transitions in the
569		Antarctic Peninsula region. BioScience 49, 393-404 (1999).
570	15.	D. Ainley, J. Russell, S. Jenouvrier, E. Woehler, P. O. Lyver, W. R. Fraser, G. L. Kooyman,
571		Antarctic penguin response to habitat change as Earth's troposphere reaches 2°C above
572		preindustrial levels. Ecol. Monogr. 80, 49-66 (2010).
573	16.	M. A. Cimino, H. J. Lynch, V. S. Saba, M. J. Oliver, Projected asymmetric response of Adélie
574		penguins to Antarctic climate change. Sci. Rep. 6, 28785 (2016).
575	17.	C. Che-Castaldo, S. Jenouvrier, C. Youngflesh, K. T. Shoemaker, G. Humphries, P. McDowall,
576		L. Landrum, M. M. Holland, Y. Li, R. Ji, H. J. Lynch, Pan-Antarctic analysis aggregating spatial
577		estimates of Adélie penguin abundance reveals robust dynamics despite stochastic noise. Nat.
578		<i>Commun.</i> <b>8</b> , 832 (2017).
579	18.	H. J. Lynch, M. A. LaRue, First global census of the Adélie Penguin. The Auk 131, 457-466
580		(2014).
581	19.	G. R. W. Humphries, C. Che-Castaldo, P. J. Bull, G. Lipstein, A. Ravia, B. Carrión, T. Bolton, A.
582		Ganguly, H. J. Lynch, Predicting the future is hard and other lessons from a population time
583		series data science competition. Ecol. Inform. 48, 1-11 (2018).
584	20.	E. J. Talis, C. Che-Castaldo, B. Şen, K. Krumhardt, H. J. Lynch, Variability, skipped breeding
585		and heavy-tailed dynamics in an Antarctic seabird. J. Anim. Ecol. 91, 2437-2450 (2022).
586	21.	D. Iles, S. Jenouvrier, "Projected population consequences of climate change" in Effects of
587		Climate Change on Birds (Oxford University Press, 2019;
588		https://oxford.university press cholarship.com/view/10.1093/oso/9780198824268.001.0001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-10001/oso-100001/oso-100000000000000000000000000000000000
589		9780198824268-chapter-12), pp. 147–164.
590	22.	M. Frederiksen, M. P. Harris, S. Wanless, Inter-population variation in demographic parameters:
591		a neglected subject? Oikos 111, 209-214 (2005).
592	23.	V. Grosbois, O. Gimenez, JM. Gaillard, R. Pradel, C. Barbraud, J. Clobert, A. P. Møller, H.
593		Weimerskirch, Assessing the impact of climate variation on survival in vertebrate populations.
594		<i>Biol. Rev.</i> <b>83</b> , 357–399 (2008).
595	24.	D. Irons, A. Petersen, T. Anker-Nilssen, Y. Artukhin, R. Barrett, D. Boertmann, M. Gavrilo, H.
596		Gilchrist, E. Hansen, M. Hario, K. Kuletz, M. Mallory, F. Merkel, A. Mosbech, A. Labansen, B.

- 597 Olsen, H. Österblom, J. Reid, G. Robertson, H. Strøm, *Circumpolar Seabird Monitoring Plan*598 (2015).
- 599 25. A. Nicol-Harper, C. P. Doncaster, G. M. Hilton, K. A. Wood, T. H. G. Ezard, Conservation
  600 implications of a mismatch between data availability and demographic impact. *Ecol. Evol.* 13,
  601 e10269 (2023).
- 602 26. B.-E. Sæther, Ø. Bakke, Avian Life History Variation and Contribution of Demographic Traits to
  603 the Population Growth Rate. *Ecology* 81, 642–653 (2000).
- C. Southwell, L. Emmerson, J. McKinlay, K. Newbery, A. Takahashi, A. Kato, C. Barbraud, K.
   DeLord, H. Weimerskirch, Spatially Extensive Standardized Surveys Reveal Widespread, Multi Decadal Increase in East Antarctic Adélie Penguin Populations. *PLOS ONE* 10, e0139877 (2015).

28. P. J. Kappes, K. M. Dugger, A. Lescroël, D. G. Ainley, G. Ballard, K. J. Barton, P. O. Lyver, P.
R. Wilson, Age-related reproductive performance of the Adélie penguin, a long-lived seabird
exhibiting similar outcomes regardless of individual life-history strategy. *J. Anim. Ecol.* 90, 931–
942 (2021).

- 611 29. J. T. Hinke, S. G. Trivelpiece, W. Z. Trivelpiece, Adélie penguin (Pygoscelis adeliae) survival
  612 rates and their relationship to environmental indices in the South Shetland Islands, Antarctica.
  613 *Polar Biol.* 37, 1797–1809 (2014).
- 30. J. T. Hinke, S. G. Trivelpiece, W. Z. Trivelpiece, Variable vital rates and the risk of population
  declines in Adélie penguins from the Antarctic Peninsula region. *Ecosphere* 8, e01666 (2017).
- 616 31. L. Emmerson, C. Southwell, Environment-triggered demographic changes cascade and compound
  617 to propel a dramatic decline of an Antarctic seabird metapopulation. *Glob. Change Biol.* 28
  618 (2022).
- 32. S. Jenouvrier, C. Barbraud, H. Weimerskirch, Sea ice affects the population dynamics of Adélie
  penguins in Terre Adélie. *Polar Biol.* 29, 413–423 (2006).
- 33. T. Ballerini, G. Tavecchia, S. Olmastroni, F. Pezzo, S. Focardi, Nonlinear effects of winter sea ice
  on the survival probabilities of Adélie penguins. *Oecologia* 161, 253–265 (2009).
- 623 34. L. Emmerson, C. Southwell, Ade lie penguin survival: age structure, temporal variability and624 environmental influences. 15 (2011).
- 35. A. E. Schmidt, A. Lescroël, S. Lisovski, M. Elrod, D. Jongsomjit, K. M. Dugger, G. Ballard, Sea
  ice concentration decline in an important Adélie penguin molt area. *Proc. Natl. Acad. Sci.* 120,

627 e2306840120 (2023).

- 36. S. R. Rintoul, S. L. Chown, R. M. DeConto, M. H. England, H. A. Fricker, V. Masson-Delmotte,
  T. R. Naish, M. J. Siegert, J. C. Xavier, Choosing the future of Antarctica. *Nature* 558, 233–241
  (2018).
- 37. B. Reid, An interpretation of the age structure and breeding status of an Adélie penguin
  population. *Notornis Q. J. Ornithol. Soc. N. Z.*, 193 (1968).
- 633 38. D. G. Ainley, *The Adélie Penguin: Bellwether of Climate Change* (Columbia University Press,
  634 New York, 2002).
- 39. J. A. Badia-Boher, J. Real, J. L. Riera, F. Bartumeus, F. Parés, J. M. Bas, A. Hernández-Matías,
  Joint estimation of survival and dispersal effectively corrects the permanent emigration bias in
  mark-recapture analyses. *Sci. Rep.* 13, 6970 (2023).
- 40. K. M. Dugger, G. Ballard, D. G. Ainley, K. J. Barton, Effects of Flipper Bands on Foraging
  Behavior and Survival of Adélie Penguins (Pygoscelis Adeliae). *The Auk* 123, 858–869 (2006).
- 41. C. Saraux, C. Le Bohec, J. M. Durant, V. A. Viblanc, M. Gauthier-Clerc, D. Beaune, Y.-H. Park,
  N. G. Yoccoz, N. C. Stenseth, Y. Le Maho, Reliability of flipper-banded penguins as indicators
  of climate change. *Nature* 469, 203–206 (2011).
- 42. W. R. Fraser, WayneZ. Trivelpiece, D. G. Ainley, SusanG. Trivelpiece, Increases in Antarctic
  penguin populations: reduced competition with whales or a loss of sea ice due to environmental
  warming? *Polar Biol.* 11, 525–531 (1992).
- 43. P. Wilson, D. Ainley, N. Nur, S. Jacobs, K. Barton, G. Ballard, J. Comiso, Adélie penguin
  population change in the pacific sector of Antarctica: relation to sea-ice extent and the Antarctic
  Circumpolar Current. *Mar. Ecol. Prog. Ser.* 213, 301–309 (2001).
- 44. J. T. Hinke, K. Salwicka, S. G. Trivelpiece, G. M. Watters, W. Z. Trivelpiece, Divergent
  responses of Pygoscelis penguins reveal a common environmental driver. *Oecologia* 153, 845–
  855 (2007).
- 45. J. T. Hinke, G. M. Watters, C. S. Reiss, J. A. Santora, M. M. Santos, Acute bottlenecks to the
  survival of juvenile Pygoscelis penguins occur immediately after fledging. *Biol. Lett.* 16,
  20200645 (2020).
- 46. S. L. Cox, M. Authier, F. Orgeret, H. Weimerskirch, C. Guinet, High mortality rates in a juvenile
  free-ranging marine predator and links to dive and forage ability. *Ecol. Evol.* 10, 410–430 (2020).

- 47. J.-M. Gaillard, C. Toïgo, Temporal variation in fitness component and population dynamics of
  large herbivores. 31, 367–393 (2000).
- 48. B. Naef-Daenzer, M. U. Grüebler, Post-fledging survival of altricial birds: ecological
  determinants and adaptation. *J. Field Ornithol.* 87, 227–250 (2016).
- 49. C. L. McIntyre, M. W. Collopy, M. S. Lindberg, Survival Probability and Mortality of Migratory
  Juvenile Golden Eagles from Interior Alaska. J. Wildl. Manag. 70, 717–722 (2006).
- 50. R. Massom, P. Reid, S. Stammerjohn, B. Raymond, A. Fraser, S. Ushio, Change and Variability
  in East Antarctic Sea Ice Seasonality, 1979/80–2009/10. *PLoS ONE* 8, e64756 (2013).
- 51. C. Eayrs, D. Holland, D. Francis, T. Wagner, R. Kumar, X. Li, Understanding the Seasonal Cycle
  of Antarctic Sea Ice Extent in the Context of Longer-Term Variability. *Rev. Geophys.* 57, 1037–
  1064 (2019).
- 52. M. A. Morales Maqueda, A. J. Willmott, N. R. T. Biggs, Polynya Dynamics: a Review of
  Observations and Modeling. *Rev. Geophys.* 42 (2004).
- 53. K. R. Arrigo, G. L. van Dijken, A. L. Strong, Environmental controls of marine productivity hot
  spots around Antarctica. *J. Geophys. Res. Oceans* 120, 5545–5565 (2015).
- 54. M. R. Enstipp, C.-A. Bost, C. Le Bohec, N. Chatelain, H. Weimerskirch, Y. Handrich, The early
  life of king penguins: ontogeny of dive capacity and foraging behaviour in an expert diver. *J. Exp. Biol.* 224, jeb242512 (2021).
- 55. D. G. Ainley, G. Ballard, B. J. Karl, K. M. Dugger, Leopard seal predation rates at penguin
  colonies of different size. *Antarct. Sci.* 17, 335–340 (2005).
- 56. A. Takahashi, M. Ito, K. Nagai, J. Thiebot, H. Mitamura, T. Noda, P. Trathan, T. Tamura, Y.
  Watanabe, Migratory movements and winter diving activity of Adélie penguins in East
  Antarctica. *Mar. Ecol. Prog. Ser.* 589, 227–239 (2018).
- 57. J. Clarke, K. Kerry, C. Fowler, R. Lawless, S. Eberhard, R. Murphy, Post-fledging and winter
  migration of Adélie penguins Pygoscelis adeliae in the Mawson region of East Antarctica. *Mar. Ecol. Prog. Ser.* 248, 267–278 (2003).
- 58. D. G. Ainley, G. Ballard, K. J. Barton, B. J. Karl, G. H. Rau, C. A. Ribic, P. R. Wilson, Spatial
  and Temporal Variation of Diet Within a Presumed Metapopulation Of Adélie Penguins. *The Condor* 105, 95–106 (2003).

- 59. M. Beaulieu, A. Dervaux, A.-M. Thierry, D. Lazin, Y. Le Maho, Y. Ropert-Coudert, M. Spée, T.
  Raclot, A. Ancel, When sea-ice clock is ahead of Adélie penguins' clock. *Funct. Ecol.* 24, 93–
  102 (2010).
- 689 60. Y. Y. Watanabe, K. Ito, N. Kokubun, A. Takahashi, Foraging behavior links sea ice to breeding
  690 success in Antarctic penguins. *Sci. Adv.* 6, eaba4828 (2020).
- 691 61. C. L. Guen, A. Kato, B. Raymond, C. Barbraud, M. Beaulieu, C.-A. Bost, K. Delord, A. J. J.
  692 MacIntosh, X. Meyer, T. Raclot, M. Sumner, A. Takahashi, J.-B. Thiebot, Y. Ropert-Coudert,
  693 Reproductive performance and diving behaviour share a common sea-ice concentration optimum
  694 in Adélie penguins (Pygoscelis adeliae). *Glob. Change Biol.* 24, 5304–5317 (2018).
- 695 62. D. G. Ainley, K. M. Dugger, M. L. Mesa, G. Ballard, K. J. Barton, S. Jennings, B. J. Karl, A.
  696 Lescroël, P. O. Lyver, A. Schmidt, P. Wilson, Post-fledging survival of Adélie penguins at
  697 multiple colonies: chicks raised on fish do well. *Mar. Ecol. Prog. Ser.* 601, 239–251 (2018).
- 63. E. Chapman, E. Hofmann, D. Patterson, C. Ribic, W. Fraser, Marine and terrestrial factors
  affecting Adélie-penguin Pygoscelis adeliae chick growth and recruitment off the western
  Antarctic Peninsula. *Mar. Ecol. Prog. Ser.* 436, 273–289 (2011).
- 64. M. A. Cimino, J. A. Conroy, E. Connors, J. Bowman, A. Corso, H. Ducklow, W. Fraser, A.
  Friedlaender, H. H. Kim, G. D. Larsen, C. Moffat, R. Nichols, L. Pallin, D. Patterson-Fraser, D.
  Roberts, M. Roberts, D. K. Steinberg, P. Thibodeau, R. Trinh, O. Schofield, S. Stammerjohn,
  Long-term patterns in ecosystem phenology near Palmer Station, Antarctica, from the perspective
  of the Adélie penguin. *Ecosphere* 14, e4417 (2023).
- 65. B.-E. Sæther, T. Coulson, V. Grøtan, S. Engen, R. Altwegg, K. B. Armitage, C. Barbraud, P. H.
  Becker, D. T. Blumstein, F. S. Dobson, M. Festa-Bianchet, J.-M. Gaillard, A. Jenkins, C. Jones,
  M. A. C. Nicoll, K. Norris, M. K. Oli, A. Ozgul, H. Weimerskirch, How Life History Influences
  Population Dynamics in Fluctuating Environments. *Am. Nat.* 182, 743–759 (2013).
- 66. C. Barbraud, K. Delord, C. A. Bost, A. Chaigne, C. Marteau, H. Weimerskirch, Population trends
  of penguins in the French Southern Territories. *Polar Biol.* 43, 835–850 (2020).
- 712 67. Y. Ropert-Coudert, A. Kato, K. Shiomi, C. Barbraud, F. Angelier, K. Delord, T. Poupart, P.
  713 Koubbi, T. Raclot, Two Recent Massive Breeding Failures in an Adélie Penguin Colony Call for
  714 the Creation of a Marine Protected Area in D'Urville Sea/Mertz. *Front. Mar. Sci.* 5 (2018).
- 68. J. Forcada, P. N. Trathan, K. Reid, E. J. Murphy, J. P. Croxall, Contrasting population changes in
  sympatric penguin species in association with climate warming. *Glob. Change Biol.* 12, 411–423

717 (2006).

- 69. W. Z. Trivelpiece, J. T. Hinke, A. K. Miller, C. S. Reiss, S. G. Trivelpiece, G. M. Watters,
  Variability in krill biomass links harvesting and climate warming to penguin population changes
  in Antarctica. *Proc. Natl. Acad. Sci.* 108, 7625–7628 (2011).
- 70. H. J. Lynch, R. Naveen, P. N. Trathan, W. F. Fagan, Spatially integrated assessment reveals
  widespread changes in penguin populations on the Antarctic Peninsula. *Ecology* 93, 1367–1377
  (2012).
- 71. M. Wethington, C. Flynn, A. Borowicz, H. J. Lynch, Adélie penguins north and east of the
  'Adélie gap' continue to thrive in the face of dramatic declines elsewhere in the Antarctic
  Peninsula region. *Sci. Rep.* 13, 2525 (2023).
- 727 72. J. Turner, G. J. Marshall, K. Clem, S. Colwell, T. Phillips, H. Lu, Antarctic temperature
  728 variability and change from station data. *Int. J. Climatol.* 40, 2986–3007 (2020).
- 729 73. N. Lecomte, The great melt will shape unprotected ecosystems. *Nature* **620**, 499–500 (2023).

730 74. K. M. Dugger, D. G. Ainley, P. O. Lyver, K. Barton, G. Ballard, Survival differences and the
rstation effect of environmental instability on breeding dispersal in an Adélie penguin meta-population. *Proc. Natl. Acad. Sci.* 107, 12375–12380 (2010).

- 733 75. V. Morandini, K. M. Dugger, A. E. Schmidt, A. Varsani, A. Lescroël, G. Ballard, P. O. Lyver, K.
  734 Barton, D. G. Ainley, Sex-specific recruitment rates contribute to male-biased sex ratio in Adélie
  735 penguins. *Ecol. Evol.* 14, e10859 (2024).
- 736 76. M. Gauthier-Clerc, J. P. Gendner, C. A. Ribic, W. R. Fraser, E. J. Woehler, S. Descamps, C.
  737 Gilly, C. Le Bohec, Y. Le Maho, Long-term effects of flipper bands on penguins. *Proc. R. Soc. B*738 *Biol. Sci.* 271, S423–S426 (2004).
- 739 77. K. Kerry, J. Clarke, G. Else, The use of an automated weighing and recording system for the
  740 study of the biology of Adélie penguins. *Proc NIPR Symp Polar Biol* 6, 62–75 (1993).
- 741 78. S. Olmastroni, S. Corsolini, F. Pezzo, S. Focardi, K. Kerry, The first five years of the Italian742 Australian joint programme on the Adélie Penguin: An overview. *Ital. J. Zool.* 67, 141–145
  743 (2000).
- 744 79. A. Lescroël, G. Ballard, D. Grémillet, M. Authier, D. G. Ainley, Antarctic Climate Change:
  745 Extreme Events Disrupt Plastic Phenotypic Response in Adélie Penguins. *PLOS ONE* 9, e85291
  746 (2014).

- 747 80. J.-P. Gendner, M. Gauthier-Clerc, C. Le Bohec, S. Descamps, Y. Le Maho, A new application for
  748 transponders in studying penguins. *J. Field Ornithol.* 76, 138–142 (2005).
- 81. G. Bardon, R. Cristofari, A. Winterl, T. Barracho, M. Benoiste, C. Ceresa, N. Chatelain, J.
  Courtecuisse, F. A. N. Fernandes, M. Gauthier-Clerc, J.-P. Gendner, Y. Handrich, A. Houstin, A.
  Krellenstein, N. Lecomte, C.-E. Salmon, E. Trucchi, B. Vallas, E. M. Wong, D. P. Zitterbart, C.
  Le Bohec, RFIDeep: Unfolding the potential of deep learning for radio-frequency identification. *Methods Ecol. Evol.* 14, 2814–2826 (2023).
- 82. N. Dehnhard, K. Ludynia, M. Poisbleau, L. Demongin, P. Quillfeldt, Good Days, Bad Days:
  Wind as a Driver of Foraging Success in a Flightless Seabird, the Southern Rockhopper Penguin.
- 756 *PLOS ONE* **8**, e79487 (2013).
- 757 83. V. Afanasyev, S. V. Buldyrev, M. J. Dunn, J. Robst, M. Preston, S. F. Bremner, D. R. Briggs, R.
- 758 Brown, S. Adlard, H. J. Peat, Increasing Accuracy: A New Design and Algorithm for
- Automatically Measuring Weights, Travel Direction and Radio Frequency Identification (RFID)
   of Penguins. *PLOS ONE* 10, e0126292 (2015).
- 84. S. Robinson, A. Chiaradia, M. A. Hindell, The effect of body condition on the timing and success
  of breeding in Little Penguins Eudyptula minor. *Ibis* 147, 483–489 (2005).
- 763 85. E. Barreau, Y. Ropert-Coudert, K. Delord, C. Barbraud, A. Kato-Ropert, Scale matters: sea ice
  764 and breeding success of Adélie penguins. *Polar Biol.* 42, 1405–1410 (2019).
- 86. J. Lebreton, J. D. Nichols, R. J. Barker, R. Pradel, J. A. Spendelow, "Chapter 3 Modeling
  Individual Animal Histories with Multistate Capture–Recapture Models" in *Advances in Ecological Research* (Elsevier, 2009;
- 768 https://linkinghub.elsevier.com/retrieve/pii/S0065250409004036)vol. 41, pp. 87–173.
- 769 87. G. C. White, K. P. Burnham, Program MARK: survival estimation from populations of marked
  770 animals. *Bird Study* 46, S120–S139 (1999).
- 88. R. Choquet, J.-D. Lebreton, O. Gimenez, A.-M. Reboulet, R. Pradel, U-CARE: Utilities for
  performing goodness of fit tests and manipulating CApture–REcapture data. *Ecography* 32,
  1071–1074 (2009).
- 89. R. Pradel, C. M. A. Wintrebert, O. Gimenez, A Proposal for a Goodness-of-Fit Test to the
  Arnason-Schwarz Multisite Capture-Recapture Model. *Biometrics* 59, 43–53 (2003).
- 90. R. Kwok, J. C. Comiso, Southern Ocean Climate and Sea Ice Anomalies Associated with the

777 Southern Oscillation. J. Clim. 15, 487–501 (2002).

- 91. C. Sauser, K. Delord, C. Barbraud, Demography of cape petrels in response to environmental
  changes. *Popul. Ecol.* 65, 25–37 (2021).
- 780 92. F. Abadi, C. Barbraud, O. Gimenez, Integrated population modeling reveals the impact of climate
  781 on the survival of juvenile emperor penguins. *Glob. Change Biol.* 23, 1353–1359 (2017).
- 93. C. Sauser, K. Delord, C. Barbraud, Increased sea ice concentration worsens fledging condition
  and juvenile survival in a pagophilic seabird, the snow petrel. *Biol. Lett.* 14, 20180140 (2018).
- 94. J.-B. Thiebot, Y. Ropert-Coudert, T. Raclot, T. Poupart, A. Kato, A. Takahashi, Adélie penguins'
  extensive seasonal migration supports dynamic Marine Protected Area planning in Antarctica. *Mar. Policy* 109, 103692 (2019).
- 95. W. Meier, F. Fetterer, A. Windnagel, S. Stewart, NOAA/NSIDC Climate Data Record of Passive
  Microwave Sea Ice Concentration, Version 4, NSIDC (2021); https://doi.org/10.7265/EFMZ2T65.
- 96. C. Sauser, P. Blévin, O. Chastel, G. W. Gabrielsen, S. A. Hanssen, E. Lorentzen, B. Moe, S.
  Moreau, K. Sagerup, S. Descamps, Ice type matters: impacts of landfast and drift ice on body
  condition in a high Arctic seabird community. *Front. Mar. Sci.* 11, 1326658 (2024).
- 97. T. J. Maness, D. J. Anderson, Predictors of Juvenile Survival in Birds Predictores de la
  Supervivencia de Aves Juveniles. *Ornithol. Monogr.* 78, 1–55 (2013).
- 98. C. Horswill, J. Matthiopoulos, J. A. Green, M. P. Meredith, J. Forcada, H. Peat, M. Preston, P. N.
  Trathan, N. Ratcliffe, Survival in macaroni penguins and the relative importance of different
  drivers: individual traits, predation pressure and environmental variability. *J. Anim. Ecol.* 83, 1057–1067 (2014).
- 99. C. F. Dormann, J. Elith, S. Bacher, C. Buchmann, G. Carl, G. Carré, J. R. G. Marquéz, B. Gruber,
  B. Lafourcade, P. J. Leitão, T. Münkemüller, C. McClean, P. E. Osborne, B. Reineking, B.
  Schröder, A. K. Skidmore, D. Zurell, S. Lautenbach, Collinearity: a review of methods to deal
  with it and a simulation study evaluating their performance. *Ecography* 36, 27–46 (2013).
- 803 100. R Core Team (2023). \_R: A Language and Environment for Statistical Computing\_. R
  804 Foundation for Statistical Computing, Vienna, Austria. <a href="https://www.R-project.org/">https://www.R-project.org/</a>.
- 805 101. K. Matsuoka, A. Skoglund, G. Roth, J. de Pomereu, H. Griffiths, R. Headland, B. Herried, K.
  806 Katsumata, A. Le Brocq, K. Licht, F. Morgan, P. D. Neff, C. Ritz, M. Scheinert, T. Tamura, A.

Van de Putte, M. van den Broeke, A. von Deschwanden, C. Deschamps-Berger, B. Van
Liefferinge, S. Tronstad, Y. Melvær, Quantarctica, an integrated mapping environment for
Antarctica, the Southern Ocean, and sub-Antarctic islands. *Environ. Model. Softw.* 140, 105015
(2021).

811

#### **812** Acknowledgements:

We are deeply grateful to all the members of Project 137, including Benjamin Friess, Yvon Le Maho (former PI of 137-ECOPHY), Victor Planas-Bielsa, Claire Saraux, and all the wintering and summering field teams since the inception of this project in the field in 2005. We would also like to thank all the members of the missions in Dumont D'Urville since then, and the French Polar Institute-IPEV logistics team in Dumont d'Urville for their important and continuous support in the field.

819

#### 820 Funding:

This study was supported by the Institut Polaire Français Paul-Emile Victor (IPEV) within the framework of the Project 137-ANTAVIA, by the Centre Scientifique de Monaco with additional support from the LIA-647 and RTPI-NUTRESS (CSM/CNRS-UNISTRA), by the Centre National de la Recherche Scientifique (CNRS) through the Programme Zone Atelier Antarctique et Terres Australes (ZATA), and by the Deutsche Forschungsgemeinschaft (DFG) grants FA336/5-1 and ZI1525/3-1 in the framework of the priority program "Antarctic research with comparative investigations in Arctic ice areas".

828

#### 829 Author contribution:

830 Author contributions follow the CRediT guidelines. In each section, authors are listed by order

831 of appearance in the author list.

- 832
- 833 Conceptualization: TB, NL, CLB
- 834 Methodology: TB, RCH, AH, MBE, RCR, TR, NC, JC, MBU, NL, CLB
- 835 Software: TB, GB, RCH
- 836 Validation: TB, NL, CLB
- 837 Formal analysis: TB, GB, RCH, NL
- 838 Investigation: All authors
- 839 Resources: CLB
- 840 Data Curation: TB, GB, CCC, CLB

- 841 Writing original draft preparation: TB, NL, CLB
- 842 Writing review and editing: All authors
- 843 Visualization: TB
- 844 Supervision: NL, CLB
- 845 Project administration: CLB
- 846 Funding acquisition: DPZ, NL, CLB
- 847

**848 Competing interests:** 

- 849 We declare no competing interests.
- 850
- 851 Data and material availability:
- 852 Data and codes will be made available upon acceptance of the manuscript.
- 853
- **Figures and Tables:**



**Fig. 1.** Overview of the Adélie penguin individual monitoring system operating in Pointe Géologie archipelago (Adélie Land) and location of similar monitoring sites across Antarctica still active as of 2024. (A) Main panel: Satellite imagery of Pointe Géologie archipelago (Pléiades Neo, Airbus DS 2021, 12/10/2021) with locations of the mobile RFID-antennas deployment sites (red), undisturbed control colonies (blue) and the main study colony (yellow). Top-left inset: location of study sites across Antarctica where longitudinal monitoring of marked Adélie penguins is still carried out (ADMI: Admiralty Bay, BECH: Béchervaise Island, EDMO: Edmonson Point, PGEO: Pointe Géologie, ROSS: Ross Island (4 monitored colonies including one on neighboring Beaufort Island)). **B**) Study colony in Pointe Géologie where all fledged chicks have been implanted with RFID-tags since 2007. (**C**) Penguins can only access or exit through two RFID-equipped passageways (only one shown here



Fig. 2. Summary of the six locations around Antarctica where Adélie penguin juvenile survival ( $\phi$ , 0-2 years old) has been quantified. Among them, only three have investigated temporal variations. A) Survival of juvenile Adélie penguins from Pointe Géologie archipelago, Adélie Land (PGEO, electronic tagging, 2007-2020). The dots and intervals represent annual survival probabilities  $\pm$  SE. The regression line and associated SE are from a linear regression (p = 0.036,  $R^2 = 0.37$ ); B) Adélie penguin abundance (number of breeding pairs) at Pointe Géologie Archipelago (2007-2019, data reproduced from (*66*)); C-D) Juvenile survival (1982-2000 aluminum bands, 1998-2011 stainless steel bands) and abundance (1982-2011) at Admiralty Bay, King George Island, Western Antarctic Peninsula (ADMI, data reproduced from (*29*, *30*)); E-F) Juvenile survival (electronic tagging, 1992-2015) and abundance (1992-2020) at Béchervaise Island, Mac. Robertson Land (BECH, data reproduced from (*31*)); G)Only mean estimates were available for three populations in the Ross Sea: Cape Hallett, Victoria Land (HALL, metal bands, 1959-1962, data extracted from (*37*)), Cape Crozier, Ross Island (CROZ, aluminum bands, 1961-1976, data extracted from (*38*)), Edmondson Point, Victoria Land (EDMO, electronic tagging, 1994-2002, data extracted from (*33*)). Survival estimates from HALL and to a lesser extent CROZ should be compared to other estimates with caution because they were not computed using modern capture-recapture techniques accounting for imperfect detection as in the other studies. Juvenile survival time series were fitted with linear regressions when authors reported a linear trend (**A**,**C**,**E**). Abundance time series were fitted with Generalized Additive Models (GAM)





Fig. 3. Relationships between the survival ( $\phi$ ) of juvenile Adélie penguins from Pointe Géologie archipelago (Adélie Land, East Antarctica) and several sea ice metrics within 200 kilometers of the natal colony for the month of April. A) Average sea ice concentration (SIC), **B**) Fraction of the total ice area (15-100% SIC) covered by the marginal ice zone (15-80% SIC), **C**) Fraction of total ice area (15-100% SIC) covered by dense ice (80-100% SIC). **D**) Time series (2007-2021) of the fraction of total ice area (15-100% SIC) covered by dense ice (80-100% SIC). Black thick lines and associated SE (gray shadow) on each panel are from linear regressions (respectively: p = 0.009,  $R^2 = 0.51$ ; p < 0.001,  $R^2 = 0.79$ ; p = 0.049,  $R^2 = 0.33$ ; p = 0.016,  $R^2 = 0.37$ ).



Fig. 4. Model-predicted relationship between Adélie penguin juvenile survival ( $\phi$ ) and body mass at tagging. Modeling was carried out using the program MARK (87). The predicted relationship is drawn from model 34 (Table 1). Green and violet dots represent individuals, and shaded areas represent probability density for each fate.

Table 1. Comparison of the effect of covariates on Adélie penguin juvenile survival probability. Abbreviations: AICc, Akaike Information Criterion corrected for small sample sizes ;  $\Delta$ AICc, AICc difference between current model and time-dependant model ; K, number of parameters ; Dev, Model deviance ; p ANODEV, p-value of the analysis of deviance ; %DEV, percentage of the time-dependent model's deviance explained by the inclusion of covariate (23) ; LRT, Likelihood Ratio Test. Covariates abbreviations: see footnote and Table S2.

Footnote Table 1: MIZFRAC.april (April fraction of total ice area between 135°E-145°E covered by the marginal ice zone (15-80% SIC)) ; DENSEFRAC.april (April fraction of total ice area between 135°E-145°E covered by dense ice (80-100% SIC)) ; SIC.april (April average SIC in area between 135°E-145°E) ; SIC.autumn (Mar-Apr average SIC in area between 135°E-145°E) ; WCT (Windchill temperatures for Dumont D'Urville in Apr-Sept) ; SAM (Southern Annular Mode, Mar-Sept) ; SIC.march (March average SIC in area between 135°E-145°E) ; SIC.latewinter (Aug-Sept average SIC in area between 110°E-135°E) ; SOI (Southern Oscillation Index, Mar-Sept) ; SIEA (Sea Ice Extent Anomalies, Mar-Sept), SIC.winter (May-Jul average SIC in area between 110°E-135°E).

							A		Individu mod	al cov. Iels	
Index	Candidate model (Φ 0-2)	AICc	ΔΑΙϹϲ	AICc Weights	К	Dev	p ANODEV	Slope [95%Cl]	%DEV	χ2 LRT	p LRT
Annua	l covariate models	6			-				-		
13	time	12044.7	0	1	75	2303.7	-	-	-	-	-
27	MIZFRAC.april	12068.2	23.4	0	62	2353.7	<0.001	-0.97 [-0.99 , -0.94]	78.8	-	-
28	DENSEFRAC.april	12160.1	115.3	0	62	2445.6	0.005	0.84 [0.82 , 0.86]	39.7	-	-
26	SIC.april	12160.8	116.1	0	62	2446.3	0.005	0.96 [0.93 , 0.99]	39.4	-	-
30	linear	12180.5	135.7	0	62	2466.0	0.016	-0.02 [-0.03 , -0.02]	31.1	-	-
22	SIC.autumn	12187.1	142.4	0	62	2472.6	0.023	0.92 [0.89 , 0.95]	28.2	-	-
29	MFM	12201.8	157.1	0	62	2487.3	0.050	0.03 [0.03 , 0.03]	22.0	-	-
20	WCT	12215.8	171.0	0	62	2501.3	0.099	0 [-0.03 , 0.03]	16.1	-	-
21	WCT <sup>2</sup>	12217.0	172.2	0	63	2500.4	0.095	-	16.4	-	-
25	SIC.march	12220.3	175.5	0	62	2505.8	0.124	0.75 [0.74 , 0.76]	14.2	-	-
17	SOI	12232.0	187.3	0	62	2525.7	0.340	0.03 [0 , 0.07]	5.7	-	-
19	SIEA	12241.5	196.8	0	62	2527.0	0.365	0.02 [0.02 , 0.03]	5.1	-	-
31	cohort size	12243.2	198.4	0	62	2528.6	0.400	0[-0.01,0.01]	4.4	-	-
23	SIC.winter	12244.6	199.9	0	62	2530.1	0.437	0.32 [0.31 , 0.33]	3.8	-	-
18	SAM	12245.2	200.5	0	62	2530.7	0.453	-0.02 [-0.07 , 0.04]	3.6	-	-
24	SIC.latewinter	12245.7	201.0	0	62	2531.2	0.467	-0.04 [-0.05 , -0.02]	3.4	-	-
15	constant	12265.8	221.1	0	68	2539.1	-	-	0.0	-	-
Individ	lual covariate mod	els									
35	mass + time	12031.6	0.0	0.998	76	11878.0	-	-	-	15.227	< 0.001
32	time	12044.7	13.2	0.001	75	11893.2	-	-	-	-	-
35	mass x time	12048.4	16.9	0.000	89	11868.3	-	-	-	24.935	0.0352
34	mass	12219.8	188.3	0.000	69	12080.6	-	-	-	48.014	< 0.001
33	constant	12265.8	234.3	0.000	68	12128.6	-	-	-	-	-

#### SUPPLEMENTARY MATERIAL:

#### **Supplementary Text**

## Manipulating capture histories to account for near-zero recapture probability at age 1 and for Goodness-Of-Fit (GOF) testing.

#### 1. Accounting for near-zero recapture probability at age 1

In capture-recapture modeling, the capture history (CH) of an individual is coded as a suite of numbers for each capture occasion. For example, if 0 = not seen; 1 = seen in state 1; and 2 = seen in state 2, the individual A presented in the table below was tagged in state 1 in year 2, not seen in year 3, seen in state 1 in year 4 and seen in state 2 in year 5.

Year/Capture occasion	1	2	3	4	5
Capture history of individual A	0	1	0	1	2

The CH of individual A can be written as: "01012". Because recapture probabilities in the year following tagging (year 3 in the case of individual A) were near 0 in our study (only one individual was detected at one year old over our entire study period, Fig. S3), we rewrote CH of all individuals by removing the year after tagging. For example, the CH of individual A was transformed from "01012" into "0112".

This allows accounting for the fact that with so few individuals seen again one year after fledging, it is not possible to estimate survival probability ( $\phi$ ) from age 0 to age 1 ( $\phi$  0 $\rightarrow$ 1). Instead, we estimated survival from age 0 to age 2 ( $\phi$  0 $\rightarrow$ 2). This is the typical "juvenile survival" in capture-recapture studies where individuals are not available for detection before two years after fledging. Under the specific (but untestable in this case) hypothesis that  $\phi$  0 $\rightarrow$ 1 =  $\phi$  1 $\rightarrow$ 2, survival from age 0 to age 1 ( $\phi$  0 $\rightarrow$ 1) can nonetheless be estimated as sqrt( $\phi$  0 $\rightarrow$ 2).

#### 2. GOF testing

There is no direct GOF test for age-dependent, multistate capture-recapture models (88). However, the presence of transients (individuals never seen again after tagging) is a common source of age-dependence in recapture probabilities. To quantify how much of the lack of fit of a time-dependent (JollyMoVe, JMV) model can be attributed to transients, a common strategy is to test the GOF of the JMV model to a dataset where transients are removed by replacing the first capture of each individual by 0 in their capture histories. For example, in the dataset where transients are removed, the CH of individual A would be "00012".

When the GOF of the dataset without transients is satisfactory, the original dataset (i.e. including transients) can be used but recapture probabilities must be specified as age-dependent in the starting model, therefore accounting for transience.



Fig. S1. Sectors of Antarctica used for large-scale sea ice analyses. Figure extracted from (7).



#### Fig. S2.

A mobile RFID detection unit being deployed on an Adélie penguin (*Pygoscelis adeliae*) passageway near a breeding colony in Adélie Land. This system allows the passive detection of RFID-tagged individuals away from the main study site where tagging is conducted. Detection data and batteries are stored in the acquisition box (1), to which an antenna is affixed and buried in the snow on penguin passageways (2, burying in process in this picture). To increase running time up to several weeks during the summer season, the system can be fitted with solar panels (3). © Gregory Tran (main panel), Téo Barracho (inset), Institut Polaire Français - French Polar Institute. Image rights: the person depicted in the picture is the first author.



Fig. S3.

Age of first return at the colony after fledging for RFID-tagged Adélie penguins in Pointe Géologie archipelago, Adélie land, Antarctica (2007-2020)



#### Fig. S4.

**Spatial windows considered for estimating the sea ice concentrations covariates for autumn (orange), and winter/late winter (green).** The yellow line depicts the average April sea ice extent and the blue line the average maximal sea ice extent (both over the 1981-2010 period). Map produced using Quantarctica (https://www.npolar.no/quantarctica/; (101))



Fig. S5.

**Time series of breeding productivity for the study colony (2011-2024).** Breeding productivity was calculated as the number of chicks fledged divided by the annual maximum number of adults inside the colony divided by two). This proxy for breeding productivity is a slight overestimation of the actual breeding productivity, since all individuals present at the colony may not be breeders. However, it still reflects population-scale breeding productivity accurately (Fig. S5). Breeding productivity may be superior to 1 in some years because Adélie penguins can raise two chicks (38)



Fig. S6.

The annual breeding productivity (chicks per pair) of the study colony correlates with that of the whole population of Pointe Géologie archipelago, Adélie Land, Antarctica (2011-2017, r = 0.99). Data for Pointe Géologie was extracted from (85). Breeding productivity may be superior to 1 in some years because Adélie penguins can raise two chicks (38).



**Fig. S7.** 

Annual maximal number of adults (a proxy for the number of breeding pairs, weekly surveys) in three neighboring Adélie penguin colonies in Pointe Géologie archipelago, Adélie Land, Antarctica. These three colonies include the study colony (Antavia) and two control colonies (Hall Fusée and Isabelle) where no monitoring activities are conducted except for weekly photo counts.



**Fig. S8.** 

Chick body mass at the time of RFID-tagging across cohorts (2007-2021) for Adélie penguins born in the study colony (Pointe Géologie archipelago, Adélie Land). Red dots represent annual means. Tagging consistently occurred 10-15 days prior to fledging, thus making mass at tagging a reliable proxy for fledging mass (62).



#### **Fig. S9.**

Estimated recapture probabilities of known-age RFID-tagged Adélie penguins in the local (A) and distant (B) states. States refer to spatial locations of colonies where individuals are detected. The local state includes the study colony and adjacent colonies (< 200 m), while the distant state includes colonies elsewhere in Pointe Géologie archipelago (see Fig. 1 in the main text). Error bars indicate  $\pm$  SE.



#### Fig. S10.

Annual increase in the proportion of dense sea ice (80-100% sea ice concentration) in the vicinity of Pointe Géologie archipelago (~200 km) between March and May of each year (2007-2021). The gray shaded area represents the month of April, where the increase in dense ice fraction is strongest. Daily sea ice concentration data over 25×25 km grids was downloaded from the NSDIC ((95), https://nsidc.org/data/g02202/versions/4).





#### Fig. S11.

Sea ice-scape at the end of April near Pointe Géologie archipelago in two years of contrasting survival probabilities for juvenile Adélie penguins (2008, high survival probability; 2018, low survival probability. MODIS satellite images were downloaded from the NASA Worldview application (https://worldview.earthdata.nasa.gov), part of the NASA Earth Observing System Data and Information System (EOSDIS).

#### Table S1:

**Results of Goodness-of-fit tests for the multistate capture-recapture dataset used to estimate Adélie penguin juvenile survival probability at Pointe-Géologie archipelago, Adélie Land, Antarctica**). df, Degrees of freedom; ĉ, Deviance inflation factor.

Test	χ2	pval	df	ĉ =χ2/Df				
A) Original dataset								
WBWA	2.469	0.7	4	0.617				
3G.SR	1827.802	< 0.001	12	152.317				
3G.Sm	1393.189	<0.001	35	39.805				
M.ITEC	18.37	0.005	6	3.062				
M.LTEC	6.831	0.2	5	1.366				
JMV model	3248.661	<0.001	62	52.398				
B) Dataset where 1st ca	pt. set to 0 (no trans	sients)						
WBWA	3.428	0.5	4	0.857				
3G.SR	14.744	0.324	13	1.134				
3G.Sm	47.702	0.0	31	1.539				
M.ITEC	12.541	0.051	6	0.805				
M.LTEC	6.015	0.3	5	0.919				
JMV model	84.429	0.017	59	1.431				

#### Table S2.

Annual covariates considered in the present study for explaining temporal variation in Adélie penguin juvenile survival.

Variable type	Variable	Spatial scale	Temporal scale	Hypothesized mechanisms and previous results	Expected effects on Adélie penguin juvenile survival
	Southern Oscillation Index (SOI)	global	March- September	Low SOI values linked to higher winter temperatures were negatively related to adult survival in Adélie penguins in Adélie Land (32)	Negative
Global climatic index	Southern Annular Mode (SAM)	global	March- September	Positive SAM values were related to higher juvenile survival in Emperor penguins in Adélie Land, possibly through increased food availability ( <i>92</i> ). Also in Adélie Land, this relationship was negative for snow petrels, possibly because positive SAM	Positive

				increases fast ice extent and reduces food availability for this surface-feeding species (93).	
	Sea ice extent anomalies (SIEA)	regional (90°E - 150°E)	March- September	SIEA were quadratically related to Adélie penguin adult survival in the Ross sea ( <i>33</i> ) and positively related to juvenile survival in the WAP, possibly because increased ice extent provides more resting habitat ( <i>29</i> ).	Positive
Sea ice metrics	<b>SIC.autumn</b> (average SIC in area)	local (135°E - 145°E)	March-April	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival ( <i>11</i> ). Furthermore, tracking data from the Western Antarctic Peninsula (WAP) suggest a survival bottleneck occurs right	Positive

			after fledging (45).	
<b>SIC.march</b> (average SIC in area)	local (135°E - 145°E)	March	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival ( <i>11</i> ). Furthermore, tracking data from the Western Antarctic Peninsula (WAP) suggest a survival bottleneck occurs right after fledging ( <i>45</i> ).	Positive
<b>SIC.april</b> (average SIC in area)	local (135°E - 145°E)	April	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival ( <i>11</i> ). Furthermore, tracking data from the Western Antarctic Peninsula (WAP) suggest a survival	Positive

			bottleneck occurs right after fledging ( <i>45</i> ).	
<b>SIC.winter</b> (average SIC in area)	local/regional (110°E - 135°E)	May-July	Increased autumn/winter sea ice concentrations were positively related to population growth rate across Antarctica, consistent with effects on juvenile survival ( <i>11</i> ). Higher winter sea ice concentrations in the Weddell sea were positively related to juvenile survival in the WAP ( <i>29</i> ).	Positive
<b>SIC.latewinter</b> (average SIC in area)	local/regional (110°E - 135°E)	August- September	Sea ice concentrations in Aug-Sept were weakly but negatively related to Adélie penguin juvenile survival in another east Antarctica population, possibly because too much ice pushes individuals away from foraging areas and/or affects predation pressure ( <i>34</i> ).	Negative

	<b>MIZFRAC.april</b> (% of ice area (15- 100% SIC) covered by the marginal ice zone (15-80% SIC))	local (135°E - 145°E)	April	As different types of sea ice may affect seabird vital rates differently ( <i>96</i> ), we decomposed the April sea ice concentration into its loose fraction (MIZFRAC.april) and its solid fraction (DENSEFRAC.april).	Unknown
	DENSEFRAC.april (% of ice area (15- 100% SIC) covered by dense ice (80-100% SIC))	local (135°E - 145°E)	April	As different types of sea ice may affect seabird vital rates differently ( <i>96</i> ), we decomposed the April sea ice concentration into its loose fraction (MIZFRAC.april) and its solid fraction (DENSEFRAC.april)	Unknown
Local climatic and intrinsic variables	Windchill temperature (WCT)	local/regional	April-September	WCT were quadratically or positively related to Adélie penguin juvenile survival in another east Antarctica population possibly because higher temperatures decreased thermoregulatory costs	Positive

			(31).	
Cohort size	local	Annual	Density-dependent effects (positive relationship between cohort size and juvenile survival) were reported in another east Antarctica Adélie penguin population ( <i>31</i> ), possibly acting through diluted predation ( <i>55</i> ) pressure or increased social foraging.	Positive
Average chick mass at tagging	local	Annual	Higher mass at fledging buffers inexperienced fledglings against depletion of energy reserve and is a general predictor of juvenile survival in birds ( <i>97</i> ).	Positive

#### Table S3:

**Model selection for recapture, transition, and survival probabilities.** Abbreviations: AICc, Akaike Information Criterion corrected for small sample sizes ;  $\Delta$ AICc, AICc difference between current model and time-dependent model ; K, number of parameters ; Dev, Model deviance.

Submodel set	Inde x	Model	AICc	ΔAICc (for each submode l set)	AICc Weight s	K	Dev
Recapture (p)							
			12076.			10	2276.
	3	$\Phi$ (5 age classes × time) <b>p(state1: 3 age classes × time; state2: 3 age classes)</b> $\Psi$ (5 age classes × state)	5	0.0	0.902	4	1
			12081.			11	2260.
	2	$\Phi$ (5 age classes × time) <b>p(state1: 4 age classes × time; state2: 4 age classes)</b> $\Psi$ (5 age classes × state)	8	5.3	0.063	4	8
			12083.	<i></i>	0.00 <b>-</b>	12	2243.
	1	$\Phi$ (5 age classes × time) <b>p(state1: 5 age classes × time; state2: 5 age classes)</b> $\Psi$ (5 age classes × state)	0	6.5	0.035	3	4
	_		12095.	10.2	0.000	74	2356.
	5	$\Phi$ (5 age classes × time) <b>p(state1: 5 age classes + time; state2: 5 age classes)</b> $\Phi$ (5 age classes × state)	/	19.2	0.000	/4	/
	6	Φ (5 ago classos × tima) <b>n(ctato1; 4 ago classos + timo; stato2; 4 ago classos)</b> W(5 ago classos × stato)	12098.	21 7	0.000	72	2301.
	0	$\Phi$ (J age classes ~ time) <b>p(state1.</b> $\Phi$ age classes + time, state2. $\Phi$ age classes) $\mp$ (J age classes ~ state)	12135	21.7	0.000	/3	2357
	4	Φ (5 age classes x time) <b>n(state1: 2 age classes + time: state2: 2 age classes)</b> Ψ(5 age classes x state)	6	591	0.000	93	2337.
State transition	-			0,712	0.000	,,,	0
(Ψ)							
1			12076.			10	2276.
	3	$\Phi$ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (5 age classes × state)	5	0.0	0.991	4	1
			12086.			10	2294.
	8	$\Phi$ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (3 age classes × state)	9	10.3	0.006	0	7
			12087.			10	2291.
	7	$\Phi$ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (4 age classes × state)	7	11.2	0.004	2	4
			12096.				2308.
	9	$\Phi$ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (2 age classes × state)	4	19.9	0.000	98	3
	10		12098.	21.4	0.000	07	2311.
	10	$\Psi$ (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (state)	0	21.4	0.000	97	9
Survival ( $\Phi$ )							

		12044.				2303.
13	<b>Φ</b> (2 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (5 age classes × state)	7	0.0	0.997	75	7
		12056.				2294.
12	<b>Φ</b> (3 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (5 age classes × state)	1	11.4	0.003	85	7
		12064.				2282.
11	<b>Φ</b> (4 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (5 age classes × state)	6	19.9	0.000	95	7
		12076.			10	2276.
3	<b>Φ</b> (5 age classes × time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (5 age classes × state)	5	31.8	0.000	4	1
		12127.				2406.
16	<b>Φ</b> (2 age classes + time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (5 age classes × state)	1	82.3	0.000	65	4
	$\Phi$ (2 age classes: cst/time) p(state1: 3 age classes × time; state2: 3 age classes) $\Psi$ (5 age classes ×	12264.				2537.
15	state)	6	219.8	0.000	68	9
		12280.				2580.
14	<b>Φ</b> (5 age classes) $p(\text{state1: 3 age classes} \times \text{time}; \text{ state2: 3 age classes}) \Psi(5 \text{ age classes} \times \text{state})$	2	235.5	0.000	55	0