

High-precision 40Ar/39Ar dating of Australasian tektites associated with bifacial tools in the Bose Basin (Xiaomei and Fengshudao sites), South China and in Vietnam (Go Da and Roc Tung 1 sites)

Véronique Michel, Fred Jourdan, Marie-Hélène Moncel, Bernard Gratuze, Guanjun Shen, Wei Wang, Celia Mayers, Adam Frew, Dominique Cauche, Patricia Valensi, et al.

▶ To cite this version:

Véronique Michel, Fred Jourdan, Marie-Hélène Moncel, Bernard Gratuze, Guanjun Shen, et al.. High-precision 40Ar/39Ar dating of Australasian tektites associated with bifacial tools in the Bose Basin (Xiaomei and Fengshudao sites), South China and in Vietnam (Go Da and Roc Tung 1 sites). Quaternary Science Reviews, 2024, 346, pp.109065. 10.1016/j.quascirev.2024.109065. hal-04792396

HAL Id: hal-04792396 https://hal.science/hal-04792396v1

Submitted on 20 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 https://doi.org/10.1016/j.quascirev.2024.109065

2

- 3 High-precision 40Ar/39Ar dating of Australasian tektites associated with
- 4 bifacial tools in the Bose Basin (Xiaomei and Fengshudao sites), South China
- 5 and in Vietnam (Go Da and Roc Tung 1 sites).

6

- 7 Véronique Michel^{a,b,*}, Fred Jourdan^c, Marie-Hélène Moncel^d, Bernard Gratuze^e, Guanjun
- 8 Shen^f, Wei Wang^g, Celia Mayers^c, Adam Frew^c, Dominique Cauche^h, Patricia Valensi^d, Sylvain
- 9 Gallet^b, Anatoly P. Dereviankoⁱ, Alexander V. Kandybaⁱ, Sergey A. Gladyshevⁱ, Henry de
- 10 Lumley^h
- 11 ^a Université Côte d'Azur, CNRS, CEPAM, Nice, France

12

- b Université Côte d'Azur, CNRS, Observatoire de la Côte d'Azur, IRD, Géoazur, Valbonne,
- 14 France

15

- ^c Western Australian Argon Isotope Facility, School of Earth and Planetary Sciences, John de
- 17 Laeter Centre, Space Science and Technology Centre, Curtin University, Perth, WA,
- 18 Australia.

19

- 20 d UMR 7194 HNHP (MNHN-CNRS-UPVD), Département Homme et Environnement, Muséum
- 21 National d'Histoire Naturelle, 75013 Paris, France

22

- ^e UMR 7065, IRAMAT, CNRS-Université d'Orleans, Centre Ernest-Babelon, 45071 Orleans,
- 24 France

25

- ^f College of Geographical Sciences, Nanjing Normal University, Nanjing, Jiangsu 210023,
- 27 China

30 31

28

^g Institute of Culture Heritage, Shandong University, Qingdao Shandong, 266237, China

^h Institut de Paléontologie Humaine, HNHP CNRS-MNHN, 75013, Paris, France

i Institute of Archaeology and Ethnography, Siberia Branch, Russian Academy of Sciences,
Pr. Akademika Lavrentieva 17, Novosibirsk, 630090, Russia

35

- *Corresponding author.
- 38 Email address: veronique.michel@cnrs.fr (V. Michel).
- 39 Full postal address: Université Côte d'Azur, Pôle Universitaire Saint Jean d'Angély, SJA3,
- 40 CEPAM, UMR7264, CNRS, UCA, 24 avenue des Diables Bleus, 06300 Nice, France.

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

ABSTRACT

Bifacial tools discovered at about a hundred Lower Palaeolithic sites in the Bose Basin, southern China, have been previously dated to around 800,000 years ago. This age was obtained by dating four tektites directly associated with the tools at the Nalai site using the ⁴⁰Ar/³⁹Ar method (Michel et al., 2021). Similar ages on tektites from the Bogu and Yangwu sites had been previously published, albeit with limited analytical details, by Hou et al. (2000). In this study, a total of eight tektites, discovered alongside abundant lithic artefacts including bifacial tools, were dated with the 40Ar/39Ar technique, using an incremental temperature increase approach. Six of these tektites are from two Lower Palaeolithic sites in the Bose Basin; three from Xiaomei and three others from Fengshudao (China). The remaining two tektites come from two sites in Vietnam, one from Go Da and the other from Roc-Tung 1. With the exception of the site of Go Da, the tektites were buried in lateritic sediments associated with the stone tools. At the Go Da site, the tektites were found in a layer overlying the deposit containing bifacial tools. The tektites display no signs of fluvial abrasion or reworking, making them potentially strong chronological markers. Chemical analyses of major elements and rare earth elements indicate that the tektites belong to the Australasian tektite group. This is further confirmed by new 40 Ar/ 39 Ar analyses with a weighted mean age of 787.2 ± 8.2 ka (2σ) (MSWD = 0.96, P = 0.45). Therefore, when combined with the available results on Australasian tektites, this suggests a highly precise age of 788.0 \pm 2.6 ka (2 σ ; P = 0.84) for these tektites and consequently, for the bifacial tools in this part of Southeast Asia, located east of the Movius Line. The presence of bifacial tools attests to either a diffusion of the Acheulean from Africa

or a local emergence from previous occupations and traditions in Asia.

66

65

67 **Keywords:** bifaces, bifacial tools, China, Vietnam, tektite, dating, 40Ar/39Ar, step-heating, Mid-

Pleistocene climate transition

69

70

68

1. Introduction

Understanding the evolution, diffusion, behavioural changes and adaptation of hominins over 71 72 time, and the impact and role of climate cycles on the distribution of occupations, is intricately linked to defining an accurate chronological framework. Many questions remain unclear, such 73 as the precise dates of the onset of Lower Palaeolithic-type industries, including the Acheulean, 74 75 in Asia (arrivals of new populations/traditions or local evolution). The most commonly utilized methods for dating events from the earliest phases of hominin occupations and dispersion, i.e., 76 2.6-0.6 Ma, are based on radioactive decay (such as K-Ar and its improved derivative ⁴⁰Ar/³⁹Ar, 77 U-Th, and U-Pb), burial dating by isotopic decay (²⁶Al/¹⁰Be), cumulative processes (Fission 78 track FT, Electron spin resonance ESR, Luminescence), or on the recording of natural processes 79 80 (such as magnetostratigraphy), often incorporating biochronological data (Pike and Pettitt, 2003; Scaillet et al., 2008; Grün et al., 2010; Shen et al., 2012; Tribolo et al., 2013; Jourdan et 81 al., 2014; Lebatard et al., 2014; Hellstrom and Pickering, 2015; Tu et al., 2017; Carbonell et al., 82 83 2010; de Lombera-Hermida et al., 2015; Carbonell et al., 2016; Harmand et al., 2015; Semaw et al., 2003; Garcia et al., 2010; Toro-Moyano et al., 2013). 84 In Asia, the earliest evidence, based on current knowledge, was discovered in Central China, 85 and consists of four localities (Xihoudu, Renzidong, Longuppo, Longgudong) potentially 86 indicating the presence of hominins. The ages are between ~2.6 and 1.8 Ma (Shen et al., 2020; 87 88 Han et al., 2017; Wei et al., 2014; Hou and Zhao, 2010; Li et al., 2017; Bae, 2024, Fig. 1, Fig.

2, Table S1). Several other early sites located in North China date between approximately 2 and 89 90 1 Ma ago, such as Shangchen, Gongwangling or sites in the Nihewan Formation (Majuangou, Xiantai, Xiaochangliang, Feiliang, Madigou, Cenjiawan, Donggutuo) (Liu et al., 2014; Zhu et 91 92 al., 2004; Zhu et al., 2018; Tu et al., 2017; Ao et al., 2010; Ao et al., 2012; Deng et al., 2006; Keates, 2010; Liu et al., 2013; Zhu et al., 2001, 2003; Yang et al., 2016; Ao et al., 2013; Li et 93 al., 2016; Pei et al., 2019; Guan et al., 2016; Liu et al., 2013; Wang et al., 2005; Yang et al., 94 2017; Yang et al., 2020, see Table S1 for details). All these sites have yielded lithic 95 assemblages, described by some as crudely-made tools. Recent dating studies have improved 96 the chronological framework of early hominin presence in this part of Eurasia, as described in 97 98 Appendix A (Table S1, Fig. 2). The emergence and diffusion of the Acheulean (with new technological features including 99 Large Cutting Tools-LCTs) is still enigmatic. The earliest evidence is located in Africa, with 100 101 sites corresponding to the transition from the Oldowan/Mode 1 (core-and-flake industries) to the Early Acheulean/Mode 2. This occurred diachronically around 1.95-1.0 Ma (Lepre et al., 102 103 2011; Beyene et al., 2013; Mussi et al., 2023a). These African industries evolved over time into what is described as the Classical Acheulean from ~1.5 to 0.8 Ma (Mussi et al., 2023b; Galloti 104 and Mussi, 2017). Outside Africa, the Acheulean is younger, with evidence in the Levant of an 105 early "Out of Africa" dispersion for this tradition at ~1.6-1.4 Ma at Ubeidiya (Martinez-Navarro 106 107 et al., 2012; Sagi, 2005; Rink et al., 2007), possibly followed by a second wave at ~800 ka at Gesher Benot Yakov (Goren-Inbar et al., 2018), suggesting, at the least recurrent phases of 108 diffusion outside Africa. 109 Moving to Asia, two hypotheses have been put forward regarding the emergence of the 110 Acheulean culture beyond the "Movius Line" in an area where LCTs were considered to be 111 absent (Movius, 1948; Bae, 2024): either this represents the diffusion of the Acheulean from 112

Africa, as observed in the Near East, or it is a local emergence from previous occupations and traditions in Asia (Bae, 2024). It is therefore essential to accurately date sites with LCTs (bifaces/bifacial tools/handaxes according to studies and authors) in Eurasia in order to contribute to discussions on the onset of this/these new traditions(s) in Asia, and to better define the possible diffusion of what is described in Africa as the Acheulean. Only we then will gain insights into Pleistocene hominin mobility and behavioural strategies in response to a continually changing climate (Bae et al., 2018, Larick and Ciochon, 2015; Qin and Sun, 2023). For convenience, throughout our paper, we use the generic terminology of LCTs to designate lithic assemblages with bifaces, handaxes and bifacial tools. In Michel et al. (2021), tektites in close association with LCTs at the Nalai site (Bose Basin, South China) were dated to 809 ± 12 ka from 55 total fusion subsamples. A probability density distribution yielded an age centred at 797 ± 6 ka, tentatively suggesting that they might be part of the Australasian tektite strewnfield recently dated to 788 ± 3 ka $(2\sigma, P = 0.54)$ by the ⁴⁰Ar/³⁹Ar method (Jourdan et al., 2019). In this new study, our aim is to improve the accuracy of our preliminary dating result of the Bose sites (Michel et al., 2021) and obtain new dates for two sites in Central Vietnam yielding LCTs. We dated tektites via 40Ar/39Ar analysis by the "step-heating" method in order to circumvent inherited argon present in tektites during the initial steps. Furthermore, with this approach, inverse isochron ages can be calculated, which resolves and takes into account any inherited ⁴⁰Ar* present in the samples in the calculation, provided that its argon isotopic composition is homogenous, thus leading to more accurate results (Jourdan et al., 2019). 40Ar/39Ar dating analyses were conducted on tektites from two archaeological sites located in the Bose Basin in China and from two others sites located in Vietnam (An Khê): Xiaomei, Fengshudao, Go Da and Roc Tung 1, respectively (Fig. 1, Fig.

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

S1, Fig. S2). Major and trace element geochemistry was applied to tektites to assess whether they were part of the Australasian strewnfield.

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

2. Australasian tektites: origin, background chronology and chronological markers Tektites are glasses produced by the melting of the continental crust during the impact of a meteorite. This impact causing the formation of Australasian tektites is supposed to have resulted in the formation of a crater located in Indochina, or more likely, in southern Laos. According to the recent study by Sieh et al. (2020), the crater would be buried under the Bolaven volcanic field, near the northern limit of the strewnfield (Fig. 1), although firm evidence is still lacking. These tektites have been found at varying distances from the impact structure, spanning regions from Southeast Asia to Antarctica, with a chemical composition characteristic of the group of Australasian tektites (Koeberl, 1990; Heide et al., 2001; Glass and Koeberl, 2006; Schwarcz et al., 2016). Tektites closely associated with artefacts from the Bose Basin in China were initially dated using fission track dating (Guo et al., 1997). The authors obtained an age of 732 ± 39 ka for the tektites of the Bogu site (Fig. S1). Subsequently, the ⁴⁰Ar/³⁹Ar method was applied to two tektites from the Bogu site and one from the Yangwu site, but these results were published without age spectra or detailed analytical data (Hou et al., 2000) (Fig. S1). The authors obtained an overall weighted mean age of 803 ± 6 ka (2σ) from the isochron ages of three samples (Hou et al., 2000; Potts et al., 2000). This first ⁴⁰Ar/³⁹Ar dating of tektite in Bose Palaeolithic sites, presented by Hou et al. (2000) was questioned by Koeberl and Glass (2000). The latter authors cast doubt on the association of tektites with the artefacts and claimed that tektites are generally found in deposits younger than their isotopic age. They thus called for caution in using them as stratigraphic markers

(Koeberl and Glass, 2000; Keates, 2000). Potts et al. (2000) responded clearly, affirming that

the Bose tektites are *in situ* and associated with the stone tools in terrace T4, as described by Hou et al. (2000), in the same way as the stone tools found alongside tektites at the Gaolingpo site (Xie et al., 2021). They attributed the tektite age to the tools as there is no geomorphological and stratigraphical indication of the redeposition of either the tools or tektites (Potts et al., 2000). Later, Wang et al. (2014) presented a detailed morphological study of the Fengshudao site and its bifacial industry. In this site, the first excavated association of tektites, dating to ca. 803 ka, with LCTs (including bifaces), and found in situ, was described in one stratigraphic level in terrace T4, indicating that the Bose Basin handaxes most likely date to the Early to Middle Pleistocene transition (Fig. 2). However, according to Langbroek (2015), the age of the earliest bifaces in China and the close association between tektites and bifaces have not yet been demonstrated, although this was clearly established by Wang and Bae (2015), and later confirmed at the Nalai site with Australasian tektites and bifaces found together in situ (Michel et al., 2021). It is important to note that two Australasian tektites found in situ at the Sangiran area have been successfully used as chronological markers. They are consistent with the magnetic polarity stratigraphy and predate the Matuyama-Brunhes transition (MB) (Hyodo et al., 2011). The data demonstrate the near co-occurrence of the tektite level and the magnetostratigraphic constraint, arguing that the last occurrence of Sangiran hominins dates to $\sim 0.79 \text{ Ma (Fig. 2)}.$ Later, Jourdan et al. (2019) conducted precise dating on four Australasian tektites from Western Australia, Thailand, Vietnam, and southern China, including a tektite from the Bose Basin previously dated by Hou et al. (2000). The exact origin of the tektite from the Bose Basin was not specified by Jourdan et al. (2019). Age plateau and isochron diagrams yielded concordant data with a weighted-mean age of 788.1 ± 2.8 (P = 0.54) for these Australasian tektites, which aligns closely with the less precise 40 Ar/ 39 Ar age of 793 \pm 14 ka published by Schwarz et al. (2016).

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

3. Xiaomei and Fengshudao sites in Southern China: Geography, stratigraphy and lithic 187 assemblages 188 The history of Palaeolithic research in the Bose Basin has been thoroughly documented by 189 Wang et al. (2014). Approximately a hundred sites have been discovered in this basin (de 190 Lumley et al., 2020; Wang et al., 2024). In these three abovementioned studies, the 191 geography/geology of the Bose Basin with its Quaternary terrace stratigraphy (see Fig. S3, Fig. 192 S4), as well as overviews of the site's location and lithic assemblages, are also described in 193 194 detail. 195 The Xiaomei site is located in Xiaomei village, Youjiang district, Bose city, Guangxi Zhuang Autonomous Region (23°46'41.5" N, 106°42'21.9" E), on the fourth terrace of the Youjiang 196 River (Fig. S1, Fig. S3, Fig. S4, Fig. S5). It was discovered in the 1980s during an 197 archaeological survey covering an area of 350,000 m². Many lithic artefacts and a few tektites 198 were exposed on the ground surface of the laterite deposits at the site. The lithic series were 199 made up of diversified LCTs, including pick and tongue-shaped bifaces typical of the area (Fig. 200 3, n° 6), on sandstone or quartzite pebbles (Fig. 3, n° 7 and 8), associated with flake-tools 201 (scrapers), and notably a large unifacial core on a massive thick quartzite pebble (Fig. 3, n° 9) 202 with centripetal removals. A test excavation conducted in 1996 unearthed artefacts, associated 203 204 with tektites. Based on geomorphology and artefacts, the age of this site is assumed to be the same as that of Bogu site. 205 The Fengshudao site (23°57'39"N, 106°40'39"E) is situated along the south bank of the 206 Chengbihe River, a significant tributary of the Youjiang River (Fig. S1, Fig. S3, Fig. S4). The 207 208 site is located in the fourth terrace, spanning 100 metres from east to west, with a sloping terrain 209 from north to south of approximately 10 to 30 metres. Discovered in 2004, the site was excavated from 2004 to 2005, revealing nearly 100 artefacts and nine tektites in situ from an 210 excavated area of 62.6 m² (Wang et al., 2014). The lithic assemblage at the Fengshudao site 211

includes the same diversity as at Xiaomei (Fig. 3, n° 5) with in addition, evidence of the use of

bipolar debitage. The main raw materials consist of sandstone, quartzite and quartz collected

as pebbles. Both tektites and LCTs are found in exactly the same layer within the laterite (Wang

- et al., 2024) (Fig. 4).
- The bifaces exhibit the characteristic shape of the Bose Basin bifaces (Hou et al., 2000; Xie
- and Bodin, 2007; Zhang et al., 2010; Huang et al., 2012; Wang et al., 2014; Feng et al. 2018;
- Wang et al., 2024). They are notably short, broad and stocky, featuring a tongue-shaped
- 219 extremity (Fig. 3, n° 1, 2 and 3).

220

221

- 4. Go Da and Roc Tung 1 sites in Central Vietnam: Geography, stratigraphy and lithic
- 222 assemblage
- Following the discovery of sites with LCTs near the city of An Khê in Central Vietnam in 2014,
- by Vietnamese archaeologists, systematic excavations have been conducted since 2015
- 225 (Derevianko et al., 2018) (Fig. 1, Fig. S2).
- Several other similar Palaeolithic sites were identified at the Roc Tung location, situated on the
- left bank of the Ba River. Artefacts were collected on the surface or discovered *in situ* during
- excavations and test pits, with similar stratigraphic contexts. The Roc Tung 1 site (14°02'253"
- N; 108°40'822" E) is situated on a high and relatively flat area of agricultural land. The
- stratigraphy is well-documented through excavations and various test pits (Derevianko et al.,
- 2018). Three layers have been identified below the surface debris of an irrigation canal. The
- cultural layer, situated at the top of the laterite (Layer 3), has yielded several *in situ* artefacts
- associated with tektites (Fig. S6, Fig. 5).
- 234 Go Da (13°58'306" N; 108°39'136" E; 440 m a.s.l.) is located downstream, near An Khê, on a
- sloping hill, 900 m west of the Ba River (Fig. S2). The stratigraphy established during the
- excavation is somewhat different on this right bank of the Ba (Derevianko et al. (2018);

Kandyba et al. (2021) (Fig. S7). The stratigraphic sequence begins with a substantial weathering crust (layer 1) of the granite bedrock, succeeded by slope wash sediments (layer 2). The cultural horizon at Go Da formed during these erosion and slope wash processes, probably in a cooler climate than the present one. Archaeological materials are specifically situated at the top of the weathering crust and at the base of the slope wash layer (Fig. S7). Tektites found in the upper part of this layer 2 indicate that hominins were present at Go Da before the deposition of Australasian tektites. In other words, it can be asserted that the industries at Go Da predate those at Roc Tung 1, where artefacts are found in situ with tektites. The local industry is well described by Derevianko et al. (2018) and Kandyba et al. (2021). The lithic assemblage from the Roc Tung 1 site, collected on the surface but also in stratigraphy context during excavations and test pits, includes over 126 pieces (Derevianko et al. 2018), made on pebbles or boulders. This lithic industry from the left bank of the Ba River is made on local raw materials, most often quartzite cobbles or blocks, but also other raw materials of lesser quality, such as vein quartz. Many tested pebbles have been identified in the assemblage, no doubt corresponding to knappers' efforts to select good-quality blanks (Fig. 6). As in the other An Khê-tradition assemblages (Derevianko et al. 2018), the main features of the industry at Roc Tung 1 are heavy-duty-tools including numerous LCTs (bifaces, picks), choppers, and chopping-tools (Fig. 6, n^o 2 and 3) and light-duty tools, such as scrapers and pointed tools, often made on flakes. Finally, flakes and cores attest to simple reduction processes: cores mainly bear a single debitage surface with unipolar or centripetal removals and a limited prepared or cortical striking platform. The lithic assemblage at the Go Da site comprises around a hundred artefacts (Derevianko et al. 2018; Kandyba et al 2021), comparable to those from Roc Tung 1. These consist of numerous split tested pebbles, often large in size, hammerstones on rounded granite pebbles, a heavy-duty tool kit with mainly thick, short and stocky LCTs (including bifaces and picks), on

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

quartzite pebbles or slabs. Some display a tongue-shaped extremity. Others are choppers on long and massive quartzite pebbles, and pointed tools and scrapers (Fig. 6, n°1). The cores, flakes and small flakes indicate *in situ* reduction processes, mainly unifacial and unipolar removals. This assemblage displays all the characteristics of the An Khê industry, as well as regional traditions.

These two assemblages from Roc Tung 1 and Go Da share many similarities to industries from the Bose Basin in China, notably those from Fengshudao and Xiaomei, discussed in this study.

5. Materials and methods

5.1. Materials.

Our analysed tektites come from the Fengshudao and Xiaomei sites (Bose Basin, China), and Roc Tung 1 and Go Da sites (Central Vietnam). They were discovered *in situ* associated with tools (including LCTs), except for the Go Da site, where the tektites were found in a layer overlying bifacial tool-bearing deposits, providing a minimum age for the artefacts.

Three tektites (F1, F2, F3) collected from the laterite layer of terrace T4 at the Fengshudao site (Fig. 7), display a very thin profile (~0.4-0.6 g) and show no signs of fluvial abrasion. They are very angular and sharp, suggesting that no river re-depositing occurred. Additionally, three tektites, namely B1 (10.2 g), B2 (8.7 g), and B3 (6.5 g), were excavated from the same laterite layer from terrace T4 at the Xiaomei site, with a rounder morphology than the others (Fig. 7).

Two tektites from Central Vietnam; one larger tektite, GD (19.1 g), excavated at the Go Da site, and another, large and very angular tektite, RT1 (10.3 g), with no signs of fluvial abrasion, excavated at the Roc Tung 1 site, complete the sample. The GD tektite is rounder, similar to the Xiaomei tektites studied here, but neither show signs of fluvial abrasion or reworking.

285 5.2. Method

Before analysis, the eight tektite samples were cleaned in an ultrasonic water bath to remove surface-bonded residues (Fig. 7). The thinner tektites (F1, F2, F3) were reduced to pieces. For the larger ones (B1, B2, B3, GD, and RT1), a piece of each tektite was extracted from the original tektites and reduced to pieces. Some were further crushed in an agate mortar and divided into subsamples for multiple analyses. For ⁴⁰Ar/³⁹Ar dating, subsamples 400-500 µm in size, from F1, F2, B1, B2, B3, GD and RT1 tektites, were cleaned with an HNO₃ 2N ultrasonic bath for 15 minutes, in HF 2% for 20 minutes, in H2O, and then in acetone. Subsamples weighing 30-70 mg were analysed using the "step-heating" method with incremental temperature increases to produce an age spectrum and inverse isochron. The ⁴⁰Ar/³⁹Ar analyses were conducted at the Western Australian Argon Isotope Facility (Australia) using an ARGUS VI multicollection instrument and following the procedure described by Jourdan et al. (2019), used for dating the Australasian tektites. The raw data (Appendix B) were processed using the ArArCALC software (Koppers, 2002) and the ages were calculated using the FCs standards (Jourdan and Renne, 2007), for which an age of 28.294 Ma (± 0.13%) was used (Renne et al., 2011), along with the decay constants recommended by Renne et al. (2011). Our criteria for plateau determination are as follows: plateaus must include at least 70% of ³⁹Ar. The plateau should be distributed over a minimum of three consecutive steps agreeing at 95% confidence level and satisfying a probability of fit (P) of at least 0.05. Plateau ages are given at the 2σ level and are calculated using the mean of all the plateau steps, each weighted by the inverse variance of their individual analytical error. Uncertainties include analytical and J-value errors. Errors with all sources of uncertainties such as the decay constants errors increase the uncertainty by ± 0.2 ka for those samples and are indicated by a square bracket (e.g., $[\pm 3 \text{ ka}]$). The composition of major and trace elements presented in Tables S2 and S3, was determined for all tektites by Laser Ablation Inductively Coupled Plasma Mass Spectrometry at the Centre Ernest-Babelon of the IRAMAT (CNRS/Orléans University) or by Inductively Coupled Plasma

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

Optical Emission Spectrometry at SARM (Service d'Analyses des Roches et des Minéraux, CRPG Nancy, France (see additional details in SI). The composition of major elements of tektites N411, N425, N862, N959 from Nalai site in the Bose Basin, China, as reported by Michel et al. (2021), was supplemented with the analysis of rare earth elements and is presented in Table S3.

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

311

312

313

314

6. Results

6.1 Chemical composition

The major elemental compositions obtained for the tektites B1, B2, B3 (Xiaomei), F1, F2, F3 (Fengshudao), GD (Go Da) and RT1 (Roc Tung 1), are presented in Table S2. They comprise the following ranges: 70.05-74.74 % for SiO₂, 11.13-15.44 % for Al₂O₃, 4.17-4.96 % for FeO₃ 1.77-2.67 % for CaO, 2.06-2.42 % for K₂O, 1.92-2.67 % for MgO, 0.93-1.51 % for Na₂O, 0.68-0.78 % for TiO₂, 0.08-0.10 % for MnO, 0.03-0.07 % for P₂O₅. These values are very close to those of the tektites from Nalai (N411, N425, N862, N959) (Michel et al., 2021) (Table S2). In Figure 8, these values are compared with various groups of tektites according to Schwarz et al. (2016) and Wesgate et al. (2021). Figure 8 shows that our tektites belong to the group of Australasian tektites. Tektites from Vietnam (GD and RT1) are very slightly different to those from the Bose Basin. They belong to the subgroup of indochinite showing some regional compositional variation in comparison with the australite subgroup, but ultimately belong to the same Australasian group. In the same way as the Australasian tektites, our tektites exhibit light rare-earth-element-(REE)-enriched compositions with a marked negative EU anomaly, as described by Koeberl (1990), Son and Koeberl (2005), Amare and Koeberl (2006), Wesgate et al. (2021) and Schwarz et al. (2016) (Fig. 9).

334

335

 6.2^{40} Ar/ 39 Ar dating

We obtained seven plateau ages ranging from 779 \pm 19 ka (MSWD = 0.53; P=0.93) to 827 \pm 57 ka (MSWD = 0.09; P = 1.00) and including between 91 and 100 % of the total released ³⁹Ar (Fig. 10, Appendix B). Inverse isochrons yielded similar ages ranging from 774 ± 24 to $842 \pm$ 67 and with trapped 40 Ar/ 36 Ar ratios in error of the atmospheric value of 298.6 \pm 0.3 and ranging from 293.3 \pm 4.9 to 300.8 \pm 6.0 (Appendix C). The K/Ca ratio (derived from $^{39}\text{Ar}_{\text{K}}/^{37}\text{Ar}_{\text{Ca}}$) shows homogenous values across a given tektite but varies among samples with average values ranging from 0.32 ± 0.18 to 1.29 ± 0.11 . The seven plateau ages (P = 0.84) and inverse isochron ages (P = 0.25) are all indistinguishable from each other, indicating the dating of a single event, as expected for tektites originating from a single impact. Furthermore, all intercept 40Ar/36Ar values define a single population (P = 0.57) with a weighted mean of 296 \pm 3, suggesting that no inherited ⁴⁰Ar* is present in the samples analysed in this study. Therefore, we calculated a single weighted mean age of 787.2 ± 8.2 ka $(2\sigma; MSWD = 0.96; P = 0.45)$ as the best estimate for the impact event that produced these tektites. This age is indistinguishable from the previous age of 788.1 \pm 2.8 (Jourdan et al., 2019), indicating that the tektites are part of the Australasian strewnfield, as also evidenced by their chemical composition. When combined with our new age, we obtained a conclusive best age estimate of 788.0 ± 2.6 [2.8] ka (2σ ; P = 0.84) for the Australasian tektites that will be used in the rest of this study.

353

354

355

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

7. Discussion on the new results of the Chinese and Vietnamese sites regarding the

emergence and diffusion of the Acheulean in Eurasia

- 356 7.1 Emergence of the Acheulean in Africa and its initial diffusion
- The emergence of the Acheulean in Africa is dated to around 1.75 to < 1.0 Ma (i.e., in southern
- Ethiopia, Beyene et al., 2013) and around 1.76 Ma years ago at Kokiselei 4 in Kenya (Lepre et
- al., 2011) but may have occurred earlier, around 1.95 Ma years ago at Garba IV (Upper Awash,
- 360 Ethiopia, Mussi et al., 2023a). This shift was characterized among others (flaking methods,

land-use patterns) by hominin capability to produce large flakes and LCTs, including for example, handaxes, picks and sometimes cleavers. The earliest LCTs (including bifaces), outside Africa are reported at ~1.4-1.2 Ma ago in Ubeidiya in the Levant (Martinez-Navarro et al., 2012; Sagi, 2005; Rink et al., 2007), and are also dated to ~1.5 Ma in Attirampakkam, India, indicating a rapid diffusion throughout Eurasia, if we consider that these industries are related to the African Acheulean. These types of LCTs are not present, earlier than 0.9-0.7 Ma in Western Europe (Pappu et al., 2011; Vallverdu et al., 2014; Voinchet et al., 2015; Antoine et al., 2019; Moncel et al., 2020a; Moncel et al., 2020b). In Asia, the core-and-flake series (Mode 1) persist when Lower Palaeolithic/Acheulean (Mode 2) industries with LCTs (bifaces/bifacial tools/handaxes) appear in some areas around 1.5-0.8 Ma ago. The number of discovered sites increase during the Mid-Pleistocene climate transition (MPT) (1.0-0.6 Ma) suggesting that this period played a crucial role in the diffusion of hominins in Eurasia (Fig. 2, Table S1).

7.2. New insights from Chinese and Vietnamese sites

These new assemblages with LCTs in Asia are currently found in four main Chinese areas (specifically Dingcun, Bose Basin, Luonan Basin, and the Danjiangkou Reservoir Region DRR) (Li et al., 2018, Li et al., 2014). Unfortunately, dating details are limited, except for the site of Yunxian (open-air site of Xuetangliangzi, DRR region, Central China), where two *Homo erectus* crania were discovered with one handaxe and 13 LCTs collected from the surface (Li et al., 2018). This site is approximatively dated between ~1.1-0.85 Ma using both paleomagnetism, biochronology and ESR dating on teeth and quartz (de Lumley and Li, 2008; Echassoux et al., 2008; Tissoux et al., 2008; Feng, 2008) (Fig.1, Fig.2, Table S1). Many other sites with LCTs, up to 200 LCTs, were discovered in this area but further chronological research is required. They are regarded as the most significant persistence of LCTs for a long period, from the late Early Pleistocene to the early part of the Late Pleistocene (Li et al., 2018).

Our results on four Chinese and Vietnamese sites provide strong and accurate data on Asian 386 387 human occupations with LCTs dated to ~ 788 ka. Through these new data, we contribute to understanding of the chronology and diffusion of the earliest populations in Asia. 388 Early human populations thrived and expanded across mainland East Asia, from low latitudes 389 whatever their technology (e.g., the Bose Basin) to high northern latitudes (e.g., the Nihewan 390 391 Basin, Zhoukoudian). The associated industry is often roughly dated using paleomagnetism, as 392 seen in Maliang (Nihewan Formation, North China), Wutaicun (Central China), Longgangsi-2 and 3 (Central China). The determined ages are approximately ~0.78 Ma (Wang et al., 2005), 393 ~0.9 Ma and ~1.2-0.7 Ma (Sun et al., 2017) respectively. Despite approximate dating, an 394 395 overview of paleoenvironmental and archaeological data in the Nihewan Basin reveals changes in hominin presence and indicates that sustained occupation in high latitudes characterized by 396 strong seasonality was not possible until the Late Pleistocene, as evidenced by chronological 397 398 gaps (Yang et al., 2020). Model/Oldowan industries persisted in Asia at Bailong Cave with human teeth in Central China, dated between ~0.78 Ma and ~0.51 Ma, and also using multi-399 dating methods such as paleomagnetism, ²⁶Al/¹⁰Be burial on quartz and ESR/U-series on teeth 400 (Liu et al., 2015; Han et al., 2018; Kong et al., 2018). 401 Moving outside China, this lithic industry is also evidenced at the site of Kalinga between 402 403 around 1.05 and 0.63 Ma, in the Philippines (Ingicco et al., 2018), and between around 1.01 and 0.7 Ma at Mata Menge and Wolo Sege in Indonesia (Brumm et al., 2016). During the same 404 period, an Acheulean-type industry is present at Shuigou-Huixinggou, in Central China, but 405 approximatively dated to ~0.9 Ma by paleomagnetism (Li et al., 2017). Other Mode 1/Mode 2 406 sites dated by alternative radioisotopic methods provide more precise dating (Fig.2, Table S1). 407 This is the case for Zhoukoudian Locality 1, in North China, where human remains and 408 409 industries with bifacial tools and flake-tools (Derevianko, 2008) are well dated to ~0.78-0.4 Ma

using multi-dating methods, such as paleomagnetism, ²⁶Al/¹⁰Be burial and U-Th (Shen et al., 2009, Shen et al., 2016). Li et al. (2018) propose that the technological transmission hypothesis could be a suitable explanation for the emergence of Chinese LCTs technology and suggest a diffusion of the Acheulean-type tradition by hominins from west to east to China and other Asia aeras as Vietnam. Other authors propose a local onset derived from previous Mode 1/core-and-flake assemblages (convergence), giving rise to the An Khê tradition observed at the Go Da and Roc Tung Lower Palaeolithic sites in Vietnam (Derevianko et al., 2018) and the Acheulean in the Bose Lower Palaeolithic sites (Wang et al., 2014; Wang and Bae, 2015; Wang et al., 2024; Bae, 2024) (Fig.1, Fig.2, Table S1). Our results highlight that the industries at the Bose and An Khê sites are now well dated to 788.0 ± 2.6 ka (2σ) by dating in situ Australasian tektites associated with the bifaces. This significant chronological marker accurately positions the industries at the Bose and An Khê sites within the chronological sequence of industries in Asia and also indicates a well-established presence of LCTs and possibly the Acheulean-type tradition in this region of Asia.

8. Conclusion

Our dating of Australasian tektites discovered *in situ* alongside artefacts at the Bose sites in southern China and An Khê in Central Vietnam using 40 Ar/ 39 Ar gives an age of 788.0 ± 2.6 [2.8] ka. This age indicates that LCTs (including bifaces), and thus what can be described as the Acheulean, developed in this part of Asia around 788 ka during the Mid-Pleistocene climate transition, regardless of the hypothesis regarding the onset of the Acheulean (Fig. 2, Table S1). Our result can be compared to the age of the Acheulean industry also found at around 1.1-0.8 Ma at DRR sites, Yunxian and Shuigou-Huixinggou in Central China. This new LCT-making tradition is penecontemporaneous with persistent core-and-flake series (Mode 1) at Nihewan in

northern China, at Bailong, as well as Longgangsi in Central China. This persistence of assemblages without LCTs extends to the Philippines at Kalinga and Indonesia at Mata Menge and Wolo Sege, with for some, the presence of *Homo erectus* while the Acheulean-type assemblages persist at Zhoukoudian (~0.8-0.6 Ma) in North China. The Bose and An Khê sites can be considered as the earliest well-dated evidence of Acheulean-type tradition beyond the Movius Line in eastern Asia, showing that hominins in these areas were capable of making LCTs, including bifaces at ~788 ka.

Acknowledgements

This work was carried out with partial financial support from CEPAM-CNRS. Louise Byrne is acknowledged for English improvements of the manuscript. The Handling Editor, Christopher J. Bae, and the two anonymous reviewers are thanked for their constructive comments and helpful suggestions for improving this manuscript.

Figure captions

Figure 1. Portion of the Australasian tektite strewnfield (modified from Wesgate et al., 2021) with a potential crater impact in southern Laos (Sieh et al., 2020). The Movius Line is represented by the red dashed line, separating Western Asia, where LCTs were discovered from Eastern Asia, where LCTs, were considered absent (Movius, 1948). However, since then, Acheulean assemblages with LCTs have been discovered east of the line (Bae, 2024). The archaeological sites are located south of the 45°N parallel and are classified in chronological order. The sites are dated using methods such as radiometric dating and/or relative dating, including biochronology and paleomagnetism, based on current knowledge (Table S1). The distribution indicates sites with Mode 1 (core-and-flake assemblages) (marked with stars), those

with *Homo* remains (marked with circles), and those associated with Acheulean/Mode 2 (marked with triangles). Red symbols represent the studied sites, while black symbols denote other key sites (see details in Tab S1): (1) Masol, India, (2) Riwat, Pakistan, (3) Attirampakkam, India, (4) Renzidong, Central China, (5) Xihoudu, northern China (6) Longgupo, Central China, (7) Longgudong, Central China, (8) Shangchen, Central China, (9) Gongwangling (Lantian man), North China, (10) Niujianbao, Southwest China, (11) Many sites in the Nihewan Basin, northern China: e.g. Donggutuo, Maliang, Feilang Xiaochangliang, Xiantai, Majuangou, Madigou, Cenjiawan Xiaochangliang, (12) Sangiran, Bapang, Lower Lahar unit, Central Java, Indonesia, (13) Longgangsi 3-2, (14) Soa Basin, Ola Bula Formation, Mata Menge, Wolo Sege, Central Flores, eastern Indonesia (15) Yunxian, DRR sites, Central China, (16) Wutaicun, Central China, (17) Shuigou-Huiximggou, Central China, (18) Meipu, Central China, (19) Trinil, Java, Indonesia, (20) Many sites in the Bose Basin, Bogu, Yangwu; Nalai, Fengshudao, Xiaomei (this study), (21) Roc Tung 1 and Go Da, Central Vietnam (this study), (22) Kalinga, Philippines, (23) Bailong, Central China, (24) Zhoukoudian Locality 1, northern China, (25) Yiyuan, northern China.

Figure 2. Chronological sequence of Palaeolithic sites with hominin fossils and/or lithic assemblages according to geographical location and correlation with marine isotope stages and δ^{18} O stacks (Lisiecki and Raymo, 2005). The horizontal grey band denotes the Mid-Pleistocene climate transition MPT (1.0-0.6 Ma).

Figure 3. Lithic artefacts from Fengshudao site (1 to 5) and from Xiaomei site (6 to 9). 1 to 3-tongue-shaped bifaces (N007612; N007622; N007623); 4- chopper (N007775); 5- large flake (N007566); 6-tongue-shaped biface (N73 H05); 7 & 8- picks (N94 5.6; N193 G2); 9- large core (N59).

- Figure 4. Fengshudao site excavation. Photographs showing the *in situ* discovery of biface No.
- 486 Z007773 (2) in the laterite layer, in close association with tektite No. FSD-059 (1).

- 488 Figure 5. Roc Tung 1 excavation. Photograph showing an *in situ* discovery of a pebble core
- No 16.RT1.ST79 (2) in the laterite layer (Layer 3), in close association with a tektite No 14 (1).
- 490 Bifaces were excavated in the same layer.

491

- 492 **Figure 6.** Lithic artefacts from Go Da site (1) and from Roc Tung 1 site (2 and 3). 1- scraper;
- 493 2- chopper; 3- biface.

494

- 495 **Figure 7.** Photograph of tektites analysed in this study: F1, F2, F3 (~0.4-0.6 g) are three thin
- tektites excavated from Fengshudao site, Bose Basin, China; B1 (10.2 g), B2 (8.7 g), B3 (6.5
- 497 g) are three tektites excavated from Xiaomei site, Bose Basin, China; GD (19.1 g) is a tektite
- 498 excavated from Go Da site and RT1 (10.3 g) is a tektite excavated from Roc Tung 1 site,
- 499 Vietnam. GD and RT1 were cleaned with a H₂0 ultrasonic bath.

500

- Figure 8. Chemical composition of various tektite groups, based on the original figure by
- Schwarz et al. (2016) with modifications. This figure includes a comparison of compositions
- with cited references by Schwarcz et al. (2016) (Taylor, 1962; Chapman and Schreiber, 1969;
- O'Reilly et al., 1983; Koeberl et al., 1997; Meisel et al., 1997; Albin et al., 2000; Son and
- Koeberl, 2005; Amare and Koeberl, 2006), including Belize (BZ1) (Central America, with
- impact glasses formed at 804 ± 9 ka, Rochette et al., 2021), Ivory Coast tektite (Ivorite), and

Australasian tektites such as Indochinite B (Muong-Nong), Western Canada (WC1) with impact glasses formed at 785 ± 7 ka (40 Ar/ 39 Ar age, Schwarcz et al., 2016), and those with a weighted FT age of 822 ± 20 ka from Malaysia, Java, Australia (group A) (Wesgate et al., 2021). Additionally, we present the chemical compositions of tektites from Nalai, Xiaomei, Fengshudao, Roc Tung 1 and Go Da sites, demonstrating similarities to the Australasian tektite group.

Figure 9. Rare earth element abundances for tektites N411, N425, N862, N959 (Nalai site); F1, F2, F3: tektites (Fengshudao site); B1, B2, B3 (Xiaomei site); GD (Go Da site); RT1 (Roc Tung 1 site). They are chondrite-normalized rare-earth-element (REE) abundance curves (chondrite concentrations are from McDonough and Sun, 1995). The abundances are compared with various tektites, including Belize (BZ1), Ivory Coast tektite 1, and Australasian tektites, such as Indochinite B (Muong-Nong), Western Canada (WC1) (Schwarcz et al., 2016), and those from Malaysia (UT1389), Java (UT2336), Australia (UT2330, UT2334, UT1392, UT2402) (Wesgate et al., 2021). The patterns of our tektites are similar to Australasian tektites (Schwarcz et al., 2016; Wesgate et al., 2021), but distinct to the Belize and the Ivory coast tektites with their lower concentrations and different impact events (Schwarcz et al., 2016).

Figure 10. ⁴⁰Ar/³⁹Ar age spectra of tektites. Plateau age given at the 2σ level. B1, B2, B3: tektites from Xioamei site; F1, F2: tektites from Fengshudao site; RT1: tektite from Roc Tung 1; GD: tektite from Go Da.

- **Appendix A.** Supplementary information.
- 530 MC2. Ar isotopic data corrected for blank.
- 531 MC3. ³⁹Ar/⁴⁰Ar vs. ³⁶Ar/⁴⁰Ar inverse isochron plots.

532 MC4. Tables S2 and S3

533 **References**

- Albin, E. F., Norman, M. D., Rodan, M., 2000. Major and trace element compositions of
- georgiaites: clues to the source of North American tektites. Meteoritic Planetary Science 35,
- 536 795–806.

537

- Amare, K., Koeberl, C., 2006. Variation of chemical composition in Australasian tektites from
- different localities in Vietnam. Meteoritic & Planetary Science 41, 107–123.

540

- Ao, H., Deng, C., Dekkers, M.J., Liu, Q., Qin, L., Xiao, G., Chang, H., 2010. Astronomical
- dating of the Xiantai, Donggutuo and Maliang Paleolithic sites in the Nihewan Basin (North
- 543 China) and implications for early human evolution in East Asia. Palaeogeography,
- Palaeoclimatology, Palaeoecology 297, 129-137.
- Ao, H., An, Z.S., Dekkers, M.J., Wei, Q., Pei, S.W., Zhao, H., Zhao, H.L., Xiao, G.Q., Qiang,
- 546 X.K., Wu, D.C., Chang, H., 2012. High-resolution record of geomagnetic excursions in the
- Matuyama chron constrains the ages of the Feiliang and Lanpo Paleolithic sites in the Nihewan
- Basin, North China. Geochemistry, Geophysics, Geosystems 13, Q08017.
- 549 Ao, H., An, Z., Zhisheng, A., Dekkers, M.J., Li, Y., Xiao, G., Zhao, H., Qiang, X., 2013.
- Pleistocene magnetochronology of the fauna and Paleolithic sites in the Nihewan Basin:
- 551 Significance for environmental and hominin evolution in North China. Quaternary
- 552 Geochronology 18, 78-92.

553

- Antoine P., Moncel M-H., Locht J-L. Bahain J-J., Voinchet P., Herisson D., Hurel A., 2019.
- 555 The earliest record of Acheulean human occupation in North-West Europe. Nature Scientific
- 556 Reports 9, 13091.
- 557 https://doi.org/10.1038/s41598-019-49400-w

558

- Bae, C.J., 2024. The Paleoanthropology of Eastern Asia, University of Hawai'i Press,
- Honolulu.

561

- Bae, C.J., Li, F., Cheng, L., Wang, W., Hong, H., 2018. Hominin distribution and density
- patterns in Pleistocene China: Climatic influences. Palaeogeography, Palaeoclimatology,
- 564 Palaeoecology 512, 118-131.

565

- Beyene, Y., Katoh, S., WoldeGabriel, G., Hart, W.K., Uto, K., Sudo, M., Kondo, M., Hyodo,
- 567 M., Renne, P.R., Suwa, G., Asfaw, B., 2013. The characteristics and chronology of the earliest
- Acheulean at Konso, Ethiopia. PNAS 110, 1584-1591.

569

- Brumm, A., van den Bergh, G., Storey, M., Kurniawan, I., Alloway, B.V., Setiawan, R.,
- 571 Setiyabudi, E., Grün, R., Moore, M.W., Yurnaldi, D., Puspaningrum, M.R., Wibowo, U.P.,
- Insani, H., Sutisna, I., Westgate, J.A., Pearce, N.J.G., Duval, M., Meijer, H.J.M., Aziz, F.,
- 573 Sutikna, T., van der Kaars, S., Morwood, M.J., 2016. Age and context of the oldest known
- 574 hominin fossils from Flores. Nature 534, 249–253.

- 576 Carbonell, E., Sala Ramos, R., Rodriguez, X.P., Mosquera, M., Ollé, A., Vergès J.M., Martinez-
- Navarro, B., Bermudez de Castro, J.M., 2010. Early hominid dispersals: A technological
- 578 hypothesis for « out of Africa". Quaternary International 223-224, 36-44.

Carbonell, E., Barsky, D., Sala, R., Celiberti, V., 2016. Structural continuity and technological change in Lower Pleistocene toolkits. Quaternary International 393, 6-18.

582

Chapman, D.R., Scheiber, L.C., 1969. Chemical Investigation of Australasian Tektites. Journal of Geophysical Research 74, 6737-6776.

585

- 586 Deng, C.L., Wei, Q., Zhu, R.X., Wang, H.Q., Zhang, R., Ao, H., Chang, L., Pan, Y.X., 2006.
- Magnetostratigraphic age of the Xiantai Paleolithic site in the Nihewan Basin and implications
- for early human colonization of Northeast Asia. Earth Planet. Sci. Lett. 244, 336–348.

589

- 590 Derevianko, A.P., 2008. Paleoenvironment. The Stone Age. The Bifacial technique on China.
- 591 Archaeology, Ethnology & Anthropology of Eurasia 33/1, 2-32.
- 592 Derevianko, A.P., Kandyda, A.V., Khac Su, N., Gladyshev, S.A., Gia Doi, N., Lebedev, V.A.,
- 593 Chekha, A.M., Rybalko, A.G., Kharevich, V.M., Tsybankov, A.A., 2018. The Discovery of a
- Bifacial Industry in Vietnam. Archaeology, Ethnology and Anthropology of Eurasia 46/3, 3-21.

595

- 596 Echassoux, A., Moigne, A.-M., Moullé, P.-E., Li, T., Feng, X.B., Li, W., WU, Z.,
- Fauquembergue, E., Magnaldi, B., 2008. Les faunes de grands mammifères du site de l'Homme
- 598 de Yunxian. In : de Lumley H., Li T. (Eds.), Le site de l'Homme de Yunxian. CNRS éditions,
- 599 Paris, pp. 253-364.

600

- 601 Feng, X., 2008. Stratégie de débitage et mode de façonnage des industries du Paléolithique
- inférieur en Chine et en Europe entre 1 Ma et 400 000 ans. Ressemblances et différences de la
- culture de l'homme de Yunxian et Acheuléen européen. L'Anthropologie 112, 423-447.

604

- Feng, X., Qi, Y., Li, Q., Ma, X., Liu, K., 2018. The industry of the Lower Paleolithic site of
- Baigu, Bose Basin, Guangxi Zhuang autonome Province, PR China. L'Anthropologie 122, 14—
- 607 32.

608

- Gallotti, R., & Mussi, M., 2017. Two Acheuleans, two humankinds. From 1.5 to 0.85 Ma at
- Melka Kunture (Upper Awash, Ethiopian highlands). Journal of Anthropological Sciences 95,
- 611 137-181.

612

- 613 Garcia, T., Féraud, G., Falguères, C., de Lumley, H., Perrenoud, C., Lordkipanidze, D., 2010.
- Earliest human remains in Eurasia: New ⁴⁰Ar/³⁹Ar dating of the Dmanisi hominid-bearing
- levels, Georgia. Quaternary Geochronology 5, 443-451.

616

- 617 Glass, B.P., Koeberl, C., 2006. Australasian microtektites and associated impact ejecta in the
- South China Sea and the Middle Pleistocene superuption of Toba. Meteoritics & Planetary
- 619 Science 41, 305-326.

620

- 621 Goren-Inbar, N. Alperson-Alif, N., Sharon, G., Herzlinger, G., 2018. The Acheulian site of
- 622 Gesher Benot Ya 'aqov. Volume IV. The lithic Assemblages. Springer, Cham.

- 624 Grün, R., Aubert, M., Hellstrom, J., Duval, M., 2010. The challenge of direct dating old human
- fossils. Quaternary International 223-224, 87-93.
- 626
- Guan, Y., Wang, F-G, Xie, F., Pei, S-W, Zhou, Z-Y, Gao, X., 2016. Flint knapping strategies
- at Cenjiawan, an Early Paleolithic site in the Nihewan Basin, North China. Quaternary
- 629 International 400, 86-92.
- 630
- 631 Guo, S.-L., Huang, W., Hao, X.-H., Chen, B.L., 1997. Fission Track Dating of Ancient Man
- Site in Baise, China, and its signicances in space research, Paleomagnetism and Stratigraphy.
- Radiation Measurements 28, 565-570.
- 634
- Han, F., Bahain, J.-J., Deng, C., Boëda, E., Hou, Y., Wei, G., Huang, W., Garcia, T., Shao, Q.,
- He, C., Falguères, C., Voinchet, P., Yin, G., 2017. The earliest evidence of hominid settlement
- 637 in China: Combined electron spin resonance and uranium series (ESR/U-series) dating of
- 638 mammalian fossil teeth from Longgupo cave. Quaternary International 434, 75-83.
- 639
- Han, F., Shao, Q., Bahain, J.-J., Sun, X., Yin, G., 2018. Coupled ESR and U-series dating of
- Middle Pleistocene hominin site Bailongdong cave, China, Quaternary Geochronology 49, 291-
- 642 296.
- 643
- 644
 - Harmand, S., Lewis, J.E., Feibel, C.S., Lepre, C.J., Prat, S., Lenoble, A., Boës, X., Quinn, R.,
 - Brenet, M., Arroyo, A., Taylor, N., Clément, S., Daver, G., Brugal, J.-P., Leakey, L. Mortlocks,
 - R.A., Wright, J.D., Lokorodi, S., Kirwa, C., Kent, D.V., Roche, H., 2015. 3.3-million-year-old
 - stone tools from Lomekwi 3, West Turkana, Kenya. Nature 521, 310-315.

- Heide, K., Heide, G., Kloess, G., 2001. Glass chemistry of tektites. Planet. Space Sci. 49, 839–
- 651 844.
- 652
- Hellstrom, J., Pickering, R., 2015. Recent advances and future prospects of the U-Th and U-Pb
- chronometers applicable to archaeology. Journal of Archaeological Science 56, 32-40.
- 655
- Hou, Y., Potts, R., Baoyin, Y., Zhengtang, G., Deino, A., Wie, W., Clark, J., Guangmao, X.,
- Weiwen, H., 2000. Mid-Pleistocene Acheulean-like Stone Technology of the Bose Basin, South
- 658 China. Science 287, 1622–1626.

659

- Hou, Y.M., Zhao, L.X., 2010. An archaeological view for the presence of early humans in
- 661 China. Quaternary International 223-224, 10-19.

662

- Huang, S., Wang, W., Bae, C. J., Xu, G., Liu, K., 2012. Recent Paleolithic field investigations
- in Bose Basin (Guangxi, China). Quaternary International 281, 5–9.

665

- 666 Hyodo, M., Matsu'ura, S., Kamishima, Y., Kondo, M., Takeshita, Y., Kitaba, I., Danhara, T.,
- Aziz, F., Kurniawan, I., Kumai, H., 2011. High-resolution record of the Matuyama-Brunhes
- transition constrains the age of Javanese *Homo erectus* in the Sangiran dome, Indonesia. PNAS
- 669 108, 19563-19568.

- 671 Ingicco, T., van den Bergh, G.D., Jago-on, C., Bahain, J.-J., Chacon, M.G., Amano, N.,
- 672 Forestier, H., King, C., Manalo, K., Nomade, S., Pereira, A., Reyes, M.C., Sémah, A.-M., Shao,
- Q., Voinchet, P., Falguères, C., Albers, P.C.H., Lising, M., Lyras, G., Yurnaldi, D., Rochette,

- P., Baustita, A., de Vos, J., 2018. Earliest known hominin activity in the Philippines by 709 674
- thousand years ago. Nature 557, 233-237. 675

Jourdan, F., Renne, P.R., 2007. Age calibration of the Fish Canyon sanidine 40Ar/39Ar datin g 677 standard using primary K-Ar standards. Geochimica et Cosmochimica Acta 71, 387-402. 678

679

Jourdan, F., Mark, D.F., Vérati, C. (eds), 2014. Advances in 40Ar/39Ar dating: from 680 archaeological to planetary sciences-introduction. Geological Society, London, Special 681 Publications 378, 1–8.

682

- 683 Jourdan, F., Nomade, S., Wingate, M.T.D., Eroglu, E., Deino, A., 2019. Ultraprecise age and 684
- Meteoritics & Planetary Science 54, 2373-2591. 686

687

685

Kandyba, A.V., Chekha, A.M., Doi, N.G., Su, N. K., Gladyshev, S.A., Derevianko, A.P., 2021. 688

formation temperature of the Australasian tektites constrained by ⁴⁰Ar/³⁹Ar analyses.

- The early Paleolithic Go Da site and the Bifacial lithic industries of Southeast Asia. 689
- Archaeology, Ethnology & Anthropology of Eurasia 49, 3-14. 690

691

692 Keates, S.G., 2000. Tektites and the age paradox in Mid-Pleistocene China. Science 289, 507a.

693

- Keates, S.G., 2010. Evidence for the earliest Pleistocene hominid activity in the Nihewan Basin 694
- of northern China. Quaternary International 223-224, 408-417. 695

696

Koeberl, C., 1990. The geochemistry of tektites: An overview. Tectonophysics 171, 405–422. 697

698

Koeberl C., Bottomley R., Glass B.P., Storzer D., 1997. Geochemistry and age of Ivory Coast 699 700 tektites and microtektites. Geochimica et Cosmochimica Acta 61, 1745-1772.

701

702 Koeberl, C., Glass, B.P., 2000. Tektites and the age paradox in Mid-Pleistocene China. Science 289, 507a. 703

704

Kong, Y., Deng, C., Liu, W., Pei, S., Sun, L., Ge, J., Yi, L., Zhu, R., 2018. Magnetostratigraphic 705 dating of the hominin occupation of Bailong Cave, central China. Scientific Reports 8, 9699. 706

707

Koppers, A.A.P., 2002. ArArCALC-software for 40Ar/39Ar age calculations. Computers & 708 709 Geosciences 28, 605–619.

710

Langbroek, M., 2015. Do tektites really date the bifaces from the Bose (Baise) Basin, Guangxi, 711 southern China?. Journal of Human Evolution 80, 175-178. 712

713

Larick, R., Ciochon, R.L., 2015. Early hominin Biogeography in Island Southeast Asia. 714 Evolutionary Anthropology 24, 185-213. 715

716

- 717 Lebatard, A.-E., Alçiçek, M.C., Rochette, P., Khatib, S., Vialet, A., Bouldes, N., Bourlès, D.J.,
- Demory, F., Guipert, G., Mayda, S., Titov, V.V., Vidal, L., de Lumley, H., 2014. Dating the 718
- Homo erectus bearing travertine from Kocabaş (Denizli, Turkey) at at least 1.1 Ma. Earth and 719
- 720 Planetary Science Letters 390, 8-18.

- Lepre, C.J., Roche, H., Kent, D.V., Harmand, S., Quinn, R.L., Brugal, J.-P., Texier, P.-J., 722
- 723 Lenoble, A., Feigel, C.S., 2011. An earlier origin for the Acheulian. Nature 477, 81-85

- Li, H., Li, C., Kuman, K., 2014. Rethinking the "Acheulean" in East Asia: Evidence from recent
- 725 investigations in the Danjiangkou Reservoir Region, central China. Quaternary International
- 726 347, 163-175.

- Li, H., Li, C., Kuman, K., 2017. Longgudong, an Early Pleistocene site in Jianshi, South China,
- with stratigraphic association of human teeth and lithics. Science China Earth Sciences 60, 452-
- 730 462.

731

- Li, H., Kuman, K., Li, C., 2018. What is currently (un)known about the Chinese Acheulean,
- vith implications for hypotheses on the earlier dispersal of hominids. C.R. Palevol 17, 120-
- 734 130.

735

- Li, X., Pei, S., Jia, Z., Guan, Y., Niu, D., Ao, H., 2016. Paleoenvironmental conditions at
- Madigou (MDG), a newly discovered Early Paleolithic site in the Nihewan Basin, North Chian.
- 738 Quaternary International 400, 100-110.

739

- Li, X., Ao, H., Dekkers, M.J., Roberts, A.P., Zhang, P., Lin, S., Huang, W., Hou, Y., Zhang,
- 741 W., An, Z., 2017. Early Pleistocene occurrence of Acheulian technology in North China.
- 742 Quaternary Science Reviews 156, 12-22.

743

- Lisiecki, L.E, Raymo, M.E., 2005. A Pleicene-Pleistocene stack of 57 globally distributed
- benthic δ^{18} O records. Paleoceanography and Paleoclimatology 20, PA1003.

746

- Liu, Y., Hu, Y., Wei, Q., 2013. Early to Late Pleistocene human settlements and the evolution
- 748 of lithic technology in the Nihewan Basin, North China: A macroscopic perspective.
- 749 Quaternary International 295, 204-214.

750

- Liu, C.-R., Yin, G.-M., Deng, C.-L., Han, F., Song, W.-J., 2014. ESR dating of the Majuangou
- and Banshan Paleolithic sites in the Nihewan Basin, North China. Journal of Human Evolution
- 753 73, 58-63.

754

- Liu, Y., Hu, Y., Wei, Q., 2013. Early to Late Pleistocene human settlements and the evolution
- of lithic technology in the Nihewan Basin, North China: A macroscopic perspective.
- 757 Quaternary International 295, 204-214.

758

- Liu, X., Shen, G., Tu, H., Lu, C., Granger, D.E., 2015. Initial ²⁶Al/¹⁰Be burial dating of the
- 760 hominin site Bailong Cave in Hubei Province, central China. Quaternary International 389, 235-
- 761 240.

762

- 763 Lombera-Hermida de, A., Bargallo, A., Terradillos-Bernal, M., Huguet, R., Vallverdu, J.,
- Garcia-Anton, M.-D., Mosquera, M., Ollé, A., Sala, R., Carbonell, E., Rodriguez-Alvarez, X.-
- P., 2015. The lithic industry of Sima del Elefante (Atapuerca, Burgos, Spain) in the context of
- Early and Middle Pleistocene technology in Europe. Journal of Human Evolution 82, 95-106.

767

- Lumley de, H., Li, T., (Eds.), 2008. Le site de l'homme de Yunxian : Quyuanhekou, Quingqu,
- 769 Yunxian, Province de Hubei. CNRS Editions, Paris.

- Lumley de, H., Cauche, D., Pollet, G., Notter, O., Rossini-Notter, E., Xie, G., Lin, Q, Feng, X.,
- Ghen, X., Qi, Y., Wei, J., Wang, W., Huang, Q., 2020. Les industries lithiques du Paléolithique

- ancien du bassin de Bose. Province autonome du Guangxi Zhuang, Chine du Sud. CNRS
- 774 Editions, Paris, 325 p. ISBN: 978-2-271-13406-6

- Matinez-Navarro, B., Belmaker, M., Bar-Yosef, O., 2012. The Bovid assemblage (Bovidae,
- 777 Mammalia) from the Early Pleistocene site of 'Ubeidiya, Israel: Biochronological and
- paleoecological implications for the fossil and lithic bearing strata. Quaternary International
- 779 267, 78-97.

780

781 McDonough, W. F., Sun, S.-S., 1995. The composition of the Earth. Chemical Geology 120, 223-253.

783

Meisel, T., Lange, J.-M., Krähenbühl, U., 1997. The chemical variation of moldavite tektites: simple mixing of terrestrial sediments. Meteoritics Planetary Science 32, 493–502.

786

- Michel, V., Feng, X., Shen, G., Cauche, D., Moncel, M.-H., Gallet, S., Gratuze, B., Wei, J.,
- Ma, X., Liu, K., 2021. First ⁴⁰Ar/³⁹Ar analyses of Australasian tektites in close association with
- bifacially worked artifacts at Nalai site in Bose Basin, South China: The question of the early
- 790 Chinese Acheulean. Journal of Human Evolution, 153, 102953.

791

- Moncel, M-H., Santagata, C., Pereira, A., Nomade, S., Voinchet, P., Bahain, J-J., Daujeard, C.,
- Curci, A. Lemorini, C., Hardy, B., Eramo, G., Berto, C., Raynal, J-P., Arzarello, M., Mecozzi,
- B., Iannucci, A., Sardella, R., Allegretta, I., Delluniversità, E., Terzano, R., Dugas, P., Jouanic,
- 795 G., Queffelec, A., d'Andrea, A., Valentini, R., Minucci, E., Carpentiero, L., Piperno, M.,
- 796 2020a. The origin of early Acheulean expansion in Europe 700 ka ago: new findings at
- 797 Notarchirico (Italy). Nature. Scientific Reports 10(1), 1-16.
- 798 https://doi.org/10.1038/s41598-020-68617-8

799

- 800 Moncel M-H., Despriée J., Courcimaut G., Voinchet P., Bahain J-J., 2020b. La Noira site
- 801 (Centre, France) and the technological behaviours and skills of the earliest Acheulean in
- Western Europe between 700 and 600 kyrs. Journal of Paleolithic Archeology 3 (3), 255-301.
- 803 https://doi.org/10.1007/s41982-020-00049-2

804

Movius, H., 1948. The Lower Palaeolithic Cultures of Southern and Eastern Asia. Transactions of the American Philosophical Society 38, 329-420.

807

- 808 Mussi, M., Skinner, M.M., Melis, R.T., Pneras, J., Rubio-Jara, S., Davies, T.W., Geraads, D.,
- Bocherens, H., Briatico, G., Le Cabec, A., Hublin, J.-J., Gidna, A., Bonnefille, R., Di Bianco,
- 810 L., Mendez-Quinta, E., 2023a. Early *Homo erectus* lived at high altitudes and produced both
- Oldowan and Acheulean tools. Science 382, 713-718.

812

- Mussi, M., Altamura, F., Di Bianco, L., Bonnefille, R., Gaudzinski-Windheuser, S., Geraads,
- D., Melis, R.T., Panera, J., Piarulli, F., Pioli, L., Ruta, G., Sanchez-Dehesa Galn, S., Mendez -
- Quintas, E. 2023b. After the emergence of the Acheulean at Melka Kunture (Upper Awash,
- 816 Ethiopia): From Gombore IB (1.6 Ma) to Gombore Iγ (1.4 Ma), Gombore iδ (1.3 Ma) and
- Gombore II OAM test pit C (1.2 Ma). Quaternary International 657, 3-25.

818

O'Reilly, T.C., Haskin, L.A., King, E.A., 1983. Element correlations among north American tektites. 14th Luna Planet. Sci., 580-581 (abstract).

- Pappu, S., Gunnell, Y., Akhilesh, K., Braucher, R., Taieb, M., Demory, F., Thouveny, N., 2011.
- 823 Early Pleistocene Presence of Acheulian Hominins in South India. Science 331, 1596-1599.

- Pei, S., Deng, C., de la Torre, I., Jia, Z., Ma, D., Li, X., Wang, X., 2019. Magnetostratigraphic
- and archaeological records at the Early Pleistocene site complex of Madigou (Nihewan Basin):
- 827 Implications for human adaptations in North China. Palaeogeography, Palaeoclimatology,
- 828 Palaeoecology 530, 176-189.

829

- Pike, A.W.G., Pettitt, P.B., 2003. U-series Dating and Human Evolution. In Uranium Series
- 631 Geochemistry (edited by B. Bourdon, G.M. Henderson, C.C. Lundstrom, et al.). Reviews in
- 832 Mineralogy and Geochemistry 52, 607-630.

833

- Potts, R., Huang, W., Hou, Y., Deino, A., Baoyin, Y., Zhengtang, G., Clark, J., 2000. Tektites
- and the age paradox in Mid-Pleistocene China. Response. Science 289, 507a.

836

Qin, Z., Sun, X., 2023. Glacial-Interglacial Cycles and Early Human Evolution in China. Land
12, 1683.

839

- Renne, P.R., Balco, G., Ludwig, K.R., Mundil, R., Min, K., 2011. Response to the comment by
- W.H. Schwarz et al. on "Joint determination of K-40 decay constants and Ar-40*/K-40 for the
- Fish Canyon sanidine standard, and improved accuracy for Ar-40/Ar-39 geochronology" by PR
- Renne et al. (2010). Geochimica et Cosmochimica Acta 75, 5097-5100.

844

- Rink, W.J., Bartoll, J., Schwarcz, H.P., Shane, P., Bar-Osef, O., 2007. Testing the reliability of
- ESR dating of optically exposed buried quartz sediments. Radiation Measurements 42, 1618-
- 847 1626.

848

- Rochette, P., Beck, P., Bizzarro, M., Braucher, R., Cornec, J., Debaille, V., Devouard, B.,
- 850 Gattacceca, J., Jourdan, F., Moustard, F., Moynier, F., Nomade, S., Reynard, B., 2021. Impact
- 851 glasses from Belize represent tektites from the Pleistocene Pantasma impact crater in
- Nicaragua. Communications earth & environment 2, 94.

853

- 854 Sagi, A., 2005. Magnetostratigraphy of 'Ubeidiya Formation, Northern Dead Sea Transform,
- 855 Israel. M.S. Dissertation, The Hebrew University.

856

Scaillet, S., Vita-Scaillet, G., Guillou, H., 2008. Oldest human footprints dated by Ar/Ar. Earth and Planetary Science Letters 275, 320–325.

858 859

- 860 Schwarz, W.H., Trieloff, M., Bollinger, K., Gantert, N., Fernandes, V.A., Meyer, H.-P.,
- Povenmire, H., Jessberger, E.K., Guglielmino, M., Koeberl, C., 2016. Coeval ages of
- Australasian, Central American and Western Canadian tektites reveal multiple impacts 790 ka
- ago. Geochimica et Cosmochimica Acta 178, 307-319.

864

- 865 Semaw, S., Rogers, M.J., Quade, J., Renne, P.R., Butler, R.F., Dominguez-Rodrigo, M., Stout,
- D., Hart, W.S., Pickering, T., Simpson, S.W., 2003. 2.6-Million-year-old stone tools and
- associated bones from OGS-6 and OGS-7, Gona, Afar, Ethiopia. Journal of Human Evolution
- 868 45, 169-177.

- 870 Shen, C., Zhang, X., Gao, X., 2016. Zhoukoudian in transition: Research history, lithic
- 871 technologies, and transformation of Chinese Palaeolithic archaeology. Quaternary International
- 872 400, 4-13.

Shen, G.J., Gao, X., Gao, B., Granger, D.E., 2009. Age of Zhoukoudian *Homo erectus* determined with ²⁶Al/¹⁰Be burial dating. Nature 458, 198-200.

876

- 877 Shen, G.J., Michel, V., Despriée, J., Han, F., Granger, D., 2012. Datation d'enfouissement par
- 878 ²⁶Al/¹⁰Be et son application préliminaire à des sites du Paléolithique inférieur en Chine et en
- 879 France. L'Anthropologie 116, 1-11.

880

- 881 Shen, G., Wang, Y., Tu, Hua, Tong, H., Wu, Z., Kuman, K., Fink, D., Granger, D.E., 2020.
- Isochron ²⁶Al/¹⁰Be burial dating of Xiboudu: Evidence for the earliest human settlement in
- northern China. L'Anthropologie 124, 102790.

884

- 885 Sieh, K., Herrin, J., Jicha, B., Angel, D.S., Moore, J.D.P., Banerjee, P., Wiwegwin, W.,
- 886 Sihavong, V., Singer, B., Chualaowanich, T., Charusiri, P., 2020. Australasian impact crater
- buried under the Bolaven volcanic field, Southern Laos. PNAS 117, 1346-1353.

888

- 889 Son, T.H., Koeberl, C., 2005. Chemical variation within fragments of Australasian tektites.
- 890 Meteoritics & Planetary Science 40, 805-815.

891

- 892 Sun, X., Lu, H., Wang, S., Yi, L., Li, Y., Bahain, J.-J., Vionchet, P., Hu, X., Zeng L., Zhang,
- 893 W., Zhuo, H., 2017. Early human settlements in the southern Qinling Mountains, central China,
- 894 Quaternary Science Reviews 164, 168-186.

895

- 896 Taylor, S.R., 1962. The chemical composition of australites. Geochimica et Cosmochimica
- 897 Acta 26, 685-722.

898

- Tissoux, H., Bahain, J.-J., de Lumley, H., Li, T., Feng, X., Li, W., 2008. Essai de datation par
- 900 les méthodes combinées (RPE/U-Th) de dents d'herbivores et par RPE de quartz blanchis
- 901 extraits des sédiments fluviatiles provenant du site de l'Homme de Yunxian. In : de Lumley H.,
- 902 Li T. (Eds.), Le site de l'Homme de Yunxian. CNRS éditions, Paris, pp. 237-252.

903

- 904 Toro-Moyano, I., Martinez-Navarro, B., Augusti, J., Souday, C., Bermudez de Castro, J.M.,
- 905 Martinon-Torres, M., Fajardo, B., Duval, M., Falguères, C., Oms, O., Pares, J.M., Andon, P.,
- Julia, R., Garcia-Aguilar, J.M., Moigne, A.-M., Espigares, M.P., Ros-Montoya, S., Palmqvist,
- 907 P., 2013. The oldest human fossil in Europe, from Orce (Spain). Journal of Human Evolution
- 908 65, 1-9.

909

- 910 Tribolo, C., Mercier, N., Douville, E., Joron, J.-L., Reyss, J.-L., Rufer, D., Cantin, N., Lefrais,
- 911 Y., Miller, C.E., Porraz, G., Parlington, J., Rigaud, J.-P., Texier, P.-J., 2013. OSL and TL dating
- of the Middle Stone Age sequence at Diepkloof Rock Shelter (South Africa): A clarification.
- 913 Journal of Archaeological Science 40, 3401-3411.

914

- 915 Tu, H., Shen, G., Granger, D., Yang, X., Lai, Z., 2017. Isochron ²⁶Al/¹⁰Be burial dating of the
- Lantian hominin site at Gongwangling in Northwestern China. Quaternary Geochronology 41,
- 917 174-179.

- 919 Vallverdu, J., Saladié, P., S., Rosas, A., Huguet R., Caceres, I., Mosquera, M., Garcia-Taberno,
- 920 A., Estalrrich, A., Lozana-Fernandez, I., Pineda-Alcala, A., Carrancho, A., Villalain, J.J.,
- Bourlès, D., Braucher R., Lebatard, E., Vilalta, J., Esteban-Nadal, M., Bennasar M.L., Bastir,
- 922 M., Lopez-Polin L., Ollé, A., Vergés, J.M., Ros-Montoya, S., Martinez-Navarro, B., Garcia,
- A., Martinell, J., Exposito, I., Burjachs, F., Agusti, J., Carbonell, E., 2014. Age and Date for
- Early Arrival of the Acheulian in Europe (Barranc de la Boella, la Canonja, Spain). PLoS One
- 925 9, e103634.
- 926
- Voinchet, P., Moreno, D., Bahain, J-J., Tissoux, H., Tombret, O., Falguères, Moncel, M-H. C.,
- 928 Schreve, D., Candy, I., Antoine, P., Ashton, N., Beamish, M., Cliquet, D., Despriée, J., Lewis,
- 929 S., Limondin-Lozouet, N., Locht, J-L., Parfitt, S., Pope, M., 2015. New chronological data
- 930 (ESR and ESR/U-series) for the earliest Acheulean sites of north-western Europe. J. Quat, Sci.
- 931 30, 610–622.
- 932
- 933 Wang, H., Deng, C., Zhu, R., Wei, Q., Hou, Y., Boëda, E., 2005. Magnetostratigraphic dating
- of the Donggutuo and Maliang Paleolithic sites in the Nihewan Basin, North China. Quaternary
- 935 Research 64, 1-11.
- 936
- 937 Wang, W., Bae, C.J., Huang, S., Huang, X., Tian, F., Mo, J., Huang, Z., Huang, C., Xie, S., Li,
- 938 D., 2014. Middle Pleistocene bifaces from Fengshudao (Bose Basin, Guangxi, China). Journal
- 939 of Human Evolution 69, 110-122.
- 940
- Wang, W., Bae, C.J., 2015. How old are the Bose (Baise) Basin (Guangxi, southern China)
- bifaces? The Australasian tektites question revisited. Journal of Human Evolution 80, 171-174.
- 943
- Wang, W., Fan, Y., Bae, C., 2024. The Bose (South China) Handaxes: Early Palaeolithic
- 945 Bifaces and Early Hominin Cognition. In: The Oxford Handbook of Cognitive Archaeology
- 946 (eds. K.A. Overmann, F.L. Coolidge), chap. 31, p. 741-770.
- 947
- Wei, G., Huang, W., Chen, S., He, C., Pang, L., Wu, Y., 2014. Paleolithic culture of Longgupo
- and its creators. Quaternary International 354, 154-161.
- 950
- Wesgate, J.A., Pillans, B.J., Alloway, B.V., Pearce, N.J.G., Simmonds P., 2021. New fission-
- 952 track ages of Australasian tektites define two age groups: discriminating between formation and
- 953 reset ages. Quaternary Geochronology 66, 101113.
- 954
- 355 Xie, G.-M., Bodin, E., 2007. Les industries paléolithiques du Bassin de Bose (Chine du Sud).
- 956 L'Anthropologie 111, 182–206.
- 957
- 958 Xie, G., Chen, X., Li, D., Yu, M., Hu, Z., Lu, H., Huang, Q., Lin, Q., 2021. Stratigraphy and
- ohronology of the palaeolithic industry in Bose Basin, South China: Excavation of Gaolingpo.
- 960 Archaeological Research in Asia 26, 100284.
- 961
- 962 Yang, S.-X., Hou, Y.-M., Yue, J.-P., Petraglia, M.D., Deng, C.-L., Zhu, R.-X., 2016. The Lithic
- 963 Assemblages of Xiaochangliang, Nihewan Basin: Implications for Early Pleistocene Hominin
- 964 Behaviour in North China. PLoS ONE 11(5), e0155793.
- 965
- 966 Yang, S.-X., Petraglia, M.D., Hou, Y.-M., Yue, J.-P., Deng, C.-L., Zhu, R.-X., 2017. The lithic
- assemblages of Donggutuo, Nihewan basin: Knapping skills of Early Pleistocene hominins in
- 968 North China. PLoS ONE 12, e0185101.

970 Yang, S.-X., Deng, C.-L., Zhu, R.-X., Petraglia, M.D., 2020. The Paleolithic in the Nihewan 971 Basin, China: Evolutionary history of an Early to Late Pleistocene record in Eastern Asia. 972

Evolutionary Anthropology 29, 125-142.

973

974 Zhang, P., Huang, W., Wang, W., 2010. Acheulean handaxes from Fengshudao, Bose sites of South China. Quaternary International 223–224, 440–443. 975

976

Zhu, R.X., Hoffman, K.A., Potts, R., Deng, C.L., Pan, Y.X., Guo, B., Shi, C.D., Guo, Z.T., 977 Yuan, B.Y., Hou, Y.M., Huang, W.W., 2001. Earliest presence of humans in northeast Asia. 978 Nature 413, 413–417. 979

980 981

Zhu, R., An, Z., Potts, R., Hoffman, K.A., 2003. Magnetostratigraphic dating of early humans in China. Earth-Science Reviews 61, 341-359.

982 983 984

Zhu, R.X., Potts, R., Xie, F., Hoffmn, K.A., Deng, C.L., Shi, C.D., Pan, Y.X., Wang, H.Q., Shi, R.P., Wang, Y.C., Shi, G.H., Wu, N.Q., 2004. New evidence on the earliest human presence at high northern latitudes in northeast Asia. Nature 431, 559-562.

986 987 988

985

Zhu, Z., Dennell, R., Huang, W., Wu, Y., Qiu, S., Yang, S., Rao, Z., Hou, Y., Xie, J., Han, J., Ouyang, T., 2018. Hominin occupation of the Chinese Loess Plateau since about 2.1 million years ago. Nature 559, 608-612.

990 991 992

993

Figure 1




Figure 2




Figure 3




Figure 4






1072

Figure 5



Figure 6









Figure 8







Chondrite-nomalized after chondrite concentrations McDonough and Sun (1995)

