

High-precision 40Ar/39Ar dating of Australasian tektites associated with bifacial tools in the Bose Basin (Xiaomei and Fengshudao sites), South China and in Vietnam (Go Da and Roc Tung 1 sites)

Véronique Michel, Fred Jourdan, Marie-Hélène Moncel, Bernard Gratuze, Guanjun Shen, Wei Wang, Celia Mayers, Adam Frew, Dominique Cauche, Patricia Valensi, et al.

▶ To cite this version:

Véronique Michel, Fred Jourdan, Marie-Hélène Moncel, Bernard Gratuze, Guanjun Shen, et al.. Highprecision 40Ar/39Ar dating of Australasian tektites associated with bifacial tools in the Bose Basin (Xiaomei and Fengshudao sites), South China and in Vietnam (Go Da and Roc Tung 1 sites). Quaternary Science Reviews, 2024, 346, pp.109065. 10.1016/j.quascirev.2024.109065 . hal-04792396

HAL Id: hal-04792396 https://hal.science/hal-04792396v1

Submitted on 20 Nov 2024 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High-precision ⁴⁰Ar/³⁹Ar dating of Australasian tektites associated with bifacial tools in the Bose Basin (Xiaomei and Fengshudao sites), South China and in Vietnam (Go Da and Roc Tung 1 sites).

7	Véronique Michel ^{a,b,*} , Fred Jourdan ^c , Marie-Hélène Moncel ^d , Bernard Gratuze ^e , Guanjun
8	Shen ^f , Wei Wang ^g , Celia Mayers ^c , Adam Frew ^c , Dominique Cauche ^h , Patricia Valensi ^d , Sylvain
9	Gallet ^b , Anatoly P. Derevianko ⁱ , Alexander V. Kandyba ⁱ , Sergey A. Gladyshev ⁱ , Henry de
10	Lumley ^h
11 12	^a Université Côte d'Azur, CNRS, CEPAM, Nice, France
13 14 15	^b Université Côte d'Azur, CNRS, Observatoire de la Côte d'Azur, IRD, Géoazur, Valbonne, France
16 17 18 19	^c Western Australian Argon Isotope Facility, School of Earth and Planetary Sciences, John de Laeter Centre, Space Science and Technology Centre, Curtin University, Perth, WA, Australia.
20 21 22	^d UMR 7194 HNHP (MNHN-CNRS-UPVD), Département Homme et Environnement, Muséum National d'Histoire Naturelle, 75013 Paris, France
23 24 25	^e UMR 7065, IRAMAT, CNRS-Université d'Orleans, Centre Ernest-Babelon, 45071 Orleans, France
26 27 28	^f College of Geographical Sciences, Nanjing Normal University, Nanjing, Jiangsu 210023, China
29 30	^g Institute of Culture Heritage, Shandong University, Qingdao Shandong, 266237, China
31 32	^h Institut de Paléontologie Humaine, HNHP CNRS-MNHN, 75013, Paris, France
33 34 35 36	ⁱ Institute of Archaeology and Ethnography, Siberia Branch, Russian Academy of Sciences, Pr. Akademika Lavrentieva 17, Novosibirsk, 630090, Russia

- 37 *Corresponding author.
- 38 *Email address*: <u>veronique.michel@cnrs.fr</u> (V. Michel).
- 39 Full postal address: Université Côte d'Azur, Pôle Universitaire Saint Jean d'Angély, SJA3,
- 40 CEPAM, UMR7264, CNRS, UCA, 24 avenue des Diables Bleus, 06300 Nice, France.
- 41
- 42
- 43

44 ABSTRACT

Bifacial tools discovered at about a hundred Lower Palaeolithic sites in the Bose Basin, 45 southern China, have been previously dated to around 800,000 years ago. This age was obtained 46 by dating four tektites directly associated with the tools at the Nalai site using the ${}^{40}\text{Ar}/{}^{39}\text{Ar}$ 47 method (Michel et al., 2021). Similar ages on tektites from the Bogu and Yangwu sites had 48 49 been previously published, albeit with limited analytical details, by Hou et al. (2000). In this study, a total of eight tektites, discovered alongside abundant lithic artefacts including bifacial 50 tools, were dated with the ⁴⁰Ar/³⁹Ar technique, using an incremental temperature increase 51 approach. Six of these tektites are from two Lower Palaeolithic sites in the Bose Basin; three 52 from Xiaomei and three others from Fengshudao (China). The remaining two tektites come 53 from two sites in Vietnam, one from Go Da and the other from Roc-Tung 1. With the exception 54 of the site of Go Da, the tektites were buried in lateritic sediments associated with the stone 55 tools. At the Go Da site, the tektites were found in a layer overlying the deposit containing 56 57 bifacial tools. The tektites display no signs of fluvial abrasion or reworking, making them potentially strong chronological markers. Chemical analyses of major elements and rare earth 58 elements indicate that the tektites belong to the Australasian tektite group. This is further 59 confirmed by new ${}^{40}\text{Ar}/{}^{39}\text{Ar}$ analyses with a weighted mean age of 787.2 \pm 8.2 ka (2 σ) (MSWD 60 = 0.96, P = 0.45). Therefore, when combined with the available results on Australasian tektites, 61 this suggests a highly precise age of 788.0 \pm 2.6 ka (2 σ ; P = 0.84) for these tektites and 62 consequently, for the bifacial tools in this part of Southeast Asia, located east of the Movius 63

Line. The presence of bifacial tools attests to either a diffusion of the Acheulean from Africaor a local emergence from previous occupations and traditions in Asia.

66

Keywords: bifaces, bifacial tools, China, Vietnam, tektite, dating, ⁴⁰Ar/³⁹Ar, step-heating, MidPleistocene climate transition

69

70 **1. Introduction**

Understanding the evolution, diffusion, behavioural changes and adaptation of hominins over 71 72 time, and the impact and role of climate cycles on the distribution of occupations, is intricately linked to defining an accurate chronological framework. Many questions remain unclear, such 73 as the precise dates of the onset of Lower Palaeolithic-type industries, including the Acheulean, 74 75 in Asia (arrivals of new populations/traditions or local evolution). The most commonly utilized methods for dating events from the earliest phases of hominin occupations and dispersion, i.e., 76 2.6-0.6 Ma, are based on radioactive decay (such as K-Ar and its improved derivative 40 Ar/ 39 Ar, 77 U-Th, and U-Pb), burial dating by isotopic decay (²⁶Al/¹⁰Be), cumulative processes (Fission 78 track FT, Electron spin resonance ESR, Luminescence), or on the recording of natural processes 79 80 (such as magnetostratigraphy), often incorporating biochronological data (Pike and Pettitt, 2003; Scaillet et al., 2008; Grün et al., 2010; Shen et al., 2012; Tribolo et al., 2013; Jourdan et 81 al., 2014; Lebatard et al., 2014; Hellstrom and Pickering, 2015; Tu et al., 2017; Carbonell et al., 82 83 2010; de Lombera-Hermida et al., 2015; Carbonell et al., 2016; Harmand et al., 2015; Semaw et al., 2003; Garcia et al., 2010; Toro-Moyano et al., 2013). 84

In Asia, the earliest evidence, based on current knowledge, was discovered in Central China,
and consists of four localities (Xihoudu, Renzidong, Longuppo, Longgudong) potentially
indicating the presence of hominins. The ages are between ~2.6 and 1.8 Ma (Shen et al., 2020;
Han et al., 2017; Wei et al., 2014; Hou and Zhao, 2010; Li et al., 2017; Bae, 2024, Fig. 1, Fig.

2, Table S1). Several other early sites located in North China date between approximately 2 and 89 90 1 Ma ago, such as Shangchen, Gongwangling or sites in the Nihewan Formation (Majuangou, Xiantai, Xiaochangliang, Feiliang, Madigou, Cenjiawan, Donggutuo) (Liu et al., 2014; Zhu et 91 92 al., 2004; Zhu et al., 2018; Tu et al., 2017; Ao et al., 2010; Ao et al., 2012; Deng et al., 2006; Keates, 2010; Liu et al., 2013; Zhu et al., 2001, 2003; Yang et al., 2016; Ao et al., 2013; Li et 93 al., 2016; Pei et al., 2019; Guan et al., 2016; Liu et al., 2013; Wang et al., 2005; Yang et al., 94 2017; Yang et al., 2020, see Table S1 for details). All these sites have yielded lithic 95 assemblages, described by some as crudely-made tools. Recent dating studies have improved 96 the chronological framework of early hominin presence in this part of Eurasia, as described in 97 98 Appendix A (Table S1, Fig. 2).

The emergence and diffusion of the Acheulean (with new technological features including 99 Large Cutting Tools-LCTs) is still enigmatic. The earliest evidence is located in Africa, with 100 101 sites corresponding to the transition from the Oldowan/Mode 1 (core-and-flake industries) to the Early Acheulean/Mode 2. This occurred diachronically around 1.95-1.0 Ma (Lepre et al., 102 103 2011; Beyene et al., 2013; Mussi et al., 2023a). These African industries evolved over time into what is described as the Classical Acheulean from ~1.5 to 0.8 Ma (Mussi et al., 2023b; Galloti 104 and Mussi, 2017). Outside Africa, the Acheulean is younger, with evidence in the Levant of an 105 early "Out of Africa" dispersion for this tradition at ~1.6-1.4 Ma at Ubeidiya (Martinez-Navarro 106 107 et al., 2012; Sagi, 2005; Rink et al., 2007), possibly followed by a second wave at ~800 ka at Gesher Benot Yakov (Goren-Inbar et al., 2018), suggesting, at the least recurrent phases of 108 diffusion outside Africa. 109

110 Moving to Asia, two hypotheses have been put forward regarding the emergence of the 111 Acheulean culture beyond the "Movius Line" in an area where LCTs were considered to be 112 absent (Movius, 1948; Bae, 2024): either this represents the diffusion of the Acheulean from

Africa, as observed in the Near East, or it is a local emergence from previous occupations and
traditions in Asia (Bae, 2024).

It is therefore essential to accurately date sites with LCTs (bifaces/bifacial tools/handaxes 115 116 according to studies and authors) in Eurasia in order to contribute to discussions on the onset of this/these new traditions(s) in Asia, and to better define the possible diffusion of what is 117 described in Africa as the Acheulean. Only we then will gain insights into Pleistocene hominin 118 119 mobility and behavioural strategies in response to a continually changing climate (Bae et al., 2018, Larick and Ciochon, 2015; Qin and Sun, 2023). For convenience, throughout our paper, 120 we use the generic terminology of LCTs to designate lithic assemblages with bifaces, handaxes 121 and bifacial tools. 122

In Michel et al. (2021), tektites in close association with LCTs at the Nalai site (Bose Basin, 123 South China) were dated to 809 ± 12 ka from 55 total fusion subsamples. A probability density 124 distribution yielded an age centred at 797 ± 6 ka, tentatively suggesting that they might be part 125 of the Australasian textite strewnfield recently dated to 788 \pm 3 ka (2 σ , P = 0.54) by the 126 40 Ar/ 39 Ar method (Jourdan et al., 2019). In this new study, our aim is to improve the accuracy 127 of our preliminary dating result of the Bose sites (Michel et al., 2021) and obtain new dates for 128 two sites in Central Vietnam vielding LCTs. We dated tektites via ⁴⁰Ar/³⁹Ar analysis by the 129 "step-heating" method in order to circumvent inherited argon present in tektites during the 130 initial steps. Furthermore, with this approach, inverse isochron ages can be calculated, which 131 resolves and takes into account any inherited ⁴⁰Ar* present in the samples in the calculation, 132 provided that its argon isotopic composition is homogenous, thus leading to more accurate 133 results (Jourdan et al., 2019). ⁴⁰Ar/³⁹Ar dating analyses were conducted on tektites from two 134 archaeological sites located in the Bose Basin in China and from two others sites located in 135 Vietnam (An Khê): Xiaomei, Fengshudao, Go Da and Roc Tung 1, respectively (Fig. 1, Fig. 136

137 S1, Fig. S2). Major and trace element geochemistry was applied to tektites to assess whether
138 they were part of the Australasian strewnfield.

139

140 2. Australasian tektites: origin, background chronology and chronological markers

Tektites are glasses produced by the melting of the continental crust during the impact of a 141 meteorite. This impact causing the formation of Australasian tektites is supposed to have 142 143 resulted in the formation of a crater located in Indochina, or more likely, in southern Laos. According to the recent study by Sieh et al. (2020), the crater would be buried under the Bolaven 144 volcanic field, near the northern limit of the strewnfield (Fig. 1), although firm evidence is still 145 146 lacking. These tektites have been found at varying distances from the impact structure, spanning regions from Southeast Asia to Antarctica, with a chemical composition characteristic of the 147 group of Australasian tektites (Koeberl, 1990; Heide et al., 2001; Glass and Koeberl, 2006; 148 Schwarcz et al., 2016). 149

Tektites closely associated with artefacts from the Bose Basin in China were initially dated using fission track dating (Guo et al., 1997). The authors obtained an age of 732 ± 39 ka for the tektites of the Bogu site (Fig. S1). Subsequently, the 40 Ar/ 39 Ar method was applied to two tektites from the Bogu site and one from the Yangwu site, but these results were published without age spectra or detailed analytical data (Hou et al., 2000) (Fig. S1). The authors obtained an overall weighted mean age of 803 ± 6 ka (2σ) from the isochron ages of three samples (Hou et al., 2000; Potts et al., 2000).

This first ⁴⁰Ar/³⁹Ar dating of tektite in Bose Palaeolithic sites, presented by Hou et al. (2000) was questioned by Koeberl and Glass (2000). The latter authors cast doubt on the association of tektites with the artefacts and claimed that tektites are generally found in deposits younger than their isotopic age. They thus called for caution in using them as stratigraphic markers (Koeberl and Glass, 2000; Keates, 2000). Potts et al. (2000) responded clearly, affirming that

the Bose tektites are *in situ* and associated with the stone tools in terrace T4, as described by 162 163 Hou et al. (2000), in the same way as the stone tools found alongside tektites at the Gaolingpo site (Xie et al., 2021). They attributed the tektite age to the tools as there is no geomorphological 164 and stratigraphical indication of the redeposition of either the tools or tektites (Potts et al., 165 2000). Later, Wang et al. (2014) presented a detailed morphological study of the Fengshudao 166 167 site and its bifacial industry. In this site, the first excavated association of tektites, dating to ca. 803 ka, with LCTs (including bifaces), and found in situ, was described in one stratigraphic 168 level in terrace T4, indicating that the Bose Basin handaxes most likely date to the Early to 169 Middle Pleistocene transition (Fig. 2). However, according to Langbroek (2015), the age of the 170 171 earliest bifaces in China and the close association between tektites and bifaces have not yet been demonstrated, although this was clearly established by Wang and Bae (2015), and later 172 confirmed at the Nalai site with Australasian tektites and bifaces found together in situ (Michel 173 174 et al., 2021). It is important to note that two Australasian tektites found *in situ* at the Sangiran area have been successfully used as chronological markers. They are consistent with the 175 176 magnetic polarity stratigraphy and predate the Matuyama-Brunhes transition (MB) (Hyodo et al., 2011). The data demonstrate the near co-occurrence of the tektite level and the 177 magnetostratigraphic constraint, arguing that the last occurrence of Sangiran hominins dates to 178 ~ 0.79 Ma (Fig. 2). 179

Later, Jourdan et al. (2019) conducted precise dating on four Australasian tektites from Western Australia, Thailand, Vietnam, and southern China, including a tektite from the Bose Basin previously dated by Hou et al. (2000). The exact origin of the tektite from the Bose Basin was not specified by Jourdan et al. (2019). Age plateau and isochron diagrams yielded concordant data with a weighted-mean age of 788.1 \pm 2.8 (P = 0.54) for these Australasian tektites, which aligns closely with the less precise 40 Ar/ 39 Ar age of 793 \pm 14 ka published by Schwarz et al. (2016).

3. Xiaomei and Fengshudao sites in Southern China: Geography, stratigraphy and lithic assemblages

The history of Palaeolithic research in the Bose Basin has been thoroughly documented by Wang et al. (2014). Approximately a hundred sites have been discovered in this basin (de Lumley et al., 2020; Wang et al., 2024). In these three abovementioned studies, the geography/geology of the Bose Basin with its Quaternary terrace stratigraphy (see Fig. S3, Fig. S4), as well as overviews of the site's location and lithic assemblages, are also described in detail.

195 The Xiaomei site is located in Xiaomei village, Youjiang district, Bose city, Guangxi Zhuang Autonomous Region (23°46'41.5" N, 106°42'21.9" E), on the fourth terrace of the Youjiang 196 River (Fig. S1, Fig. S3, Fig. S4, Fig. S5). It was discovered in the 1980s during an 197 archaeological survey covering an area of 350,000 m². Many lithic artefacts and a few tektites 198 were exposed on the ground surface of the laterite deposits at the site. The lithic series were 199 made up of diversified LCTs, including pick and tongue-shaped bifaces typical of the area (Fig. 200 3, n° 6), on sandstone or quartzite pebbles (Fig. 3, n° 7 and 8), associated with flake-tools 201 (scrapers), and notably a large unifacial core on a massive thick quartzite pebble (Fig. 3, n° 9) 202 with centripetal removals. A test excavation conducted in 1996 unearthed artefacts, associated 203 204 with tektites. Based on geomorphology and artefacts, the age of this site is assumed to be the same as that of Bogu site. 205

The Fengshudao site $(23^{\circ}57'39''N, 106^{\circ}40'39''E)$ is situated along the south bank of the Chengbihe River, a significant tributary of the Youjiang River (Fig. S1, Fig. S3, Fig. S4). The site is located in the fourth terrace, spanning 100 metres from east to west, with a sloping terrain from north to south of approximately 10 to 30 metres. Discovered in 2004, the site was excavated from 2004 to 2005, revealing nearly 100 artefacts and nine tektites *in situ* from an excavated area of 62.6 m² (Wang et al., 2014). The lithic assemblage at the Fengshudao site includes the same diversity as at Xiaomei (Fig. 3, n° 5) with in addition, evidence of the use of bipolar debitage. The main raw materials consist of sandstone, quartzite and quartz collected as pebbles. Both tektites and LCTs are found in exactly the same layer within the laterite (Wang et al., 2024) (Fig. 4).

The bifaces exhibit the characteristic shape of the Bose Basin bifaces (Hou et al., 2000; Xie and Bodin, 2007; Zhang et al., 2010; Huang et al., 2012; Wang et al., 2014; Feng et al. 2018; Wang et al., 2024). They are notably short, broad and stocky, featuring a tongue-shaped extremity (Fig. 3, n° 1, 2 and 3).

220

4. Go Da and Roc Tung 1 sites in Central Vietnam: Geography, stratigraphy and lithic assemblage

Following the discovery of sites with LCTs near the city of An Khê in Central Vietnam in 2014,
by Vietnamese archaeologists, systematic excavations have been conducted since 2015
(Derevianko et al., 2018) (Fig. 1, Fig. S2).

226 Several other similar Palaeolithic sites were identified at the Roc Tung location, situated on the left bank of the Ba River. Artefacts were collected on the surface or discovered in situ during 227 excavations and test pits, with similar stratigraphic contexts. The Roc Tung 1 site (14°02'253" 228 229 N; 108°40'822" E) is situated on a high and relatively flat area of agricultural land. The stratigraphy is well-documented through excavations and various test pits (Derevianko et al., 230 2018). Three layers have been identified below the surface debris of an irrigation canal. The 231 cultural layer, situated at the top of the laterite (Layer 3), has yielded several in situ artefacts 232 associated with tektites (Fig. S6, Fig. 5). 233

Go Da (13°58'306" N; 108°39'136" E; 440 m a.s.l.) is located downstream, near An Khê, on a
sloping hill, 900 m west of the Ba River (Fig. S2). The stratigraphy established during the

excavation is somewhat different on this right bank of the Ba (Derevianko et al. (2018);

Kandyba et al. (2021) (Fig. S7). The stratigraphic sequence begins with a substantial weathering 237 238 crust (layer 1) of the granite bedrock, succeeded by slope wash sediments (layer 2). The cultural horizon at Go Da formed during these erosion and slope wash processes, probably in a cooler 239 240 climate than the present one. Archaeological materials are specifically situated at the top of the weathering crust and at the base of the slope wash layer (Fig. S7). Tektites found in the upper 241 242 part of this layer 2 indicate that hominins were present at Go Da before the deposition of 243 Australasian tektites. In other words, it can be asserted that the industries at Go Da predate those at Roc Tung 1, where artefacts are found *in situ* with tektites. 244

The local industry is well described by Derevianko et al. (2018) and Kandyba et al. (2021). The lithic assemblage from the Roc Tung 1 site, collected on the surface but also in stratigraphy context during excavations and test pits, includes over 126 pieces (Derevianko et al. 2018), made on pebbles or boulders. This lithic industry from the left bank of the Ba River is made on local raw materials, most often quartzite cobbles or blocks, but also other raw materials of lesser quality, such as vein quartz. Many tested pebbles have been identified in the assemblage, no doubt corresponding to knappers' efforts to select good-quality blanks (Fig. 6).

As in the other An Khê-tradition assemblages (Derevianko et al. 2018), the main features of the industry at Roc Tung 1 are heavy-duty-tools including numerous LCTs (bifaces, picks), choppers, and chopping-tools (Fig. 6, n^o 2 and 3) and light-duty tools, such as scrapers and pointed tools, often made on flakes. Finally, flakes and cores attest to simple reduction processes: cores mainly bear a single debitage surface with unipolar or centripetal removals and a limited prepared or cortical striking platform.

The lithic assemblage at the Go Da site comprises around a hundred artefacts (Derevianko et al. 2018; Kandyba et al 2021), comparable to those from Roc Tung 1. These consist of numerous split tested pebbles, often large in size, hammerstones on rounded granite pebbles, a heavy-duty tool kit with mainly thick, short and stocky LCTs (including bifaces and picks), on quartzite pebbles or slabs. Some display a tongue-shaped extremity. Others are choppers on long and massive quartzite pebbles, and pointed tools and scrapers (Fig. 6, n°1). The cores, flakes and small flakes indicate *in situ* reduction processes, mainly unifacial and unipolar removals. This assemblage displays all the characteristics of the An Khê industry, as well as regional traditions.

These two assemblages from Roc Tung 1 and Go Da share many similarities to industries from
the Bose Basin in China, notably those from Fengshudao and Xiaomei, discussed in this study.

270 **5. Materials and methods**

271 5.1. Materials.

Our analysed tektites come from the Fengshudao and Xiaomei sites (Bose Basin, China), and Roc Tung 1 and Go Da sites (Central Vietnam). They were discovered *in situ* associated with tools (including LCTs), except for the Go Da site, where the tektites were found in a layer overlying bifacial tool-bearing deposits, providing a minimum age for the artefacts.

Three tektites (F1, F2, F3) collected from the laterite layer of terrace T4 at the Fengshudao site (Fig. 7), display a very thin profile (-0.4-0.6 g) and show no signs of fluvial abrasion. They are very angular and sharp, suggesting that no river re-depositing occurred. Additionally, three tektites, namely B1 (10.2 g), B2 (8.7 g), and B3 (6.5 g), were excavated from the same laterite layer from terrace T4 at the Xiaomei site, with a rounder morphology than the others (Fig. 7).

Two tektites from Central Vietnam; one larger tektite, GD (19.1 g), excavated at the Go Da site, and another, large and very angular tektite, RT1 (10.3 g), with no signs of fluvial abrasion, excavated at the Roc Tung 1 site, complete the sample. The GD tektite is rounder, similar to the Xiaomei tektites studied here, but neither show signs of fluvial abrasion or reworking.

285 5.2. Method

Before analysis, the eight tektite samples were cleaned in an ultrasonic water bath to remove 286 287 surface-bonded residues (Fig. 7). The thinner tektites (F1, F2, F3) were reduced to pieces. For the larger ones (B1, B2, B3, GD, and RT1), a piece of each tektite was extracted from the 288 original tektites and reduced to pieces. Some were further crushed in an agate mortar and 289 divided into subsamples for multiple analyses. For ⁴⁰Ar/³⁹Ar dating, subsamples 400-500 µm 290 in size, from F1, F2, B1, B2, B3, GD and RT1 tektites, were cleaned with an HNO₃ 2N 291 ultrasonic bath for 15 minutes, in HF 2% for 20 minutes, in H₂O, and then in acetone. 292 Subsamples weighing 30-70 mg were analysed using the "step-heating" method with 293 incremental temperature increases to produce an age spectrum and inverse isochron. The 294 ⁴⁰Ar/³⁹Ar analyses were conducted at the Western Australian Argon Isotope Facility (Australia) 295 using an ARGUS VI multicollection instrument and following the procedure described by 296 297 Jourdan et al. (2019), used for dating the Australasian textites. The raw data (Appendix B) were 298 processed using the ArArCALC software (Koppers, 2002) and the ages were calculated using 299 the FCs standards (Jourdan and Renne, 2007), for which an age of 28.294 Ma ($\pm 0.13\%$) was 300 used (Renne et al., 2011), along with the decay constants recommended by Renne et al. (2011). Our criteria for plateau determination are as follows: plateaus must include at least 70% of ³⁹Ar. 301 The plateau should be distributed over a minimum of three consecutive steps agreeing at 95% 302 confidence level and satisfying a probability of fit (P) of at least 0.05. Plateau ages are given at 303 the 2σ level and are calculated using the mean of all the plateau steps, each weighted by the 304 inverse variance of their individual analytical error. Uncertainties include analytical and J-value 305 306 errors. Errors with all sources of uncertainties such as the decay constants errors increase the 307 uncertainty by ± 0.2 ka for those samples and are indicated by a square bracket (e.g., $[\pm 3 \text{ ka}]$). 308 The composition of major and trace elements presented in Tables S2 and S3, was determined for all tektites by Laser Ablation Inductively Coupled Plasma Mass Spectrometry at the Centre 309 Ernest-Babelon of the IRAMAT (CNRS/Orléans University) or by Inductively Coupled Plasma 310

311 Optical Emission Spectrometry at SARM (Service d'Analyses des Roches et des Minéraux,

312 CRPG Nancy, France (see additional details in SI). The composition of major elements of

tektites N411, N425, N862, N959 from Nalai site in the Bose Basin, China, as reported by

Michel et al. (2021), was supplemented with the analysis of rare earth elements and is presentedin Table S3.

- 316
- 317 **6. Results**
- 318 *6.1 Chemical composition*

The major elemental compositions obtained for the tektites B1, B2, B3 (Xiaomei), F1, F2, F3 319 320 (Fengshudao), GD (Go Da) and RT1 (Roc Tung 1), are presented in Table S2. They comprise the following ranges: 70.05-74.74 % for SiO₂, 11.13-15.44 % for Al₂O₃, 4.17-4.96 % for FeO, 321 1.77-2.67 % for CaO, 2.06-2.42 % for K₂O, 1.92-2.67 % for MgO, 0.93-1.51 % for Na₂O, 0.68-322 323 0.78 % for TiO₂, 0.08-0.10 % for MnO, 0.03-0.07 % for P₂O₅. These values are very close to those of the tektites from Nalai (N411, N425, N862, N959) (Michel et al., 2021) (Table S2). In 324 325 Figure 8, these values are compared with various groups of tektites according to Schwarz et al. (2016) and Wesgate et al. (2021). Figure 8 shows that our tektites belong to the group of 326 Australasian tektites. Tektites from Vietnam (GD and RT1) are very slightly different to those 327 from the Bose Basin. They belong to the subgroup of indochinite showing some regional 328 compositional variation in comparison with the australite subgroup, but ultimately belong to 329 the same Australasian group. In the same way as the Australasian tektites, our tektites exhibit 330 light rare-earth-element-(REE)-enriched compositions with a marked negative EU anomaly, as 331 described by Koeberl (1990), Son and Koeberl (2005), Amare and Koeberl (2006), Wesgate et 332 al. (2021) and Schwarz et al. (2016) (Fig. 9). 333

334

335 $6.2^{40} Ar / {}^{39} Ar dating$

We obtained seven plateau ages ranging from 779 \pm 19 ka (MSWD = 0.53; P=0.93) to 827 \pm 336 57 ka (MSWD = 0.09; P = 1.00) and including between 91 and 100 % of the total released ³⁹Ar 337 (Fig. 10, Appendix B). Inverse isochrons yielded similar ages ranging from 774 ± 24 to $842 \pm$ 338 67 and with trapped ${}^{40}\text{Ar}/{}^{36}\text{Ar}$ ratios in error of the atmospheric value of 298.6 \pm 0.3 and ranging 339 from 293.3 \pm 4.9 to 300.8 \pm 6.0 (Appendix C). The K/Ca ratio (derived from ³⁹Ar_K/³⁷Ar_{Ca}) 340 shows homogenous values across a given tektite but varies among samples with average values 341 ranging from 0.32 ± 0.18 to 1.29 ± 0.11 . The seven plateau ages (P = 0.84) and inverse isochron 342 ages (P = 0.25) are all indistinguishable from each other, indicating the dating of a single event, 343 as expected for tektites originating from a single impact. Furthermore, all intercept ⁴⁰Ar/³⁶Ar 344 345 values define a single population (P = 0.57) with a weighted mean of 296 ± 3 , suggesting that no inherited ⁴⁰Ar* is present in the samples analysed in this study. Therefore, we calculated a 346 single weighted mean age of 787.2 \pm 8.2 ka (2 σ ; MSWD = 0.96; P = 0.45) as the best estimate 347 for the impact event that produced these tektites. This age is indistinguishable from the previous 348 age of 788.1 ± 2.8 (Jourdan et al., 2019), indicating that the tektites are part of the Australasian 349 strewnfield, as also evidenced by their chemical composition. When combined with our new 350 age, we obtained a conclusive best age estimate of 788.0 \pm 2.6 [2.8] ka (2 σ ; P = 0.84) for the 351 Australasian tektites that will be used in the rest of this study. 352

353

7. Discussion on the new results of the Chinese and Vietnamese sites regarding the

355 emergence and diffusion of the Acheulean in Eurasia

356 7.1 Emergence of the Acheulean in Africa and its initial diffusion

The emergence of the Acheulean in Africa is dated to around 1.75 to < 1.0 Ma (i.e., in southern Ethiopia, Beyene et al., 2013) and around 1.76 Ma years ago at Kokiselei 4 in Kenya (Lepre et al., 2011) but may have occurred earlier, around 1.95 Ma years ago at Garba IV (Upper Awash, Ethiopia, Mussi et al., 2023a). This shift was characterized among others (flaking methods,

land-use patterns) by hominin capability to produce large flakes and LCTs, including for 361 362 example, handaxes, picks and sometimes cleavers. The earliest LCTs (including bifaces), outside Africa are reported at ~1.4-1.2 Ma ago in Ubeidiya in the Levant (Martinez-Navarro et 363 al., 2012; Sagi, 2005; Rink et al., 2007), and are also dated to ~1.5 Ma in Attirampakkam, India, 364 indicating a rapid diffusion throughout Eurasia, if we consider that these industries are related 365 to the African Acheulean. These types of LCTs are not present, earlier than 0.9-0.7 Ma in 366 Western Europe (Pappu et al., 2011; Vallverdu et al., 2014; Voinchet et al., 2015; Antoine et 367 al., 2019; Moncel et al., 2020a; Moncel et al., 2020b). In Asia, the core-and-flake series (Mode 368 1) persist when Lower Palaeolithic/Acheulean (Mode 2) industries with LCTs (bifaces/bifacial 369 370 tools/handaxes) appear in some areas around 1.5-0.8 Ma ago. The number of discovered sites increase during the Mid-Pleistocene climate transition (MPT) (1.0-0.6 Ma) suggesting that this 371 372 period played a crucial role in the diffusion of hominins in Eurasia (Fig. 2, Table S1).

373

374 7.2. New insights from Chinese and Vietnamese sites

375 These new assemblages with LCTs in Asia are currently found in four main Chinese areas (specifically Dingcun, Bose Basin, Luonan Basin, and the Danjiangkou Reservoir Region 376 DRR) (Li et al., 2018, Li et al., 2014). Unfortunately, dating details are limited, except for the 377 site of Yunxian (open-air site of Xuetangliangzi, DRR region, Central China), where two Homo 378 erectus crania were discovered with one handaxe and 13 LCTs collected from the surface (Li 379 et al., 2018). This site is approximatively dated between ~1.1-0.85 Ma using both 380 paleomagnetism, biochronology and ESR dating on teeth and quartz (de Lumley and Li, 2008; 381 Echassoux et al., 2008; Tissoux et al., 2008; Feng, 2008) (Fig.1, Fig.2, Table S1). Many other 382 sites with LCTs, up to 200 LCTs, were discovered in this area but further chronological research 383 is required. They are regarded as the most significant persistence of LCTs for a long period, 384 from the late Early Pleistocene to the early part of the Late Pleistocene (Li et al., 2018). 385

386 Our results on four Chinese and Vietnamese sites provide strong and accurate data on Asian 387 human occupations with LCTs dated to ~ 788 ka. Through these new data, we contribute to 388 understanding of the chronology and diffusion of the earliest populations in Asia.

Early human populations thrived and expanded across mainland East Asia, from low latitudes 389 whatever their technology (e.g., the Bose Basin) to high northern latitudes (e.g., the Nihewan 390 391 Basin, Zhoukoudian). The associated industry is often roughly dated using paleomagnetism, as 392 seen in Maliang (Nihewan Formation, North China), Wutaicun (Central China), Longgangsi-2 and 3 (Central China). The determined ages are approximately ~0.78 Ma (Wang et al., 2005), 393 ~0.9 Ma and ~1.2-0.7 Ma (Sun et al., 2017) respectively. Despite approximate dating, an 394 395 overview of paleoenvironmental and archaeological data in the Nihewan Basin reveals changes in hominin presence and indicates that sustained occupation in high latitudes characterized by 396 strong seasonality was not possible until the Late Pleistocene, as evidenced by chronological 397 398 gaps (Yang et al., 2020). Mode1/Oldowan industries persisted in Asia at Bailong Cave with human teeth in Central China, dated between ~0.78 Ma and ~0.51 Ma, and also using multi-399 dating methods such as paleomagnetism, ²⁶Al/¹⁰Be burial on quartz and ESR/U-series on teeth 400 (Liu et al., 2015; Han et al., 2018; Kong et al., 2018). 401

401 (Liu et al., 2015, Hall et al., 2016, Kolig et al., 2016).

Moving outside China, this lithic industry is also evidenced at the site of Kalinga between 402 403 around 1.05 and 0.63 Ma, in the Philippines (Ingicco et al., 2018), and between around 1.01 and 0.7 Ma at Mata Menge and Wolo Sege in Indonesia (Brumm et al., 2016). During the same 404 period, an Acheulean-type industry is present at Shuigou-Huixinggou, in Central China, but 405 approximatively dated to ~0.9 Ma by paleomagnetism (Li et al., 2017). Other Mode 1/Mode 2 406 sites dated by alternative radioisotopic methods provide more precise dating (Fig.2, Table S1). 407 This is the case for Zhoukoudian Locality 1, in North China, where human remains and 408 409 industries with bifacial tools and flake-tools (Derevianko, 2008) are well dated to ~0.78-0.4 Ma using multi-dating methods, such as paleomagnetism, ²⁶Al/¹⁰Be burial and U-Th (Shen et al.,
2009, Shen et al., 2016).

Li et al. (2018) propose that the technological transmission hypothesis could be a suitable 412 413 explanation for the emergence of Chinese LCTs technology and suggest a diffusion of the Acheulean-type tradition by hominins from west to east to China and other Asia aeras as 414 415 Vietnam. Other authors propose a local onset derived from previous Mode 1/core-and-flake 416 assemblages (convergence), giving rise to the An Khê tradition observed at the Go Da and Roc Tung Lower Palaeolithic sites in Vietnam (Derevianko et al., 2018) and the Acheulean in the 417 Bose Lower Palaeolithic sites (Wang et al., 2014; Wang and Bae, 2015; Wang et al., 2024; Bae, 418 419 2024) (Fig.1, Fig.2, Table S1).

Our results highlight that the industries at the Bose and An Khê sites are now well dated to 788.0 \pm 2.6 ka (2 σ) by dating *in situ* Australasian tektites associated with the bifaces. This significant chronological marker accurately positions the industries at the Bose and An Khê sites within the chronological sequence of industries in Asia and also indicates a wellestablished presence of LCTs and possibly the Acheulean-type tradition in this region of Asia.

425

426 **8. Conclusion**

Our dating of Australasian tektites discovered in situ alongside artefacts at the Bose sites in 427 southern China and An Khê in Central Vietnam using ${}^{40}\text{Ar}/{}^{39}\text{Ar}$ gives an age of 788.0 ± 2.6 428 [2.8] ka. This age indicates that LCTs (including bifaces), and thus what can be described as 429 the Acheulean, developed in this part of Asia around 788 ka during the Mid-Pleistocene climate 430 transition, regardless of the hypothesis regarding the onset of the Acheulean (Fig. 2, Table S1). 431 Our result can be compared to the age of the Acheulean industry also found at around 1.1-0.8 432 Ma at DRR sites, Yunxian and Shuigou-Huixinggou in Central China. This new LCT-making 433 tradition is penecontemporaneous with persistent core-and-flake series (Mode 1) at Nihewan in 434

northern China, at Bailong, as well as Longgangsi in Central China. This persistence of
assemblages without LCTs extends to the Philippines at Kalinga and Indonesia at Mata Menge
and Wolo Sege, with for some, the presence of *Homo erectus* while the Acheulean-type
assemblages persist at Zhoukoudian (~0.8-0.6 Ma) in North China. The Bose and An Khê sites
can be considered as the earliest well-dated evidence of Acheulean-type tradition beyond the
Movius Line in eastern Asia, showing that hominins in these areas were capable of making
LCTs, including bifaces at ~788 ka.

442

443 Acknowledgements

This work was carried out with partial financial support from CEPAM-CNRS. Louise Byrne is
acknowledged for English improvements of the manuscript. The Handling Editor, Christopher
J. Bae, and the two anonymous reviewers are thanked for their constructive comments and
helpful suggestions for improving this manuscript.

448

449 **Figure captions**

Figure 1. Portion of the Australasian tektite strewnfield (modified from Wesgate et al., 2021) 450 with a potential crater impact in southern Laos (Sieh et al., 2020). The Movius Line is 451 represented by the red dashed line, separating Western Asia, where LCTs were discovered from 452 Eastern Asia, where LCTs, were considered absent (Movius, 1948). However, since then, 453 454 Acheulean assemblages with LCTs have been discovered east of the line (Bae, 2024). The archaeological sites are located south of the 45°N parallel and are classified in chronological 455 order. The sites are dated using methods such as radiometric dating and/or relative dating, 456 including biochronology and paleomagnetism, based on current knowledge (Table S1). The 457 distribution indicates sites with Mode 1 (core-and-flake assemblages) (marked with stars), those 458

with Homo remains (marked with circles), and those associated with Acheulean/Mode 2 459 460 (marked with triangles). Red symbols represent the studied sites, while black symbols denote other key sites (see details in Tab S1): (1) Masol, India, (2) Riwat, Pakistan, (3) Attirampakkam, 461 India, (4) Renzidong, Central China, (5) Xihoudu, northern China (6) Longgupo, Central China, 462 (7) Longgudong, Central China, (8) Shangchen, Central China, (9) Gongwangling (Lantian 463 man), North China, (10) Niujianbao, Southwest China, (11) Many sites in the Nihewan Basin, 464 465 northern China: e.g. Donggutuo, Maliang, Feilang Xiaochangliang, Xiantai, Majuangou, Madigou, Cenjiawan Xiaochangliang, (12) Sangiran, Bapang, Lower Lahar unit, Central Java, 466 Indonesia, (13) Longgangsi 3-2, (14) Soa Basin, Ola Bula Formation, Mata Menge, Wolo Sege, 467 468 Central Flores, eastern Indonesia (15) Yunxian, DRR sites, Central China, (16) Wutaicun, Central China, (17) Shuigou-Huiximggou, Central China, (18) Meipu, Central China, (19) 469 Trinil, Java, Indonesia, (20) Many sites in the Bose Basin, Bogu, Yangwu; Nalai, Fengshudao, 470 471 Xiaomei (this study), (21) Roc Tung 1 and Go Da, Central Vietnam (this study), (22) Kalinga, Philippines, (23) Bailong, Central China, (24) Zhoukoudian Locality 1, northern China, (25) 472 473 Yiyuan, northern China.

474

Figure 2. Chronological sequence of Palaeolithic sites with hominin fossils and/or lithic assemblages according to geographical location and correlation with marine isotope stages and δ^{18} O stacks (Lisiecki and Raymo, 2005). The horizontal grey band denotes the Mid-Pleistocene climate transition MPT (1.0-0.6 Ma).

479

Figure 3. Lithic artefacts from Fengshudao site (1 to 5) and from Xiaomei site (6 to 9). 1 to 3tongue-shaped bifaces (N007612; N007622; N007623); 4- chopper (N007775); 5- large flake
(N007566); 6-tongue-shaped biface (N73 H05); 7 & 8- picks (N94 5.6; N193 G2); 9- large core
(N59).

19

484

Figure 4. Fengshudao site excavation. Photographs showing the *in situ* discovery of biface No.
Z007773 (2) in the laterite layer, in close association with tektite No. FSD-059 (1).

487

Figure 5. Roc Tung 1 excavation. Photograph showing an *in situ* discovery of a pebble core
No 16.RT1.ST79 (2) in the laterite layer (Layer 3), in close association with a tektite No 14 (1).
Bifaces were excavated in the same layer.

491

492 Figure 6. Lithic artefacts from Go Da site (1) and from Roc Tung 1 site (2 and 3). 1- scraper;
493 2- chopper; 3- biface.

494

Figure 7. Photograph of tektites analysed in this study: F1, F2, F3 (~0.4-0.6 g) are three thin
tektites excavated from Fengshudao site, Bose Basin, China; B1 (10.2 g), B2 (8.7 g), B3 (6.5
g) are three tektites excavated from Xiaomei site, Bose Basin, China; GD (19.1 g) is a tektite
excavated from Go Da site and RT1 (10.3 g) is a tektite excavated from Roc Tung 1 site,
Vietnam. GD and RT1 were cleaned with a H₂0 ultrasonic bath.

500

Figure 8. Chemical composition of various tektite groups, based on the original figure by Schwarz et al. (2016) with modifications. This figure includes a comparison of compositions with cited references by Schwarcz et al. (2016) (Taylor, 1962; Chapman and Schreiber, 1969; O'Reilly et al., 1983; Koeberl et al., 1997; Meisel et al., 1997; Albin et al., 2000; Son and Koeberl, 2005; Amare and Koeberl, 2006), including Belize (BZ1) (Central America, with impact glasses formed at 804 ± 9 ka, Rochette et al., 2021), Ivory Coast tektite (Ivorite), and Australasian tektites such as Indochinite B (Muong-Nong), Western Canada (WC1) with impact glasses formed at 785 ± 7 ka (40 Ar/ 39 Ar age, Schwarcz et al., 2016), and those with a weighted FT age of 822 ± 20 ka from Malaysia, Java, Australia (group A) (Wesgate et al., 2021). Additionally, we present the chemical compositions of tektites from Nalai, Xiaomei, Fengshudao, Roc Tung 1 and Go Da sites, demonstrating similarities to the Australasian tektite group.

513

Figure 9. Rare earth element abundances for tektites N411, N425, N862, N959 (Nalai site); F1, 514 F2, F3: tektites (Fengshudao site); B1, B2, B3 (Xiaomei site); GD (Go Da site); RT1 (Roc Tung 515 516 1 site). They are chondrite-normalized rare-earth-element (REE) abundance curves (chondrite concentrations are from McDonough and Sun, 1995). The abundances are compared with 517 various tektites, including Belize (BZ1), Ivory Coast tektite 1, and Australasian tektites, such 518 519 as Indochinite B (Muong-Nong), Western Canada (WC1) (Schwarcz et al., 2016), and those from Malaysia (UT1389), Java (UT2336), Australia (UT2330, UT2334, UT1392, UT2402) 520 521 (Wesgate et al., 2021). The patterns of our tektites are similar to Australasian tektites (Schwarcz et al., 2016; Wesgate et al., 2021), but distinct to the Belize and the Ivory coast tektites with 522 their lower concentrations and different impact events (Schwarcz et al., 2016). 523

524

Figure 10. ⁴⁰Ar/³⁹Ar age spectra of tektites. Plateau age given at the 2σ level. B1, B2, B3:
tektites from Xioamei site; F1, F2: tektites from Fengshudao site; RT1: tektite from Roc Tung
1; GD: tektite from Go Da.

528

- 529 Appendix A. Supplementary information.
- 530 MC2. Ar isotopic data corrected for blank.
- 531 MC3. 39 Ar/ 40 Ar vs. 36 Ar/ 40 Ar inverse isochron plots.

532 MC4. Tables S2 and S3

533 **References**

Albin, E. F., Norman, M. D., Rodan, M., 2000. Major and trace element compositions of
georgiaites: clues to the source of North American tektites. Meteoritic Planetary Science 35,
795–806.

- 537
- Amare, K., Koeberl, C., 2006. Variation of chemical composition in Australasian tektites from
 different localities in Vietnam. Meteoritic & Planetary Science 41, 107–123.
- 540

Ao, H., Deng, C., Dekkers, M.J., Liu, Q., Qin, L., Xiao, G., Chang, H., 2010. Astronomical
dating of the Xiantai, Donggutuo and Maliang Paleolithic sites in the Nihewan Basin (North
China) and implications for early human evolution in East Asia. Palaeogeography,
Palaeoclimatology, Palaeoecology 297, 129-137.

- Ao, H., An, Z.S., Dekkers, M.J., Wei, Q., Pei, S.W., Zhao, H., Zhao, H.L., Xiao, G.Q., Qiang,
- 546 X.K., Wu, D.C., Chang, H., 2012. High-resolution record of geomagnetic excursions in the
- 547 Matuyama chron constrains the ages of the Feiliang and Lanpo Paleolithic sites in the Nihewan
- 548 Basin, North China. Geochemistry, Geophysics, Geosystems 13, Q08017.
- Ao, H., An, Z., Zhisheng, A., Dekkers, M.J., Li, Y., Xiao, G., Zhao, H., Qiang, X., 2013.
 Pleistocene magnetochronology of the fauna and Paleolithic sites in the Nihewan Basin:
 Significance for environmental and hominin evolution in North China. Quaternary
 Geochronology 18, 78-92.
- 553
- Antoine P., Moncel M-H., Locht J-L. Bahain J-J., Voinchet P., Herisson D., Hurel A., 2019.
 The earliest record of Acheulean human occupation in North-West Europe. Nature Scientific
- 556 Reports 9, 13091.
- 557 https://doi.org/10.1038/s41598-019-49400-w
- 558
- Bae, C.J., 2024. The Paleoanthropology of Eastern Asia, University of Hawai'i Press,
 Honolulu.
- 561
- Bae, C.J., Li, F., Cheng, L., Wang, W., Hong, H., 2018. Hominin distribution and density
 patterns in Pleistocene China: Climatic influences. Palaeogeography, Palaeoclimatology,
 Palaeoecology 512, 118-131.
- 565
- Beyene, Y., Katoh, S., WoldeGabriel, G., Hart, W.K., Uto, K., Sudo, M., Kondo, M., Hyodo,
 M., Renne, P.R., Suwa, G., Asfaw, B., 2013. The characteristics and chronology of the earliest
 Acheulean at Konso, Ethiopia. PNAS 110, 1584-1591.
- 569
- Brumm, A., van den Bergh, G., Storey, M., Kurniawan, I., Alloway, B.V., Setiawan, R.,
 Setiyabudi, E., Grün, R., Moore, M.W., Yurnaldi, D., Puspaningrum, M.R., Wibowo, U.P.,
 Insani, H., Sutisna, I., Westgate, J.A., Pearce, N.J.G., Duval, M.,Meijer, H.J.M., Aziz, F.,
 Sutikna, T., van der Kaars, S., Morwood, M.J., 2016. Age and context of the oldest known
 hominin fossils from Flores. Nature 534, 249–253.
- 575

- 576 Carbonell, E., Sala Ramos, R., Rodriguez, X.P., Mosquera, M., Ollé, A., Vergès J.M., Martinez-
- 577 Navarro, B., Bermudez de Castro, J.M., 2010. Early hominid dispersals: A technological
- 578 hypothesis for « out of Africa". Quaternary International 223-224, 36-44.
- 579
 580 Carbonell, E., Barsky, D., Sala, R., Celiberti, V., 2016. Structural continuity and technological
 581 change in Lower Pleistocene toolkits. Quaternary International 393, 6-18.
- 582
- Chapman, D.R., Scheiber, L.C., 1969. Chemical Investigation of Australasian Tektites. Journal
 of Geophysical Research 74, 6737-6776.
- 585
- Deng, C.L., Wei, Q., Zhu, R.X., Wang, H.Q., Zhang, R., Ao, H., Chang, L., Pan, Y.X., 2006.
 Magnetostratigraphic age of the Xiantai Paleolithic site in the Nihewan Basin and implications
 for early human colonization of Northeast Asia. Earth Planet. Sci. Lett. 244, 336–348.
- 590 Derevianko, A.P., 2008. Paleoenvironment. The Stone Age. The Bifacial technique on China.
 591 Archaeology, Ethnology & Anthropology of Eurasia 33/1, 2-32.
- 592 Derevianko, A.P., Kandyda, A.V., Khac Su, N., Gladyshev, S.A., Gia Doi, N., Lebedev, V.A.,
- 593 Chekha, A.M., Rybalko, A.G., Kharevich, V.M., Tsybankov, A.A., 2018. The Discovery of a
- Bifacial Industry in Vietnam. Archaeology, Ethnology and Anthropology of Eurasia 46/3, 3-21.
- 595
- Echassoux, A., Moigne, A.-M., Moullé, P.-E., Li, T., Feng, X.B., Li, W., WU, Z.,
 Fauquembergue, E., Magnaldi, B., 2008. Les faunes de grands mammifères du site de l'Homme
 de Yunxian. In : de Lumley H., Li T. (Eds.), Le site de l'Homme de Yunxian. CNRS éditions,
 Paris, pp. 253-364.
- 600
- Feng, X., 2008. Stratégie de débitage et mode de façonnage des industries du Paléolithique
 inférieur en Chine et en Europe entre 1 Ma et 400 000 ans. Ressemblances et différences de la
 culture de l'homme de Yunxian et Acheuléen européen. L'Anthropologie 112, 423-447.
- 604
- Feng, X., Qi, Y., Li, Q., Ma, X., Liu, K., 2018. The industry of the Lower Paleolithic site of
 Baigu, Bose Basin, Guangxi Zhuang autonome Province, PR China. L'Anthropologie 122, 14–
 32.
- 608
 609 Gallotti, R., & Mussi, M., 2017. Two Acheuleans, two humankinds. From 1.5 to 0.85 Ma at
 610 Melka Kunture (Upper Awash, Ethiopian highlands). Journal of Anthropological Sciences 95,
 611 137-181.
- 612
- Garcia, T., Féraud, G., Falguères, C., de Lumley, H., Perrenoud, C., Lordkipanidze, D., 2010.
 Earliest human remains in Eurasia: New ⁴⁰Ar/³⁹Ar dating of the Dmanisi hominid-bearing
 levels, Georgia. Quaternary Geochronology 5, 443-451.
- 616
- Glass, B.P., Koeberl, C., 2006. Australasian microtektites and associated impact ejecta in the
 South China Sea and the Middle Pleistocene superuption of Toba. Meteoritics & Planetary
 Science 41, 305-326.
- 620
- 621 Goren-Inbar, N. Alperson-Alif, N., Sharon, G., Herzlinger, G., 2018. The Acheulian site of
- 622 Gesher Benot Ya 'aqov. Volume IV. The lithic Assemblages. Springer, Cham.
- 623

Grün, R., Aubert, M., Hellstrom, J., Duval, M., 2010. The challenge of direct dating old human 624 fossils. Quaternary International 223-224, 87-93. 625 626 Guan, Y., Wang, F-G, Xie, F., Pei, S-W, Zhou, Z-Y, Gao, X., 2016. Flint knapping strategies 627 at Cenjiawan, an Early Paleolithic site in the Nihewan Basin, North China. Quaternary 628 629 International 400, 86-92. 630 Guo, S.-L., Huang, W., Hao, X.-H., Chen, B.L., 1997. Fission Track Dating of Ancient Man 631 Site in Baise, China, and its signicances in space research, Paleomagnetism and Stratigraphy. 632 Radiation Measurements 28, 565-570. 633 634 Han, F., Bahain, J.-J., Deng, C., Boëda, E., Hou, Y., Wei, G., Huang, W., Garcia, T., Shao, Q., 635 636 He, C., Falguères, C., Voinchet, P., Yin, G., 2017. The earliest evidence of hominid settlement in China: Combined electron spin resonance and uranium series (ESR/U-series) dating of 637 mammalian fossil teeth from Longgupo cave. Quaternary International 434, 75-83. 638 639 Han, F., Shao, Q., Bahain, J.-J., Sun, X., Yin, G., 2018. Coupled ESR and U-series dating of 640 Middle Pleistocene hominin site Bailongdong cave, China, Quaternary Geochronology 49, 291-641 642 296. 643 644 645 Harmand, S., Lewis, J.E., Feibel, C.S., Lepre, C.J., Prat, S., Lenoble, A., Boës, X., Quinn, R., Brenet, M., Arroyo, A., Taylor, N., Clément, S., Daver, G., Brugal, J.-P., Leakey, L. Mortlocks, 646 R.A., Wright, J.D., Lokorodi, S., Kirwa, C., Kent, D.V., Roche, H., 2015. 3.3-million-year-old 647 648 stone tools from Lomekwi 3, West Turkana, Kenya. Nature 521, 310-315. 649 650 Heide, K., Heide, G., Kloess, G., 2001. Glass chemistry of tektites. Planet. Space Sci. 49, 839-844. 651 652 Hellstrom, J., Pickering, R., 2015. Recent advances and future prospects of the U-Th and U-Pb 653 chronometers applicable to archaeology. Journal of Archaeological Science 56, 32-40. 654 655 Hou, Y., Potts, R., Baoyin, Y., Zhengtang, G., Deino, A., Wie, W., Clark, J., Guangmao, X., 656 Weiwen, H., 2000. Mid-Pleistocene Acheulean-like Stone Technology of the Bose Basin, South 657 China. Science 287, 1622–1626. 658 659 Hou, Y.M., Zhao, L.X., 2010. An archaeological view for the presence of early humans in 660 China. Quaternary International 223-224, 10-19. 661 662 Huang, S., Wang, W., Bae, C. J., Xu, G., Liu, K., 2012. Recent Paleolithic field investigations 663 664 in Bose Basin (Guangxi, China). Quaternary International 281, 5-9. 665 Hyodo, M., Matsu'ura, S., Kamishima, Y., Kondo, M., Takeshita, Y., Kitaba, I., Danhara, T., 666 667 Aziz, F., Kurniawan, I., Kumai, H., 2011. High-resolution record of the Matuyama-Brunhes transition constrains the age of Javanese Homo erectus in the Sangiran dome, Indonesia. PNAS 668 108, 19563-19568. 669 670 671 Ingicco, T., van den Bergh, G.D., Jago-on, C., Bahain, J.-J., Chacon, M.G., Amano, N., Forestier, H., King, C., Manalo, K., Nomade, S., Pereira, A., Reyes, M.C., Sémah, A.-M., Shao, 672 673 Q., Voinchet, P., Falguères, C., Albers, P.C.H., Lising, M., Lyras, G., Yurnaldi, D., Rochette,

- P., Baustita, A., de Vos, J., 2018. Earliest known hominin activity in the Philippines by 709
 thousand years ago. Nature 557, 233-237.
- 676
- Jourdan, F., Renne, P.R., 2007. Age calibration of the Fish Canyon sanidine ⁴⁰Ar/³⁹Ar datin g
 standard using primary K-Ar standards. Geochimica et Cosmochimica Acta 71, 387-402.
- 679
- Jourdan, F., Mark, D.F., Vérati, C. (eds), 2014. Advances in ⁴⁰Ar/³⁹Ar dating: from
 archaeological to planetary sciences-introduction. Geological Society, London, Special
 Publications 378, 1–8.
- 683
- Jourdan, F., Nomade, S., Wingate, M.T.D., Eroglu, E., Deino, A., 2019. Ultraprecise age and
 formation temperature of the Australasian tektites constrained by ⁴⁰Ar/³⁹Ar analyses.
 Meteoritics & Planetary Science 54, 2373-2591.
- Kandyba, A.V., Chekha, A.M., Doi, N.G., Su, N. K., Gladyshev, S.A., Derevianko, A.P., 2021.
 The early Paleolithic Go Da site and the Bifacial lithic industries of Southeast Asia.
 Archaeology, Ethnology & Anthropology of Eurasia 49, 3-14.
- Keates, S.G., 2000. Tektites and the age paradox in Mid-Pleistocene China. Science 289, 507a.
- Keates, S.G., 2010. Evidence for the earliest Pleistocene hominid activity in the Nihewan Basinof northern China. Quaternary International 223-224, 408-417.
- Koeberl, C., 1990. The geochemistry of tektites: An overview. Tectonophysics 171, 405–422.
- Koeberl C., Bottomley R., Glass B.P., Storzer D., 1997. Geochemistry and age of Ivory Coast
 tektites and microtektites. Geochimica et Cosmochimica Acta 61, 1745-1772.
- Koeberl, C., Glass, B.P., 2000. Tektites and the age paradox in Mid-Pleistocene China. Science289, 507a.
- Kong, Y., Deng, C., Liu, W., Pei, S., Sun, L., Ge, J., Yi, L., Zhu, R., 2018. Magnetostratigraphic
 dating of the hominin occupation of Bailong Cave, central China. Scientific Reports 8, 9699.
- 707
 708 Koppers, A.A.P., 2002. ArArCALC-software for ⁴⁰Ar/³⁹Ar age calculations. Computers &
 709 Geosciences 28, 605–619.
- 710
- Langbroek, M., 2015. Do tektites really date the bifaces from the Bose (Baise) Basin, Guangxi,
 southern China?. Journal of Human Evolution 80, 175-178.
- 713
- Larick, R., Ciochon, R.L., 2015. Early hominin Biogeography in Island Southeast Asia.
 Evolutionary Anthropology 24, 185-213.
- 716
- Lebatard, A.-E., Alçiçek, M.C., Rochette, P., Khatib, S., Vialet, A., Bouldes, N., Bourlès, D.J.,
- Demory, F., Guipert, G., Mayda, S., Titov, V.V., Vidal, L., de Lumley, H., 2014. Dating the *Homo erectus* bearing travertine from Kocabaş (Denizli, Turkey) at at least 1.1 Ma. Earth and
 Planetary Science Letters 390, 8-18.
- 721
- Lepre, C.J., Roche, H., Kent, D.V., Harmand, S., Quinn, R.L., Brugal, J.-P., Texier, P.-J., Lenoble, A., Feigel, C.S., 2011. An earlier origin for the Acheulian. Nature 477, 81-85

- Li, H., Li, C., Kuman, K., 2014. Rethinking the "Acheulean" in East Asia: Evidence from recent
 investigations in the Danjiangkou Reservoir Region, central China. Quaternary International
 347, 163-175.
- 727

Li, H., Li, C., Kuman, K., 2017. Longgudong, an Early Pleistocene site in Jianshi, South China,
with stratigraphic association of human teeth and lithics. Science China Earth Sciences 60, 452462.

- 731
- Li, H., Kuman, K., Li, C., 2018. What is currently (un)known about the Chinese Acheulean,
 with implications for hypotheses on the earlier dispersal of hominids. C.R. Palevol 17, 120130.
- 735

743

Li, X., Pei, S., Jia, Z., Guan, Y., Niu, D., Ao, H., 2016. Paleoenvironmental conditions at
Madigou (MDG), a newly discovered Early Paleolithic site in the Nihewan Basin, North Chian.
Quaternary International 400, 100-110.

- Li, X., Ao, H., Dekkers, M.J., Roberts, A.P., Zhang, P., Lin, S., Huang, W., Hou, Y., Zhang,
 W., An, Z., 2017. Early Pleistocene occurrence of Acheulian technology in North China.
 Quaternary Science Reviews 156, 12-22.
- ⁷⁴⁴ Lisiecki, L.E, Raymo, M.E., 2005. A Pleicene-Pleistocene stack of 57 globally distributed ⁷⁴⁵ benthic δ^{18} O records. Paleoceanography and Paleoclimatology 20, PA1003. ⁷⁴⁶
- Liu, Y., Hu, Y., Wei, Q., 2013. Early to Late Pleistocene human settlements and the evolution
 of lithic technology in the Nihewan Basin, North China: A macroscopic perspective.
 Quaternary International 295, 204-214.
- Liu, C.-R., Yin, G.-M., Deng, C.-L., Han, F., Song, W.-J., 2014. ESR dating of the Majuangou and Banshan Paleolithic sites in the Nihewan Basin, North China. Journal of Human Evolution 73, 58-63.
- Liu, Y., Hu, Y., Wei, Q., 2013. Early to Late Pleistocene human settlements and the evolution
 of lithic technology in the Nihewan Basin, North China: A macroscopic perspective.
 Quaternary International 295, 204-214.
- 758
- Liu, X., Shen, G., Tu, H., Lu, C., Granger, D.E., 2015. Initial ²⁶Al/¹⁰Be burial dating of the
 hominin site Bailong Cave in Hubei Province, central China. Quaternary International 389, 235240.
- Lombera-Hermida de, A., Bargallo, A., Terradillos-Bernal, M., Huguet, R., Vallverdu, J.,
 Garcia-Anton, M.-D., Mosquera, M., Ollé, A., Sala, R., Carbonell, E., Rodriguez-Alvarez, X.P., 2015. The lithic industry of Sima del Elefante (Atapuerca, Burgos, Spain) in the context of
 Early and Middle Pleistocene technology in Europe. Journal of Human Evolution 82, 95-106.
- 767
- Lumley de, H., Li, T., (Eds.), 2008. Le site de l'homme de Yunxian : Quyuanhekou, Quingqu,
 Yunxian, Province de Hubei. CNRS Editions, Paris.
- Lumley de, H., Cauche, D., Pollet, G., Notter, O., Rossini-Notter, E., Xie, G., Lin, Q, Feng, X.,
 Ghen, X., Qi, Y., Wei, J., Wang, W., Huang, Q., 2020. Les industries lithiques du Paléolithique

- ancien du bassin de Bose. Province autonome du Guangxi Zhuang, Chine du Sud. CNRS
 Editions, Paris, 325 p. ISBN: 978-2-271-13406-6
- 775
- Matinez-Navarro, B., Belmaker, M., Bar-Yosef, O., 2012. The Bovid assemblage (Bovidae,
 Mammalia) from the Early Pleistocene site of 'Ubeidiya, Israel: Biochronological and
 paleoecological implications for the fossil and lithic bearing strata. Quaternary International
 267, 78-97.
- 780
- McDonough, W. F., Sun, S.-S., 1995. The composition of the Earth. Chemical Geology 120,
 223-253.
- 783
- Meisel, T., Lange, J.-M., Krähenbühl, U., 1997. The chemical variation of moldavite tektites:
 simple mixing of terrestrial sediments. Meteoritics Planetary Science 32, 493–502.
- 786
- Michel, V., Feng, X., Shen, G., Cauche, D., Moncel, M.-H., Gallet, S., Gratuze, B., Wei, J.,
 Ma, X., Liu, K., 2021. First ⁴⁰Ar/³⁹Ar analyses of Australasian tektites in close association with
 bifacially worked artifacts at Nalai site in Bose Basin, South China: The question of the early
 Chinese Acheulean. Journal of Human Evolution, 153, 102953.
- 791
- Moncel, M-H., Santagata, C., Pereira, A., Nomade, S., Voinchet, P., Bahain, J-J., Daujeard, C.,
 Curci, A. Lemorini, C., Hardy, B., Eramo, G., Berto, C., Raynal, J-P., Arzarello, M., Mecozzi,
 B., Iannucci, A., Sardella, R., Allegretta, I., Delluniversità, E., Terzano, R., Dugas, P., Jouanic,
 G., Queffelec, A., d'Andrea, A., Valentini, R., Minucci, E., Carpentiero, L., Piperno, M.,
 2020a. The origin of early Acheulean expansion in Europe 700 ka ago: new findings at
 Notarchirico (Italy). Nature. Scientific Reports 10(1), 1-16.
- 798 https://doi.org/10.1038/s41598-020-68617-8 799
- Moncel M-H., Despriée J., Courcimaut G., Voinchet P., Bahain J-J., 2020b. La Noira site
 (Centre, France) and the technological behaviours and skills of the earliest Acheulean in
 Western Europe between 700 and 600 kyrs. Journal of Paleolithic Archeology 3 (3), 255-301.
 https://doi.org/10.1007/s41982-020-00049-2
- 804
 805 Movius, H., 1948. The Lower Palaeolithic Cultures of Southern and Eastern Asia. Transactions
 806 of the American Philosophical Society 38, 329-420.
- 807

Mussi, M., Skinner, M.M., Melis, R.T., Pneras, J., Rubio-Jara, S., Davies, T.W., Geraads, D.,
Bocherens, H., Briatico, G., Le Cabec, A., Hublin, J.-J., Gidna, A., Bonnefille, R., Di Bianco,

- L., Mendez-Quinta, E., 2023a. Early *Homo erectus* lived at high altitudes and produced both
 Oldowan and Acheulean tools. Science 382, 713-718.
- 812

Mussi, M., Altamura, F., Di Bianco, L., Bonnefille, R., Gaudzinski-Windheuser, S., Geraads,
D., Melis, R.T., Panera, J., Piarulli, F., Pioli, L., Ruta, G., Sanchez-Dehesa Galn, S., Mendez Quintas, E. 2023b. After the emergence of the Acheulean at Melka Kunture (Upper Awash,

- Ethiopia): From Gombore IB (1.6 Ma) to Gombore Iγ (1.4 Ma), Gombore iδ (1.3 Ma) and
- 817 Gombore II OAM test pit C (1.2 Ma). Quaternary International 657, 3-25.
- 818819 O'Reilly, T.C., Haskin, L.A., King, E.A., 1983. Element correlations among north American
- tektites. 14th Luna Planet. Sci., 580-581 (abstract).
- 821

- Pappu, S., Gunnell, Y., Akhilesh, K., Braucher, R., Taieb, M., Demory, F., Thouveny, N., 2011.
- Early Pleistocene Presence of Acheulian Hominins in South India. Science 331, 1596-1599.
- 824
- Pei, S., Deng, C., de la Torre, I., Jia, Z., Ma, D., Li, X., Wang, X., 2019. Magnetostratigraphic
 and archaeological records at the Early Pleistocene site complex of Madigou (Nihewan Basin):
 Implications for human adaptations in North China. Palaeogeography, Palaeoclimatology,
 Palaeoecology 530, 176-189.
- Pike, A.W.G., Pettitt, P.B., 2003. U-series Dating and Human Evolution. In Uranium Series
 Geochemistry (edited by B. Bourdon, G.M. Henderson, C.C. Lundstrom, et al.). Reviews in
 Mineralogy and Geochemistry 52, 607-630.
- 833
- Potts, R., Huang, W., Hou, Y., Deino, A., Baoyin, Y., Zhengtang, G., Clark, J., 2000. Tektites
 and the age paradox in Mid-Pleistocene China. Response. Science 289, 507a.
- Qin, Z., Sun, X., 2023. Glacial-Interglacial Cycles and Early Human Evolution in China. Land
 12, 1683.
- 839
- 840 Renne, P.R., Balco, G., Ludwig, K.R., Mundil, R., Min, K., 2011. Response to the comment by
- W.H. Schwarz et al. on "Joint determination of K-40 decay constants and Ar-40*/K-40 for the
 Fish Canyon sanidine standard, and improved accuracy for Ar-40/Ar-39 geochronology" by PR
- 843 Renne et al. (2010). Geochimica et Cosmochimica Acta 75, 5097-5100.
 - 844
 - Rink, W.J., Bartoll, J., Schwarcz, H.P., Shane, P., Bar-Osef, O., 2007. Testing the reliability of
 ESR dating of optically exposed buried quartz sediments. Radiation Measurements 42, 16181626.
 - 848
 - Rochette, P., Beck, P., Bizzarro, M., Braucher, R., Cornec, J., Debaille, V., Devouard, B.,
 Gattacceca, J., Jourdan, F., Moustard, F., Moynier, F., Nomade, S., Reynard, B., 2021. Impact
 glasses from Belize represent tektites from the Pleistocene Pantasma impact crater in
 Nicaragua. Communications earth & environment 2, 94.
 - Sagi, A., 2005. Magnetostratigraphy of 'Ubeidiya Formation, Northern Dead Sea Transform,
 Israel. M.S. Dissertation, The Hebrew University.
 - 856
 - Scaillet, S., Vita-Scaillet, G., Guillou, H., 2008. Oldest human footprints dated by Ar/Ar. Earth
 and Planetary Science Letters 275, 320–325.
 - 859
 - Schwarz, W.H., Trieloff, M., Bollinger, K., Gantert, N., Fernandes, V.A., Meyer, H.-P.,
 Povenmire, H., Jessberger, E.K., Guglielmino, M., Koeberl, C., 2016. Coeval ages of
 Australasian, Central American and Western Canadian tektites reveal multiple impacts 790 ka
 ago. Geochimica et Cosmochimica Acta 178, 307-319.
 - 864
 - Semaw, S., Rogers, M.J., Quade, J., Renne, P.R., Butler, R.F., Dominguez-Rodrigo, M., Stout,
 D., Hart, W.S., Pickering, T., Simpson, S.W., 2003. 2.6-Million-year-old stone tools and
 associated bones from OGS-6 and OGS-7, Gona, Afar, Ethiopia. Journal of Human Evolution
 45, 169-177.
- 869

- Shen, C., Zhang, X., Gao, X., 2016. Zhoukoudian in transition: Research history, lithic
 technologies, and transformation of Chinese Palaeolithic archaeology. Quaternary International
 400, 4-13.
- 873
- Shen, G.J., Gao, X., Gao, B., Granger, D.E., 2009. Age of Zhoukoudian *Homo erectus* determined with ²⁶Al/¹⁰Be burial dating. Nature 458, 198-200.
- 876
- 877 Shen, G.J., Michel, V., Despriée, J., Han, F., Granger, D., 2012. Datation d'enfouissement par
- 26 Al/¹⁰Be et son application préliminaire à des sites du Paléolithique inférieur en Chine et en
- 879 France. L'Anthropologie 116, 1-11.880
- Shen, G., Wang, Y., Tu, Hua, Tong, H., Wu, Z., Kuman, K., Fink, D., Granger, D.E., 2020.
 Isochron ²⁶Al/¹⁰Be burial dating of Xiboudu: Evidence for the earliest human settlement in northern China. L'Anthropologie 124, 102790.
- Sieh, K., Herrin, J., Jicha, B., Angel, D.S., Moore, J.D.P., Banerjee, P., Wiwegwin, W.,
 Sihavong, V., Singer, B., Chualaowanich, T., Charusiri, P., 2020. Australasian impact crater
 buried under the Bolaven volcanic field, Southern Laos. PNAS 117, 1346-1353.
- Son, T.H., Koeberl, C., 2005. Chemical variation within fragments of Australasian tektites.
 Meteoritics & Planetary Science 40, 805-815.
- 891

884

888

- Sun, X., Lu, H., Wang, S., Yi, L., Li, Y., Bahain, J.-J., Vionchet, P., Hu, X., Zeng L., Zhang,
 W., Zhuo, H., 2017. Early human settlements in the southern Qinling Mountains, central China,
 Quaternary Science Reviews 164, 168-186.
- 895
- Taylor, S.R., 1962. The chemical composition of australites. Geochimica et CosmochimicaActa 26, 685-722.
- 898
- Tissoux, H., Bahain, J.-J., de Lumley, H., Li, T., Feng, X., Li, W., 2008. Essai de datation par les méthodes combinées (RPE/U-Th) de dents d'herbivores et par RPE de quartz blanchis extraits des sédiments fluviatiles provenant du site de l'Homme de Yunxian. In : de Lumley H.,
 Li T. (Eds.), Le site de l'Homme de Yunxian. CNRS éditions, Paris, pp. 237-252.
- 903
- Toro-Moyano, I., Martinez-Navarro, B., Augusti, J., Souday, C., Bermudez de Castro, J.M.,
 Martinon-Torres, M., Fajardo, B., Duval, M., Falguères, C., Oms, O., Pares, J.M., Andon, P.,
 Julia, R., Garcia-Aguilar, J.M., Moigne, A.-M., Espigares, M.P., Ros-Montoya, S., Palmqvist,
 P., 2013. The oldest human fossil in Europe, from Orce (Spain). Journal of Human Evolution
 65, 1-9.
- 909
- Tribolo, C., Mercier, N., Douville, E., Joron, J.-L., Reyss, J.-L., Rufer, D., Cantin, N., Lefrais,
 Y., Miller, C.E., Porraz, G., Parlington, J., Rigaud, J.-P., Texier, P.-J., 2013. OSL and TL dating
 of the Middle Stone Age sequence at Diepkloof Rock Shelter (South Africa): A clarification.
 Journal of Archaeological Science 40, 3401-3411.
- 914
- Tu, H., Shen, G., Granger, D., Yang, X., Lai, Z., 2017. Isochron ²⁶Al/¹⁰Be burial dating of the
 Lantian hominin site at Gongwangling in Northwestern China. Quaternary Geochronology 41,
 174-179.
- 918

Vallverdu, J., Saladié, P., S., Rosas, A., Huguet R., Caceres, I., Mosquera, M., Garcia-Taberno,
A., Estalrrich, A., Lozana-Fernandez, I., Pineda-Alcala, A., Carrancho, A., Villalain, J.J.,
Bourlès, D., Braucher R., Lebatard, E., Vilalta, J., Esteban-Nadal, M., Bennasar M.L., Bastir,
M., Lopez-Polin L., Ollé, A., Vergés, J.M., Ros-Montoya, S., Martinez-Navarro, B., Garcia,
A., Martinell, J., Exposito, I., Burjachs, F., Agusti, J., Carbonell, E., 2014. Age and Date for
Early Arrival of the Acheulian in Europe (Barranc de la Boella, la Canonja, Spain). PLoS One
9, e103634.

926

Voinchet, P., Moreno, D., Bahain, J-J., Tissoux, H., Tombret, O., Falguères, Moncel, M-H. C.,
Schreve, D., Candy, I., Antoine, P., Ashton, N., Beamish, M., Cliquet, D., Despriée, J., Lewis,
S., Limondin-Lozouet, N., Locht, J-L., Parfitt, S., Pope, M., 2015. New chronological data
(ESR and ESR/U-series) for the earliest Acheulean sites of north-western Europe. J. Quat, Sci.
30, 610–622.

932

Wang, H., Deng, C., Zhu, R., Wei, Q., Hou, Y., Boëda, E., 2005. Magnetostratigraphic dating
of the Donggutuo and Maliang Paleolithic sites in the Nihewan Basin, North China. Quaternary
Research 64, 1-11.

936

Wang, W., Bae, C.J., Huang, S., Huang, X., Tian, F., Mo, J., Huang, Z., Huang, C., Xie, S., Li,
D., 2014. Middle Pleistocene bifaces from Fengshudao (Bose Basin, Guangxi, China). Journal
of Human Evolution 69, 110-122.

940

Wang, W., Bae, C.J., 2015. How old are the Bose (Baise) Basin (Guangxi, southern China)
bifaces? The Australasian tektites question revisited. Journal of Human Evolution 80, 171-174.

Wang, W., Fan, Y., Bae, C., 2024. The Bose (South China) Handaxes: Early Palaeolithic
Bifaces and Early Hominin Cognition. In: The Oxford Handbook of Cognitive Archaeology
(eds. K.A. Overmann, F.L. Coolidge), chap. 31, p. 741-770.

947

954

Wei, G., Huang, W., Chen, S., He, C., Pang, L., Wu, Y., 2014. Paleolithic culture of Longgupo
and its creators. Quaternary International 354, 154-161.

Wesgate, J.A., Pillans, B.J., Alloway, B.V., Pearce, N.J.G., Simmonds P., 2021. New fissiontrack ages of Australasian tektites define two age groups: discriminating between formation and
reset ages. Quaternary Geochronology 66, 101113.

Xie, G.-M., Bodin, E., 2007. Les industries paléolithiques du Bassin de Bose (Chine du Sud).
L'Anthropologie 111, 182–206.

- Xie, G., Chen, X., Li, D., Yu, M., Hu, Z., Lu, H., Huang, Q., Lin, Q., 2021. Stratigraphy and
 chronology of the palaeolithic industry in Bose Basin, South China: Excavation of Gaolingpo.
 Archaeological Research in Asia 26, 100284.
- 961

Yang, S.-X., Hou, Y.-M., Yue, J.-P., Petraglia, M.D., Deng, C.-L., Zhu, R.-X., 2016. The Lithic
Assemblages of Xiaochangliang, Nihewan Basin: Implications for Early Pleistocene Hominin
Behaviour in North China. PLoS ONE 11(5), e0155793.


965


Yang, S.-X., Petraglia, M.D., Hou, Y.-M., Yue, J.-P., Deng, C.-L., Zhu, R.-X., 2017. The lithic
assemblages of Donggutuo, Nihewan basin: Knapping skills of Early Pleistocene hominins in
North China. PLoS ONE 12, e0185101.


- 969
- Yang, S.-X., Deng, C.-L., Zhu, R.-X., Petraglia, M.D., 2020. The Paleolithic in the Nihewan
 Basin, China: Evolutionary history of an Early to Late Pleistocene record in Eastern Asia.
 Evolutionary Anthropology 29, 125-142.
- 973
- Zhang, P., Huang, W., Wang, W., 2010. Acheulean handaxes from Fengshudao, Bose sites of
 South China. Quaternary International 223–224, 440–443.
- 976
- 977 Zhu, R.X., Hoffman, K.A., Potts, R., Deng, C.L., Pan, Y.X., Guo, B., Shi, C.D., Guo, Z.T.,
- Yuan, B.Y., Hou, Y.M., Huang, W.W., 2001. Earliest presence of humans in northeast Asia.
 Nature 413, 413–417.
- 979 980


983

- Zhu, R., An, Z., Potts, R., Hoffman, K.A., 2003. Magnetostratigraphic dating of early humans
 in China. Earth-Science Reviews 61, 341-359.
- Zhu, R.X., Potts, R., Xie, F., Hoffmn, K.A., Deng, C.L., Shi, C.D., Pan, Y.X., Wang, H.Q., Shi,
 R.P., Wang, Y.C., Shi, G.H., Wu, N.Q., 2004. New evidence on the earliest human presence at
 high northern latitudes in northeast Asia. Nature 431, 559-562.
- 287
 288 Zhu, Z., Dennell, R., Huang, W., Wu, Y., Qiu, S., Yang, S., Rao, Z., Hou, Y., Xie, J., Han, J.,
 2018. Hominin occupation of the Chinese Loess Plateau since about 2.1 million
 2019. years ago. Nature 559, 608-612.
- 991 992
- 993 Figure 1









F3

B3

90

Ivorite

Moldavite

