



HAL
open science

Effects of Vibrance Gold® on soil parameters during wheat development

Séverine Piutti, Hugues Clivot, Virginie Braun, Olivier Borde, S Slezack

► **To cite this version:**

Séverine Piutti, Hugues Clivot, Virginie Braun, Olivier Borde, S Slezack. Effects of Vibrance Gold® on soil parameters during wheat development. European Root Health Forum, Mar 2015, Berlin, Germany. hal-04796689

HAL Id: hal-04796689

<https://hal.science/hal-04796689v1>

Submitted on 21 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effects of Vibrance Gold® on soil parameters during wheat development

European
Root Health
Forum
2015

S. Piutti^{a,b}, H. Clivot^{a,b}, V. Braun^c, O. Borde^c, S. Slezack-Deschaumes^{a,b}

^a Université de Lorraine, Laboratoire Agronomie et Environnement, UMR 1121, TSA 40602, 54518 Vandœuvre cedex, France

^b INRA, Laboratoire Agronomie et Environnement, UMR 1121, TSA 40602, 54518 Vandœuvre cedex, France

^c Syngenta France SAS, 1 avenue des Prés, 78286 Guyancourt, France

Introduction :


Nitrogen is a limiting nutrient for vegetative growth supporting leaf area development and photosynthesis. Vibrance Gold® applied as seed treatment has a beneficial effect on wheat growth (Slezack-Deschaumes et al.). One hypothesis is that Vibrance Gold®, by

modulating root architecture traits related to nutrient acquisition, could improve root N uptake. A second hypothesis is that seed treatment acts as a biostimulant that induced some soil microbiological functions such as organic matter decomposition and then, improve the bioavailability of nutrients for plant uptake.

Material and methods :

At each growth stages (3-4 tillers, ear 1cm and 1-2 nodes), rhizosphere soils were sampled for Redigo (100 g l⁻¹ prothioconazol) and Vibrance Gold® (50g l⁻¹ sedaxane) treatments. The following soil parameters were measured :

- Hot water extractable carbon (HWC) and nitrogen (HWN)
- Microbial biomass carbon (MBC) and nitrogen (MBN)
- Enzyme activities linked to biogeochemical cycles (carbon, nitrogen, phosphorus and sulfur)



Results and discussion :

HWC was significantly reduced (-74%) at the 1-2 nodes stage in Vibrance Gold® treatment compared to Redigo treatment (Figure 1a). This soil C pool constitutes a source of C and energy for soil heterotrophic microbial biomass. As this C pool is linked to rhizodeposition, the largest decrease of HWC in the Vibrance Gold® treatment could result from a greater reduction in the rhizodeposition compared to Redigo treatment. No effect of the seed treatment was evidenced at any development stage on MBC (data not shown).





Figure 1 : Evolution of soil extractable carbon (HWC, figure a) and soils extractable nitrogen (HWN, figure b) content (mean value \pm SE, n=4) in Redigo and Vibrance Gold® treatments. (Significant levels : * p<0.05; ** p<0.02; *** p<0.001)

This greater N immobilisation in microbial biomass in the Vibrance Gold® treatment explained the significant decrease of the C:N ratio of microbial biomass at the last development stage (Figure 3). No positive effect of Vibrance Gold® treatment was evidenced on the microbial enzyme activities related with C, N, P and S cycles.

Conclusion :

Soil parameters evolved during wheat development. The strong decrease of soil carbon and nitrogen contents in




Figure 2 : Evolution of microbial biomass nitrogen (mean value \pm SE, n=4) in Redigo and Vibrance Gold® treatments (Significant levels : * p<0.05; ** p<0.02; *** p<0.001)

HWN (soluble organic and inorganic N forms) decreased significantly in the Vibrance Gold® treatment during stem elongation (Figure 1b), suggesting a greater uptake of soil mineral N by plants and/or a larger microbial N immobilisation (Figure 2).




Figure 3 : Evolution of the C:N ratio of microbial biomass (mean value \pm SE, n=4) in Redigo and Vibrance Gold® treatments (Significant levels : * p<0.05; ** p<0.02; *** p<0.001)

Vibrance Gold® treatment suggest that : (i) the root soluble C content is reduced due to changes in plant C allocation, (ii) the acquisition of soil N by the plant is more efficient due to changes in the root morphology.