



HAL
open science

Vocal communication: The enigmatic production of low-frequency purrs in cats

David Reby, Andrey Anikin

► To cite this version:

David Reby, Andrey Anikin. Vocal communication: The enigmatic production of low-frequency purrs in cats. *Current Biology*, 2023, 33 (23), pp.R1236-R1237. <10.1016/j.cub.2023.10.033>. <hal-04798364>

HAL Id: hal-04798364

<https://hal.science/hal-04798364v1>

Submitted on 11 Mar 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

DISPATCH

Vocal communication: the enigmatic production of low-frequency purrs in cats

David Reby^{1,2,*} and Andrey Anikin^{1,3}

¹ENES Bioacoustics Research Laboratory, CRNL, CNRS, Inserm, University of Saint-Etienne, 42000 Saint-Etienne, France

²Institut universitaire de France, 75005 Paris, France.

³Division of Cognitive Science, Lund University, SE-221 00 Lund, Sweden

*Correspondence: dreby@mac.com (D.R.)

Cat purring, the unusual, pulsed vibration that epitomizes comfort, enjoys a special status in the world of vocal communication research. Indeed, it has long been flagged as a rare exception to the dominant theory of voice production in mammals. A new study presents histological and biomechanical evidence that purring can occur passively, without needing muscle vibration in the larynx controlled by an independent neural oscillator.

In mammals, including humans, vocal signals are produced by gestures that are mostly internal, which makes voice production a bit of a mystery and its investigation particularly difficult. Perhaps as a consequence, it was not until Ferrein's¹ experiments with cadavers published in 1741 that scientists began to understand the nature of voice production: phonation results from the vibration inside the larynx of vocal 'ribbons', under the effect of the airflow coming from the lungs. And it took two more centuries of investigation for scientists to arrive at the now-dominant *source-filter* theory of voice production^{2,3}, which states that vocal signals are generated through the combination of the vibration of vocal folds (which are more like lips than strings!) and subsequent filtering by the vocal tract (the throat, mouth, and nasal cavity). Crucially, the vibration of the vocal folds occurs as the air is forced out of the lungs (or, as we will see, also sometimes *into* the lungs), without nerve induced muscle contraction at these frequencies.

Yet, seeds of doubt sprouted again in the 1950s and 60s, when a new battle raged in the voice-science field about the forces that drive human phonation. Phonation involves lateral, oscillatory movements of the two vocal folds within the larynx⁴. As the two vocal folds repeatedly open and close the respiratory airway, this creates a series of regular air 'puffs', whose rate of repetition corresponds to the fundamental frequency or f_0 of the produced sound and determines its pitch. But why do the vocal folds oscillate? As we have seen, these vocal-fold movements were traditionally thought to be driven aerodynamically by the flow of air from the lungs (reminiscent of a flag flapping in the wind), but the new theory introduced by Husson⁵ argued that they were driven actively, by very rapid nerve impulses and muscle contractions that matched the rate of vocal-fold oscillations. Husson's theory was disproved for the human voice, since the f_0 s of human speech (about 100–300 oscillations per second in adults⁴), and especially of singing (over 1,000 oscillations per second in sopranos⁶), greatly exceed the possible rate of motor-nerve firing or muscle contraction, and because *ex-vivo* experiments have

confirmed that excised human larynges can still phonate normally when air is blown through them.

Thus, the 'myo-elastic aerodynamic theory' triumphed for the human voice⁷, and it has since been extended to most vertebrate vocalizations, whose f_0 ranges from 10 Hz in infrasonic elephant rumbles⁸ to well over 100 kHz in ultrasonic echolocation calls of bats⁹. Yet, one intriguing exception remained: cat purring. Because f_0 in purring is so low (20–30 Hz), despite the relatively small size of cats and their larynges, purrs were widely thought to be driven by active muscle contraction^{10,11}. Could this exception finally fall, too? In a recent paper published in *Current Biology*, Herbst *et al.*¹² show that the vocal folds in excised cat larynges can vibrate at the frequencies characteristic of purring when they are driven solely by air flow, and all nerve firing or muscle contraction is rendered impossible experimentally. Ex-vivo experiments using excised larynges work as follows: larynges from animals that died of natural causes are excised and mounted on an experimental set-up that allows the experimenters to control the air pressure below the glottis and to monitor its effect on the onset and periodicity of phonation¹³. In this case, the authors showed that the vocal folds in cat larynges can vibrate in the absence of muscle activity at frequencies ranging from 15–200 Hz, which encompasses the frequencies of purring. At these frequencies, the vocal folds behave in a way reminiscent of what happens in human vocal fry — a low, pulse-like vocal register, also known as strohbass¹⁴, in which the glottis remains closed for the majority of the oscillatory cycle.

The authors also conducted anatomical investigations that confirmed the presence of thick *pads* in the vocal folds. Such adaptations can have the effect of slowing down the rate of oscillation of vocal folds by increasing the effective mass in vibration¹⁵. Similar adaptations have previously been identified in studies of other felids, namely lions and tigers¹⁶, where they are also believed to support the efficient production of high-amplitude calls with low f_0 s.

To conclude, this study shows that the cat's vocal folds, despite their small size, are indeed capable of vibrating solely under the effect of airflow from the lungs, at a rate that corresponds to that observed in purrs, suggesting that active muscle vibration may not be necessary, after all. However, the authors acknowledge that the anatomical adaptation that enables passive vocal-fold vibration at low frequencies might also have evolved to enhance the effects of active muscular vibration. It is indeed much more energy efficient to make an oscillator vibrate at its natural frequency — a phenomenon known as resonance, which makes it possible to achieve very high amplitude oscillations with minimal energy¹⁷. Thus, Remmers and Gautier¹⁰ may still have been correct about cats generating each pulse of purring with an active muscular contraction, but at a very specific, anatomically optimal frequency close to the natural oscillatory frequency of the vocal folds.

Future work should investigate what happens during the inhalation phase as cat purrs are continuous vocalisations that take place on both inhalation and exhalation^{10,11}. In particular, it would be interesting to investigate why the transition between exhalation

and inhalation appears to have so little effect on the rate and amplitude of oscillations in the larynx (Figure 1). Finally, *in vivo* investigations should revisit the presence or absence of neural firing and muscle activity, and test for a possible coupling between the muscle- and airflow-induced oscillations documented here.

This paper has the merit of challenging a theory that may have been accepted too hastily, perhaps because we love an exception that confirms the rule. It calls for more investigations of this intriguing phenomena using the modern tools at our disposal. Science at its best!

Declaration of interests

The authors declare no competing interests.

References

1. Ferrein, M. (1741). De la formation de la voix de l'homme. *Hist. Acad. Roy. Sci.* 3, 409–432.
2. Chiba, T., and Kajiyama, M. (1958). The vowel, its nature and structure (Tokyo: Phonetic society of Japan).
3. Fant, G. (1960). Acoustic theory of speech production: with calculations based on X-ray studies of Russian articulations (The Hague: Mouton & Co. N.V., Publishers).
4. Titze, I.R. (2000). Principles of voice production. Second printing (Iowa City: National Center for Voice and Speech).
5. Husson, R. (1950). Etude des phénomènes physiologiques et acoustiques fondamentaux de la voix chantée. Thèse Fac. Sci., Paris (17 juin 1950). *Revue Scientifique*, Paris (1951).
6. Echternach, M., Burk, F., Köberlein, M., Döllinger, M., Burdumy, M., Richter, B., Titze, I., Elemans, C., and Herbst, C. (2023). Biomechanical sound production in high-pitched classical singing—the “Queen of the Night” does not whistle. *Research Square* <https://doi.org/10.21203/rs.3.rs-3222892/v1>
7. Van den Berg, J. (1958). Myoelastic-aerodynamic theory of voice production. *J. Speech Hear. Res.* 1, 227–244.
8. Herbst, C.T., Stoeger, A.S., Frey, R., Lohscheller, J., Titze, I.R., Gumpenberger, M., and Fitch, W.T. (2012). How low can you go? Physical production mechanism of elephant infrasonic vocalizations. *Science* 337, 595–599.
9. Håkansson, J., Mikkelsen, C., Jakobsen, L., and Elemans, C.P. (2022). Bats expand their vocal range by recruiting different laryngeal structures for echolocation and social communication. *Plos Biology* 20, e3001881.

10. Remmers, J., and Gautier, H. (1972). Neural and mechanical mechanisms of feline purring. *Respir. Physiol.* 16, 351–361.
11. Sissom, D.E.F., Rice, D., and Peters, G. (1991). How cats purr. *J. Zool.* 223, 67–78.
12. Herbst, C.T., Prigge, T., Garcia, M., Hampala, V., Hofer, R., Weissengruber, G.E., Svec, J.G., and Fitch, W.T. (2023). Domestic cat larynges can produce purring frequencies without neural input. *Curr. Biol.* 33, XXXX–XXXX.
13. Herbst, C.T. (2016). Biophysics of vocal production in mammals. In *Vertebrate Sound Production and Acoustic Communication*, R.A. Suthers, W.T. Fitch, R.R. Fay, and A.N. Popper, eds. (Cham, Switzerland: Springer), pp. 159–189.
14. Henrich, D.N. (2006). Mirroring the voice from Garcia to the present day: Some insights into singing voice registers. *Logoped. Phoniatr. Vocol.* 31, 3–14.
15. Kime, N.M., Ryan, M.J., and Wilson, P.S. (2019). Modelling the production of complex calls in the túngara frog (*Physalaemus pustulosus*). *Bioacoustics* 28, 345–363.
16. Titze, I.R., Fitch, W.T., Hunter, E.J., Alipour, F., Montequin, D., Armstrong, D.L., McGee, J., and Walsh, E.J. (2010). Vocal power and pressure–flow relationships in excised tiger larynges. *J. Exp. Biol.* 213, 3866–3873.
17. Feynman, R.P., Leighton, R.B., and Sands, M. (2011). *The Feynman lectures on physics, Vol. I: The new millennium edition: mainly mechanics, radiation, and heat* (New York City: Basic books).




Figure 1. Waveform and spectrogram of a bout of four purrs extracted from a long purring sequence recorded at the cat's nostrils.

Purring sequences consist of alternating calls produced during exhalation and inhalation. The fundamental frequency (f_0) of the purr is very low (here 30 Hz) and remains highly regular throughout the sequence, which has led to a debate on whether it is produced by nerve induced muscular vibration or by passive vibration as a consequence of the airflow. The study by Herbst *et al.* shows that vibration at these low frequencies does not require muscular activity.

In Brief

Cat purring, the unusual, pulsed vibration that epitomizes comfort, enjoys a special status in the world of vocal communication research. Indeed, it has long been flagged as a rare exception to the dominant theory of voice production in mammals. A new study presents histological and biomechanical evidence that purring can occur passively, without needing muscle vibration in the larynx controlled by an independent neural oscillator.