



Complete and circularized genome sequences of five nitrogen-fixing *Bradyrhizobium* sp. strains isolated from root nodules of peanut, *Arachis hypogaea*, cultivated in Tunisia

Besma Bouznif, Amira Boukherissa, Yan Jaszczyszyn, Mohamed Mars,
Tatiana Timchenko, Jacqui A Shykoff, Benoît Alunni

► To cite this version:

Besma Bouznif, Amira Boukherissa, Yan Jaszczyszyn, Mohamed Mars, Tatiana Timchenko, et al.. Complete and circularized genome sequences of five nitrogen-fixing *Bradyrhizobium* sp. strains isolated from root nodules of peanut, *Arachis hypogaea*, cultivated in Tunisia. *Microbiology Resource Announcements*, 2024, 13, 10.1128/mra.01078-23 . hal-04798785

HAL Id: hal-04798785

<https://hal.science/hal-04798785v1>

Submitted on 22 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Complete and circularized genome sequences of five nitrogen-fixing *Bradyrhizobium* sp. strains isolated from root nodules of peanut, *Arachis hypogaea*, cultivated in Tunisia

Besma Bouznif,^{1,2,3,4} Amira Boukherissa,^{1,2} Yan Jaszczyszyn,¹ Mohamed Mars,³ Tatiana Timchenko,¹ Jacqui A. Shykoff,² Benoît Alunni^{1,5}

AUTHOR AFFILIATIONS See affiliation list on p. 3.

ABSTRACT This manuscript reports the complete and circularized Oxford Nanopore Technologies (ONT) long read-based genome sequences of five nitrogen-fixing symbionts belonging to the genus *Bradyrhizobium*, isolated from root nodules of peanut (*Arachis hypogaea*) grown on soil samples collected from Tunisia.

KEYWORDS symbiosis, nitrogen fixation, *Bradyrhizobium*, peanut, Tunisia

Legume plants have evolved the capacity to harbor symbiotic nitrogen-fixing soil bacteria within root nodules. This symbiosis provides these plants the ability to grow in diverse habitats and to become major players in agricultural sustainability (1, 2). Bacteria belonging to the genus *Bradyrhizobium* are known as predominant peanut (*Arachis hypogaea*) symbionts (3–6).

In this announcement, we report the complete genome sequence of five bacterial strains isolated from peanut root nodules and belonging to the genus *Bradyrhizobium* based on average nucleotide identity (ANI) value analysis (Table 1). The soils were sampled from five sites in Tunisia (Table 1). Then, Tunisian traditional varieties of peanuts were grown in pots containing the sampled soils for bacterial trapping. After 7 weeks of greenhouse cultivation (28°C, 16 h photoperiod, 160 μmol·m⁻²·s⁻¹), nodules were collected and surface-sterilized in 96% ethanol for 1 min, then in 3% sodium hypochlorite solution for 3 min before being rinsed three times with sterile distilled water and individually crushed, plated on yeast extract mannitol (YM) agar medium and grown for 5 to 10 days at 28°C (7, 8). The strains were purified by successive streaking and single-colony picking and then stored in YM medium with 20% (vol/vol) glycerol at -80°C.

Total genomic DNA was extracted from the five strains grown in the YM medium for 4 days at 28°C using the MasterPure complete DNA and RNA Purification Kit (Epicentre). DNA libraries were prepared from intact genomic DNA with the Native Barcoding Kit 24 V14 (SQK-NBD114.24), and sequencing was performed on a MinION flow cell (R10.4.1). Basecalling was performed with the Super Accuracy model in Guppy 5.3.1. Nanopore library adaptors were trimmed using Porechop v2.0.4 (<https://github.com/rrwick/Porechop>). The read quality was assessed using NanoPlot v1.41.0 (9). The long reads were then assembled using Flye 2.8.3 (10), which generated a single contig assembly for four strains and two contigs for the strain BWA-3-5 (<https://doi.org/10.6084/m9.figshare.24547348>). The presence of plasmids was determined with plASGraph2 v1.0.0 (11). The assemblies were polished with Medaka v1.7.2 (model r1041_e82_260bps_sup_g632) (<https://github.com/nanoporetech/medaka>) using the Oxford Nanopore Technology (ONT) reads to create a consensus sequence. The assemblies of the chromosomes were then rotated with Circlator all at the DnaA gene (12).

Editor Leighton Pritchard, University of Strathclyde, Glasgow, United Kingdom

Address correspondence to Benoît Alunni, benoit.alunni@inrae.fr.

The authors declare no conflict of interest.

See the funding table on p. 3.

Received 13 November 2023

Accepted 15 April 2024


Published 15 May 2024

Copyright © 2024 Besma et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

TABLE 1 Genome features of the five *Bradyrhizobium* sp. strains

Strain characteristic(s)	Data for:				
	<i>Bradyrhizobium</i> sp. BEA-2-5	<i>Bradyrhizobium</i> sp. BWA-3-5	<i>Bradyrhizobium</i> sp. BWC-3-1	<i>Bradyrhizobium</i> sp. NDS-1	<i>Bradyrhizobium</i> sp. sBnM-33
Sampling site	Béja, Elmarja	Béja, Wechtata	Béja, Wechtata	Nabeul, Dar Allouch	Ben Gardane
Peanut variety	Arbi	Arbi	Chounfakhi	Siniya	Massriya
No. of reads (post QC)	91,274	118,365	141,671	99,886	80,650
No. of contigs	1	2	1	1	1
No. of genes	7,837	7,729	8,367	7,134	8,642
Genome length (bp)	8,406,336	8,041,125	8,767,305	7,645,890	9,184,954
		(chromosome: 7,897,608; plasmid: 143,517)			
Coverage (x)	148	188	161	190	146
N50 (bp)	8,406,336	7,897,608	8,767,305	7,645,890	9,184,954
GC content (%)	63.92	62.51	62.92	64.01	61.62
BUSCO score (%)	99.1	99.9	99.5	99.4	99.2
Closest taxonomic assignment—accession	<i>Bradyrhizobium</i> <i>pachyrhizi</i> —	<i>Bradyrhizobium</i> <i>hereditatis</i> —	<i>Bradyrhizobium</i> <i>canariense</i> —	<i>Bradyrhizobium</i> <i>frederickii</i> —	<i>Bradyrhizobium</i> <i>retamae</i> —
(ANI value to closest species [%])	GCF_029714545.1 (95.89)	GCF_020329435.1 (89.14)	GCF_019402665.1 (94.70)	GCF_004570865.1 (90.13)	GCF_001440415.1 (92.53)
BioSample ID	SAMN37684786	SAMN37684780	SAMN37683990	SAMN37684784	SAMN37684830
SRA ID	SRR26337671	SRR26335727	SRR26335726	SRR26337672	SRR26336054
GenBank ID	CP136629	CP136626-	CP136625	CP136628	CP136624
		CP136627			

The plasmid of the strain BWA-3-5 was rotated with Prodigal v2.6.2 at position 73241 (13). The statistics of the assembly were obtained with QUAST v5.2.0 (<https://github.com/ablab/quast>), and the completeness of the genomes was assessed using BUSCO v5.6.1 (14) (Table 1). Functional annotation of the genomes was performed using the NCBI

**FIG 1** Heatmap of ANI values between the five *Bradyrhizobium* sp. strains. ANI values for pairwise comparisons of these five strains are in the range between 81.2% and 87.7%, below the species threshold of 95%, and therefore appear to belong to five distinct *Bradyrhizobium* species.

Prokaryotic Genome Annotation Pipeline v6.3 (15, 16). ANI calculation was carried out using ANIclustermap v1.2.0 (Fig. 1)(17).

ACKNOWLEDGMENTS

B.B. benefited from support from the Programme Hubert Curien UTIQUE Project 17G0918, the Government of Tunisia, for a mobility grant (Bourse d'Alternance) and a doctoral partner scholarship from the French Ministry of Foreign and European Affairs. A.B. benefited from a Ph.D. contract in the frame of the CNRS 80|PRIME—2021 program. We acknowledge the sequencing and bioinformatics expertise of the I2BC High-throughput sequencing facility, supported by France Génomique (funded by the French National Program "Investissement d'Avenir" ANR-10-INBS-09). This work has benefited from a French State grant (Saclay Plant Sciences, reference no. ANR-17-EUR-0007, EUR SPS-GSR) under a France 2030 program (reference no. ANR-11-IDEX-0003).

AUTHOR AFFILIATIONS

¹Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), Gif-sur-Yvette, France

²IDEEV—Laboratoire Ecologie, Systématique et Evolution, CNRS, AgroParisTech Université Paris-Saclay, Gif-sur-Yvette, France

³Research Unit Biodiversity and Valorization of Arid Areas Bioresources (VBAA), Faculty of Sciences, Gabès, Tunisia

⁴Université de Picardie Jules Verne, UMR BioEcoAgro 1158-INRAE, BIOPI, Amiens, France

⁵Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), Versailles, France

AUTHOR ORCIDs

Benoît Alunni  <http://orcid.org/0000-0002-3616-0669>

FUNDING

Funder	Grant(s)	Author(s)
Programme Hubert Curien	UTIQUE Project 17G0918	Bouznif Besma Mars Mohamed Shykoff Jacqui
Centre National de la Recherche Scientifique (CNRS)	80 Prime	Boukherissa Amira Benoît Alunni
Agence Nationale de la Recherche (ANR)	ANR-10-INBS-09	Jaszczyszyn Yan Benoît Alunni
Agence Nationale de la Recherche (ANR)	ANR-17-EUR-0007	Timchenko Tatiana Benoît Alunni
Agence Nationale de la Recherche (ANR)	ANR-11-IDEX-0003	Timchenko Tatiana Benoît Alunni

DATA AVAILABILITY

The raw reads and complete genomes of the five *Bradyrhizobium* sp. strains were deposited at GenBank under the BioProject no. PRJNA1023532. The BioSample numbers, raw reads, and assembly GenBank accession numbers are given in Table 1.

REFERENCES

1. Sprent JI. 2007. Evolving ideas of legume evolution and diversity: a taxonomic perspective on the occurrence of nodulation. *New Phytol* 174:11–25. <https://doi.org/10.1111/j.1469-8137.2007.02015.x>
2. Liu A, Ku Y-S, Contador CA, Lam H-M. 2020. The impacts of domestication and agricultural practices on legume nutrient acquisition through symbiosis with rhizobia and arbuscular mycorrhizal fungi. *Front Genet* 11:583954. <https://doi.org/10.3389/fgene.2020.583954>
3. Muñoz V, Ibañez F, Tonelli ML, Valetti L, Anzuay MS, Fabra A. 2011. Phenotypic and phylogenetic characterization of native peanut *Bradyrhizobium* isolates obtained from Córdoba, Argentina. *Syst Appl Microbiol* 34:446–452. <https://doi.org/10.1016/j.syapm.2011.04.007>
4. Jaiswal SK, Msimbira LA, Dakora FD. 2017. Phylogenetically diverse group of native bacterial symbionts isolated from root nodules of groundnut (*Arachis hypogaea L.*) in South Africa. *Syst Appl Microbiol* 40:215–226. <https://doi.org/10.1016/j.syapm.2017.02.002>
5. Shao S, Chen M, Liu W, Hu X, Wang E-T, Yu S, Li Y. 2020. Long-term monoculture reduces the symbiotic rhizobial biodiversity of peanut. *Syst Appl Microbiol* 43:126101. <https://doi.org/10.1016/j.syapm.2020.126101>
6. Bouznif B, Alunni B, Mars M, Shykoff JA, Timchenko T, Rodríguez de la Vega RC. 2021. Draft genome sequences of nitrogen-fixing *Bradyrhizobia* isolated from root nodules of peanut, *Arachis hypogaea*, cultivated in Southern Tunisia. *Microbiol Resour Announc* 10:e0043421. <https://doi.org/10.1128/MRA.00434-21>
7. Rodríguez-Echeverría S, Pérez-Fernández MA, Vlaar S, Finan TM. 2003. Analysis of the legume–rhizobia symbiosis in shrubs from central Western Spain. *J Appl Microbiol* 95:1367–1374. <https://doi.org/10.1046/j.1365-2672.2003.02118.x>
8. Vincent JM. 1970. A manual for the practical study of the root-nodule bacteria. *Man Pract Study Root-Nodule Bact.*
9. De Coster W, Rademakers R. 2023. NanoPack2: population-scale evaluation of long-read sequencing data. *Bioinformatics* 39:btad311. <https://doi.org/10.1093/bioinformatics/btad311>
10. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol* 37:540–546. <https://doi.org/10.1038/s41587-019-0072-8>
11. Sielemann J, Sielemann K, Brejová B, Vinař T, Chauve C. 2023. PIAS-graph2: using graph neural networks to detect plasmid contigs from an assembly graph. *Front Microbiol* 14:1267695. <https://doi.org/10.3389/fmich.2023.1267695>
12. Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>
13. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>
14. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>
15. Li W, O'Neill KR, Haft DH, DiCuccio M, Chetvernin V, Badretdin A, Coulouris G, Chitsaz F, Derbyshire MK, Durkin AS, Gonzales NR, Gwadz M, Lanczycki CJ, Song JS, Thanki N, Wang J, Yamashita RA, Yang M, Zheng C, Marchler-Bauer A, Thibaud-Nissen F. 2021. RefSeq: expanding the prokaryotic genome annotation pipeline reach with protein family model curation. *Nucleic Acids Res* 49:D1020–D1028. <https://doi.org/10.1093/nar/gkaa1105>
16. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>
17. Lisa M, Bot H. 2017. My research software (2.0.4)