



HAL
open science

Correction: Toxicity of metal–organic framework nanoparticles: from essential analyses to potential applications

Romy Ettliger, Ulrich Lächelt, Ruxandra Gref, Patricia Horcajada, Twan Lammers, Christian Serre, Patrick Couvreur, Russell E Morris, Stefan Wuttke

► To cite this version:

Romy Ettliger, Ulrich Lächelt, Ruxandra Gref, Patricia Horcajada, Twan Lammers, et al.. Correction: Toxicity of metal–organic framework nanoparticles: from essential analyses to potential applications. *Chemical Society Reviews*, 2023, 52 (3), 10.1039/d3cs90014b . hal-04799043

HAL Id: hal-04799043

<https://hal.science/hal-04799043v1>

Submitted on 22 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Cite this: *Chem. Soc. Rev.*, 2023, 52, 1156

Correction: Toxicity of metal–organic framework nanoparticles: from essential analyses to potential applications

Romy Ettlinger,^{*a} Ulrich Lächelt,^{bc} Ruxandra Gref,^d Patricia Horcajada,^e Twan Lammers,^f Christian Serre,^g Patrick Couvreur,^h Russell E. Morris^a and Stefan Wuttke^{*ij}

DOI: 10.1039/d3cs90014b

Correction for 'Toxicity of metal–organic framework nanoparticles: from essential analyses to potential applications' by Romy Ettlinger *et al.*, *Chem. Soc. Rev.*, 2022, 51, 464–484, <https://doi.org/10.1039/D1CS00918D>.

rsc.li/chem-soc-rev

The authors regret that the information presented in Fig. 12 and discussed in the text on p. 478 may have been unclear, or inconsistent with the information derived from ref. 26. The corrected version of Fig. 12 is presented here. The corresponding sentence on p. 478, beginning, "Herein, endogenous fumaric acid...", should also be revised as follows:

"Herein, nitro- and tetramethyl-terephthalic acid modifications turned out to be the most biofriendly organic linkers, followed by exogenous trimesic acid, amino-terephthalic acid, and terephthalic acid, and the endogenous fumaric acid had the lowest assessed IC₅₀ value."




Fig. 12 Ranking of different organic linkers, based on their reported IC₅₀ toxicity data.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a School of Chemistry, University of St. Andrews, St. Andrews, UK. E-mail: rle6@st-andrews.ac.uk

^b Department of Pharmacy and Center for NanoScience (CeNS), Ludwig-Maximilians-Universität München, Munich, Germany

^c Division of Pharmaceutical Technology and Biopharmaceutics, University of Vienna, Vienna, Austria

^d Institut de Sciences Moléculaires d'Orsay, Université Paris Saclay, Paris, France

^e Madrid Institute for Advanced Studies, Madrid, Spain

^f Institute for Experimental Molecular Imaging, RWTH Aachen University Clinic, Aachen, Germany

^g Département de Chimie, Ecole Normale Supérieure de Paris, Paris, France

^h Institut Galien Paris-Sud, Université Paris Saclay, Paris, France

ⁱ Ikerbasque, Basque Foundation for Science, Bilbao, Spain

^j Basque Center for Materials, UPV/EHU Science Park, Leioa, Spain. E-mail: stefan.wuttke@bcmaterials.net

