



# The inducible KPC:APC mouse organoid model for colon carcinogenesis studies

Mélissandre Gomot, Nora Essakhi, Clément Codan, Quentin Pascal,  
Guillaume Vares

## ► To cite this version:

Mélissandre Gomot, Nora Essakhi, Clément Codan, Quentin Pascal, Guillaume Vares. The inducible KPC:APC mouse organoid model for colon carcinogenesis studies. Engineering in vitro Human Physiology - Tissues, Organoids and Organs-on-Chip for Clinical Applications -, Nov 2024, Paris, France.  
hal-04799476

HAL Id: hal-04799476

<https://hal.science/hal-04799476v1>

Submitted on 22 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

# The inducible KPC:APC mouse organoid model for colon carcinogenesis studies

Mélissandre Gomot<sup>1\*</sup>, Nora Essakhi<sup>1</sup>, Clément Codan<sup>1</sup>, Quentin Pascal<sup>2</sup>, Guillaume Vares<sup>1</sup>

1 Institut de Radioprotection et de Sûreté Nucléaire (IRSN), PSE-SANTE/SESANE/LRTOX, Fontenay-aux-Roses, France

2 Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA), IMVA-HB/IDMIT, Fontenay-aux-Roses, France

## Introduction


Every day, 500,000 CT-scans are performed worldwide, and this number is on the rise<sup>1</sup>.

Abdominal and pelvic CT-scans have the highest LAR (Lifetime Attributable Risk) of solid cancer. They deliver 8-31 mGy per scan on average<sup>2,3</sup>. An increasing number of patients are exposed to doses >100 mSv<sup>1</sup>.

### Ionizing radiation and colon cancer

Epidemiological studies show an increased risk of colorectal cancer after low dose medical radiation exposure (LDR) but benefits still outweigh the risk of radio-induced cancer<sup>4</sup>. However, LDR increase tumor multiplicity and interact with carcinogenesis-related pathways through gene expression modulation, DNA hypomethylation or protein interactions<sup>5,6,7</sup>.

The Apc<sup>min</sup> mouse is a widely used model of colorectal cancer but inducible models are more suitable for colon carcinogenesis studies.



**Objectives →** To develop and characterize suitable *in vivo* and organoid models to study colon carcinogenesis after medical diagnostic radiation exposure.


## Methods

**KPC : APC**  
**Inducible colon cancer mouse model**  
B6.Cg-Kras<sup>tm1Tij</sup> Apc<sup>tm1Tno</sup> Tg(CDX2-cre/ERT2)752Erf/Maraj<sup>8</sup>  

- o CreLox system
- o Under control of the CDX2 promoter

Kras<sup>+/f</sup> Apc<sup>ff</sup> Cre<sup>ERT2+</sup> (ApcKras)  
 Kras<sup>+/f</sup> Apc<sup>ff</sup> Cre<sup>ERT2++</sup> (Apc)  
 Kras<sup>+/f</sup> Apc<sup>ff</sup> (Control)


Organoids were grown in complemented Matrigel (B27, N2, ROCK inhibitor, EGF and N-acetylcysteine). ROCK inhibitor was added to the medium for 3 days after culture initiation and organoid passaging. Organoids were treated with 4-OHT (0,125-1µM) for 24h two days after culture initiation at passage 0 or 1 to induce Apc and/or Kras mutations.




## Results

### Treatment on the second week of culture p(1)


Dose effect at p(2)



Organoid size




Proportion of spheroids and organoids for each genotype




N=2 ; n=3 - Dunnett's multiple comparison test, each treated group was compared to the corresponding untreated genotype, \* : p-value < 0,05

### Treatment on the first week of culture p(0)


Dose effect at p(0)




Organoid size




Proportion of spheroids and organoids for each genotype




Characterization after treatment with 0,125µM of 4-OHT




Stem cell markers



Wnt pathway activation



RNA methylases and demethylases




n=3, Standard deviation, Two-way ANOVA with Tukey multiple comparison.  
\* : p-value < 0,05

## Application


Organoids were irradiated with 25 or 250mGy X-rays. The irradiation was performed on the 5<sup>th</sup> day of p(0) (early irradiation), or on the 5<sup>th</sup> day of p(1) (late irradiation).

### Effect of early irradiation on organoids size



N>2 ; n=8 - Dunnett's multiple comparison test, \* : p-value < 0,05

### Effect of late irradiation on organoids size



\* Contact: melissandre.gomot@irsn.fr

## Conclusions

We have developed a new KPC:APC dynamic organoid model which allows the induction and study of colon carcinogenesis *in vitro*. After inducing Kras and/or Apc mutations, the organoids exhibited increased size and a characteristic cystic morphology. Markers for proliferation, stemness and Wnt pathway activation were also increased, suggesting that this model may be able to recapitulate colon carcinogenesis *in vitro*.

Organoids derived from KPC:APC mice therefore provide the opportunity to perform parallel *in vivo* histopathological and *in vitro* mechanistic studies to advance research on colon carcinogenesis and to investigate the effects of low-dose radiation exposures.

## References

- IRSN, Exposition de la population aux rayonnements ionisants - Rapport EXPRI, 2017.
- S.Bagherzadeh et al., Evaluation of radiation dose and radiation-induced cancer risk associated with routine CT scan examinations, Radiation Physics and Chemistry, 2023.
- J. Albert, Radiation Risk From CT: Implications for Cancer Screening, American Journal of Roentgenology, 2013.
- A. Gonzales, Radiation-Related Cancer Risks From CT Colonography Screening: A Risk-Benefit Analysis, American Journal of Roentgenology, 2012.
- D. Trani, Sex-dependent Differences in Intestinal Tumorigenesis Induced in Apc1638N/+ Mice by Exposure to γ Rays, International Journal of Radiation Biology, 2012.
- S. Soleymaniard et al., Effects of radiation dose on the stemness -related genes expression in colorectal cancer cell line ; International Journal of Radiation Biology, 2021.
- J.-H. Bae, Identification of radiation-induced aberrant hypomethylation in colon cancer, BMC Genomics, 2015.
- R. Maitra et al., Development and Characterization of a Genetic Mouse Model of KRAS Mutated Colorectal Cancer, International Journal of Molecular Sciences, 2019.