

Implementation of a platform for the identification of broad-spectrum antiviral compounds based on 2D/3D cultures of the central nervous system

François Piumi, Berry Noémie, Chaillot Valentine, Magniez Aurélie, Gorna Kamila, Coulpier Muriel

▶ To cite this version:

François Piumi, Berry Noémie, Chaillot Valentine, Magniez Aurélie, Gorna Kamila, et al.. Implementation of a platform for the identification of broad-spectrum antiviral compounds based on 2D/3D cultures of the central nervous system. Journées d'animation scientifique du département santé animale de l'INRAE, Sep 2024, Seignosse, France. hal-04803577

HAL Id: hal-04803577 https://hal.science/hal-04803577v1

Submitted on 25 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Implementation of a platform for the identification of broad-spectrum antiviral compounds based on 2D/3D cultures of the central nervous system

François PIUMI, Noémie BERRY, Valentine CHAILLOT, Aurélie MAGNIEZ, Kamila GORNA, Muriel COULPIER


UMR 1161 Virologie, Anses (Laboratoire de Santé Animale), INRAe, Ecole Nationale Vétérinaire d'Alfort, Université Paris-Est, Maisons-Alfort corresponding author: francois.piumi@vet-alfort.fr

INTRODUCTION

Viral infections of the nervous system are often very serious affections, and can be responsible for fatal encephalitis or severely disabling neurological sequelae. In humans, many viruses are capable of infecting the nervous system; some of them are also responsible for encephalitis in horses. However, there are currently very few antiviral treatments available, and none to block the replication of neurotropic arboviruses (viruses transmitted by mosquitoes and ticks) and limit the damage caused to the brain. Our aim is to identify broad-spectrum antivirals that will be effective against several viral families, in both humans and horses.

This poster will show the different infection models based on 2D/3D cultures of central nervous system cells that we have developed in recent years, and our strategy for identifying broad-spectrum antiviral compounds active in the nervous system.

VIRUSES OF INTEREST





Endemic, emerging or at-risk arboviruses responsible for serious neurological diseases


Viral infection of hNGC, hNPC and eNPC


Neurons loss


PHYSIOLOGICALLY RELEVANT 2D MODEL TO MIMIC VIRAL INFECTION














TBEV: Tick-Borne Encephalitis Virus **WNV**: West Nile Virus **BoDV**: Borna Disease Virus **EEEV**: Eastern Equine Encephalitis Virus **Sars-Cov2:** severe acute respiratory syndrome coronavirus 2


eNPCs

DEVELOPMENT OF A PHENOTYPIC SCREEN BASED ON hNGCs AND IMAGING



COMPARISON OF THE ANTIVIRAL ACTIVITY OF SELECTED **MOLECULES**

- 8 molecules selected from the literature (already described for their anti-flavivirus activity).
- Cells: hNGCs + two others in vitro models with different levels of relevance (A549: human alveolar basal epithelial cells; hNPCs: human neural progenitor cells) – all permissive to TBEV infection.
- Read-out: % of infected cells and % cell viability by an automatized imaging approach.




→ Importance to choose the most physiologically relevant in vitro model to screen antiviral drugs.


→ Our pipeline allows to test the antiviral potential of chemical compounds on a panel of physiologically relevant in vitro models representative of the human and equine infected brain.

CONCLUSION


- Tytotoxicity of molecules for human/equine brain cells can be assessed and their selectivity index can be calculated.
- → Hits identification is statistically determined.
- The diversity of our models allows the identification of broad-spectrum antiviral compounds.
- → Whether the compounds act on neurons and/or astrocytes (antiviral effect/cytotoxicity) maybe resoluted.

Statistical analysis for hits identification






Hits confirmation by quantification of viral RNA/titer




USING HUMAN CEREBRAL ORGANOIDS (3D MODEL) TO **VALIDATE OUR HITS**

TBEV infects and replicates in h. cerebral organoids WNV infects and replicates in h. cerebral organoids (See poster A. Magniez)



Validation of the « hits » using TBEV-infected human cerebral organoids :



→ Molecule γ strongly inhibits TBEV infection in human cerebral organoids.





