



**HAL**  
open science

## Synergy of fine-resolution regional model and SWOT measurements to study coastal dynamics

Nushrat Yeasmin, Audrey Hyeans, Laurent Testut, Valérie Ballu

► **To cite this version:**

Nushrat Yeasmin, Audrey Hyeans, Laurent Testut, Valérie Ballu. Synergy of fine-resolution regional model and SWOT measurements to study coastal dynamics. 30 Years of Progress in radar altimetry symposium, Sep 2024, Montpellier, France. hal-04803664

**HAL Id: hal-04803664**

**<https://hal.science/hal-04803664v1>**

Submitted on 25 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Introduction

SWOT (Surface Water and Ocean Topography) altimetry mission has made remarkable advancements in resolution, coverage and overall water surface observability. Although SWOT's primary objective is to monitor both open ocean and inland water bodies at an unprecedented spatial resolution, its potential application in coastal regions shows great promise. However, fine scale SWOT measurements in the coast faces major challenges regarding fine scale correction of tide and coastal dynamic processes. In this context, the objective of this study is to assess the relevance of high resolution coastal model and its application in analyzing high resolution SWOT observations in a macrotidal region.

### Study Area

- Coastal region of Pertuis Charentais (Bay of Biscay, France).
- Semi-enclosed macro-tidal coastal regime with tidal range 1.5 m to 5.5 m.
- Located under 2 SWOT passes (348, 419) along with several nadir altimetry mission.
- Covered by a network of tide gauges and pressure sensors.

### Model Description SCHISM (Semi-Implicit Cross-scale Hydroscience Integrated System Model)

- A semi-implicit cross-scale hydrodynamic model which solves Navier-Stokes equations (in hydrostatic form) in 3D unstructured grid (Zhang et al., 2016).
- SCHISM in depth-averaged barotropic mode with 2DH configuration developed by Tranchant et al., (2021).
- 57,828 grid nodes with 3 km resolution towards the open ocean and 50 m resolution near the coast.
- Boundary forcing → tide (Tranchant et al., 2021), sea level pressure and wind (ECMWF-ERA5).




Fig 1: SCHISM model mesh with SWOT passes overlay and available tide gauges in the study area

## SWOT ( Surface Water and Ocean Topography)

- Wide-swath 2D altimetry mission using Ka-band Radar Interferometry (KaRIn)**
- Joint project of NASA, CNES, the Canadian Space Agency and UK Space Agency
- Launched on **16<sup>th</sup> December, 2022**
- Mission duration: **3 years**
- Repeatability:
  - Calibration phase: **1 day** ( 5<sup>th</sup> April to 11<sup>th</sup> July of 2023)
  - Science phase: **~21 days** ( 26<sup>th</sup> July, 2023 to onward)




Fig 2. Diagram of SWOT Data Collection  
<https://swot.jpl.nasa.gov/resources/104/diagram-of-swot-data-collection/>

### Data type and observability

#### Ocean

- Low Resolution (LR)
- 250 m x 250 m
- Characterizing meso- and sub-mesoscale 2D ocean circulations

#### Inland

- High Resolution (HR)
- 5 m (along-track) x 10-70 m
- Observation of water bodies with width larger than 100 m

#### Coast

- No dedicated product
- Both LR and HR coverage
- Possibility of observing fine scale coastal dynamics, surges etc.

## Model Performance

### Tidal constituents considered for model performance validation

Astronomical	K1	O1	M2	N2	S2	K2	MU2	NU2	M3
Shallow-water	M4	M6	MN4	MS4					




Fig 3: Model performance evaluation using complex error ( $\sigma$ ) between observed and modelled tidal constituents. The estimation of  $\sigma$  accounts for both amplitude and phase difference.




Fig 4: Comparison of modelled shallow water constituents with observation. Top panels show the amplitude differences, mid panels show the phase differences and bottom panels show the complex differences between observation and FES 2014 (blue bars), FES 2022 (violet bars) and Tranchant et al., (2021) (red bars).

## Conclusion and Perspective

- In the coastal region, high resolution SWOT measurements can bring crucial information regarding fine scale coastal dynamics.
- High resolution regional models for geophysical corrections are essential to fully use SWOT's potential.
- The regional model used in this study is performing well compared to both FES 2022 and FES 2014 global model. Further improvement in representation of shallow water constituents are ongoing.
- Current model will be coupled with Wind Wave Model (WWM) to include short wave dynamics.
- Dedicated multi-sensor field campaigns are currently being conducted over the study area to complement the modelling (c.f. Poster by Audrey Hyeans) in assessing SWOT measurements in coasts.

## Recommendation

- Dedicated SWOT data products for coastal region are required to utilize the unique dataset of SWOT at its full potential in the coastal region.
- Observed permanent artifacts in the SWOT measurements near coasts should be further investigated by the SWOT processing team.

## Result

### High resolution coastal model: a necessity for SWOT !!




Fig 5: Tide and dynamic atmospheric correction (dac) from regional model (a) and from SWOT LR level 2 product (b) and their comparison.

### SWOT observed coastal gradient




Fig 6: Coastal SSHA gradient from model (a) and SWOT (b, c) relative to Aix island tide gauge location (red star). SSHA in (b) and (c) are computed using Eqn 1. Mean dynamic topography (mdt) correction used in (b) is from SWOT L2 product (d) and correction used in (c) is computed with 15 cycles of SWOT data (e) using Eqn 2, which includes mdt and stationary SWOT artifacts.

$$SSH A_{SWOT} = ssh_{karin\_2} + xover\ cor(crossover\ correction) - geoid - mdt - solid\ earth\ tide - load\ tide - pole\ tide - internal\ tide \quad Eqn. 1$$

$$mdt_{SWOT} = \frac{1}{N} \sum_{cycle=N} ssh_{karin\_2} + xover\ cor - geoid - (ocean\ tide + dac)_{model} - (solid\ earth\ tide + load\ tide + pole\ tide + internal\ tide)_{L2\ product} \quad Eqn. 2$$

- With appropriate geophysical correction SWOT can observe coastal sea surface gradient.
- SWOT data derived mdt includes some stationary artifacts and ocean dynamics in SWOT measurements.

## References

- Tranchant, Y.-T., Testut, L., Chupin, C., Ballu, V., & Bonnefond, P. (2021). Near-Coast Tide Model Validation Using GNSS Unmanned Surface Vehicle (USV), a Case Study in the Pertuis Charentais (France). *Remote Sensing*, 13(15), Article 15. <https://doi.org/10.3390/rs13152886>
- Zhang, Y. J., Ye, F., Stanev, E. V., & Grashorn, S. (2016). Seamless cross-scale modeling with SCHISM. *Ocean Modelling*, 102, 64–81. <https://doi.org/10.1016/j.ocemod.2016.05.002>
- Surface Water Ocean Topography (SWOT). 2024. SWOT Level 2 KaRin Low Rate Sea Surface Height Data Product, Version 2.0. Ver. 2.0. PO.DAAC, CA, USA. Dataset accessed [2024-08-22] at <https://doi.org/10.5067/SWOT-SSH-2.0>
- SWOT Level-2 KaRin Low Rate SSH Unsmoothed (v2.0) [Data set]. CNES. <https://doi.org/10.24400/527896/A01-2023-016>.