



HAL
open science

The Fractional Poisson Process and Other Limit Point Processes for Rare Events in Infinite Ergodic Theory

Dylan Bansard-Tresse

► **To cite this version:**

Dylan Bansard-Tresse. The Fractional Poisson Process and Other Limit Point Processes for Rare Events in Infinite Ergodic Theory. 2024. hal-04811660

HAL Id: hal-04811660

<https://hal.science/hal-04811660v1>

Preprint submitted on 29 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

The Fractional Poisson Process and Other Limit Point Processes for Rare Events in Infinite Ergodic Theory

Dylan Bansard-Tresse

CPHT, CNRS, École polytechnique,
Institut Polytechnique de Paris, 91120 Palaiseau, France

dylan.bansard-tresse@polytechnique.edu

<https://sites.google.com/view/bansard-tresse>

November 28, 2024

Abstract

We study the process of suitably normalized successive return times to rare events in the setting of infinite-measure preserving dynamical systems. Specifically, we consider small neighborhoods of points whose measure tends to zero. We obtain two types of results. First, we conduct a detailed study of a class of interval maps with a neutral fixed point and we fully characterize the limit processes for all points, highlighting a trichotomy and the emergence of the fractional (possibly compound) Poisson process. This is the first time that these processes have been explicitly identified in this context. Second, we prove an abstract result that offers an explanation for the emergence of the fractional Poisson process, as the unique fixed point of a functional equation, drawing a parallel with the well-established behavior of the Poisson process in finite-measure preserving dynamical systems.

Contents

1	Introduction	2
2	Statement of results	4
2.1	Preliminaries	4
2.2	Abstract results	6
2.2.1	The general relationship between hitting and return REPPs	6
2.2.2	The fractional Poisson process as a common limit law	8
2.3	Rare event point processes in maps with an indifferent fixed point	10
2.3.1	Points that are not preimages of the indifferent fixed point	11
2.3.2	The indifferent fixed point and its preimages	11
2.3.2.1	Barely infinite case $p = 1$	13
3	Return and Hitting Rare Event Point Processes and the Fractional Poisson Process	13
3.1	Equivalence between Hitting and Return Point Processes	13
3.1.1	Proof of Theorems 2.1-2.2	13
3.1.2	Proof of Proposition 2.4	18
3.2	Sufficient conditions for convergence towards FPP_α and $CFPP_\alpha$	19
4	Quantitative recurrence for maps with an indifferent fixed point	22
4.1	Additional properties	22
4.2	Proof of Theorems 2.4-2.5	24
4.2.0.1	Proof of Theorem 2.4: Non periodic points.	24
4.2.0.2	Proof of Theorem 2.5: Periodic points.	25
4.3	Proofs of Theorems 2.7 and 2.9	26
4.3.1	The case $x = 1/2$	26
4.3.2	Preimages of $1/2$	27
4.3.3	Proof of Theorem 4.1	27
4.3.4	Proof of Theorems 2.7 and 2.9	33
4.3.5	Proof of Proposition 2.7	34
5	Fractional Poisson processes in infinite ergodic theory: a concluding perspective	34
	References	36

A	Additional proofs	38
A.1	Proof of Corollary 2.6	38
A.2	Proof of Lemma 2.8	39

1 Introduction

We consider a measure-theoretic dynamical system (X, \mathcal{B}, μ, T) where T acts on the phase space X and preserves the measure μ which can be finite or infinite. We are interested in “asymptotically rare events”, meaning that we consider sequences $(B_n)_{n \geq 0}$ of measurable sets such that $\mu(B_n) \rightarrow 0$ as $n \rightarrow \infty$. Denoting $r_{B_n}^{(k)}$ the k -th return to B_n , we want to find the asymptotic behavior of the point process

$$N_{B_n}^\gamma := \sum_{k \geq 1} \delta_{\gamma(\mu(B_n)) r_{B_n}^{(k)}}$$

where γ is an appropriate scaling function. Such point processes are often called “Rare Event Point Processes” (REPP for short). When μ is a probability measure, extensive research has focused on the limiting distribution of the first hitting or return time, with contributions from many authors (see, *e.g.*, [Sau09, Hay13] and references therein), or the whole sequence of return times seen as a process (see [LFF+16] and references therein). For differentiable dynamical systems, the measure μ is either absolutely continuous with respect to the Lebesgue measure or, more broadly, is an SRB (Sinai-Ruelle-Bowen) measure; see, *e.g.*, [CC13, SB22]. Due to Kač’s theorem, the right scaling consists in taking $\gamma = \text{id}$. In this context, if mixing is sufficiently strong or correlations decay sufficiently fast, when the sets B_n are either balls or cylinder sets shrinking to a point x in the phase space and x is μ -generic, then $N_{B_n}^{\text{id}}$ converges towards the Poisson point process (PPP for short), with this convergence understood when $N_{B_n}^{\text{id}}$ is considered as a random variable on the probability space (X, \mathcal{B}, μ) or on the restricted probability spaces $(B_n, \mathcal{B} \cap B_n, \mu_{B_n})$. These cases are commonly referred to as “hitting REPP” and “return REPP”, respectively.

Hitting and return REPPs are intrinsically tied, converging to the same limit (see [HLV05] for the first hitting/return time relationship and [Mar17, Zwe16] for the point process version). The Poisson point process is the unique fixed point of the equation connecting the two limits, thereby confirming its prominent role as the anticipated limit point process for sufficiently mixing systems.

Yet, not all dynamical systems of interest preserve a probability measure, and some indeed preserve a σ -finite measure with infinite mass. Examples include null-recurrent Markov chains and Markov shifts [Sar01], interval maps with indifferent fixed points [Aar97, Tha80], \mathbb{Z}^d -extensions of probability preserving systems or billiards with cusps. For conservative ergodic systems (see Section 2.1 for precise definitions), every set of positive measure is visited infinitely often by almost every orbit. Consequently, $N_{B_n}^\gamma$ is well defined, and the question of its limiting behavior arises naturally. Research on infinite measure systems has been more limited but has seen significant growth in recent years. According to Kač’s formula, the mean return time to any finite measure set is infinite, making the identity scaling $\gamma = \text{id}$ inappropriate. Nevertheless, suitable scalings can still be identified [RZ20]. The convergence of first hitting and return times has been investigated, revealing non-exponential limiting laws for various infinite measure preserving dynamical systems and natural targets [BZ01, PS10, RZ20, Yas18, Yas24], with fewer studies considering the whole sequence of successive returns [PS17]. Recent work [PS24] has shed light on the limiting behavior of point processes for \mathbb{Z} -extensions of subshifts of finite type.

In this paper, we encompass both an in-depth analysis of a paradigmatic interval map with a neutral fixed point and a general theoretical framework. We will now present our results in a fairly informal way, deferring the precise statements until afterwards.

We choose to focus on the following class of interval maps with a neutral fixed point as it allows for a clear illustration of our main results without the need for overly technical considerations. Namely, we consider $X = [0, 1]$ and

$$T_p(x) = \begin{cases} x + 2^p x^{p+1}, & 0 \leq x < 1/2, \\ 2x - 1, & 1/2 \leq x \leq 1, \end{cases} \quad (1.1)$$

where 0 is the neutral fixed point, and p is a nonnegative parameter. This map preserves an absolutely continuous measure which is finite for $p < 1$ and infinite for $p \geq 1$ (see more details below). This map is commonly referred to as Manneville-Pomeau map or LSV map in the literature, following Liverani, Saussol and Vaienti [LSV99]. This paradigmatic one-parameter family of maps displays a rich spectrum of statistical behaviors in both finite (see *e.g.* [Gou04]) and infinite (see *e.g.* [Tha80, Tha83]) contexts.

Our first result establishes the *fractional Poisson process* as the limiting point process for cylinders shrinking to *generic points* of the invariant measure in the infinite measure case (see Theorem 2.4, p.11 for the precise statement).

Theorem A. For the map (1.1) with $p > 1$, and for cylinders B_n shrinking to a point x which is generic for the absolutely continuous invariant measure, the point process $N_{B_n}^\gamma$ converges in law towards a fractional Poisson process (parametrized by p), both when one starts in B_n or off B_n (return and hitting REPPs, respectively).

The fractional Poisson process, which we will define below, was developed to extend the framework of the Poisson process to systems exhibiting long-term temporal correlations, which prevent exponential behavior [Las03] (see also [MS19] for related applications). The following result offers an abstract explanation for the emergence of the fractional Poisson process, drawing a parallel with the well-established behavior of the Poisson process in finite-measure preserving dynamical systems, as previously mentioned. It generalizes the results of [Zwe16] to the infinite measure situation and builds up on [RZ20] where only the first return is considered.

Theorem B (Abstract result). Consider a dynamical system (X, \mathcal{B}, μ, T) preserving an infinite measure, and let $(B_n)_{n \geq 0}$ be a sequence of asymptotically rare events lying in a “good” subset of X . Then, $N_{B_n}^\gamma$ when one starts off B_n (hitting REPP) converges in law if and only if $N_{B_n}^\gamma$ when one starts in B_n (return REPP) converges in law. If convergence takes place, both limits determine one another through a functional equation whose unique fixed point is the fractional Poisson process (with parameter related to γ).

For the precise statement, see Theorem 2.1, p.6. It is worth noting that restricting to “good” subsets is an intrinsic limitation arising in infinite ergodic theory, and that the scaling function γ is directly linked to the regularly varying function defining the correct normalizing sequence in the analogue of the ergodic theorem for the infinite measure case, see Section 2.1 for further details.

In the context of finite-measure preserving dynamical systems, the study of non-generic points, especially periodic points, has gained significant attention in recent years, particularly in the context of extreme value theory. It is not surprising that these periodic points give rise to clustering phenomena leading to a compound Poisson limit process for hitting and return times. The intensity and multiplicity of Poisson compound processes are characterized by a parameter θ , lying between 0 and 1, known as the “extremal index” (see [LFF⁺16] and references therein). Besides the generic points of the invariant measure under consideration and periodic points, there also exist generic points for other invariant measures, as well as points that are not generic for any measure. For several classes of probability-preserving dynamical systems with sufficiently rapid decay of correlations, a *dichotomy* arises in the asymptotic behavior of points: periodic points lead to a compound Poisson limit process, while *all* other points give rise to a Poisson limit process; see *e.g.*, [LFF⁺16] (Rychlik and Gibbs-Markov maps), [FFTV16] (for the map (1.1) with $p < 1$) and [BTF23, DT23] (quadratic maps with Misiurewicz parameters).

In the context of infinite measure preserving dynamical systems, the only existing study of certain non-generic points is done for hyperbolic periodic points of prototypical null-recurrent interval maps [RZ20]. In this paper, we conduct a thorough investigation of the application of the interval (1.1) and demonstrate that the asymptotic behavior of successive return times exhibits not a dichotomy but a trichotomy, involving fractional Poisson processes, compound fractional Poisson processes, and a third type of process (for which no standard nomenclature has been established). This result represents a novel contribution to the theory of infinite-measure dynamical systems.

Theorem C. Consider the map (1.1) with $p =: 1/\alpha > 1$ with its (infinite) absolutely continuous invariant measure. Let $x \in (0, 1]$ and (B_n) be the cylinders shrinking to x . Then we have the following trichotomy:

- If x is a periodic point, then $N_{B_n}^\gamma$ converges in law towards a compound fractional Poisson process CFPP $_\alpha$ with an extremal index $\theta \in (0, 1)$ depending only on x (see Theorem 2.5, p.11).
- If x is a preimage of 0, then $N_{B_n}^\gamma$ converges in law towards a point process N_x depending on the point x (see Theorem 2.7, p.12).
- Otherwise, $N_{B_n}^\gamma$ converges in law towards the fractional Poisson process FPP $_\alpha$ (see Theorem 2.4, p.11).

This theorem generalizes Theorem A. As in that theorem, the parameter γ is determined by α , which in turn is determined by p . In the case of finite measure, *i.e.*, $p < 1$, prior results [FFTV16, Zwe19] have shown that two types of limiting behavior emerge: a compound Poisson process appears around periodic points, while a standard Poisson process describes almost all other points except for the fixed point at 0. This point, being indifferent, is distinguished by an extremal index $\theta = 0$, producing an infinite cluster. Interestingly, for shrinking neighborhoods around 0, a different scaling allows the hitting time distribution to converge to the exponential distribution. However, for $p > 1$, the limiting distribution deviates from the exponential law [Zwe08]. Notably, unlike the finite measure case, as established in Theorem C, preimages of 0 give rise to limit point processes that do not align with the expected fractional Poisson process FPP $_\alpha$. Instead, these processes are obtained through thinning and rescaling of a specific renewal process.

Finally, in the “barely infinite case” ($p = 1$), we prove a similar dichotomy to that found in the finite-measure case. Our approach parallels the method for $p > 1$, with the essential distinction that the preimages of

0 demand specific handling, differing from that of other points. Here, however, we benefit from the persistence of the exponential distribution as the limiting law, which continues to apply for neighborhoods around 0 that are scaled down [CG93].

Theorem D. *Consider the map (1.1) with $p = 1$ with its (infinite) Let $x \in (0, 1]$ and (B_n) be the cylinders shrinking to x . Then we have the following dichotomy:*

- *If x is a periodic point, then $N_{B_n}^\gamma$ converges in law towards a compound Poisson point process with an extremal index $\theta \in (0, 1)$ depending only on x (see Theorem 2.5, p.11).*
- *If x is not a periodic point, then $N_{B_n}^\gamma$ converges in law towards the fractional Poisson process (see Theorems 2.4 p.11 and 2.9 p.13).*

This paper is structured as follows: Section 2 states our main results, Section 3 provides proofs of the abstract theorems relating hitting and return REPPs and establishes convergence to fractional and compound fractional Poisson processes, Section 4 focuses on convergence for neighborhoods of all points for the map (1.1) and Section 5 discusses related work and future research directions.

2 Statement of results

2.1 Preliminaries

Let (X, \mathcal{B}, μ, T) be a measure-theoretic dynamical system. This means that (X, \mathcal{B}, μ) is a measure space and the self-map $T : X \rightarrow X$ leaves the measure μ invariant (*i.e.*, the push forward $T_\# \mu$ of μ by T is equal to μ). Assume that μ is σ -finite and $\mu(X) = +\infty$. The transfer operator $\widehat{T} : L^1(\mu) \rightarrow L^1(\mu)$ of the system is defined via the following identity : $\forall f \in L^1(\mu), \forall g \in L^\infty(\mu), \int f \cdot (g \circ T) d\mu = \int (\widehat{T}f) \cdot g d\mu$. We say that (X, \mathcal{B}, μ, T) is a ‘‘conservative ergodic measure preserving transformation’’ (CEMPT for short) if $\sum_{k \geq 0} \widehat{T}^k u = +\infty$ μ - a.e. for all $u \in L^1_+(\mu) := \{u \in L^1(\mu) \mid u \geq 0, \int u d\mu > 0\}$ or, equivalently, if this is true for all $u \in \mathcal{D}(\mu) := \{u \in L^1(\mu) \mid u \geq 0, \int u d\mu = 1\}$ (see [Aar97, Propostion 1.3.2]).

For $A \in \mathcal{B}$ and $x \in X$, let $r_A(x) = r_A^{(1)}(x) = \inf\{n \geq 1 \mid T^n x \in A\}$ be the first time the orbits of x hits A , and for $k \geq 1$, define inductively the $(k + 1)$ -th return time to A , namely $r_A^{(k+1)}(x) := \inf\{n > r_A^{(k)}(x) \mid T^n x \in A\}$, with the convention that $r_A^{(k)}(x) = +\infty$ if $A = \emptyset$. If (X, \mathcal{B}, μ, T) is a CEMPT and $\mu(A) > 0$, then $r_A^{(k)}$ is finite μ -almost everywhere, for each $k \geq 1$.

Remark 2.1. *It is also possible to work with inter-arrival times. Set $r_A^{\{1\}} := r_A = r_A^{(1)}$ and $r_A^{\{k+1\}} = r_A \circ T^{r_A^{(k)}}$ for $k \geq 1$. By construction, we have $r_A^{\{k\}} = r_A^{(k)} - r_A^{(k-1)}$ with the convention that $r_A^{(0)} = 0$. To distinguish between inter-arrival and return times, we use the super-scripts $\{k\}$ and (k) , respectively. Again, if (X, \mathcal{B}, μ, T) is a CEMPT and $\mu(A) > 0$, then $r_A^{\{k\}}$ is finite for every $k \geq 1$, μ -almost everywhere.*

Remark 2.2. *Sometimes, we will consider return times on an induced system. For example, if $A \subset Y \subset X$, we will write $r_A^{Y, (k)}$ for the k -th return time to A in the induced dynamical system $(Y, \mathcal{B} \cap Y, T_Y, \mu_Y)$ where $T_Y(x) := T^{r_Y(x)}(x)$ for $x \in X$ and $\mu_Y = \mu(\cdot \cap Y)/\mu(Y)$. In particular, on Y ,*

$$r_A = \sum_{k=0}^{r_A^Y - 1} r_Y \circ T_Y^k = r_Y^{(r_A^Y)}.$$

We are now able to define the objects that we are going to study along this article.

Definition 2.1 (Process of hitting/return times). For $A \in \mathcal{B}$, let

$$\Phi_A := (r_A^{(1)}, r_A^{(2)}, \dots).$$

Furthermore, for every $d \geq 1$, we will write $\Phi_A^{[d]}$ for $(r_A^{(1)}, r_A^{(2)}, \dots, r_A^{(d)})$.

Remark 2.3. *For all $A \in \mathcal{B}$ and $x \in X$, $\Phi_A(x) \in (\overline{\mathbb{R}}_+)^{\mathbb{N}}$ where $\overline{\mathbb{R}}_+ := [0, +\infty]$. If $\mu(A) > 0$ and (X, \mathcal{B}, μ, T) is a CEMPT, then $\Phi_A \in (\overline{\mathbb{R}}_+)^{\mathbb{N}}$ μ -almost everywhere. However, it is more convenient to see it as a function taking values in the compact space $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$ (see [Zwe16] for instance). On $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$ one can take the product metric $d((x_n)_{n \geq 1}, (y_n)_{n \geq 1}) = \sum_{n \geq 0} d_1(x_n, y_n)/(2^n(1 + d_1(x_n, y_n)))$ where $d_1(s, t) := |e^{-s} - e^{-t}|$ is a standard distance on $\overline{\mathbb{R}}_+$. Furthermore, as we consider successive return times, Φ_A is taking values inside the compact subset \mathcal{W} of $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$ where*

$$\mathcal{W} := \left\{ (\phi^{(i)})_{i \geq 1} \in (\overline{\mathbb{R}}_+)^{\mathbb{N}} \mid \forall i \geq 0, \phi^{(i)} \leq \phi^{(i+1)} \right\}.$$

For a non-decreasing function $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and a process $\Phi = (\phi_k)_{k \geq 1}$ taking values in $(\overline{\mathbb{R}_+})^{\mathbb{N}}$, we write $f(\Phi)$ the process $(f(\phi_k))_{k \geq 1}$ with the convention that $f(+\infty) = +\infty$. Note that in this case, $f(\mathcal{W}) \subseteq \mathcal{W}$.

Definition 2.2 (Rare Event Point Process). For a set $A \in \mathcal{B}$ and a function $\gamma : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, the Rare Event Point Process (REPP) is defined by

$$N_A^\gamma := \sum_{k \geq 1} \delta_{\gamma(\mu(A)) r_A^{(k)}}.$$

For a CEMPT, if $\mu(A) > 0$, N_A^γ is well defined for almost every point (all the points such that $r_A^{(k)} < +\infty$ for all $k \geq 1$ or equivalently when $\Phi_A \in (\mathbb{R}_+)^{\mathbb{N}}$) and belongs to the set $\mathcal{M}_{\text{atom}}^{\text{Rad}}(\mathbb{R}_+)$ of Radon atomic measures on \mathbb{R}_+ . $\mathcal{M}_{\text{atom}}^{\text{Rad}}(\mathbb{R}_+)$ is endowed with the topology of vague convergence.

When $\mu(A) < +\infty$ and N_A^γ is treated as a random variable on the probability space $(A, \mathcal{B} \cap A, \mu_A)$, we refer to it as the return REPP. Conversely, when N_A^γ is considered as a random variable on (X, \mathcal{B}, ν) , where ν is a probability measure absolutely continuous with respect to μ , it is termed the hitting REPP. For a fixed set A , the hitting REPP depends on the specific choice of the probability measure ν . However, as our interest lies in the behavior of N_A^γ as $\mu(A) \rightarrow 0$, [Zwe07b, Corollary 6] ensures that any limiting distribution, if it exists, is independent of ν . This result justifies the use of the term ‘‘hitting REPP’’ without dependence on the particular choice of ν .

For a CEMPT preserving a σ -finite measure μ with infinite mass, it is well-established that a direct analogue of the Birkhoff theorem is unattainable, as no normalization exists such that the time average along an orbit converges almost surely [Aar97, Theorem 2.4.1]. Nonetheless, many such systems exhibit a related property, which provides insight into their asymptotic behavior.

Definition 2.3 (Pointwise dual ergodicity). A CEMPT (X, \mathcal{B}, μ, T) is said to be pointwise dual ergodic (PDE) if there exists a sequence $(a_n)_{n \in \mathbb{N}}$ such that

$$\frac{1}{a_n} \sum_{k=0}^{n-1} \widehat{T}^k u \xrightarrow[n \rightarrow +\infty]{\mu\text{-a.e.}} \int u \, d\mu, \quad \forall u \in L^1(\mu). \quad (2.1)$$

In this case $(a_n)_{n \in \mathbb{N}}$ is called a normalizing sequence for (X, \mathcal{B}, μ, T) .

For instance, examples include the Boole map [Aar97, §3.7], interval maps with a finite number of indifferent fixed points [Zwe00] or null recurrent Markov shifts [Sar01]. In fact, the PDE property is equivalent to the existence of a uniform subset on which the convergence is stronger. Such sets are of paramount importance in the study of quantitative recurrence.

Definition 2.4 (Uniform set). A set $Y \in \mathcal{B}$ with $\mu(Y) > 0$ is said to be f -uniform for $f \in L^1(\mu)$ if there exists a sequence $(a_n)_{n \geq 0}$ such that

$$\frac{1}{a_n} \sum_{k=0}^{n-1} \widehat{T}^k f \xrightarrow[n \rightarrow +\infty]{L^\infty(\mu_Y)} \int f \, d\mu.$$

We say that Y is uniform if it is f -uniform for some $f \in L^1(\mu)$.

The existence of uniform sets from the PDE property is immediate by Egorov’s theorem. The proof of the reciprocal can be found in [Aar97, Proposition 3.7.5].

Additionally, in order to get convergence results, we usually require more information on the normalizing sequence through regular variation properties. A measurable function $a : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is said to be regularly varying of index $\alpha \in \mathbb{R}$ at infinity if, for all $y \in \mathbb{R}_+$,

$$\lim_{x \rightarrow +\infty} \frac{a(xy)}{a(x)} = y^\alpha.$$

The notion of regular variation at infinity can be generalized for sequences $(u_n)_{n \geq 0}$ by looking at the function $u : x \mapsto u_{\lfloor x \rfloor}$. In both cases, we will write $a \in \text{RV}(\alpha)$ or $(u_n)_{n \geq 0} \in \text{RV}(\alpha)$. We also say that a function $b : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is regularly varying of index α at 0 if $x \mapsto b(1/x)$ is regularly varying of parameter α at infinity. We write $\text{RV}_{0+}(\alpha)$ the set of such functions or simply $\text{RV}(\alpha)$ when the limit 0 or $+\infty$ is clear from the context.

Given a normalizing sequence $(a(n))_{n \geq 0} \in \text{RV}(\alpha)$, its asymptotic inverse b is defined by the property $b(a(s)) \sim a(b(s)) \sim s$ (see [BGT89] for more on regular variation). Furthermore, we define the scaling function

$$\gamma : s \mapsto \frac{1}{b(1/s)}, \quad \forall s > 0. \quad (2.2)$$

If $a \in \text{RV}(\alpha)$, then $\gamma \in \text{RV}_{0+}(1/\alpha)$. The function γ will be used to scale return times.

We end this section with several notations.

Notations. We write $\xrightarrow{\mu}$ for the convergence in law under the law μ . When the sequence of random variables is defined on different probability spaces with probability measures μ_n , we will write $\xrightarrow{\mu_n}$ for the convergence in law. For example, if X_n is a sequence of random variables defined on $(\Omega_n, \mathcal{B}_n, \mu_n)$ and X is another random variable, $X_n \xrightarrow{\mu_n} X$ means that $(X_n)_{\#}\mu_n$ converges weakly towards the law of X . Finally, when the measure μ is infinite, we cannot draw random variables from it. However, we can still set a convergence, that we will write $\xrightarrow{\mathcal{L}(\mu)}$ if we have $\xrightarrow{\nu}$ for every probability ν that is absolutely continuous with respect to μ . In this case, we say that we have strong convergence in law (see [Aar97, §3.6] for more details).

Notations. Every set equality is understood as an equality up to a set of 0 measure. Furthermore, we use \sqcup for a disjoint union. We will write $(a_n)_{n \geq 0}$ or $(a(n))_{n \geq 0}$ for sequence of real numbers, depending on the context.

We are now prepared to formally present our results. Section 2.2 is devoted to general, abstract findings on quantitative recurrence for pointwise dual ergodic CEMPTs. Section 2.3 examines a specific family of maps with an indifferent fixed point, where we determine the limiting behavior of shrinking targets around every point in the phase space.

2.2 Abstract results

2.2.1 The general relationship between hitting and return REPPs

We say that a sequence $(B_n)_{n \geq 0} \in \mathcal{B}^{\mathbb{N}}$ is a sequence of asymptotically rare events with respect to a measure μ if $\mu(B_n) \xrightarrow{n \rightarrow +\infty} 0$. From now on, we shall consider that asymptotically rare events fulfill the following hypothesis.

(A0) $_{\alpha}$ There exists a uniform set Y with normalizing sequence $(a_n)_{n \geq 0} \in \text{RV}(\alpha)$ such that $B_n \subset Y$ for all $n \geq 0$.

We start by showing that, if (A0) $_{\alpha}$ is satisfied, then the convergence of the hitting REPP implies the convergence of the return REPP and reciprocally, each limit being determined by the other. In particular it generalizes [RZ20, Theorem 4.3] where the authors only consider the first hitting or return.

Theorem 2.1. *Let (X, \mathcal{B}, μ, T) be a PDE CEMPT with $\mu(X) = +\infty$. Let $(B_n)_{n \geq 0}$ be a sequence of asymptotically rare events satisfying (A0) $_{\alpha}$. Let $\Psi, \tilde{\Psi}$ be stochastic processes in $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$. Then, we have*

$$\gamma(\mu(E_n)) \Phi_{E_n} \xrightarrow{n \rightarrow +\infty} \tilde{\Psi} \quad \text{if and only if} \quad \gamma(\mu(E_n)) \Phi_{E_n} \xrightarrow{n \rightarrow +\infty} \Psi.$$

Moreover, the distributions of Ψ and $\tilde{\Psi}$ uniquely determine each other in the following way. For all $d \geq 1$, denoting $F^{[d]}$ (respectively $\tilde{F}^{[d]}$) the distribution function of the d first coordinates of Ψ (respectively $\tilde{\Psi}$), we have, for all $0 \leq t_1 \leq \dots \leq t_d$,

$$F^{[d]}(t_1, \dots, t_d) = \alpha \int_0^{t_1} \left(\tilde{F}^{[d-1]}(t_2 - t_1 + x, \dots, t_d - t_1 + x) - \tilde{F}^{[d]}(x, t_2 - t_1 + x, \dots, t_d - t_1 + x) \right) (t_1 - x)^{\alpha-1} dx, \quad (2.3)$$

with the convention $\tilde{F}^{[0]} = 1$.

Remark 2.4. *Since, for all $d \geq 1$, $F^{[d]} \leq 1$ and $\alpha \int_0^{t_1} (t_1 - x)^{\alpha-1} dx = t_1^{\alpha}$, for all $t_2, \dots, t_d = t_1$, we necessarily have*

$$\lim_{x \rightarrow +\infty} \tilde{F}^{[d]}(x, \dots, x) = \lim_{x \rightarrow +\infty} F^{[d-1]}(x, \dots, x) = 1,$$

justifying the fact that $\tilde{\Psi}$ belongs to $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$ almost surely.

Remark 2.5. *Theorem 2.1 considers stochastic processes of hitting/return times taking values in $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$. However, point processes are more natural in our context. If the limits Ψ and $\tilde{\Psi}$ belong to a natural subset of $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$, it will be enough to ensure the result for REPPs. Let*

$$\mathcal{W}' := \left\{ (\phi^{(i)})_{i \geq 0} \in (\overline{\mathbb{R}}_+)^{\mathbb{N}} \mid \phi^{(i)} \leq \phi^{(i+1)} \text{ and } \lim \phi^{(i)} = +\infty \right\}.$$

and define the following map

$$\Xi : \begin{cases} \mathcal{W} & \longrightarrow \mathcal{M}_{\text{atom}}^{\text{Rad}}(\mathbb{R}_+) \\ \Psi = (\psi^{(i)})_{i \geq 1} & \longmapsto \begin{cases} \sum_{i \geq 1} \delta_{\psi^{(i)}} & \text{if } \Psi \in \mathcal{W}' \\ 0 & \text{otherwise.} \end{cases} \end{cases}$$

Then, Ξ is continuous on \mathcal{W}' (see also [Zwe22, Remark 3.5]). Thus, by the extended continuous mapping theorem, we immediately get the following corollary.

Corollary 2.1. *Under the same assumptions as in Theorem 2.1. Assume furthermore that $\Psi, \tilde{\Psi} \in \mathcal{W}'$ almost surely. Then, for $N = \Xi(\Psi)$ and $\tilde{N} = \Xi(\tilde{\Psi})$, we have*

$$N_{E_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} \tilde{N} \quad \text{and} \quad N_{E_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} N.$$

The law of N determines the law of \tilde{N} and reciprocally.

Remark 2.6. *Here, and in the following, we have chosen to change the renormalization by taking γ . We could have chosen to keep the normalization by the measure, but instead change the return random variable. As we assumed $\alpha \in (0, 1]$, both result are equivalent (see [BZ01, Lemma 1] for example). The main difference is when $\alpha = 0$ (see [RZ20, Propostion 4.1]) but we do not consider this case here.*

With the choice of scaling the stochastic process instead of scaling the measures, it gives the following theorem, equivalent to Theorem 2.1.

Theorem 2.2. *Under the same assumptions as in Theorem 2.1, we have*

$$\mu(E_n) a(\Phi_{E_n}) \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} \tilde{\Phi} \quad \text{if and only if} \quad \mu(E_n) a(\Phi_{E_n}) \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \Phi.$$

Furthermore, the laws of Φ and $\tilde{\Phi}$ determine one another through the distribution function of their marginals by a change of variable in (2.3).

In particular, Theorem 2.1 gives back the equivalence result for the first hitting and return times if we take the projection on the first component.

Corollary 2.2. [RZ20, Theorem 4.2] *Under the same assumptions as in Theorem 2.1, we have for R, \tilde{R} random variables in \mathbb{R}_+ ,*

$$\gamma(\mu(B_n)) r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} R \quad \text{if and only if} \quad \gamma(\mu(B_n)) r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \tilde{R}.$$

If convergence takes place, then

$$\mathbb{P}(R \leq t) = \alpha \int_0^t (1 - \mathbb{P}(\tilde{R} \leq u))(t - u)^{\alpha-1} du, \quad \forall t \geq 0. \quad (2.4)$$

In terms of the Laplace transform, this equation can be rewritten as

$$\mathbb{E}[e^{-sR}] = \frac{\Gamma(1 + \alpha)}{s^\alpha} \left(1 - \mathbb{E}[e^{-s\tilde{R}}]\right), \quad \forall s > 0, \quad (2.5)$$

where $\Gamma : z \mapsto \int_0^{+\infty} t^{z-1} e^{-t} dt$ is the standard Gamma function.

More generally, some properties of a point process N on \mathbb{R}_+ are defined throughout their evolution equation. In our case, (2.3) can be used to find the evolution equation of the limit point process N from the one of the point process \tilde{N} . For a point process N and $d \geq 1$, let $P_N(d, t) := \mathbb{P}(N[0, t] = d)$. We see $P_N(d, \cdot)$ as a function from \mathbb{R}_+ to $[0, 1]$.

Before stating the results, we need to recall some basics notions of fractional calculus. For every $\beta > 0$, we define the Riemann-Liouville integral I^β as follows. For every Riemann integrable f and $t \in \mathbb{R}$

$$(I^\beta f)(t) := (I_{0+}^\beta f)(t) := \frac{1}{\Gamma(\beta)} \int_0^t f(x)(t - x)^{\beta-1} dx. \quad (2.6)$$

Associated to this integral, we can go backwards and define the Caputo derivative for differentiable functions f by

$$\text{Cap} D^\beta f(t) := (I^{1-\beta} f')(t), \quad \forall t \in \mathbb{R}, \quad 0 < \beta < 1. \quad (2.7)$$

Then, Theorem 2.1 implies the following result.

Corollary 2.3. *Assume the same hypothesis as in Corollary 2.1. Then, we have*

$$\text{Cap} D^\alpha P_N(d, \cdot) = \Gamma(1 + \alpha)(P_{\tilde{N}}(d - 1, \cdot) - P_{\tilde{N}}(d, \cdot)), \quad \forall d \geq 0, \quad (2.8)$$

with the convention that $P_{\tilde{N}}(-1, t) := 0$ for all $t \geq 0$ and $P_N(0, 0) = 1$.

2.2.2 The fractional Poisson process as a common limit law

We now focus on characterizing point processes that will serve as potential limits for hitting and return REPP. As a preliminary step, we first revisit key concepts from renewal theory and present some properties of the Fractional Poisson Process.

Definition 2.5 (Renewal point process). Let W be a non negative random variable such that $\mathbb{P}(W > 0) > 0$, and $(W_i)_{i \geq 1}$ i.i.d. random variables having the same law as W . The renewal point process with waiting times $(W_i)_{i \geq 1}$ is defined by

$$\text{RPP}(W) \stackrel{\text{law}}{=} \sum_{i=1}^{+\infty} \delta_{T_i},$$

where $T_{i+1} - T_i = W_{i+1}$ for all $i \geq 0$, with the convention $T_0 = 0$. Note that $\text{RPP}(W) \in \mathcal{M}_{\text{atom}}^{\text{Rad}}(\mathbb{R}_+)$.

Definition 2.6 (Fractional Poisson Process). The Fractional Poisson Process $\text{FPP}_\alpha(\lambda)$ of parameters $\alpha \in (0, 1]$ and $\lambda > 0$ is the renewal point process $\text{RPP}(H_\alpha(\lambda))$ where $H_\alpha(\lambda)$ is a Mittag-Leffler law of the first type characterized by its Laplace transform

$$\mathbb{E}[e^{-sH_\alpha(\lambda)}] := \frac{\lambda}{\lambda + s^\alpha}, \quad \forall s \geq 0.$$

Remark 2.7.

- (i) When $\lambda = 1$, we will simply write FPP_α and H_α .
- (ii) We can easily see that $H_\alpha(\lambda) \stackrel{\text{law}}{=} \frac{1}{\lambda^{1/\alpha}} H_\alpha$.
- (iii) If $\alpha = 1$, then $H_1(\lambda)$ is the exponential law of parameter λ and $\text{FPP}_1(\lambda)$ is the homogeneous Poisson point process of parameter λ .
- (iv) The first type Mittag-Leffler law $H_\alpha(\lambda)$ should not be confused with the second type Mittag-Leffler law Y_α commonly called Mittag-Leffler law in the context of infinite measure dynamical systems (see [Aar97, §3.6] for a definition).¹
- (v) When $0 < \alpha < 1$, $\mathbb{E}[H_\alpha(\lambda)] = +\infty$.

Remark 2.8. The fractional Poisson process $\text{FPP}_\alpha(\lambda)$ was introduced as a fractional generalization of the standard Poisson process through its Kolmogorov-Feller equation [Las03]. In particular, a process N having the law of $\text{FPP}_\alpha(\lambda)$ is characterized by the independence of its waiting times and the generalized Kolmogorov-Feller evolution equation

$$\text{Cap} D^\alpha P_N(d, \cdot) = \lambda(P_N(d-1, \cdot) - P_N(d, \cdot)), \quad \forall d \geq 0, \quad (2.9)$$

with the convention $P_N(-1, \cdot) = 0$ (see [Las03, Equation (19)] or [MS19, Equation (7.10) p. 207]).

For the majority of well-behaved rare events, we anticipate that the hitting REPP and return REPP converge to the same limiting point process. Consequently, the point processes of primary interest are the fixed points of (2.3). Here, we see that (2.9) is exactly (2.8) when N and \tilde{N} have the same law and $\lambda = \Gamma(1 + \alpha)$ pointing $\text{FPP}_\alpha(\Gamma(1 + \alpha))$ as a potential fixed point. The following proposition confirms that this is indeed a fixed point and further ensures its uniqueness.

Proposition 2.4. The fractional Poisson process $\text{FPP}_\alpha(\Gamma(1 + \alpha))$ is the only process such that the distribution functions of its finite-dimensional marginals are the fixed points of (2.3).

Remark 2.9. In Proposition 2.4, we took a slight liberty in stating that the fractional Poisson process is the fixed point of the equation. It is in fact $\Phi_{\text{FPP}_\alpha(\Gamma(1+\alpha))} := (\phi^{(i)})_{i \geq 1}$ taking values in $\overline{\mathbb{R}}_+$ where for every $i \geq 1$, $\phi^{(i)} = \sum_{k=1}^i X_k$ where $(X_k)_{k \geq 1}$ are i.i.d. with common law that of $H_\alpha(\Gamma(1 + \alpha))$. However, if

$$\mathcal{W}'' := \{(\phi^{(i)})_{i \geq 0} \in (\mathbb{R}_+)^{\mathbb{N}} \mid \forall i \geq 0, \phi^{(i)} < \phi^{(i+1)} \text{ and } \lim_{i \rightarrow +\infty} \phi^{(i)} = +\infty\},$$

we then have $\Phi_{\text{FPP}_\alpha(\Gamma(1+\alpha))} \in \mathcal{W}''$ almost surely, so the law of $\Phi_{\text{FPP}_\alpha(\Gamma(1+\alpha))}$ and $\text{FPP}_\alpha(\Gamma(1 + \alpha))$ are uniquely defined by one another. In the sequel, we will not distinguish between the point process, which takes values in $\mathcal{M}_{\text{atom}}^{\text{Rad}}(\mathbb{R}_+)$, and the stochastic process, which takes values in $(\mathbb{R}_+)^{\mathbb{N}}$, hence both will be denoted by FPP_α .

¹Both are called Mittag-Leffler because they are defined from the Mittag-Leffler function $E_\alpha(z) := \sum_{k=0}^{+\infty} \frac{z^k}{\Gamma(\alpha k + 1)}$, but we have $\mathbb{E}[e^{zY_\alpha}] = E_\alpha(\Gamma(1 + \alpha)z)$ for $z \in \mathbb{R}$, whereas $\mathbb{P}(H_\alpha > t) = E_\alpha(-t^\alpha)$ for $t \geq 0$.

While the Fractional Point Process will act as the limit in most cases, other point processes naturally emerge. To accommodate situations where the first waiting time differs from subsequent ones or the mass associated to each point is random, we introduce delayed renewal point processes and compound point processes.

Definition 2.7 (Delayed renewal point process). Let V, W be two non negative random variables such that $\mathbb{P}(V > 0), \mathbb{P}(W > 0) > 0$, and let $(W_i)_{i \geq 1}$ be i.i.d. random variables with the same law as W . The delayed renewal point process with delay V and waiting times $(W_i)_{i \geq 1}$ is defined by

$$\text{DRPP}(V, W) \stackrel{\text{law}}{=} \sum_{i=1}^{+\infty} \delta_{T_i},$$

where $T_{i+1} - T_i = W_{i+1}$ for all $i \geq 1$ and $T_1 = V$. Note that $\text{DRPP}(V, W) \in \mathcal{M}_{\text{atom}}^{\text{Rad}}(\mathbb{R}_+)$.

Definition 2.8 (Compound point process). For a simple point process $P = \sum_{i=1}^{+\infty} \delta_{T_i}$, we define the associated compound point process $c(P)(\pi)$ of multiplicity π (where π is a probability distribution on \mathbb{N}) as

$$c(P)(\pi) \stackrel{\text{law}}{=} \sum_{i=1}^{+\infty} X_i \delta_{T_i},$$

where $(X_i)_{i \geq 1}$ are i.i.d. random variables distributed according to π and independent of $(T_i)_{i \geq 1}$.

Remark 2.10. *This is not the standard approach to defining a compound process. Here, we utilize the structure of \mathbb{R} and the fact that the multiplicity π takes integer values. However, compound point processes are typically constructed in a more general framework, particularly the compound Poisson process, as described in detail in [LP18, Chapter 15].*

Definition 2.9 (Compound Fractional Poisson Process). We define $\text{CFPP}_\alpha(\lambda, \pi)$ as the compound process with multiplicity π associated to the fractional Poisson process $\text{FPP}_\alpha(\lambda)$.

Remark 2.11. *In our setting, we do not impose the waiting time of a renewal process to be positive almost surely, so compound processes can also be characterized as renewal processes. For instance, let $H_\alpha(\lambda)$ be the waiting time of the fractional Poisson process. Let $\theta \in (0, 1]$ and let $W_{\alpha, \theta}(\lambda)$ be a non-negative random variable with distribution function*

$$\mathbb{P}(W_{\alpha, \theta}(\lambda) \leq t) = 1 - \theta + \theta \mathbb{P}(H_\alpha(\lambda) \leq t). \quad (2.10)$$

It means that with probability $1 - \theta$, $W_{\alpha, \theta}(\lambda) = 0$ and $(W_{\alpha, \theta}(\lambda) | W_{\alpha, \theta}(\lambda) > 0)$ has the same law as $H_\alpha(\lambda)$. Then,

$$\text{DRPP}(H_\alpha(\lambda), W_{\alpha, \theta}(\lambda)) \stackrel{\text{law}}{=} \text{CFPP}_\alpha(\lambda, \text{Geo}(\theta)),$$

where $\text{Geo}(\theta)$ is the positive geometric law, i.e. if $Y \stackrel{\text{law}}{=} \text{Geo}(\theta)$, for all $k \geq 1$, $\mathbb{P}(Y = k) = \theta(1 - \theta)^{k-1}$.

To prove convergence of the hitting REPP and the return REPPs, further conditions on the sequence of targets $(B_n)_{n \geq 0}$ are necessary.

Assumptions $(A)_\alpha$.

A sequence $(B_n) \in \mathcal{B}^{\mathbb{N}}$ of asymptotically rare events satisfies $(A)_\alpha$ if it satisfies $(A0)_\alpha$ and the following conditions :

(A1) $_\alpha$ For every $n \geq 1$, we can write $B_n = U(B_n) \sqcup Q(B_n)$, and $\lim_{n \rightarrow +\infty} \mu(Q(B_n))/\mu(B_n) = \theta \in (0, 1]$ (the extremal index).

(A2) $_\alpha$ There exists a sequence of measurable functions $\tau_n : B_n \rightarrow \mathbb{N}$ and a compact subset \mathcal{U} of $L^1(\mu)$ such that

$$\widehat{T^{\tau_n}} \left(\frac{\mathbf{1}_{Q(B_n)}}{\mu(Q(B_n))} \right) \in \mathcal{U}, \forall n \geq 1.$$

(A3) $_\alpha$ The sequence $(\tau_n)_{n \geq 0}$ satisfies $\gamma(\mu(B_n)) \tau_n \xrightarrow[n \rightarrow +\infty]{\mu B_n} 0$, where γ is defined from $(a_n)_{n \geq 0}$ by (2.2).

(A4) $_\alpha$ The sequence $(Q(B_n))_{n \geq 0}$ is such that $\mu_{Q(B_n)}(r_{B_n} < \tau_n) \xrightarrow[n \rightarrow +\infty]{} 0$.

Furthermore, if $U(B_n) \neq \emptyset$, we have

(A5) $_\alpha$ The sequence $(U(B_n))_{n \geq 0}$ is such that $\mu_{U(B_n)}(r_{B_n} > \tau_n) \xrightarrow[n \rightarrow +\infty]{} 0$

(A6) $_{\alpha}$ We have the following limit

$$\left\| \widehat{T}_{B_n} \left(\frac{\mathbf{1}_{U(B_n)}}{\mu(U(B_n))} \right) - \frac{1}{\mu(B_n)} \mathbf{1}_{B_n} \right\|_{L^{\infty}(\mu_{B_n})} \xrightarrow{n \rightarrow +\infty} 0.$$

Let us analyze these conditions. The assumptions (A) $_{\alpha}$ are identical to those outlined in [RZ20, Theorem 7.2], where only the first return is considered. We show that these assumptions are sufficiently robust to establish the convergence of the point process as well. Assumption (A0) $_{\alpha}$ ensures that Theorem 2.1. can be applied effectively.

Condition (A1) $_{\alpha}$ serves two purposes: it identifies points that are likely to form clusters within $U(B_n)$ and points that escape the target $Q(B_n)$, while also guaranteeing that the extremal index θ is well-defined.

Condition (A2) $_{\alpha}$ specifies an appropriate waiting time τ_n such that the density within the target set transforms into a desirable structure under the dynamics. However, τ_n must remain asymptotically negligible, a requirement addressed by (A3) $_{\alpha}$.

Cluster compatibility is handled through (A4) $_{\alpha}$ and (A5) $_{\alpha}$, ensuring that τ_n aligns with clustering behavior.

Finally, Condition (A6) $_{\alpha}$ explains the geometric distribution of multiplicities. While this condition could be generalized to allow for other multiplicity distributions, we retain it as stated, since the geometric law is the only distribution arising in our examples.

We can now state our main theorem, which establishes that Assumptions (A) $_{\alpha}$ are sufficient conditions for convergence to the Compound Poisson Process.

Theorem 2.3. *Let (X, \mathcal{B}, μ, T) be a PDE CEMPT with $\mu(X) = +\infty$. Let $(B_n)_{n \geq 0}$ be a sequence of asymptotically rare events satisfying (A) $_{\alpha}$. Then,*

$$N_{B_n}^{\gamma} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{CFPP}_{\alpha}(\theta\Gamma(1 + \alpha), \text{Geo}(\theta))$$

and

$$N_{B_n}^{\gamma} \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1 + \alpha))).$$

In particular, if $\theta = 1$, we have

$$N_{B_n}^{\gamma} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{FPP}_{\alpha}(\Gamma(1 + \alpha)) \quad \text{and} \quad N_{B_n}^{\gamma} \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{FPP}_{\alpha}(\Gamma(1 + \alpha)).$$

The primary goal in the following is to apply Theorem 2.3 to interval maps with a single indifferent fixed point, using shrinking cylinders as rare events. However, the theorem is broadly applicable to a wider range of maps and asymptotically rare events, making it a versatile tool for many other contexts.

2.3 Rare event point processes in maps with an indifferent fixed point

We now focus on the specific case of interval maps with an indifferent fixed point, as defined by (1.1). Recall that $x \in [0, 1]$ and

$$T_p(x) = \begin{cases} x + 2^p x^{p+1}, & 0 \leq x < 1/2, \\ 2x - 1, & 1/2 \leq x \leq 1. \end{cases}$$

We define $\alpha := p^{-1}$. Once p is fixed, for simplicity, we omit its dependence in the index and we will simply write T instead of T_p . Let T_1 and T_2 be the two diffeomorphic branches of T (more precisely their extensions to $[0, 1/2]$ and $[1/2, 1]$ respectively) and set $c_n := T_1^{-n} 1$. In particular, $c_0 = 1$ and $c_1 = 1/2$.

For $0 \leq p < 1$, the absolutely continuous invariant measure μ is unique (up to scaling) and normalized to a probability measure. In contrast, for $p \geq 1$, μ remains unique but becomes infinite. Since we are interested by the infinite case, we assume $p \geq 1$ and fix a scaling so that $\mu([1/2, 1]) = 1$. The density $\rho = d\mu/d\text{Leb}$ is the following:

$$\rho(x) = h_0(x) \frac{x}{x - T_1^{-1}x}, \quad x \in [0, 1], \quad (2.11)$$

for some continuous function h_0 bounded away from 0 and $+\infty$. For more details about such dynamical systems, see [LSV99], [You99] or [Alv20, Section 3.5] or [Tha80, Tha83].

The dynamical system $([0, 1], T, \mu)$ is a PDE CEMPT with normalizing sequence $(a_n)_{n \in \mathbb{N}} \in \text{RV}(\alpha)$ and every interval $I = [c, 1]$ with $c > 0$ is uniform (see [Zwe00] for example).

Set $Y := [c_1, c_0] = [1/2, 1]$ and define $\xi := \{[c_{k+1}, c_k], k \geq 0\} = \{Y\} \cup \{Y^c \cap \{r_Y = k\}, n \geq 1\}$ the measurable partition of $[0, 1]$ defined from return times to Y . Then, for every $n \geq 1$, let $\xi_n = \bigvee_{i=0}^{n-1} T^{-i}\xi$. The partition ξ dynamically generates the Borel σ -algebra \mathcal{B} and for a point $x \in [0, 1]$, we write $\xi_n(x)$ the element of ξ_n that contains x . If x is not a preimage of 0, it is easy to see that $\xi_n(x)$ is well defined for every $n \geq 1$.

In this case, $\xi_n(x)$ shrinks to $\{x\}$ as n goes to $+\infty$ and will serve as our asymptotically rare events in the study of the hitting and return REPP.

For the preimages of 0, such a definition for asymptotically rare events is not possible anymore. However, if $T^k x = 0$, the connected component C_x^n of $T^{-k}[0, c_n]$ containing x is a union of elements in ξ_k for all $n \geq 1$. As n goes to $+\infty$, C_x^n shrinks towards x and thus $(C_x^n)_{n \geq 1}$ will serve as our sequence of asymptotically rare events associated to x in this case.

2.3.1 Points that are not preimages of the indifferent fixed point

If x is not a preimage of 0, as determined by the chosen partition ξ , we define the asymptotically rare events as $B_n := \xi_n(x)$ for all $n \geq 1$. This setup results in two distinct behaviors depending on whether x is periodic. If x is not periodic, both the hitting and return REPPs converge to the fractional Poisson process.

Theorem 2.4. *Let $p \geq 1$. Assume $x \in [0, 1]$ is not periodic and not a preimage of 0. For all $n \geq 1$, set $B_n := \xi_n(x)$. Then,*

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{FPP}_\alpha(\Gamma(1 + \alpha)) \quad \text{and} \quad N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{FPP}_\alpha(\Gamma(1 + \alpha)).$$

When x is q -periodic, clusters of hittings appear and thus we get a compound fractional Poisson process in the limit.

Theorem 2.5. *Let $p \geq 1$. Assume $x \in (0, 1]$ is periodic of prime period q . For all $n \geq 1$, set $B_n := \xi_n(x)$. Then,*

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{CFPP}_\alpha(\theta\Gamma(1 + \alpha), \text{Geo}(\theta)) \quad \text{and} \quad N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1 + \alpha))),$$

where the extremal index $\theta = 1 - |(T^q)'(x)|^{-1}$.

The framework closely mirrors that of the finite measure case, with the significant difference being that the Poisson point process is replaced by the fractional Poisson process. Notably, in the barely infinite case where $p = 1$, the limits are instead the standard Poisson point process and the compound Poisson point process, characterized by parameters θ and multiplicity distributed as $\text{Geo}(\theta)$.

2.3.2 The indifferent fixed point and its preimages

If $p > 0$, the neutral fixed point 0 always requires a special treatment, because its extremal index $\theta = 0$ as the neutrality gives a cluster of infinite length on average. In the finite measure context, for shrinking neighborhoods $(B_n)_{n \geq 1}$ of 0, $\mu(B_n)r_{B_n}$ converges almost surely to $+\infty$ when one starts in B_n and to 0 when one starts off B_n . However, the exponential law can be recovered with the scaling $\mu(Q(B_n))$ instead of $\mu(B_n)$, where $Q(B_n) := B_n \setminus T^{-1}B_n$.

Proposition 2.5. [FFTV16, Theorem 2], [Zwe19, Theorem 5.1] *Assume $0 < p < 1$ and let $(B_n)_{n \in \mathbb{N}}$ be a nested sequence of intervals neighborhoods of 0. Then, we have*

$$\mu(Q(B_n))r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mu} \mathcal{E},$$

where \mathcal{E} is an exponential random variable.

This highlights the robustness of the exponential law in the finite measure case. The key idea of [Zwe19, Theorem 5.1] is to relate returns to 0 with returns to $1/2$, the only other preimage of 0. Unlike 0, the point $1/2$ is more favorable for analysis since it is not a fixed point and lies within the common inducing set $Y = [1/2, 1]$. Within this set, standard inducing techniques can be employed to establish the convergence of the return process to the standard homogeneous Poisson point process for shrinking neighborhoods of $1/2$. The result for shrinking neighborhoods of 0 is then derived as a consequence of this approach.

When $p \geq 1$, there are no established techniques to directly address the point $1/2$. However, it is possible to analyze hitting times for neighborhoods of 0 by viewing them as the apex of the tower constructed via the induction $Y = [1/2, 1]$. At this stage, it becomes necessary to distinguish between two scenarios: the ‘‘proper infinite case’’ ($p > 1$) and the ‘‘barely infinite case’’ ($p = 1$).

For $p > 1$, [Zwe08, Theorem 2] demonstrates that a non-degenerate limit can be achieved with an appropriately chosen scaling. However, the resulting limit does not align with the exponential law observed in the finite measure case or the Mittag-Leffler law $H_\alpha(\Gamma(1 + \alpha))$ typical for generic points.

Theorem 2.6. [Zwe08, Theorem 2] Let $p > 1$. Let $B_n := [0, \eta_n]$ such that $\eta_n \xrightarrow[n \rightarrow +\infty]{} 0$. Then,

$$\frac{1}{I(\eta_n)} r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \mathcal{J}_\alpha,$$

where \mathcal{J}_α is characterized by its Laplace transform

$$\mathbb{E} [e^{-s\mathcal{J}_\alpha}] = \frac{1}{e^{-s} + s \int_0^1 y^{-\alpha} e^{-sy} dy}, \quad \forall s \geq 0,$$

and

$$I(\eta_n) := \int_{\eta_n}^1 \frac{dx}{x - T_1^{-1}(x)}.$$

Using Lemma A.1, we can rewrite this result in our framework, using the scaling γ defined in (2.2).

Corollary 2.6. Under the same assumptions as in Theorem 2.6. Then,

$$\gamma(\mu(T_2^{-1}(B_n))) r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \mathfrak{J}_\alpha,$$

where $\mathfrak{J}_\alpha := d_\alpha \mathcal{J}_\alpha$ and $d_\alpha := (\Gamma(1 + \alpha)\Gamma(1 - \alpha))^{-1/\alpha} = (\sin(\pi\alpha)/(\pi\alpha))^{1/\alpha}$.

Using (2.5), we can construct a random variable $\tilde{\mathfrak{J}}_\alpha$ associated to \mathfrak{J}_α . In particular, the Laplace function of $\tilde{\mathfrak{J}}_\alpha$ is the following

$$\mathbb{E} [e^{-s\tilde{\mathfrak{J}}_\alpha}] = 1 - \frac{s^\alpha}{\Gamma(1 + \alpha)} \left(e^{-sd_\alpha} + sd_\alpha \int_0^1 y^{-\alpha} e^{-d_\alpha sy} dy \right)^{-1}, \quad s \geq 0. \quad (2.12)$$

The random variable $\tilde{\mathfrak{J}}_\alpha$ plays a central role in characterizing the limiting point processes obtained for shrinking targets around points that include 0 in their orbit. Specifically, any such limit can be represented as a thinning and rescaling of the renewal point process $\text{RPP}(\tilde{\mathfrak{J}}_\alpha)$. To formalize the concept of thinning and rescaling a point process, we rely on the following definition.

Definition 2.10 (Thinning and rescaling). For a point process $N = \sum_{i=1}^{+\infty} \delta_{T_i}$, $\tau > 0$ and $v > 0$, we call τ -thinning and v -rescaling of N the point process

$$N^{(\tau, v)} \stackrel{\text{law}}{=} \sum_{i=1}^{+\infty} X_i \delta_{(vT_i)},$$

where $(X_i)_{i \geq 1}$ are i.i.d. random variables, independent from N and $X_1 \sim \text{Bernoulli}(\tau)$.

We are now able to state our theorem for shrinking neighborhoods of preimages of 0.

Theorem 2.7. Let $p > 1$. Let $k \geq 0$ and assume $x \in T^{-(k+1)}\{0\}$. Let $B_n := C_x^n$ be the connected component of $T^{-(k+1)}[0, c_n]$ containing x . Then,

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{RPP}(\tilde{\mathfrak{J}}_\alpha)^{(\mathbb{Q}_k(x), \mathbb{Q}_k(x)^{1/\alpha})} \quad \text{and} \quad N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{DRPP}(\mathfrak{J}_\alpha, \tilde{\mathfrak{J}}_\alpha)^{(\mathbb{Q}_k(x), \mathbb{Q}_k(x)^{1/\alpha})},$$

where

$$\mathbb{Q}_k(x) := \frac{\rho(x)}{\rho(1/2)(T^k)'(x)}.$$

Remark 2.12. For every $k \geq 0$, \mathbb{Q}_k is a probability on $T^{-(k+1)}\{0\}$ because ρ is a fixed point of the transfer operator \hat{T} , whence

$$\rho(1/2) = (\hat{T}^k \rho)(1/2) = \sum_{T^k y = \frac{1}{2}} \frac{\rho(y)}{(T^k)'(y)}.$$

Theorem 2.7 reveals that points whose orbits include 0 exhibit markedly different behavior compared to the cases outlined in Theorems 2.4 and 2.5, where the transition from the finite measure setting is more straightforward. Specifically, in the finite measure case, preimages of 0 conform to the standard Poisson process. However, in the infinite measure scenario, the interplay between extended excursions and visits to neighborhoods of 0's preimages disrupts similar outcomes. That said, points located farther from 0 are expected to exhibit a diminishing dependence on 0. This phenomenon is explored further in the next proposition.

Proposition 2.7. For every sequence of $(x_k)_{k \geq 1}$, with $x_k \in T^{-(k+1)}\{0\}$, we have

$$\text{RPP}(\tilde{\mathfrak{J}}_\alpha)^{(\mathbb{Q}_k(x_k), \mathbb{Q}_k(x_k)^{1/\alpha})} \xrightarrow[k \rightarrow +\infty]{} \text{FPP}_\alpha(\Gamma(1 + \alpha)),$$

and

$$\text{DRPP}(\mathfrak{J}_\alpha, \tilde{\mathfrak{J}}_\alpha)^{(\mathbb{Q}_k(x_k), \mathbb{Q}_k(x_k)^{1/\alpha})} \xrightarrow[k \rightarrow +\infty]{} \text{FPP}_\alpha(\Gamma(1 + \alpha)).$$

For periodic points, the longer the period, the closer the extremal index θ approaches 1, resulting in the limiting point process converging more closely to the fractional Poisson process. Similarly, Proposition 2.7 guarantees a parallel trend for preimages of 0: the farther a point lies from the neutral fixed point 0 (in the sense of time iterations needed to hit 0), the more closely its limiting behavior aligns with that of the fractional Poisson process.

2.3.2.1 Barely infinite case $p = 1$. When, $p = 1$, the picture is different. Indeed, for neighborhoods of 0 and a well-chosen scaling, the limit law obtained is again the exponential, as in the finite measure case.

Theorem 2.8. [CG93, Theorem 5], [CI95, Theorem 3.3] Let $p = 1$ and $B_n = [0, c_n]$ for $n \geq 0$. Then,

$$\mathbb{E}_{\mu_Y}[r_{B_n}]^{-1} r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \mathcal{E}$$

where \mathcal{E} is the exponential law.

As for Corollary 2.6 for $p > 1$, we need to make sure that the result can be written in our setting to make the scaling gamma appear. This is ensured by the following lemma.

Lemma 2.8. With the same notation as in Theorem 2.8, we have

$$\gamma(\mu(T_2^{-1}B_n)) \underset{n \rightarrow +\infty}{\sim} \mathbb{E}_{\mu_Y}[r_{B_n}]^{-1}.$$

Using this and applying the same approach as for $p > 1$, we get the following theorem

Theorem 2.9. Let $p = 1$, $k \geq 0$ and $x \in T^{-k+1}\{0\}$. Let $B_n := C_x^n$. Then,

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{PPP}(1) \quad \text{and} \quad N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{PPP}(1).$$

Therefore, in the special case $p = 1$, the preimages of 0 have the same limit behavior as any other non periodic point and thus we only have a dichotomy.

3 Return and Hitting Rare Event Point Processes and the Fractional Poisson Process

3.1 Equivalence between Hitting and Return Point Processes

This part is devoted to the proofs of Theorems 2.1 and 2.2 and their applications.

3.1.1 Proof of Theorems 2.1-2.2

As stated in Remark 2.6, Theorems 2.1 and 2.2 are equivalent since $\alpha \in (0, 1]$. We are going in fact to prove Theorem 2.2 and study the sequence of processes $(\mu(E_n) a(\Phi_{E_n}^{[d]}))_{n \geq 0}$. For that, we introduce some more notation. For $A \in \mathcal{B}$, $u \in L^1(\mu)$, $u \geq 0$, $d \geq 1$ and $t_1, \dots, t_d > 0$, let

$$G_{u,A}^{[d]}(t_1, \dots, t_d) := \mu_u \left(\mu(A) a(\Phi_A^{[d]}) \leq (t_1, \dots, t_d) \right),$$

where μ_u is the measure absolutely continuous with respect to μ with density u . In particular, if $\int u d\mu = 1$, $G_{u,A}^{[d]}$ is the distribution function of the random variable $\mu(A) a(\Phi_A^{[d]})$ under the probability μ_u . We also define

$$\tilde{G}_A^{[d]}(t_1, \dots, t_d) = \mu_A \left(\mu(A) a(\Phi_A^{[d]}) \leq (t_1, \dots, t_d) \right) \quad \forall d \geq 1, t_1, \dots, t_d \geq 0,$$

as the distribution function of $\mu(A) a(\Phi_A^{[d]})$ under μ_A . Before delving into the proof, we first establish the following lemma, an adaptation of [RZ20, Lemma 5.1], which provides a decomposition for multiple return events.

Lemma 3.1. For all $A \in \mathcal{B}$, $n_1, \dots, n_d \in \mathbb{N}$, we have the following decomposition:

$$\left\{ \Phi_A^{[d]} \leq (n_1, \dots, n_d) \right\} = \bigsqcup_{\ell=1}^{n_1} T^{-\ell} \left(A \cap \left\{ \Phi_A^{[d-1]} \leq (n_2 - \ell, \dots, n_d - \ell) \right\} \cap \left\{ \Phi_A^{[d]} \leq (n_1 - \ell, \dots, n_d - \ell) \right\}^c \right).$$

Remark 3.1. In fact, we also have the easier decomposition

$$\left\{ \Phi_A^{[d]} \leq (n_1, \dots, n_d) \right\} = \bigcup_{\ell=1}^{n_1} T^{-\ell} \left(A \cap \left\{ \Phi_A^{[d-1]} \leq (n_2 - \ell, \dots, n_d - \ell) \right\} \right),$$

but we will later take advantage of the disjointness.

Proof (of Lemma 3.1). For every $\ell \geq 1$, $T^{-\ell} \left(A \cap \left\{ \Phi_A^{[d-1]} \leq (n_2 - \ell, \dots, n_d - \ell) \right\} \right) \subset \left\{ \Phi_A^{[d]} \leq (n_1, \dots, n_d) \right\}$, hence we have the inclusion of the right member into the left one.

Now, let $x \in \left\{ \Phi_A^{[d]} \leq (n_1, \dots, n_d) \right\}$. Consider $\ell := \max \{ p \leq n_1 \mid x \in T^{-p} (A \cap \{ \Phi_A^{[d-1]} \leq (n_2 - p, \dots, n_d - p) \}) \}$. The maximum ℓ is well-defined, as the set is non-empty with $r_A(x)$ included as an element. By construction, $x \in T^{-\ell} (A \cap \{ \Phi_A^{[d-1]} \leq (n_2 - \ell, \dots, n_d - \ell) \})$. If $x \in T^{-\ell} \{ \Phi_A^{[d]} \leq (n_1 - \ell, \dots, n_d - \ell) \}$, then $\ell + r_A(T^\ell x)$ is also in the set which is a contradiction by definition of ℓ .

The same approach shows that the union is disjoint. For $1 \leq \ell_1 < \ell_2 \leq n_1$, we have

$$\begin{aligned} & T^{-\ell_1} \left(A \cap \left\{ \Phi_A^{[d-1]} \leq (n_2 - \ell_1, \dots, n_d - \ell_1) \right\} \right) \cap \left\{ \Phi_A^{[d]} \leq (n_1 - \ell_1, \dots, n_d - \ell_1) \right\}^c \\ & \quad \cap T^{-\ell_2} \left(A \cap \left\{ \Phi_A^{[d-1]} \leq (n_2 - \ell_2, \dots, n_d - \ell_2) \right\} \right) \cap \left\{ \Phi_A^{[d]} \leq (n_1 - \ell_2, \dots, n_d - \ell_2) \right\}^c \\ & \subset T^{-\ell_1} \left(A \cap \left\{ \Phi_A^{[d]} \leq (n_1 - \ell_1, \dots, n_d - \ell_1) \right\}^c \right. \\ & \quad \left. \cap T^{-(\ell_2 - \ell_1)} (A \cap \{ \Phi_A^{[d-1]} \leq (n_2 - \ell_2, \dots, n_d - \ell_2) \}) \right) \\ & \subset T^{\ell_1} \left(A \cap \left\{ \Phi_A^{[d]} \leq (n_1 - \ell_1, \dots, n_d - \ell_1) \right\}^c \cap \left\{ \Phi_A^{[d]} \leq (\ell_2 - \ell_1, n_2 - \ell_1, \dots, n_d - \ell_1) \right\} \right) \\ & \subset T^{-\ell_1} \left(A \cap \left\{ \Phi_A^{[d]} \leq (n_1 - \ell_1, \dots, n_d - \ell_1) \right\}^c \cap \left\{ \Phi_A^{[d]} \leq (n_1 - \ell_1, n_2 - \ell_1, \dots, n_d - \ell_1) \right\} \right) = \emptyset. \end{aligned}$$

□

We are now ready to go on with the proof of Theorem 2.2.

Proof (of Theorem 2.2). Set $R_n := \mu(E_n) a(\Phi_{E_n})$ and for all $t \geq 0$, let $p_n^{[t]}$ be the integer such that

$$\left\{ \mu(E_n) a(r_{E_n}) \leq t \right\} = \left\{ r_{E_n} \leq p_n^{[t]} \right\}.$$

Conversely, define for all $0 \leq \ell \leq p_n^{[t]}$, the number $\vartheta_{n,\ell}^{[t]}$ such that

$$\left\{ \mu(E_n) a(r_{E_n}) \leq \vartheta_{n,\ell}^{[t]} \right\} = \left\{ r_{E_n} \leq p_n^{[t]} - \ell \right\}.$$

Thus, for all $d \geq 1$ and $t_1, \dots, t_d \geq 0$ we get

$$\left\{ R_n^{[d]} \leq (t_1, \dots, t_d) \right\} = \left\{ \Phi_{E_n}^{[d]} \leq (p_n^{[t_1]}, \dots, p_n^{[t_d]}) \right\}. \quad (3.1)$$

and

$$\left\{ R_n^{[d]} \leq (\vartheta_{n,\ell}^{[t_1]}, \dots, \vartheta_{n,\ell}^{[t_d]}) \right\} = \left\{ \Phi_{E_n}^{[d]} \leq (p_n^{[t_1]} - \ell, \dots, p_n^{[t_d]} - \ell) \right\}. \quad (3.2)$$

The quantities $p_n^{[t]}$ and $\vartheta_{n,\ell}^{[t]}$ can be computed using b the asymptotic inverse of a and a itself. We get

$$p_n^{[t]} := b(t/\mu(E_n)) \quad \text{and} \quad \vartheta_{n,\ell}^{[t]} = \mu(E_n) a(p_n^{[t]} - \ell).$$

First, assume the convergence of the return process, that is to say assume that $R_n \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} \tilde{\Phi}$. This is equivalent to the convergence of $R_n^{[d]}$ for every $d \geq 1$ and thus assume that $\tilde{G}_{E_n}^{[d]}(t_1, \dots, t_d) \Rightarrow \tilde{G}^{[d]}(t_1, \dots, t_d)$ at every point of continuity (t_1, \dots, t_d) of $\tilde{G}^{[d]}$ such that $t_1((t_2/t_1)^{1/\alpha} - m/M)^\alpha$ for all $0 \leq m \leq M$ are also continuity points of $\tilde{G}^{[d]}$. Without loss of generality, we can assume that $t_1 \leq \dots \leq t_d$ as the sequence of returns

is increasing. In the following, we write $\widetilde{G}_n^{[d]}$ instead of $\widetilde{G}_{E_n}^{[d]}$ to ease the notations and we will do the same for $G_u^{[d]}$.

Now, we have

$$\begin{aligned}
G_{u,n}^{[d]}(t_1, \dots, t_d) &= \int_{\{R_n^{[d]} \leq (t_1, \dots, t_d)\}} u \, d\mu = \int_{\{\Phi_{E_n}^{[d]} \leq (p_n^{[t_1]}, \dots, p_n^{[t_d]})\}} u \, d\mu \quad (\text{by (3.1)}) \\
&= \sum_{\ell=1}^{p_n^{[t_1]}} \left(\int_{E_n \cap \{\Phi_{E_n}^{[d-1]} \leq (p_n^{[t_2]} - \ell, \dots, p_n^{[t_d]} - \ell)\} \cap \{\Phi_{E_n}^{[d]} \leq (p_n^{[t_1]} - \ell, \dots, p_n^{[t_d]} - \ell)\}^c} \widehat{T}^\ell u \, d\mu \right) \quad (\text{by Lemma 3.1}) \\
&= \sum_{\ell=1}^{p_n^{[t_1]}} \left(\int_{E_n \cap \{\Phi_{E_n}^{[d-1]} \leq (p_n^{[t_2]} - \ell, \dots, p_n^{[t_d]} - \ell)\}} \widehat{T}^\ell u \, d\mu \right. \\
&\quad \left. - \int_{E_n \cap \{\Phi_{E_n}^{[d]} \leq (p_n^{[t_1]} - \ell, \dots, p_n^{[t_d]} - \ell)\}} \widehat{T}^\ell u \, d\mu \right) \quad (\text{because } p_n^{[t_i]} \leq p_n^{[t_{i+1}]}) \\
&= \sum_{m=0}^{M-1} \sum_{\ell=\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor + 1}^{\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor} \left(\int_{E_n \cap \{R_n^{[d-1]} \leq (\vartheta_{n,\ell}^{[t_2]}, \dots, \vartheta_{n,\ell}^{[t_d]})\}} \widehat{T}^\ell u \, d\mu \right. \\
&\quad \left. - \int_{E_n \cap \{R_n^{[d]} \leq (\vartheta_{n,\ell}^{[t_1]}, \dots, \vartheta_{n,\ell}^{[t_d]})\}} \widehat{T}^\ell u \, d\mu \right) \quad (\text{by (3.2)}) \\
&\leq \sum_{m=0}^{M-1} \sum_{\ell=\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor + 1}^{\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor} \left(\int_{E_n \cap \left\{ R_n^{[d-1]} \leq \left(\vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_2]}, \dots, \vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right\}} \widehat{T}^\ell u \, d\mu \right. \\
&\quad \left. - \int_{E_n \cap \left\{ R_n^{[d]} \leq \left(\vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_1]}, \dots, \vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right\}} \widehat{T}^\ell u \, d\mu \right) \\
&\leq \sum_{m=0}^{M-1} \int_{E_n \cap \left\{ R_n^{[d-1]} \leq \left(\vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_2]}, \dots, \vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right\}} \sum_{\ell=\lfloor \frac{m}{M} p_n^{[t_1]} \rfloor + 1}^{\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor} \widehat{T}^\ell u \, d\mu \\
&\quad - \int_{E_n \cap \left\{ R_n^{[d]} \leq \left(\vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_1]}, \dots, \vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right\}} \sum_{\ell=\lfloor \frac{m}{M} p_n^{[t_1]} \rfloor + 1}^{\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor} \widehat{T}^\ell u \, d\mu.
\end{aligned}$$

At this point, we can take advantage of the fact that Y is uniform for u . In particular, it means that of all $0 \leq c_1 < c_2$, we have

$$\sum_{\ell=c_1 k}^{c_2 k-1} \widehat{T}^\ell u \sim (c_2^\alpha - c_1^\alpha) a_k \quad \text{uniformly mod } \mu \text{ on } Y,$$

and thus

$$\sum_{\ell=\lfloor \frac{m}{M} p_n^{[t_1]} \rfloor + 1}^{\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor} \widehat{T}^\ell u \sim \left(\left(\frac{m+1}{M} \right)^\alpha - \left(\frac{m}{M} \right)^\alpha \right) a_{p_n^{[t_1]}} \quad \text{uniformly mod } \mu \text{ on } Y,$$

meaning that for all $\varepsilon > 0$ and n large enough,

$$\sum_{\ell=\lfloor \frac{m}{M} p_n^{[t_1]} \rfloor + 1}^{\lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor} \widehat{T}^\ell u \leq (1 + \varepsilon) \left(\left(\frac{m+1}{M} \right)^\alpha - \left(\frac{m}{M} \right)^\alpha \right) a_{p_n^{[t_1]}}.$$

By definition, $a_{p_n^{[t_1]}} = a(b(t_1/\mu(E_n))) \sim t_1/\mu(E_n)$ and by the mean value theorem $((m+1)/M)^\alpha - (m/M)^\alpha \leq \alpha m^{\alpha-1}/M^\alpha$. Hence, for n large enough, we get

$$G_{u,n}^{[d]}(t_1, \dots, t_d) \leq (1 + \varepsilon) \alpha t_1 \sum_{m=0}^{M-1} \frac{1}{M} \left(\frac{m}{M} \right)^{\alpha-1} \left[\mu_{E_n} \left(R_n^{[d-1]} \leq \left(\vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_2]}, \dots, \vartheta_{k, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right) \right]$$

$$\begin{aligned}
& -\mu_{E_n} \left(R_n^{[d]} \leq \left(\vartheta_{k, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_1]}, \dots, \vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right) \\
& \leq (1 + \varepsilon) \alpha t_1 \sum_{m=0}^{M-1} \frac{1}{M} \left(\frac{m}{M} \right)^{\alpha-1} \left[\tilde{G}_k^{[d-1]} \left(\vartheta_{k, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_2]}, \dots, \vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right. \\
& \quad \left. - \tilde{G}_n^{[d]} \left(\vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_1]}, \dots, \vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \right]. \tag{3.3}
\end{aligned}$$

The lower bound is obtained similarly. Now, $\forall t_2 \geq t_1 > 0, \forall 0 < c < 1$, we have

$$\begin{aligned}
\vartheta_{n, c p_n^{[t_1]}}^{[t_2]} &= \mu(E_n) a(p_n^{[t_2]} - c p_n^{[t_1]}) = \mu(E_n) a \left(b \left(\frac{t_2}{\mu(E_n)} \right) - c b \left(\frac{t_1}{\mu(E_n)} \right) \right) \\
&= \mu(E_n) a \left(b \left(\frac{t_1}{\mu(E_n)} \right) \left(b(t_2/\mu(E_n))/b(t_1/\mu(E_n)) - c \right) \right) \\
&\sim \mu(E_n) a \left(b(t_1/\mu(E_n)) \left(\left(\frac{t_2}{t_1} \right)^{1/\alpha} - c \right) \right) \\
&\sim t_1 \left(\left(\frac{t_2}{t_1} \right) - c \right)^\alpha,
\end{aligned}$$

where we use the hypothesis $a \in \text{RV}(\alpha)$ which implies $b \in \text{RV}(\alpha^{-1})$. Since we assumed the convergence for the return time process, it yields

$$\begin{aligned}
& \tilde{G}_n^{[d-1]} \left(\vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_2]}, \dots, \vartheta_{n, \lfloor \frac{m}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \\
& \xrightarrow{n \rightarrow +\infty} \tilde{G}^{[d-1]} \left(t_1 \left(\left(\frac{t_2}{t_1} \right)^{1/\alpha} - \frac{m}{M} \right)^\alpha, \dots, t_1 \left(\left(\frac{t_d}{t_1} \right)^{1/\alpha} - \frac{m}{M} \right)^\alpha \right)
\end{aligned}$$

and

$$\begin{aligned}
& \tilde{G}_n^{[d]} \left(\vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_1]}, \dots, \vartheta_{n, \lfloor \frac{m+1}{M} p_n^{[t_1]} \rfloor}^{[t_d]} \right) \\
& \xrightarrow{n \rightarrow +\infty} \tilde{G}^{[d]} \left(t_1 \left(1 - \frac{m+1}{M} \right)^\alpha, \dots, t_1 \left(\left(\frac{t_d}{t_1} \right)^{1/\alpha} - \frac{m+1}{M} \right)^\alpha \right).
\end{aligned}$$

Thus, together with (3.3), recognizing a Riemann integral and letting ε go to 0, we get

$$\begin{aligned}
G_{u,n}^{[d]}(t_1, \dots, t_d) & \xrightarrow{k \rightarrow +\infty} \alpha t_1 \int_0^1 \left[\tilde{G}^{[d-1]} \left(\left(t_2^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha, \dots, \left(t_d^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha \right) \right. \\
& \quad \left. - \tilde{G}^{[d]} \left(t_1 (1-r)^\alpha, \left(t_2^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha, \dots, \left(t_d^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha \right) \right] r^{\alpha-1} dr.
\end{aligned}$$

This means that we have convergence of the normalized hitting time point process with distribution functions

$$\begin{aligned}
G^{[d]}(t_1, \dots, t_d) &:= \alpha t_1 \int_0^1 \left[\tilde{G}^{[d-1]} \left(\left(t_2^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha, \dots, \left(t_d^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha \right) \right. \\
& \quad \left. - \tilde{G}^{[d]} \left(t_1 (1-r)^\alpha, \left(t_2^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha, \dots, \left(t_d^{1/\alpha} - t_1^{1/\alpha} r \right)^\alpha \right) \right] r^{\alpha-1} dr. \tag{3.4}
\end{aligned}$$

Now, assume conversely the convergence of the hitting process. By Helly selection theorem in \mathbb{R}^d (see [Kal02, Theorem 5.19]) and diagonal extraction, we can find a subsequence (n_k) such that $\tilde{G}_{n_k}^{[d]}$ converges pointwise towards $\tilde{G}_*^{[d]}$ for every $d \geq 0$. But, in this case, alongside this subsequence we have $G_{u, n_k}^{[d]}$ that converges pointwise towards $G_*^{[d]}$ defined by (3.4) from $\tilde{G}_*^{[d]}$. Since the equation defines uniquely $\tilde{G}_*^{[d]}$ from $G_*^{[d]} = G^{[d]}$ by Lemma 3.2, it gives the wanted convergence. \square

It remains to prove the following Lemma ensuring that the relationship found determines uniquely the law of one limit process from the laws of the other.

Lemma 3.2. *Let Φ and $\tilde{\Phi}$ be two processes on the phase space $\bar{\mathbb{R}}_+$. For all $d \geq 1$, write $G^{[d]}$ and $\tilde{G}^{[d]}$ the distribution functions of their marginals. By convention let $G^{[0]} = \tilde{G}^{[0]} = 1$. Then, the equation $\forall d \geq 1, \forall 0 \leq t_1 \leq \dots \leq t_d$,*

$$G^{[d]}(t_1, \dots, t_d) := \alpha t_1 \int_0^1 \left[\tilde{G}^{[d-1]} \left(\left(t_2^{1/\alpha} - t_1^{1/\alpha} s \right)^\alpha, \dots, \left(t_d^{1/\alpha} - t_1^{1/\alpha} s \right)^\alpha \right) \right.$$

$$- \tilde{G}^{[d]} \left(t_1 (1-s)^\alpha, \left(t_2^{1/\alpha} - t_1^{1/\alpha} s \right)^\alpha, \dots, \left(t_d^{1/\alpha} - t_1^{1/\alpha} s \right)^\alpha \right) s^{\alpha-1} ds,$$

defines uniquely the law of Φ from the law of $\tilde{\Phi}$ and conversely.

Remark 3.2. When $\alpha \in (0, 1]$, instead of looking at the distribution function $G^{[d]}$, we can look at $F^{[d]}$ defined by

$$F^{[d]}(t_1, \dots, t_d) := G^{[d]}(t_1^\alpha, \dots, t_d^\alpha), \forall t_1, \dots, t_d > 0.$$

In fact, F is exactly the family of distribution functions of the limit point processes Ψ and $\tilde{\Psi}$ of Theorem 2.1 by Remark 2.6. Thus, Lemma 3.2 is equivalent to Lemma 3.3 (below), and the equation in Lemma 3.3 gives (2.3), concluding the proof of Theorems 2.1-2.2.

Lemma 3.3. Let Φ and $\tilde{\Phi}$ be two process on the phase space $\overline{\mathbb{R}}_+$. For all $d \geq 1$, write $F^{[d]}$ and $\tilde{F}^{[d]}$ the distribution of their marginals. By convention, let $F^{[0]} = \tilde{F}^{[0]} = 1$. Then, the equations

$$F^{[d]}(t_1, \dots, t_d) = \alpha \int_0^{t_1} \left(\tilde{F}^{[d-1]}(t_2 - t_1 + x, \dots, t_d - t_1 + x) - \tilde{F}^{[d]}(x, t_2 - t_1 + x, \dots, t_d - t_1 + x) \right) (t_1 - x)^{\alpha-1} dx,$$

where $d \geq 1$ and one takes any d -uplet $0 \leq t_1 \leq \dots \leq t_d$, uniquely determines the law of Φ from the law of $\tilde{\Phi}$, and vice versa.

Proof (of Lemma 3.3). It is immediate that the law of Φ is uniquely determined by the law of $\tilde{\Phi}$. Now, assume that there are two stochastic processes $\tilde{\Phi}_1$ and $\tilde{\Phi}_2$ leading to the same Φ . We will argue by induction. The case $d = 1$ corresponds to the equivalence between hitting and return for the first return and has already been dealt in [RZ20, Remark 4.2.b)]. For $d \geq 1$, assuming the distribution functions are equal up to $d - 1$, it means that for all $0 \leq t_1 \leq \dots \leq t_d$,

$$\begin{aligned} & \alpha \int_0^{t_1} \tilde{F}_1^{[d]}(x, t_2 - t_1 + x, \dots, t_d - t_1 + x) (t_1 - x)^{\alpha-1} dx \\ &= \alpha \int_0^{t_1} \tilde{F}_2^{[d]}(x, t_2 - t_1 + x, \dots, t_d - t_1 + x) (t_1 - x)^{\alpha-1} dx. \end{aligned}$$

Taking s_2, \dots, s_d such that $t_i - t_1 = s_i$ for $2 \leq i \leq d$, we obtain, for all $t_1, s_2, \dots, s_d \geq 0$,

$$\begin{aligned} & \alpha \int_0^{t_1} \tilde{F}_1^{[d]}(x, x + s_2, \dots, x + s_d) (t_1 - x)^{\alpha-1} dx \\ &= \alpha \int_0^{t_1} \tilde{F}_2^{[d]}(x, x + s_2, \dots, x + s_d) (t_1 - x)^{\alpha-1} dx. \end{aligned}$$

Thus, if we define

$$f : x \mapsto \tilde{F}_1^{[d]}(x, x + s_2, \dots, x + s_d) - \tilde{F}_2^{[d]}(x, x + s_2, \dots, x + s_d),$$

we then have

$$\int_0^{t_1} f(x) (t_1 - x)^{\alpha-1} dx = 0, \quad \forall t_1 \geq 0.$$

Thus, by definition of the Riemann-Liouville integral (2.6), we have $I^\alpha f = 0$ which implies that $f = 0$, meaning that $\tilde{F}_1^{[d]} = \tilde{F}_2^{[d]}$. It completes the induction and the proof. \square

We turn now to the proof of Corollaries 2.2-2.3

Proof (of Corollary 2.2). This is immediate by looking at the projection on the first coordinate, *i.e.* $d = 1$, in (2.3). With $\tilde{F}^{[0]} = 1$, we get

$$F^{[1]}(t) = \alpha \int_0^1 (1 - \tilde{F}^{[1]}(u) (t - u)^{\alpha-1}) du, \quad t \geq 0,$$

which is exactly (2.4). Equation (2.5) can be found by the same method as in [PSZ13, Lemma 7]. \square

Proof (of Corollary 2.3). Recall the definition of $P_N(d, t) = \mathbb{P}(N[0, t] = d)$ for a point process N , $d \geq 0$ and $t \in \mathbb{R}$. Let $d \geq 1$ and $t \geq 0$. Then, (2.3) with $t_1, \dots, t_d = t$ is

$$F^{[d]}(t, \dots, t) = \alpha \int_0^t \left(\tilde{F}^{[d-1]}(x, \dots, x) - \tilde{F}^{[d]}(x, \dots, x) \right) (t-x)^{\alpha-1} dx.$$

Since $F^{[d]}(t, \dots, t) = \mathbb{P}(N[0, t] \geq d)$ and $\tilde{F}^{[d]}(t, \dots, t) = \mathbb{P}(\tilde{N}[0, t] \geq d)$ for all $d \geq 0$, it yields

$$\mathbb{P}(N[0, t] \geq d) = \alpha \int_0^t \mathbb{P}(\tilde{N}[0, t] = d-1) (t-x)^{\alpha-1} dx.$$

Thus, for every $d \geq 1$, we get

$$1 - \sum_{k=0}^d P_N(k, \cdot) = \Gamma(1+\alpha) I^\alpha (P_{\tilde{N}}(d, \cdot)),$$

whence

$$P_N(d, \cdot) = \Gamma(1+\alpha) I^\alpha (P_{\tilde{N}}(d, \cdot) - P_{\tilde{N}}(d-1, \cdot))$$

which is the integral formulation of (2.8). Similarly, for $d = 0$, we obtain

$$P_N(0, \cdot) = 1 - \Gamma(1+\alpha) I^\alpha (P_{\tilde{N}}(0, \cdot)).$$

□

3.1.2 Proof of Proposition 2.4

As suggested in Remark 2.9, we actually prove the characterization of the law and its uniqueness for the stochastic process. It entails the uniqueness of the point process via the composition by Ξ . The process $\Phi_{\text{FPP}_\alpha(\Gamma(1+\alpha))} \stackrel{\text{law}}{=} (\phi^{(i)})_{i \geq 1}$ with $\phi^{(i)} = \sum_{k=1}^i X_k$ and $(X_k)_{k \geq 1}$ i.i.d with common law $H_\alpha(\Gamma(1+\alpha))$, is such that $\Xi(\Phi_{\text{FPP}_\alpha(\Gamma(1+\alpha))}) \stackrel{\text{law}}{=} \text{FPP}_\alpha(\Gamma(1+\alpha))$.

We can do the same with $\Phi_{\text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha)))} = (\psi^{(i)})_{i \geq 1}$ where $\psi^{(i)} = \sum_{k=1}^i W_k$ and $(W_k)_{k \geq 1}$ are i.i.d having as a common law $W_{\alpha, \theta}(\theta\Gamma(1+\alpha))$ (see (2.10) for the definition of these random variables). Then, $\Xi(\Phi_{\text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha)))}) \stackrel{\text{law}}{=} \text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha)))$.

Proof (of Proposition 2.4). We will actually prove that for $\theta \in (0, 1]$, $\Phi_{\text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha)))}$ is the only stochastic process such that for all $d \geq 1$ and $t_1 \leq \dots \leq t_d$,

$$\begin{aligned} \tilde{F}^{[d]}(t_1, \dots, t_d) &= (1-\theta) \tilde{F}^{[d-1]}(t_1, \dots, t_d) \\ &+ \theta \alpha \int_0^{t_1} \left(\tilde{F}^{[d-1]}(t_2 - t_1 + x, \dots, t_d - t_1 + x) \right. \\ &\left. - \tilde{F}^{[d]}(x, t_2 - t_1 + x, \dots, t_d - t_1 + x) \right) (t_1 - x)^{\alpha-1} dx. \end{aligned} \quad (3.5)$$

In particular, when $\theta = 1$, we will get that $\text{FPP}_\alpha(\Gamma(1+\alpha))$ is the only process such that its finite-dimensional marginals are fixed points of the transformation from return times to hitting times.

As in Lemma 3.3, we show by induction on d that there is at most a one fixed distribution $F^{[d]}$. For $d = 1$, (2.5) in Corollary 2.2 clearly ensures that there is a unique fixed law and using the Laplace transform, $W_{\alpha, \theta}(\theta\Gamma(1+\alpha))$ can be identified as the fixed law².

Now assume $d \geq 2$ and that the result is true for $d-1$. Let $F_1^{[d]}$ and $F_2^{[d]}$ be two distribution functions compatible with $F_1^{[d-1]}$ and $F_2^{[d]}$ satisfying (3.5). Since $F_1^{[d-1]} = F_2^{[d-1]}$ by hypothesis, for all $t_1, s_2, \dots, s_d \geq 0$, we have

$$\begin{aligned} &F_1^{[d]}(t_1, t_1 + s_2, \dots, t_1 + s_d) - F_2^{[d]}(t_1, t_1 + s_2, \dots, t_1 + s_d) \\ &= \alpha \int_0^{t_1} \left(F_2^{[d]}(x, x + s_2, \dots, x + s_d) - F_1^{[d]}(x, x + s_2, \dots, x + s_d) \right) (t_1 - x)^{\alpha-1} dx. \end{aligned}$$

²This result corresponds to [RZ20, Lemma 7.1] and $W_{\alpha, \theta}(\theta\Gamma(1+\alpha))$ corresponds to the law $\tilde{H}_{\alpha, \theta}$ in the article. However, in [RZ20], the constant $\Gamma(1+\alpha)$ has been forgotten in the identification.

Letting

$$h : x \mapsto F_1^{[d]}(x, x + s_2, \dots, x + s_d) - F_2^{[d]}(x, x + s_2, \dots, x + s_d)$$

the equation can be rewritten as

$$h = -\Gamma(1 + \alpha) I^\alpha h.$$

Going into the Laplace domain (which is possible because h is locally integrable as its absolute value is bounded by 1), we get $h = 0$ and thus $F_1^{[d]} = F_2^{[d]}$.

Thus, there can be at most one process whose distribution functions are fixed points of (3.5). It remains to demonstrate the existence of at least one fixed point for this equation. Fortunately, the theory applies to certain examples of null-recurrent Markov chains for which results concerning first hitting and return times are well-established (see [BZ01, PSZ17] for the case $\theta = 1$, and [RZ20] for the general case).

By leveraging the strong Markov property, these results extend to the point process level in cases where the waiting times are independent. In such scenarios, the process $\Phi_{\text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha)))}$ emerges naturally and must satisfy (3.5). This demonstrates that a fixed point indeed exists.

Consequently, the fixed point is both unique and well-defined³. \square

3.2 Sufficient conditions for convergence towards FPP_α and CFPP_α

In this section, we prove Theorem 2.3. We first need to recall the following lemma giving a uniform control of the convergence to 0 for the average of the iterations by the transfer operator for functions in a compact subset of $L^1(\mu)$.

Lemma 3.4. [Zwe07a, Theorem 3.1] *Let (X, \mathcal{B}, μ, T) be a CEMPT and \mathcal{U} a compact subset of $L^1(\mu)$ such that $\int u d\mu = 1$ for all $u \in \mathcal{U}$. Then, uniformly in $u, u^* \in \mathcal{U}$, we have*

$$\left\| \frac{1}{M} \sum_{j=0}^{M-1} \hat{T}^j u - \frac{1}{M} \sum_{j=0}^{M-1} \hat{T}^j u^* \right\|_{L^1(\mu)} \xrightarrow{M \rightarrow +\infty} 0.$$

Now, we can dive into the proof of Theorem 2.3.

Proof (of Theorem 2.3). As discussed in Remark 2.9 and in the proof of Proposition 2.4, we will actually show the convergence of the process. Then, the application of Ξ and the extended continuous mapping theorem give the convergence of the point process.

We consider the sequence of stochastic processes $(\gamma(\mu(B_n))\Phi_{B_n})_{n \geq 0}$ on $\overline{\mathbb{R}}_+$ and the sequence of probability spaces (B_n, μ_{B_n}) , *i.e.* we look at the successive return times. As $(\overline{\mathbb{R}}_+)^{\mathbb{N}}$ is compact, this sequence is tight so up to a subsequence and that the convergence of the stochastic process is characterized by the convergence of its finite-dimensional marginals, we can assume that the family of distribution functions $(\tilde{F}_{B_n}^{[d]})_{d \geq 1}$ converges towards the family of functions $(\tilde{F}^{[d]})_{d \geq 0}$ (meaning that we have pointwise convergence for each one of them at the continuity points of $\tilde{F}^{[d]}$). We are going to show that the only possible limits are the distribution functions of the successive return times of the stochastic process $\Phi_{\text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha)))}$, which is enough to get the convergence towards this process. By Theorem 2.1, for any given density u , let $(F_{B_n, v}^{[d]})_{d \geq 1}$ denote the family of distribution functions corresponding to the finite-dimensional marginals of the stochastic process $(\gamma(\mu(B_n))\Phi_{B_n})$ on the probability space (X, μ_v) , where μ_v is a probability measure absolutely continuous with respect to μ , having density v . Then, the family of renormalized distribution functions $(\tilde{F}_{B_n, v}^{[d]})_{d \geq 1}$ converges to the family of functions $(F^{[d]})_{d \geq 1}$. Both $(\tilde{F}_{B_n, v}^{[d]})_{d \geq 1}$ and $(F^{[d]})_{d \geq 1}$ satisfy the relationship given in (2.3).

For all $d \geq 1$ and $0 \leq t_1 \leq \dots \leq t_d$ such that (t_1, \dots, t_d) is a continuity point of $\tilde{F}^{[d]}$, we have

$$\begin{aligned} \tilde{F}_{B_n}^{[d]}(t_1, \dots, t_d) &= \frac{\mu(U(B_n))}{\mu(B_n)} \mu_{U(B_n)}(\gamma(\mu_{B_n}) r_{B_n} \leq t_1, \dots, \gamma(\mu_{B_n}) r_{B_n}^{(d)} \leq t_d) \\ &\quad + \frac{\mu(Q(B_n))}{\mu(B_n)} \mu_{Q(B_n)}(\gamma(\mu_{B_n}) r_{B_n} \leq t_1, \dots, \gamma(\mu_{B_n}) r_{B_n}^{(d)} \leq t_d). \end{aligned} \quad (3.6)$$

We are going to study the two members separately, in Lemma 3.5 and Lemma 3.6. When $U(B_n) = \emptyset$ the first term is trivially equal to 0 by convention and we can jump to Lemma 3.6 directly.

³In [RZ20], the first-return case confirms this fixed point, although it omits the constant $\Gamma(1 + \alpha)$.

Lemma 3.5. Assume $U(B_n) \neq \emptyset$. For all $d \geq 1$ and $0 \leq t_1 \leq \dots \leq t_d$ such that (t_2, \dots, t_d) is a continuity point of $\tilde{F}^{[d-1]}$, we have

$$\frac{\mu(U(B_n))}{\mu(B_n)} \mu_{U(B_n)}(\gamma(\mu(B_n)) r_{B_n} \leq t_1, \dots, \gamma(\mu(B_n)) r_{B_n}^{(d)} \leq t_d) \xrightarrow{n \rightarrow +\infty} (1 - \theta) \tilde{F}^{[d-1]}(t_2, \dots, t_d).$$

Proof (of Lemma 3.5). By $(A1)_\alpha$, we have $\mu(U(B_n))/\mu(B_n) \xrightarrow{n \rightarrow +\infty} 1 - \theta$. It vanishes when $\theta = 1$ and else we have

$$\begin{aligned} & \mu_{U(B_n)}(\gamma(\mu(B_n)) r_{B_n} \leq t_1, \dots, \gamma(\mu(B_n)) r_{B_n}^{(d)} \leq t_d) \\ & \leq \mu_{U(B_n)} \left(\bigcap_{i=1}^{d-1} \left\{ \gamma(\mu(B_n)) r_{B_n}^{(i)} \circ T_{B_n} \leq t_{i+1} \right\} \right) \\ & \leq \mu(B_n) \int_{B_n} \frac{\mathbf{1}_{U(B_n)}}{\mu(U(B_n))} \left(\prod_{i=1}^{d-1} \mathbf{1}_{\{\gamma(\mu(B_n)) r_{B_n}^{(i)} \leq t_{i+1}\}} \right) \circ T_{B_n} d\mu_{B_n} \\ & \leq \mu(B_n) \int_{B_n} \widehat{T}_{B_n} \left(\frac{\mathbf{1}_{U(B_n)}}{\mu(U(B_n))} \right) \left(\prod_{i=1}^{d-1} \mathbf{1}_{\{\gamma(\mu(B_n)) r_{B_n}^{(i)} \leq t_{i+1}\}} \right) d\mu_{B_n} \\ & \xrightarrow{n \rightarrow +\infty} \int_{B_n} \prod_{i=1}^{d-1} \mathbf{1}_{\{\gamma(\mu(B_n)) r_{B_n}^{(i)} \leq t_{i+1}\}} d\mu_{B_n} = \tilde{F}^{[d-1]}(t_2, \dots, t_d) \quad (\text{by } (A6)_\alpha). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} & \mu_{U(B_n)}(\gamma(\mu(B_n)) r_{B_n} \leq t_1, \dots, \gamma(\mu(B_n)) r_{B_n}^{(d)} \leq t_d) \\ & = \mu_{U(B_n)} \left(\left\{ \gamma(\mu(B_n)) r_{B_n} \leq t_1 \right\} \cap \bigcap_{i=1}^{d-1} \left\{ \gamma(\mu(B_n)) r_{B_n}^{(i)} \circ T_{B_n} \leq t_d - \gamma(\mu(B_n)) r_{B_n} \right\} \right) \end{aligned}$$

and for all $\varepsilon > 0$ such that $(t_2 - \varepsilon, \dots, t_d - \varepsilon)$ is a continuity point of $\tilde{F}^{[d-1]}$, we have

$$\begin{aligned} & \mu_{U(B_n)}(\gamma(\mu(B_n)) r_{B_n} \leq t_1, \dots, \gamma(\mu(B_n)) r_{B_n}^{(d)} \leq t_d) \\ & \geq \mu_{U(B_n)} \left(\bigcap_{i=1}^{d-1} \left\{ \gamma(\mu(B_n)) r_{B_n}^{(i)} \circ T_{B_n} \leq t_{i+1} - \varepsilon \right\} \right) \\ & \quad - \mu_{U(B_n)}(\gamma(\mu(B_n)) r_{B_n} \geq t_1) - \mu(\gamma(\mu(B_n)) r_{B_n} \geq \varepsilon). \end{aligned}$$

Again by $(A6)_\alpha$, the first term converges towards $\tilde{F}^{[d]}(t_2 - \varepsilon, \dots, t_d - \varepsilon)$. For the two other terms, by $(A3)_\alpha$ we have $\mu_{B_n}(\gamma(\mu(B_n)) \tau_n \geq \varepsilon) \xrightarrow{n \rightarrow +\infty} 0$ and thus $\mu_{U_n}(\gamma(\mu(B_n)) \tau_n \geq \varepsilon) \xrightarrow{n \rightarrow +\infty} 0$ because $\theta < 1$. Together with $(A5)_\alpha$ it ensures the convergence to 0. Since (t_1, \dots, t_d) is a continuity point of $\tilde{F}^{[d-1]}$, letting $\varepsilon \rightarrow 0$ gives the result. \square

Lemma 3.6. For every $d \geq 1$ and $0 \leq t_1 \leq \dots \leq t_d$ such that (t_1, \dots, t_d) is a continuity point of $F^{[d]}$, we have

$$\frac{\mu(Q(B_n))}{\mu(B_n)} \mu_{Q(B_n)}(\gamma(\mu(B_n)) r_{B_n} \leq t_1, \dots, \gamma(\mu(B_n)) r_{B_n}^{(d)} \leq t_d) \xrightarrow{n \rightarrow +\infty} \theta F^{[d]}(t_1, \dots, t_d).$$

Proof (of Lemma 3.6). This time $(A1)_\alpha$ gives $\mu(Q(B_n))/\mu(B_n) \xrightarrow{n \rightarrow +\infty} \theta$. By $(A0)_\alpha$, there exist a function $u \in L^1(\mu)$ such that Y is u -uniform (without loss of generality we assume $\int u d\mu = 1$). We consider μ_{v_n} the probability absolutely continuous with respect to μ and of density $v_n := \widehat{T}^{\tau_n}(\mathbf{1}_{Q(B_n)})/\mu(Q(B_n))$ and write $(F_{B_n, v_n}^{[d]})_{d \geq 1}$ the family of distribution functions of $\gamma(\mu(B_n)) \Phi_{B_n}$ drawn from μ_{v_n} . By $(A2)_\alpha$, for all $n \geq 1$, $v_n \in \mathcal{U}$.

By definition of v_n , we have $(\gamma(\mu(B_n)) \Phi_{B_n}) \# \mu_{v_n} = (\gamma(\mu(B_n)) \Phi_{B_n} \circ T^{\tau_n}) \# \mu_{Q(B_n)}$. On $\{r_{B_n} > \tau_n\}$, we have $\gamma(\mu(B_n)) \Phi_{B_n} = \gamma(\mu(B_n)) \Phi_{B_n} \circ T^{\tau_n} + \gamma(\mu(B_n)) \tau_n$, so by $(A3)_\alpha$ - $(A4)_\alpha$ we get $d(\gamma(\mu(B_n)) \Phi_{B_n}, \gamma(\mu(B_n)) \Phi_{B_n} \circ T^{\tau_n}) \xrightarrow{n \rightarrow +\infty} 0$. In particular, for all $d \geq 1$, we have

$$\mu_{Q(B_n)}(\gamma(\mu(B_n)) r_{B_n} \leq t_1, \dots, \gamma(\mu(B_n)) r_{B_n}^{(d)} \leq t_d) - F_{B_n, v_n}^{[d]}(t_1, \dots, t_d) \xrightarrow{n \rightarrow +\infty} 0. \quad (3.7)$$

Now, we show that the stochastic processes $(\gamma(\mu(B_n))\Phi_{B_n})$ drawn from μ_{v_n} and μ_v share the same limit. We denote $(F_{B_n, v_n}^{[d]})_{d \geq 1}$ and $(F_{B_n, v}^{[d]})_{d \geq 1}$ their respective family of distribution functions of the finite-dimensional marginals. For every $d \geq 1$, consider a bounded Lipschitz function $\psi : \mathbb{R}^d \rightarrow \mathbb{R}^d$. We are going to show that

$$\int \psi \circ (\gamma(\mu(B_n))\Phi_{B_n}^{[d]}) u \, d\mu - \int \psi \circ (\gamma(\mu(B_n))\Phi_{B_n}^{[d]}) u^* \, d\mu \xrightarrow{n \rightarrow +\infty} 0 \quad \text{uniformly in } u, u^* \in \mathcal{U}.$$

Let $\varepsilon > 0$ and consider M large enough so that the quantity in Lemma 3.4 is smaller than ε . It yields,

$$\begin{aligned} \left| \int \psi \circ (\gamma(\mu(B_n))\Phi_{B_n}^{[d]}) \cdot \frac{1}{M} \sum_{j=0}^{M-1} (\hat{T}^j u - \hat{T}^j u^*) \, d\mu \right| &\leq \sup |\psi| \left\| \frac{1}{M} \sum_{j=0}^{M-1} (\hat{T}^j u - \hat{T}^j u^*) \right\|_{L^1(\mu)} \\ &\leq \varepsilon \sup |\psi|. \end{aligned}$$

Furthermore, we also have

$$\begin{aligned} &\left| \int \psi \circ (\gamma(\mu(B_n))\Phi_{B_n}^{[d]}) \left(u - \frac{1}{M} \sum_{j=0}^{M-1} \hat{T}^j u \right) \, d\mu \right| \\ &\leq \frac{1}{M} \sum_{j=0}^{M-1} \int \left| \psi \circ (\gamma(\mu(B_n))\Phi_{B_n}^{[d]}) - \psi \circ (\gamma(\mu(B_n))\Phi_{B_n}^{[d]}) \circ T^j \right| u \, d\mu \\ &\leq \frac{1}{M} \sum_{j=0}^{M-1} \left(2 \sup |\psi| \int_{\{r_{B_n} \leq j\}} u \, d\mu + \text{Lip}(\psi) \gamma(\mu(B_n)) j \right) \\ &\leq 2 \sup |\psi| \int_{\{r_{B_n} \leq M\}} u \, d\mu + \text{Lip}(\psi) \gamma(\mu(B_n)) M \\ &\leq \varepsilon \text{ for } n \text{ large enough and uniformly on } u \in \mathcal{U} \text{ since } \mathcal{U} \text{ is uniformly integrable.} \end{aligned}$$

Thus, by Portemanteau theorem, for every $d \geq 1$

$$F_{B_n, v_n}^{[d]}(t_1, \dots, t_d) - F_{B_n, v}^{[d]}(t_1, \dots, t_d) \xrightarrow{n \rightarrow +\infty} 0. \quad (3.8)$$

Since $F_{B_n, v}^{[d]}(t_1, \dots, t_d)$ converges towards $F^{[d]}(t_1, \dots, t_d)$, (3.7) and (3.8) together give the desired result. \square

Now, going back to (3.6) and with Lemmas 3.5 and 3.6, we get, for all $d \geq 1$ and $t_1 \leq \dots \leq t_d$ such that (t_1, \dots, t_d) and (t_2, \dots, t_d) are continuity points of $\tilde{F}^{[d]}$, $F^{[d]}$ and $\tilde{F}^{[d-1]}$ we have

$$\tilde{F}_{B_n}^{[d]}(t_1, \dots, t_d) \xrightarrow{n \rightarrow +\infty} (1 - \theta) \tilde{F}^{[d-1]}(t_2, \dots, t_d) + \theta F^{[d]}(t_1, \dots, t_d).$$

But we assumed that $\tilde{F}_{B_n}^{[d]}(t_1, \dots, t_d) \xrightarrow{n \rightarrow +\infty} \tilde{F}^{[d]}(t_1, \dots, t_d)$. Hence, we get the following equality

$$\tilde{F}^{[d]}(t_1, \dots, t_d) = (1 - \theta) \tilde{F}^{[d-1]}(t_2, \dots, t_d) + \theta F^{[d]}(t_1, \dots, t_d).$$

Using Theorem 2.1, we can express $F^{[d]}$ from $\tilde{F}^{[d]}$ and $\tilde{F}^{[d-1]}$ by (2.3) we get

$$\begin{aligned} \tilde{F}^{[d]}(t_1, \dots, t_d) &= (1 - \theta) \tilde{F}^{[d-1]}(t_1, \dots, t_d) \\ &\quad + \theta \alpha \int_0^{t_1} \left(\tilde{F}^{[d-1]}(t_2 - t_1 + x, \dots, t_d - t_1 + x) \right. \\ &\quad \left. - \tilde{F}^{[d]}(x, t_2 - t_1 + x, \dots, t_d - t_1 + x) \right) (t_1 - x)^{\alpha-1} \, dx, \end{aligned}$$

which is exactly (3.5). In the proof of Proposition 2.4, we showed that the only process such that the distribution functions of its marginals satisfy (3.5) is $\Phi_{\text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha)))}$. Thus, we get the convergence

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{RPP}(W_{\alpha, \theta}(\theta\Gamma(1+\alpha))).$$

By (2.3) and Corollary 2.1, we also obtain

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{CFPP}_\alpha(\theta\Gamma(1+\alpha), \text{Geo}(\theta)).$$

\square

4 Quantitative recurrence for maps with an indifferent fixed point

We now focus on the family of maps with one indifferent fixed point with LSV parameters and prove the results presented in Section 2.3.

4.1 Additional properties

We recall the basic properties of the map T defined in (1.1) where $p \geq 1$. Recall that there is a unique absolutely continuous invariant measure μ and $\mu([0, 1]) = +\infty$ if and only if $p \geq 1$. Recall the definitions of $c_n = T_1^{-n}1$, where T_1 is the first branch of T , and of the partitions $\xi = \{[c_{n+1}, c_n], n \geq 0\}$ and $\xi_k = \bigvee_{j=0}^{k-1} T^{-j}\xi$.

Although we will not directly use the coding of the map by a renewal shift, we will take advantage of this perspective and use symbolic notations in what follows.

For all $n \geq 0$, let $[n] := [c_{n+1}, c_n]$. In particular, $\xi = \{[n], n \geq 0\}$. By definition of T , for $n \geq 1$, $T|_{[n]} : [n] \rightarrow [n-1]$ and $T|_{[0]} : [0] \rightarrow [0, 1]$ are diffeomorphisms. Similarly, the partition ξ_k can be expressed with cylinder notations.

Lemma 4.1. *For all $k \geq 0$, we have*

$$\xi_k = \bigvee_{j=0}^{k-1} T^{-j}\xi = \{[a_0^{k-1}] \mid (a_0^{k-1}) \in \mathbb{N}^k, \forall 0 \leq i \leq k-2, a_{i+1} = a_i - 1 \text{ or } a_i = 0\},$$

where $[a_0^{k-1}] = \{x \in [0, 1] \mid T^j(x) \in [a_j] \forall 0 \leq j \leq k-1\}$.

Using the terminology of symbolic dynamics, we will say that (a_0^{k-1}) is admissible if $[a_0^{k-1}] \neq \emptyset$.

Proof (of Lemma 4.1). This can be easily seen by induction. When $k = 1$, this is the definition of ξ . Now, $\xi_{k+1} = \xi_k \vee T^{-1}\xi_k$. Consider an element $[a_0^{k-1}]$ in ξ_k . Then, $T^{-1}[a_0^{k-1}] = T_1^{-1}[a_0^{k-1}] \sqcup T_2^{-1}[a_0^{k-1}]$. By definition $T_2^{-1}[a_0^{k-1}] = [0a_0^{k-1}]$ and since $T_1^{-1}[a_0] = [a_0 + 1]$, we have $T_1^{-1}[a_0^{k-1}] = [(a_0 + 1)a_0^{k-1}]$ and $(0a_0^{k-1})$ and $((a_0 + 1)a_0^{k-1})$ are admissible. Reciprocally, for an admissible sequence, we have

$$[a_0^{k-1}] = T_{\sigma(a_0)}^{-1} \cdots T_{\sigma(a_{k-2})}^{-1}[a_{k-1}] \in \xi_k, \quad (4.1)$$

where $\sigma(0) := 2$ and $\sigma(k) := 1$ for all $k \geq 1$. □

Remark 4.1. *Due to the Markov property of the partition, for all $k \geq 1$ and all $(a_0^{k-1}) \in \mathbb{N}^k$ admissible, $T^j : [a_0^{k-1}] \rightarrow [a_j^{k-1}]$ is a homeomorphism. Furthermore, we always have $[a_0^k] = [a_0^k(a_k - 1)(a_k - 2) \dots 0]$.*

For all $0 \leq n \leq q \leq \infty$, we define the sets

$$[[n, q]] := \bigsqcup_{n \leq k \leq q} [k] = [c_{q+1}, c_n].$$

When $q = \infty$, we simply write $[(\geq n)] := [[n, \infty]] = [0, c_n]$ and when $n = 0$, $[(<= q)] := [[0, q]] = [c_{q+1}, 1]$.

Lemma 4.2. *For all $k \geq 1$ and $(a_0^{k-1}) \in \mathbb{N}^k$ admissible, $[a_0^{k-1}]$ is an interval. If furthermore $a_{k-1} = 0$, then, for all $0 \leq n \leq q \leq \infty$, $[a_0^{k-1}[n, q]]$ is also an interval.*

Proof (of Lemma 4.2). This is immediate from (4.1) and because $[[n, q]]$ are intervals for every $0 \leq n \leq q \leq \infty$. □

For every $j \geq 0$, it is known that the induced map on $Y := [c_{j+1}, 1]$ has a Gibbs-Markov structure. Indeed, define

$$\xi^Y := \xi_1^Y = \{[i], 1 \leq i \leq j\} \sqcup \{[0i], i \geq 0\},$$

and for all $k \geq 1$

$$\xi_k^Y := \bigvee_{i=0}^{k-1} T_Y^{-i}\xi_1^Y.$$

Then, (Y, T_Y, μ_Y) is Gibbs-Markov with respect to the partition ξ^Y . We can also define

$$\xi_0^Y := T_Y \xi^Y = \xi \cap Y = \{[i], 0 \leq i \leq j\} = \{[c_{i+1}, c_i], 0 \leq i \leq j\}.$$

Remark 4.2. If $j = 0$, i.e. $Y = [1/2, 1]$, the system is full-branched Gibbs-Markov. Otherwise, it is Gibbs-Markov with the “big image” property.

We also recall some useful estimates. This can be found in [You99] or [LSV99] for example.

Lemma 4.3. We have the following asymptotic results

$$\mu[0(\geq n)] \asymp \frac{1}{n^\alpha} \quad (4.2)$$

and

$$\mu[0n] \asymp \frac{1}{n^{\alpha+1}}, \quad (4.3)$$

where $a_n \asymp b_n$ means that there exists $C > 1$ such that $C^{-1}b_n \leq a_n \leq Cb_n$, for all $n \geq 1$.

Finally, we recall the following lemma (see [PSZ17] p.31 and [Tha80, Tha83]) giving a distortion bound for good subsets (this lemma is true for the bigger class of AFN-maps).

Lemma 4.4. Let Y be of the form $[c_{j+1}, 1]$ for some $j \geq 0$. Then, if $V, W \subset Y$ are intervals such that there exists $m \in \mathbb{N}$ with $T^m : V \rightarrow W$ being a homeomorphism onto W , then $\widehat{T}^m \mathbf{1}_V \in \mathcal{C}_C(W)$ for some $C > 0$ depending only on Y , where

$$\mathcal{C}_C(W) := \{f : Y \rightarrow [0, +\infty) \mid |f(x)/f(y) - 1| \leq Cd(x, y) \forall x, y \in W\}.$$

From this lemma we are able to derive the following corollary that is going to be of crucial importance in the proof of Theorem 2.7.

Corollary 4.5. There exists a constant $C > 0$ such that for all $k, \ell, m \geq 0$, for all $(a_0^{k-1}), (b_0^{\ell-1}), (c_0^{m-1})$ admissible (with the possibility that $b_{\ell-1}$ and c_{m-1} are of the form $[i, j]$ for some $0 \leq i \leq j \leq \infty$) such that $[0a_0^{k-1}0b_0^{\ell-1}] \neq \emptyset$ and $[0a_0^{k-1}0c_0^{m-1}] \neq \emptyset$, we get

$$\frac{\mu[0a_0^{k-1}0b_0^{\ell-1}]}{\mu[0a_0^{k-1}0c_0^{m-1}]} = (1 \pm C \text{diam}([0b_0^{\ell-1}] \cup [0c_0^{m-1}])) \frac{\mu[0b_0^{\ell-1}]}{\mu[0c_0^{m-1}]},$$

where $x = (1 \pm C)y$ means that $(1 - C)y \leq x \leq (1 + C)y$.

Proof (of Corollary 4.5). Consider the constant C from Lemma 4.4 with $Y = [1/2, 1] = [0]$. Let $j := \min\{k \geq 0 \mid b_k \neq c_k\}$ with the convention that $j := \min\{\ell, m\} - 1$ if $b_0^{\min\{\ell, m\}-1} = c_0^{\min\{\ell, m\}-1}$. In particular, we have $\text{diam}([0b_0^{\ell-1}] \cup [0c_0^{m-1}]) = \text{diam}([0b_0^{j-1}])$. By a property of the map T , we know that $T^{k+1} : [0a_0^{k-1}0b_0^{j-1}] \mapsto [0b_0^{j-1}]$ is a homeomorphism with both $[0a_0^{k-1}0b_0^{j-1}]$ and $[0b_0^{j-1}]$ being intervals of Y . Thus, by Lemma 4.4, we have $\widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]} \in \mathcal{C}_C([0b_0^{j-1}])$. Consequently,

$$\begin{aligned} \mu[0a_0^{k-1}0b_0^{\ell-1}] &= \int \mathbf{1}_{[0a_0^{k-1}0b_0^{\ell-1}]} d\mu = \int \widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{\ell-1}]} d\mu \\ &= \int \mathbf{1}_{[0b_0^{\ell-1}]} \cdot \widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]} d\mu \\ &= \widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]}(y) \int \mathbf{1}_{[0b_0^{\ell-1}]}(x) \cdot \frac{\widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]}(x)}{\widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]}(y)} d\mu \quad (\forall y \in [0c_0^{m-1}]) \\ &= (1 \pm Cd(x, y)) \widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]}(y) \int \mathbf{1}_{[0b_0^{\ell-1}]} d\mu \quad \forall y \in [0c_0^{m-1}] \\ &= \left(1 \pm C \text{diam}([0b_0^{j-1}])\right) \mu[0b_0^{\ell-1}] \mathbb{E}_{\mu|_{[0c_0^{m-1}]}} \left[\widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]}\right] \\ &= \left(1 \pm C \text{diam}([0b_0^{j-1}])\right) \frac{\mu[0b_0^{\ell-1}]}{\mu[0c_0^{m-1}]} \int \mathbf{1}_{[0c_0^{m-1}]} \widehat{T}^{k+1} \mathbf{1}_{[0a_0^{k-1}0b_0^{j-1}]} d\mu \\ &= \left(1 \pm C \text{diam}([0b_0^{j-1}])\right) \frac{\mu[0b_0^{\ell-1}]}{\mu[0c_0^{m-1}]} \mu[0a_0^{k-1}0c_0^{m-1}]. \end{aligned}$$

□

Remark 4.3. Although we are using notations coming from the symbolic dynamics, we are still working on the interval with its topology. For example, $\text{diam}([0(\geq p)]) \xrightarrow{p \rightarrow +\infty} 0$ in our case, whereas $\text{diam}([0(\geq p)]) = 1/4$ for all $p \geq 0$ with the usual geometric topology of countable Markov shifts.

4.2 Proof of Theorems 2.4-2.5

Once $x \in [0, 1] \setminus \bigcup_{k \geq 0} T^{-k}\{0\}$ is chosen, we fix Y of the form $[c_{j+1}, 1]$ such that $x \in Y$. Furthermore, there exists a sequence of integers $(a_k)_{k \geq 0}$ such that (a_0^{n-1}) is admissible for all n and $B_n = \xi_n(x) = [a_0^{n-1}]$. Set $s_n := S_{n-1}\mathbf{1}_Y(x)$ the number of visits to Y before n (note that $s_n \geq 1$ for $n \geq 2$ since we assumed $x \in Y$).

Lemma 4.6. *For all $n \geq 0$, we have*

$$B_n = \xi_{s_n}^Y(x).$$

Proof (of Lemma 4.6). We have $B_1 = \xi(x) = \xi_0^Y(x)$ and $s_1 = S_0\mathbf{1}_Y(x) = 0$. Assume that the result is true for some $n \geq 1$. We have $B_n = \xi_n(x) = [a_0^{n-1}]$ and $B_{n+1} = \xi_{n+1}(x) = [a_0^n]$. There are three cases, if $a_{n-1} > j$, $B_{n+1} = [a_0^{n-1}(a_{n-1} - 1)] = [a_0^{n-1}] = B_n$ et $s_{n+1} = S_n\mathbf{1}_Y(x) = S_{n-1}\mathbf{1}_Y(x) = s_n$ meaning that $\xi_{s_{n+1}}^Y(x) = \xi_{s_n}^Y(x) = B_n = B_{n+1}$. If $1 \leq a_{n-1} \leq j$, $s_{n+1} = s_n + 1$ but $\xi_{s_{n+1}}^Y(x) = \xi_{s_n}^Y(x)$ and again $\xi_{s_{n+1}}^Y(x) = \xi_{s_n}^Y(x) = B_n = B_{n+1}$.

Finally, if $a_{n-1} = 0$, then $s_{n+1} = s_n + 1$ and $\xi_{s_{n+1}}^Y(x) = [a_0^n] = B_{n+1}$. \square

4.2.0.1 Proof of Theorem 2.4: Non periodic points.

Proof (of Theorem 2.4). Let $x \in [0, 1]$ be non periodic and not a preimage of 0. Let $\tau_n := \min\{i \geq n-1 \mid T^i x \in Y\}$. Thus $s_n = S_{n-1}\mathbf{1}_Y(x) = S_{\tau_n}\mathbf{1}_Y(x)$ and $B_n = \xi_n(x) = \xi_{s_n}^Y(x)$ for $n \geq 1$ by Lemma 4.6. Since (Y, T_Y, μ_Y, ξ^Y) is Gibbs-Markov, there exists $\kappa, c > 0$ such that $\mu(B_n) \leq \kappa e^{-cs_n}$.

We will use Theorem 2.3 to get the result. Thus, we need to prove conditions $(A0)_\alpha$ - $(A6)_\alpha$. $(A0)_\alpha$ is immediate since Y is a uniform set. Since there is no appearance of clusters in the non periodic case, we set $U(B_n) = \emptyset$ and $Q(B_n) = 1$ for $(A1)_\alpha$. It remains to show $(A2)_\alpha$ - $(A4)_\alpha$ for our particular deterministic choice τ_n . We start with condition $(A3)_\alpha$. In fact, we will even show the stronger result $\tau_n \mu(B_n) \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} 0$. For that there are two cases, either x is a "well-behaved" point, meaning it will return to Y often enough, or x does not come back sufficiently often to Y . In the first case, the exponential decay of the cylinder measures for the induced map is enough to get the result. In the second case, since x is a special point, the measure of its cylinder will already be small enough. Consider a sequence (n_k) such that $\lim_{k \rightarrow +\infty} \tau_{n_k} \mu(B_{n_k}) = \limsup_{n \rightarrow +\infty} \tau_n \mu(B_n)$. Without loss of generality, we can assume that $(S_{\tau_{n_k}}\mathbf{1}_Y(x) / \log(\tau_{n_k}))_{k \geq 0}$ is either bounded or diverges to $+\infty$. If it diverges to $+\infty$, we have

$$\tau_{n_k} \mu(B_{n_k}) \leq \tau_{n_k} \kappa e^{-cs_{n_k}} = \tau_{n_k} \kappa e^{-cS_{\tau_{n_k}}\mathbf{1}_Y(x)} \xrightarrow[k \rightarrow +\infty]{} 0.$$

Otherwise, if the sequence is bounded by $K > 0$, we have $S_{\tau_{n_k}}\mathbf{1}_Y(x) \leq K \log(\tau_{n_k})$ and thus, there exists some $j \leq K \log(\tau_{n_k})$ such that $r_Y^{\{j\}}(x) \geq \tau_{n_k} / (K \log(\tau_{n_k}))$. It yields,

$$\begin{aligned} \tau_{n_k} \mu(B_{n_k}) &\leq \tau_{n_k} \mu(Y \cap T_Y^{-j+1}\{r_Y = r_Y^{\{j\}}(x)\}) \leq \tau_{n_k} \mu(Y \cap \{r_Y = r_Y^{\{j\}}(x)\}) \\ &\leq C \tau_{n_k} \frac{1}{r_Y^{\{j\}}(x)^{\alpha+1}} \leq CK^\alpha \frac{\tau_{n_k} \log(\tau_{n_k})^{\alpha+1}}{\tau_{n_k}^{\alpha+1}} \xrightarrow[k \rightarrow +\infty]{} 0 \quad (\text{using (4.3)}). \end{aligned}$$

Hence condition $(A3)_\alpha$ is satisfied. For condition $(A4)_\alpha$, we can use $(A3)_\alpha$ and the fact that we have return time statistics towards the exponential law for the induced map (because it is Gibbs-Markov). We get

$$\begin{aligned} \mu_{B_n}(r_{B_n} < \tau_n) &\leq \mu_{B_n}(r_Y^{(r_{B_n}^Y)} < \tau_n) \leq \mu_{B_n}(S_{\tau_n}\mathbf{1}_Y > r_{B_n}^Y) \\ &\leq \mu_{B_n}(\mu_Y(B_n) r_{B_n}^Y < \mu_Y(B_n) S_{\tau_n}\mathbf{1}_Y) \\ &\leq \mu_{B_n}(\mu_Y(B_n) r_{B_n}^Y < t) + \mu_{B_n}(\mu_Y(B_n) S_{\tau_n}\mathbf{1}_Y > t), \end{aligned}$$

for all $t > 0$. Fix $\varepsilon > 0$. Since B_n has return time statistics for the induced map, we can take t small enough so that $\limsup_{n \rightarrow +\infty} \mu_{B_n}(\mu_Y(B_n) r_{B_n}^Y < t) \leq \varepsilon$. On the other hand, $S_{\tau_n}\mathbf{1}_Y \leq \tau_n$ and using $\mu(B_n) \tau_n \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} 0$, it yields

$$\limsup_{n \rightarrow +\infty} \mu_{B_n}(r_{B_n} < \tau_n) \leq \varepsilon,$$

proving $(A4)_\alpha$.

It remains to show $(A2)_\alpha$. Since $B_n = \xi_{s_n}^Y(x)$, we have

$$\widehat{T^{\tau_n}}(\mathbf{1}_{B_n} / \mu(B_n)) = \widehat{T_Y^{s_n}}(\mathbf{1}_{\xi_{s_n}^Y(x)} / \mu(\xi_{s_n}^Y(x))).$$

As the return map to Y is Gibbs-Markov, by [Aar97, §4.7] (see also [Zwe22, Section 10.2]), there exists some \mathcal{U} compact in $L^1(\mu)$ such that $\widehat{T^{\tau_n}}(\mathbf{1}_{B_n}/\mu(B_n)) \in \mathcal{U}$ for every $n \geq 1$.

Thus, since every condition of Theorem 2.3 in the case $U(B_n) = \emptyset$ is satisfied, it gives us the wanted result. \square

4.2.0.2 Proof of Theorem 2.5: Periodic points.

Proof (of Theorem 2.5). Assume now that $x \in [0, 1] \setminus \bigcup_{k \geq 0} T^{-k}\{0\}$ is periodic of prime period q . Again, we want to use Theorem 2.3 and thus we have to show $(A0)_{\alpha}^-(A6)_{\alpha}$. $(A0)_{\alpha}$ is immediate since Y is a uniform set. Recall that $B_n = \xi_n(x) = \xi_{s_n}^Y(x)$ with $s_n = S_{n-1}\mathbf{1}_Y(x)$. Let $Q(B_n) := B_n \cap T^{-q}B_n^c$ be the escape annulus and $U(B_n) = B_n \setminus Q(B_n) = B_n \cap T^{-q}B_n$. We start by conditions $(A1)_{\alpha}$ and $(A6)_{\alpha}$. We have

$$\mu(U(B_n)) = \mu(B_n \cap T^{-q}B_n) = \int \mathbf{1}_{B_n} \cdot \mathbf{1}_{B_n} \circ T^q d\mu = \int \mathbf{1}_{B_n} \cdot \mathbf{1}_{B_n} \circ T^q \rho d\text{Leb}.$$

For n large enough such that $T^j B_n \cap B_n = \emptyset$ for $1 \leq j < q$ and by construction of $U(B_n)$, $T_{B_n} = T^q$ on $U(B_n)$ and $T^q : U(B_n) \rightarrow B_n$ is a diffeomorphism. Thus, for every $y \in B_n$

$$\widehat{T_{B_n}} \mathbf{1}_{U(B_n)}(y) = \widehat{T^q} \mathbf{1}_{U(B_n)}(y) \mathbf{1}_{B_n}(y) = \frac{1}{(T^q)'(T_{U(B_n)}^{-q}y)}$$

where $T_{U(B_n)}^{-q}y$ is the only q -preimage of y in $U(B_n)$. Since T' is continuous at x (and $T'(x) > 0$), $\|1/(T' \circ T_{U(B_n)}^{-q}) - 1/(T'(x))\|_{L^\infty(\mu_{B_n})} \xrightarrow{n \rightarrow +\infty} 0$. Thus, using that $\widehat{T_{B_n}} \left(\frac{\mathbf{1}_{U(B_n)}}{\mu(B_n)} \right)$ and $\frac{\mathbf{1}_{B_n}}{\mu(B_n)}$ are probability densities, we have

$$\left\| \widehat{T_{B_n}} \left(\frac{\mathbf{1}_{U(B_n)}}{\mu(B_n)} \right) - \frac{\mathbf{1}_{B_n}}{\mu(B_n)} \right\|_{L^\infty(\mu_{B_n})} \xrightarrow{n \rightarrow +\infty} 0$$

and

$$\frac{\mu(U(B_n))}{\mu(B_n)} \xrightarrow{n \rightarrow +\infty} \frac{1}{(T^q)'(x)},$$

proving conditions $(A1)_{\alpha}$ and $(A6)_{\alpha}$ with

$$\theta = 1 - \frac{1}{(T^q)'(x)}.$$

For $(A5)_{\alpha}$, by construction, we have $U(B_n) = B_{n+q} = \xi_{s_{n+q}}^Y(x)$ and we set $\tau_n := \min\{i \geq n+q-1 \mid T^i x \in Y\}$ so that $s_{n+q} = S_{\tau_n} \mathbf{1}_Y(x)$. First, since $r_{B_n} = q$ on $U(B_n)$, we immediately get

$$\mu_{U(B_n)}(r_{B_n} > \tau_n) = \mu_{U(B_n)}(q > \tau_n) \xrightarrow{n \rightarrow +\infty} 0.$$

Furthermore, since x is a periodic point, $S_n \mathbf{1}_Y(x) / \log(n) \xrightarrow{n \rightarrow +\infty} +\infty$. Thus, since the measure of cylinders in Gibbs-Markov maps decays exponentially and $s_{n+q} \leq s_n + q$

$$\tau_n \mu(B_n) \leq \tau_n \kappa e^{-cs_n} \leq \tau_n \kappa e^q e^{-cs_{n+q}} = (\kappa e^q) \tau_n e^{S_{\tau_n} \mathbf{1}_Y(x)} \xrightarrow{n \rightarrow +\infty} 0,$$

which proves $(A3)_{\alpha}$. For $(A4)_{\alpha}$, as in the non periodic case, we can take advantage of the return time statistics for the induced Gibbs-Markov system. This time, the convergence of $\mu_Y(B_n) r_{B_n}^Y$ is not to an exponential law but to a law with distribution function $t \mapsto (1 - \theta) + \theta(1 - e^{-\theta t})$. However, starting from $Q(B_n)$ instead of B_n implies that we do not have the cluster at 0 and a convergence to the same law as for the hitting time, that is to say the law with distribution function $t \mapsto 1 - e^{-\theta t}$ [Zwe22, Theorem 3.2]. Thus we can use the same decomposition as before to get

$$\mu_{Q(B_n)}(r_{B_n} < \tau_n) \leq \mu_{Q(B_n)}(\mu_Y(B_n) r_{B_n}^Y < t) + \mu_{Q(B_n)}(\mu_Y(B_n) S_{\tau_n} \mathbf{1}_Y > t)$$

for every $t > 0$. Fix $\varepsilon > 0$. By the convergence for the induced map, the first term will be smaller than ε if t is chosen small enough. For the second term, $S_{\tau_n} \mathbf{1}_Y \leq \tau_n$ and we already proved that $\tau_n \mu(B_n) \xrightarrow{n \rightarrow +\infty} 0$ and thus, for all $t > 0$,

$$\begin{aligned} \mu_{Q(B_n)}(\mu(B_n) \tau_n > t) &= \frac{1}{\mu(Q(B_n))} \int_{Q(B_n)} \mathbf{1}_{\{\mu(B_n) \tau_n > t\}} d\mu \\ &\leq \frac{\mu(B_n)}{\mu(Q(B_n))} \mu_{B_n}(\mu(B_n) \tau_n > t) \xrightarrow{n \rightarrow +\infty} \theta^{-1} \times 0 = 0. \end{aligned}$$

For the remaining condition $(A2)_{\alpha}$, we can again use [Aar97, §4.7] (see also [Zwe22, Section 10.2]), to ensure the existence of some \mathcal{U} compact (and convex) in $L^1(\mu)$ such that $\widehat{T^{\tau_n}}(\mathbf{1}_{Q(B_n)}/\mu(Q(B_n))) \in \mathcal{U}$ for every $n \geq 1$ because $Q(B_n)$ is a finite union of $\xi_{s_{n+q}}$ -cylinders. \square

4.3 Proofs of Theorems 2.7 and 2.9

4.3.1 The case $x = 1/2$

Proposition 4.7. *Let $B_n := [0(\geq n)] = [1/2, \delta_n]$ be a sequence of nested balls shrinking towards $1/2$, where $\delta_n = (1 + c_n)/2$. Then,*

$$\gamma(\mu(B_n)) r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \mathfrak{J}_\alpha \quad (4.4)$$

and

$$\gamma(\mu(B_n)) r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \tilde{\mathfrak{J}}_\alpha. \quad (4.5)$$

Proof (of Proposition 4.7). Since B_n is uniformly bounded away from 0, it lies inside a uniform set for all n and thus Corollary 2.2-2) ensures that (4.4) and (4.5) are equivalent. Furthermore, with the strong distributional convergence [RZ20, Lemma 4.1], (4.4) is equivalent to $\gamma(\mu(B_n)) r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mu_{[3/4, 1]}} \mathfrak{J}_\alpha$. Notice that $T(B_n) = [0, c_n]$ is a right neighborhood of 0 and that on $[3/4, 1]$, we have $r_{B_n} = r_{T(B_n)} - 1$. Using 2.6, we obtain

$$\gamma(\mu(T_2^{-1}(T(B_n)))) r_{T(B_n)} \xrightarrow[n \rightarrow +\infty]{\mu_{[3/4, 1]}} \mathfrak{J}_\alpha.$$

Since $T_2^{-1}(T(B_n)) = B_n$, it concludes the proof. \square

Proposition 4.8. *Let $B_n := [1/2, \delta_n]$ be as in Proposition 4.7. Then,*

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{DRPP}(\mathfrak{J}_\alpha, \tilde{\mathfrak{J}}_\alpha), \quad (4.6)$$

and

$$N_{B_n}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{B_n}} \text{RPP}(\tilde{\mathfrak{J}}_\alpha). \quad (4.7)$$

Proof (of Proposition 4.8). To show the convergence, it is enough to prove that the stochastic process of the waiting times converges towards the stochastic process $(W_k)_{k \geq 1}$ of independent waiting times with $W_1 \stackrel{\text{law}}{=} \mathfrak{J}_\alpha$ and $W_i \stackrel{\text{law}}{=} \tilde{\mathfrak{J}}_\alpha$ for $i \geq 2$ when the convergence is under $\mathcal{L}(\mu)$ and the i.i.d sequence of $\tilde{\mathfrak{J}}_\alpha$ when the convergence is under μ_{B_n} . This is enough to get the convergence of the point processes, as discussed in Remark 2.9. We will focus only on the return REPP as the proof is similar for the hitting REPP. It follows the argument used in [PSZ17, (xii)-(xiii)] and we proceed by induction on successive returns. The initialization is exactly Proposition 4.7. Consider now $d \geq 1$. Our goal is to show that the distribution of the $(d + 1)$ -interarrival time is independent of the previous interarrivals and the law is the same. Let $t_0, \dots, t_{d-1}, t_d \in \mathbb{R}_+^*$ and let $M_n := B_n \cap \bigcap_{i=0}^{d-1} \{\gamma(\mu(B_n)) r_{B_n} \circ T_{B_n}^i \leq t_i\}$. We have

$$\begin{aligned} \mu_{B_n}(M_n \cap \{\gamma(\mu(B_n)) r_{B_n} \circ T_{B_n}^d \leq t_d\}) &= \frac{\mu(M_n \cap \{\gamma(\mu(B_n)) r_{B_n} \circ T_{B_n}^d \leq t_d\})}{\mu(B_n)} \\ &= \mu_{B_n}(M_n) \mu_{M_n}(\gamma(\mu(B_n)) r_{B_n} \circ T_{B_n}^d \leq t_d). \end{aligned}$$

By assumption, we know the limit of $\mu_{B_n}(M_n)$. Hence, we need to prove the convergence of $\gamma(\mu(B_n)) r_{B_n} \circ T_{B_n}^d$ under μ_{M_n} . We have

$$\begin{aligned} \mu_{M_n}(\gamma(\mu(B_n)) r_{B_n} \circ T_{B_n}^d \leq t_d) &= \frac{\mu(B_n)}{\mu(M_n)} \int_{B_n} \mathbf{1}_{M_n} \mathbf{1}_{\{\gamma(\mu(B_n)) r_{B_n} \leq t_d\}} \circ T_{B_n}^d \, d\mu_{B_n} \\ &= \int_{B_n} \mu(B_n) \hat{T}_{B_n}^d \left(\frac{\mathbf{1}_{M_n}}{\mu(M_n)} \right) \mathbf{1}_{\{\gamma(\mu(B_n)) r_{B_n} \leq t_d\}} \, d\mu_{B_n}. \end{aligned}$$

Since $B_n = [0(\geq n)]$ is an interval and a union of cylinders build from ξ , the induced map is piecewise and a partition ξ_{B_n} of B_n can naturally be defined. On each element of ξ_{B_n} , r_{B_n} is constant. Furthermore, for all $i \geq 1$, we can define $\xi_{B_n, i} = \bigvee_{k=0}^{i-1} T_{B_n}^{-k} \xi_{B_n}$. In particular, M_n is $\xi_{B_n, d-1}$ measurable. We write κ_n for the element of $\xi_{B_n, d-1}$ contained in M_n . On each $V \in \kappa_n$, there exists some m_V such that $T_{B_n}^d|_V = T^{m_V} : V \rightarrow B_n$ is a homeomorphism and thus, $\hat{T}^{m_V} \mathbf{1}_V \in C_r(B_n)$ by Lemma 4.4. Hence,

$$\hat{T}_{B_n}^d \left(\frac{\mathbf{1}_{M_n}}{\mu(M_n)} \right) = \frac{1}{\mu(M_n)} \sum_{V \in \kappa_n} \hat{T}_{B_n}^d \mathbf{1}_V = \frac{1}{\mu(M_n)} \sum_{V \in \kappa_n} \hat{T}^{m_V} \mathbf{1}_V \in C_r(B_n).$$

Now, writing $u_n := \mu(B_n) \widehat{T}_{B_n}^d \left(\frac{\mathbf{1}_{M_n}}{\mu(M_n)} \right)$ and remarking that $\int u_n d\mu = \mu(B_n)$, it gives, since $\text{supp}(u_n) \subset B_n$, that $\inf_{B_n} u_n \leq 1 \leq \sup_{B_n} u_n$. Since $u_n \in C_r(B_n)$ we have, for every $x, y \in B_n$,

$$u_n(y)(1 - r \text{diam}(B_n)) \leq u_n(x) \leq u_n(y)(1 + r \text{diam}(B_n)).$$

Hence,

$$1 - r \text{diam}(B_n) \leq \inf_{B_n} u_n \leq \sup_{B_n} u_n \leq 1 + r \text{diam}(B_n),$$

which, since $\text{diam}(B_n) \xrightarrow[n \rightarrow +\infty]{} 0$, implies

$$\left\| \mu(B_n) \widehat{T}_{B_n}^d \left(\frac{\mathbf{1}_{M_n}}{\mu(M_n)} \right) - \mathbf{1}_{B_n} \right\|_{L^\infty(\mu_{B_n})} \xrightarrow[n \rightarrow +\infty]{} 0. \quad (4.8)$$

Thus

$$\mu_{M_n}(\gamma(\mu(B_n)) r_{B_n} \circ T_{B_n}^d \leq t_d) \underset{n \rightarrow +\infty}{\sim} \int \mathbf{1}_{\{\gamma(\mu(B_n)) r_{B_n} \leq t_d\}} d\mu_{B_n} \xrightarrow[n \rightarrow +\infty]{} \mathbb{P}(\tilde{\mathfrak{J}}_\alpha \leq t_d),$$

where we used again Proposition 4.7 to get the convergence and this concludes the proof. \square

In particular, Proposition 4.8 is Theorem 2.7 for $1/2$ or, otherwise stated, for $k = 0$. We will now capitalize on the convergence for $1/2$ and go backwards for the further preimages of 0 .

4.3.2 Preimages of $1/2$

Fix $Y = [1/2, 1]$ and for all $n \geq 1$, let $E_n = T_2^{-1}[0, c_n] = [0(\geq n)] = [1/2, \delta_n]$ where $\delta_n := (1 + c_n)/2$. In particular, with the notations chosen, we have for all $0 \leq p \leq n$, $[0[p, n]] = [\delta_{n+1}, \delta_p] = Y \cap \{p+1 \leq r_Y \leq n+1\}$. Fix $k \geq 1$ and let $B_n := T^{-k}E_n$. By invariance of μ we have $\mu(E_n) = \mu(B_n)$. By definition of the map T , for n large enough, we have

$$B_n = \bigsqcup_{z \in T^{-k}\{1/2\}} B_{z,n},$$

where $B_{z,n} = [z, z + \eta_{z,n}]$. With symbolic notations, we have $B_{z,n} = [z_0^{k-1}0(\geq n)]$, where $z = T_{\sigma(a_0)}^{-1} \cdots T_{\sigma(a_{k-1})}^{-1} 1/2$. So, to an element $z \in T^{-k}\{1/2\}$, we can associate a unique sequence (z_0^{k-1}) such that $(z_0^{k-1}0)$ is admissible and reciprocally, for every admissible $(z_0^{k-1}0)$ we can associate $z \in T^{-k}\{1/2\}$. We define the localization map $\phi_n : B_n \rightarrow T^{-k}\{1/2\}$ by $\phi_n(x) = z$ if $x \in B_{z,n}$. In fact, if $n_0 \geq 1$ is such that $(B_{z,n_0})_z$ are disjoint, then $\phi_n = \phi_{n_0}|_{B_n}$ for all $n \geq n_0$ and thus we can consider $\phi := \phi_{n_0}$.

Let $\Psi_n^{\gamma,k}$ be a point process on $\mathbb{R} \times T^{-k}\{1/2\}$ defined by

$$\Psi_n^{\gamma,k} := \sum_{j \geq 1} \delta_{(\gamma(\mu(B_n)) r_{B_n}^{(j)}, \phi \circ T_{B_n}^j)}.$$

Theorem 4.1. *Let $k \geq 1$ and $\Psi_n^{\gamma,k}$ be defined as previously, then we have*

$$\Psi_n^{\gamma,k} \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} P_k,$$

where P_k is an independent \mathbb{Q}_k -marking of $\text{RPP}(\tilde{\mathfrak{J}}_\alpha)$.

Recall that \mathbb{Q}_k is the probability on $T^{-k}\{1/2\}$ such that for ρ the density of μ and all $z \in T^{-k}\{1/2\}$,

$$\mathbb{Q}_k(z) = \frac{\rho(z)}{\rho(1/2)(T^k)'(z)} \quad \forall z \in T^{-k}\{1/2\}.$$

4.3.3 Proof of Theorem 4.1

We start with the following easy observations.

Lemma 4.9. *For all $k \geq 1$, $z \in T^{-k}\{1/2\}$ and all $d \geq 0$, we have $\mu_{B_n}(\phi \circ T_{B_n}^d = z) \sim \mathbb{Q}_k(z)$.*

Proof (of Lemma 4.9). By invariance of the induced measure on B_n , for all $d \geq 0$, we have

$$\begin{aligned} \mu_{B_n}(\phi \circ T_{B_n}^d = z) &= \mu_{B_n}(\phi = z) = \mu_{B_n}(B_{z,n}) = \frac{\mu(B_{z,n})}{\mu(E_n)} \\ &\underset{n \rightarrow +\infty}{\sim} \frac{\rho(z) \text{Leb}(B_{z,n})}{\rho(1/2) \text{Leb}(E_n)} \\ &\underset{n \rightarrow +\infty}{\sim} \frac{\rho(z)}{\rho(1/2)(T^k)'(z)}. \end{aligned}$$

□

Lemma 4.10. *We have*

$$\gamma(\mu(B_n)) r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} \tilde{\mathfrak{J}}_\alpha.$$

Proof (of Lemma 4.10). On E_n , by definition of B_n , we have $r_{B_n} = r_{E_n} - k$. Thus, for all $t > 0$ and by Proposition 4.7,

$$\begin{aligned} \mu_{E_n}(\gamma(\mu(B_n)) r_{B_n} \leq t) &= \mu_{E_n}(\gamma(\mu(B_n)) r_{E_n} \leq t) + \mu_{E_n}(t \leq \gamma(\mu(B_n)) r_{E_n} \leq t + k\gamma(\mu(B_n))) \\ &\xrightarrow[n \rightarrow +\infty]{} \mathbb{P}(\tilde{\mathfrak{J}}_\alpha \leq t). \end{aligned}$$

□

To establish Theorem 4.1, the strategy is to analyze the last visit to the interval $[0[p, n]] = [\delta_n, \delta_p]$ before the system returns to E_n . Leveraging the bounded distortion property (Corollary 4.5), we obtain accurate estimates for the limiting process. Throughout this analysis, it is essential to ensure that the errors introduced by approximations diminish in the limit. This concern is addressed through a series of supporting lemmas and the pivotal Proposition 4.13. To proceed systematically, we begin by introducing the necessary notations.

Fix $k \geq 1$. Let $t > 0$ and take $p \geq 1$ large enough so that $[1/2, \delta_p] \cap \bigcup_{j=1}^k T^{-j}\{1/2\} = \emptyset$. In particular, it implies that for all $z \in T^{-k}\{1/2\}$ and $0 \leq i \leq k-1$, $z_i < p$ (recall that (z_0^{k-1}) is the path in symbolic notations that corresponds to z as a k -preimage of $1/2$). Let

$$D_n^p(t) := E_n \cap \left\{ r_{[\delta_n, \delta_p]} \leq \frac{t}{\gamma(\mu(B_n))} \right\} \cap \{r_{[\delta_n, \delta_p]} \leq r_{E_n}\},$$

and let $\kappa_n^p(t)$ be a collection of branches belonging to $D_n^p(t)$ defined as follows

$$\begin{aligned} \kappa_n^p(t) &:= \{V \subset W \in \xi_m \cap E_n \mid T^m(V) = [\delta_n, \delta_p], m \leq t/\gamma(\mu(B_n)) \text{ and } \mu(T^j V \cap E_n) = 0 \forall 0 \leq j \leq m\} \\ &= \{[0a_0^{m-2}0[p, n]] \mid 2 \leq m \leq t/\gamma(\mu(B_n)), a_0 \geq n, \text{ and } \exists! 0 \leq j \leq m-2, a_j = n\}. \end{aligned}$$

For $V \in \kappa_n^p(t)$, we write m_V its associated m coming from the definition of V and $\kappa_n^p(t)$. Note that $\kappa_n^p(t)$ is not a partition of $D_n^p(t)$ as multiple returns to $[\delta_n, \delta_p]$ are possible but we can build a partition of the sets

$$D_{n,z}^p(t) = D_n^p(t) \cap \left\{ \max_{k \geq 1} \left\{ r_{[\delta_n, \delta_p]}^{(k)} \mid r_{[\delta_n, \delta_p]}^{(k)} < r_{E_n} \right\} \leq \frac{t}{\gamma(\mu(B_n))} \right\} \cap \{r_{E_n} = r_{B_{n,z}} + k\}, \quad z \in T^{-k}\{1/2\}.$$

Indeed, for $V \in \kappa_n^p(t)$ and $z \in T^{-k}\{1/2\}$, define

$$V_z := \{x \in V \mid r_{B_{n,z}}(T^{m_V} x) = r_{B_n}(T^{m_V} x) < r_{[\delta_n, \delta_p]}(T^{m_V} x)\}.$$

Then $\kappa_{n,z}^p(t) := \{V_z, V \in \kappa_n^p(t)\}$ is a partition of $D_{n,z}^p(t)$. To be more precise, for all $V = [0a_0^{m_V-2}0[p, n]] \in \kappa_n^p(t)$, we have

$$V_z = \bigsqcup_{\underline{b} \in \mathcal{I}_z^p} [0a_0^{m_V-2}0\underline{b}z_0^{k-1}0(\geq n)]$$

with $\mathcal{I}_z^p := \{\underline{b} := (b_0^{\ell-1}) \mid \ell \geq 1, (0b_0^{\ell-1}z_0) \text{ admissible}, b_0 \in [p, n] \text{ and } \exists! 0 \leq j < \ell, b_j = p\}$.

The following lemma ensures that the measure of the image $T^{m_V} V_z$ will be comparable to the one of $B_{n,z}$ for all $V_z \in \kappa_{n,z}^p(t)$.

Lemma 4.11. For all $n > p$, $z \in T^{-k}\{1/2\}$ and $V_z \in \kappa_{n,z}^p(t)$,

$$\mu(T^{mv} V_z) = \mu(B_{n,z}) + O(\mu(B_n)^2)$$

Furthermore, $(T^{mv} V_z)_z$ are disjoint and for $V_1 := \bigsqcup_{z \in T^{-k}\{1/2\}} V_z$, we have

$$\mu(T^{mv} V_1) = \mu(B_n) + O(\mu(B_n)^2).$$

Proof (of Lemma 4.11). For all $n > p$ and $z \in T^{-k}\{1/2\}$, we have by ergodicity and conservativity

$$\begin{aligned} \mu(B_{n,z}) &= \sum_{k=0}^{+\infty} \mu([\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} > k\} \cap T^{-k} B_{n,z}) \\ &= \sum_{k=0}^{+\infty} \sum_{\ell=k+1}^{+\infty} \mu([\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} = \ell\} \cap T^{-k} B_{n,z}) \\ &= \sum_{\ell=1}^{+\infty} \sum_{k=0}^{\ell-1} \mu([\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} = \ell\} \cap T^{-k} B_{n,z}). \end{aligned}$$

On another hand, we have

$$\begin{aligned} \mu(T^{mv} V_z) &= \mu([\delta_n, \delta_p] \cap \{r_{B_n} = r_{B_{n,z}} < r_{[\delta_n, \delta_p]}\}) \\ &= \mu([\delta_n, \delta_p] \cap \{r_{B_{n,z}} < r_{[\delta_n, \delta_p]}\}) - \mu([\delta_p, \delta_n] \cap \{r_{B_n} < r_{B_{n,z}} < r_{[\delta_p, \delta_n]}\}) \\ &= \sum_{\ell=1}^{+\infty} \mu\left(\bigcup_{k=0}^{\ell-1} [\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} = \ell\} \cap T^{-k} B_{n,z}\right) - \mu([\delta_p, \delta_n] \cap \{r_{B_n} < r_{B_{n,z}} < r_{[\delta_p, \delta_n]}\}). \end{aligned}$$

We have

$$\mu([\delta_p, \delta_n] \cap \{r_{B_n} < r_{B_{n,z}} < r_{[\delta_p, \delta_n]}\}) \leq \mu([\delta_p, \delta_n] \cap \{r_{B_n}^{(2)} < r_{[\delta_n, \delta_p]}\}).$$

Consider the partition of $[\delta_p, \delta_n] \cap \{r_{B_n}^{(2)} < r_{[\delta_p, \delta_n]}\} = [\delta_p, \delta_n] \cap \{r_{E_n}^{(2)} < r_{[\delta_p, \delta_n]}\}$ defined as follows

$$\begin{aligned} G_n^p &:= \{W \subset [\delta_p, \delta_n] \mid \exists q \geq 0, T^q W = [1/2, \delta_n], \\ &\quad \exists! 1 \leq j < q, T^j W \in [1/2, \delta_n], \forall 1 \leq \ell \leq q, T^\ell W \cap [\delta_n, \delta_p] = \emptyset\} \\ &= \{[0b_0^{\ell-1} 0c_0^{m-1} 0(\geq n)] \mid \ell, m \geq 1, b_0 \in [p, n), \\ &\quad c_0 \geq n, \exists! 0 \leq j < \ell b_j = p, \exists! 0 \leq i < m c_i = p\}. \end{aligned}$$

Then, for every $W = [0b_0^{\ell-1} 0c_0^{m-1} 0(\geq n)] \in G_n^p$, we can use Corollary 4.5 to get

$$\begin{aligned} \mu(W) &= \mu[0b_0^{\ell-1} 0c_0^{m-1} 0(\geq n)] \leq C \frac{\mu[0b_0^{\ell-1} 0c_0^{m-1} 0] \mu[0(\geq n)]}{\mu[0]} \\ &\leq C^2 \mu(B_n) \frac{\mu[0b_0^{\ell-1} 0c_{i+p+1}^{m-1} 0] \mu[0c_0^{m-1} 0]}{\mu[0c_{i+p+1}^{m-1} 0]} \\ &\leq C^2 \mu(B_n) \frac{\mu[0b_0^{\ell-1} 0c_{i+p+1}^{m-1} 0]}{\mu[0c_{i+p+1}^{m-1} 0]} C \frac{\mu[0c_{i+p+1}^{m-1} 0] \mu[0c_0^{i+p-1} 0]}{\mu[0]} \\ &\leq C^3 \mu(B_n) \mu[0b_0^{\ell-1} 0c_{i+p+1}^{m-1} 0] \mu[0c_0^{i+p-1} 0]. \end{aligned}$$

Thus, summing on $W \in G_n^p$ we get

$$\begin{aligned} \mu([\delta_n, \delta_p] \cap \{r_{B_n}^{(2)} < r_{[\delta_n, \delta_p]}\}) &= \sum_{W \in G_n^p} \mu(W) \\ &\leq \sum C^3 \mu(B_n) \mu[0b_0^{\ell-1} 0c_{i+p+1}^{m-1} 0] \mu[0c_0^{i+p-1} 0] \\ &\leq C^3 \mu(B_n)^2 \mu([\delta_n, \delta_p]) = O(\mu(B_n)^2). \end{aligned}$$

Going back to the computation of $\mu(B_{n,z}) - \mu(T^{mv} V_z)$, we have

$$\mu(B_{n,z}) - \mu(T^{mv} V_z) = \sum_{\ell=1}^{+\infty} \mu\left(\bigcup_{0 \leq j < k \leq \ell-1} [\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} = \ell\} \cap T^{-j} B_{n,z} \cap T^{-k} B_{n,z}\right) + O(\mu(B_n)^2)$$

$$= \mu \left([\delta_n, \delta_p] \cap \{r_{B_{n,z}}^{(2)} < r_{[\delta_n, \delta_p]}\} \right) + O(\mu(B_n)^2) = O(\mu(B_n)^2),$$

using the same argument and Corollary 4.5 as before.

Now, by definition and by the Markov property, we have for all $V = [0a_0^{m-2}0[p, n]] \in \kappa_n^p(t)$

$$T^{mV}V_z = \bigsqcup_{b \in \mathcal{I}_z^p} [0bz_0^{k-1}0(\geq n)].$$

It is easy to see that $(T^{mV}V_z)_z$ are all disjoint and thus

$$\begin{aligned} \mu(T^{mV}V_1) &= \sum_{z \in T^{-k}\{1/2\}} \mu(T^{mV}V_z) = \sum_{z \in T^{-k}\{1/2\}} \mu(B_{n,z}) + O(\mu(B_n)^2) \\ &= \mu(B_n) + O(\mu(B_n)^2). \end{aligned}$$

□

Now, let $M_{n,z}(t) := E_n \cap \{r_{B_n} \leq t/\gamma(\mu(B_n))\} \cap T_{B_n}^{-1}(B_{n,z})$ be the quantity we want to control. The next lemma ensures that we can approximate it well by $D_{n,z}^p(t)$.

Lemma 4.12. *For all $t > 0$, $p \geq 1$ large enough so that $[1/2, \delta_p] \cap \bigcup_{j=1}^k T^{-j}\{1/2\}$ and $z \in T^{-k}\{1/2\}$ we have*

$$\mu(D_{n,z}^p(t) \triangle M_{n,z}(t)) + o(\mu(B_n)).$$

Proof (of Lemma 4.12). We have the following partition $\eta_{n,z}(t)$ of our set $M_{n,z}(t)$

$$\begin{aligned} \eta_{n,z}(t) &:= \left\{ V \subset W \in \xi_{m+1} \cap E_n \mid m \leq t/\gamma(\mu(B_n)), \right. \\ &\quad \left. T^m V = B_{n,z}, \mu(T^k V \cap [1/2, \delta_n]) = 0 \forall 1 \leq k < m \right\} \\ &= \left\{ [0a_0^{m-2}z_0^{k-1}0(\geq n)] \mid m \leq t/\gamma(\mu(B_n)), \right. \\ &\quad \left. a_0 \geq n, (a_0^{m-2}z_0) \text{ admissible and } \exists! 0 \leq j \leq m-2, a_j = n \right\}. \end{aligned}$$

On the other side, recall that

$$\begin{aligned} \kappa_{n,z}^p(t) &= \left\{ [0a_0^{m-2}0[p, n]] \cap T^{-m}\{r_{B_{n,z}} = r_{B_n} < r_{[0[p, n]]}\} \mid \right. \\ &\quad \left. m \leq t/\gamma(\mu(B_n)), a_0 \geq n \text{ and } \exists! 0 \leq j \leq m-2, a_j = n \right\}. \end{aligned}$$

We start by controlling $M_{n,z}(t) \setminus D_{n,z}^p(t)$. For $V = [0a_0^{m-2}z_0^{k-1}0(\geq n)] \in \eta_{n,z}(t)$ such that $r_{[\delta_n, \delta_p]} < r_{B_n} = r_{B_{n,z}} = m$, let $j = \max\{i \leq m \mid T^i x \in [\delta_n, \delta_p]\}$. Then $V \subset W = [0a_0^{j-2}0[p, n]] \cap T^{-j}\{r_{B_{n,z}} = r_{B_n} < r_{[\delta_n, \delta_p]}\} \in \kappa_{n,z}^p(t)$. Thus, it only remains the intervals for which $r_{[\delta_n, \delta_p]} > r_{B_n}$, *i.e.*

$$M_{n,z}(t) \setminus D_{n,z}^p(t) \subset M_{n,z}(t) \cap \{r_{B_n} < r_{[\delta_n, \delta_p]}\} \subset E_n \cap \{r_{E_n} < r_{[\delta_n, \delta_p]}\}.$$

We already know that $\mu_{E_n}(r_{E_n} < r_{[\delta_n, \delta_p]}) = o(1)$, so

$$\mu(M_{n,z}(t) \setminus D_{n,z}^p(t)) = o(\mu(B_n)).$$

Now, we need to control $D_{n,z}^p(t) \setminus M_{n,z}(t)$. This is more difficult as we allow for more branches in $D_{n,z}^p(t)$ because we do not impose any time control for returns to B_n . For every $V_z = [0a_0^{m-2}0[p, n]] \cap T^{-m}\{r_{B_{n,z}} = r_{B_n} < r_{[0[p, n]]}\} \in \kappa_{n,z}^p(t)$, we can build a partition $\kappa_{n,z}^{p,V_z}(t)$ of V_z taking cylinders up to the return time to $B_{n,z}$.

$$\begin{aligned} \kappa_{n,z}^{p,V_z}(t) &:= \left\{ [0a_0^{m-2}0b_0^{j-1}z_0^{k-1}0(\geq n)] \mid m \leq \frac{t}{\gamma(\mu(B_n))}, \right. \\ &\quad \left. b_0 \in [p, n), (b_0^{j-1}z_0^{k-1}) \text{ admissible, } \exists! 0 \leq \ell < j, b_\ell = p \right\}. \end{aligned}$$

Observe that $W \in \kappa_{n,z}^{p,V_z}(t)$ is an interval included in E_n . Then, it gives another partition $\kappa_{n,z}^{p,ex}(t)$ of $D_{n,z}^p(t)$ by

$$\kappa_{n,z}^{p,ex}(t) := \bigcup_{V_z \in \kappa_{n,z}^p(t)} \kappa_{n,z}^{p,V_z}(t).$$

Consider the subset $\kappa_{n,z}^{p,long}(t)$ of $\kappa_{n,z}^{p,ex}(t)$ defined by

$$\kappa_{n,z}^{p,long}(t) := \left\{ W = [0a_0^{m-2}0b_0^{j-1}z_0^{k-1}0(\geq n)] \in \kappa_{n,z}^{p,ex}(t) \mid m+j+1 > t/\gamma(\mu(B_n)) \right\}.$$

By construction, we have

$$D_{n,z}^p(t) \setminus M_{n,z}(t) = \bigsqcup_{W \in \kappa_{n,z}^{p,long}(t)} W.$$

For every sequence $(y_n)_n$ going to $+\infty$ such that $y_n = o(\gamma(\mu(B_n))^{-1})$, we have

$$\begin{aligned} \mu(D_{n,z}^p(t) \setminus M_{n,z}(t)) &= \sum_{W \in \kappa_{n,z}^{p,long}(t), r_{E_n} - m > y_n} \mu(W) + \sum_{W \in \kappa_{n,z}^{p,long}(t), r_{E_n} - m \leq y_n} \mu(W) \\ &\leq \sum_{W \in \kappa_{n,z}^{p,long}(t), r_{E_n} - m > y_n} \mu[0a_0^{m-2}0b_0^{j-1}z_0^{k-1}0(\geq n)] \\ &\quad + \mu(E_n \cap \{t/\gamma(\mu(B_n)) \leq r_{E_n} \leq t/\gamma(\mu(B_n)) + y_n\}) \\ &\leq C \sum_{W \in \kappa_{n,z}^{p,long}(t), r_{E_n} - m > y_n} \frac{\mu[0a_0^{m-2}0(\geq n)]\mu[0b_0^{j-1}z_0^{k-1}0(\geq n)]}{\mu[0(\geq n)]} + o(\mu(B_n)), \end{aligned}$$

where we used Corollary 4.5 and Proposition 4.7. Again with Corollary 4.5, we get

$$\begin{aligned} \mu(D_{n,z}^p(t) \setminus M_{n,z}(t)) &\leq C^2 \sum_{W \in \kappa_{n,z}^{p,long}(t), r_{E_n} - m > y_n} \frac{\mu[0a_0^{m-2}0(\geq n)]}{\mu[0(\geq n)]} \frac{\mu[0b_0^{j-1}z_0^{k-1}0] \mu[0(\geq n)]}{\mu[0]} + o(\mu(B_n)) \\ &\leq C^2 \sum_{W \in \kappa_{n,z}^{p,long}(t), r_{E_n} - m > y_n} \mu[0a_0^{m-2}0(\geq n)] \mu[0b_0^{j-1}z_0^{k-1}0] + o(\mu(B_n)). \end{aligned}$$

At this point, we can split the sum into two sums on the possible a_0^{m-2} and the possible b_0^{j-1} . (b_0^{j-1}) is such that $b_0 \in [p, n)$, ($b_0^{j-1}z_0^{k-1}$) is admissible and $\exists! 0 \leq \ell < j$, $b_\ell = p$. Since $r_{E_n} - m > y_n$, this implies

$$\sum_{b_0^{j-1}} \mu[0b_0^{j-1}z_0^{k-1}0] \leq \mu([\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} > y_n\}).$$

Yet, for all $n > p$, $\mu([\delta_n, \delta_p]) > \mu([\delta_{p+1}, \delta_p]) > 0$ and thus [RZ20, Proposition 2.1 b)] ensures that $\mu([\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} > y_n\}) = o(1)$ as $y_n \xrightarrow{n \rightarrow +\infty} +\infty$.

On the other side, we have (a_0^{m-2}) that must be such that ($a_0^{m-2}0$) is admissible, $m \leq t/\gamma(\mu(B_n))$, $a_0 \geq n$ and $\exists! 0 \leq \ell < m-1$, $a_\ell = n$. Thus,

$$\sum_{a_0^{m-2}} \mu[0a_0^{m-2}0(\geq n)] \leq \mu\left(E_n \cap \left\{r_{E_n} \leq \frac{t}{\gamma(\mu(B_n))}\right\}\right).$$

By Proposition 4.7, since $t > 0$, we know that

$$\mu\left(E_n \cap \left\{r_{E_n} \leq \frac{t}{\gamma(\mu(B_n))}\right\}\right) \lesssim \mu(E_n) = \mu(B_n).$$

Hence, we have

$$\begin{aligned} \mu(D_{n,z}^p(t) \setminus M_{n,z}(t)) &\lesssim C^2 \mu([\delta_n, \delta_p] \cap \{r_{[\delta_n, \delta_p]} > y_n\}) \mu(B_n) + o(\mu(B_n)) \\ &= o(\mu(B_n)) \end{aligned}$$

and it concludes the proof of the lemma. \square

Let $M_n(t) := E_n \cap \{\gamma(\mu(B_n)) r_{B_n} \leq t\} = \bigsqcup_{z \in T^{-k}\{1/2\}} M_{n,z}(t)$.

Proposition 4.13. *For $z \in T^{-k}\{1/2\}$ and all $t > 0$, we have*

$$\mu_{M_n(t)}(T_{B_n}^{-1} B_{n,z}) = \frac{\mu(M_{n,z}(t))}{\mu(M_n(t))} \xrightarrow{n \rightarrow +\infty} \mathbb{Q}_k(z).$$

Proof (of Proposition 4.13). For every $V \in \kappa_n^p(t)$, let $V_1 := \bigsqcup_{z \in T^{-k}\{1/2\}} V_z$. By Lemma 4.12, for all $p \geq 1$ large enough, $n \geq p$ and $z \in T^{-k}\{1/2\}$, we have

$$\begin{aligned} \mu(M_{n,z}(t)) + o(\mu(B_n)) &= \mu(D_{n,z}^p(t)) = \sum_{V \in \kappa_n^p(t)} \mu(V_z) \\ &= (1 \pm C \text{diam}([\delta_n, \delta_p])) \sum_{V \in \kappa_n^p(t)} \mu(V_1) \frac{\mu(T^{mv} V_z)}{\mu(T^{mv} V_1)}, \end{aligned}$$

where we use the bounded distortion Corollary 4.5 for the last inequality. Hence, using Lemma 4.11, for all $z \in T^{-k}\{1/2\}$

$$\mu(M_{n,z}(t)) + o(\mu(B_n)) = (1 \pm C \text{diam}([\delta_n, \delta_p])) \sum_{V \in \kappa_n^p(t)} \mu(V_1) \frac{\mu(B_{n,z}) + O(\mu(B_n)^2)}{\mu(B_n) + O(\mu(B_n)^2)}$$

and

$$\begin{aligned} \sum_{V \in \kappa_n^p(t)} \mu(V_1) &= \sum_{z' \in T^{-k}\{1/2\}} \sum_{V_{z'} \in \kappa_{n,z'}^p(t)} \mu(V_{z'}) \\ &= \sum_{z' \in T^{-k}\{1/2\}} \mu(D_{n,z'}^p(t)) \\ &= \sum_{z' \in T^{-k}\{1/2\}} (\mu(M_{n,z'}(t)) + o(\mu(B_n))) \\ &= (\mu(M_n(t)) + o(\mu(B_n))). \end{aligned}$$

Thus, with Lemma 4.9, for all p large enough, we have

$$\mu(M_{n,z}(t))/\mu(M_n(t)) + o(1) = (1 \pm C \text{diam}([\delta_n, \delta_p])) \frac{\mu(B_{n,z}) + O(\mu(B_n)^2)}{\mu(B_n) + O(\mu(B_n)^2)} (1 + o(1))$$

using Lemma 4.10 to ensure that $o(\mu(B_n)) = o(\mu(M_n(t)))$.

Hence, for all $t > 0$, we have

$$\limsup_{n \rightarrow +\infty} \frac{M_{n,z}(t)}{M_n(t)} \leq (1 + C \text{diam}([1/2, \delta_p])) \lim_{n \rightarrow +\infty} \frac{\mu(B_{n,z})}{\mu(B_n)} \leq (1 + C \text{diam}([1/2, \delta_p])) \mathbb{Q}_k(z).$$

We have a similar control for the lim inf. Since it was true for p large enough, taking $p \rightarrow +\infty$ concludes the proof of Proposition 4.13. \square

Building up on all these lemmas, we are now able to prove Theorem 4.1.

Proof (of Theorem 4.1). To get the convergence of the marking process, it is enough to get the convergence of the finite dimensional marginal of the stochastic process

$$((\gamma(\mu(B_n))) r_{B_n} \circ T_{B_n}^i, \phi \circ T_{B_n}^{i+1})_{i \geq 0}.$$

Let $d \geq 1$. We want to show

$$((\gamma(\mu(B_n))) r_{B_n} \circ T_{B_n}^i, \phi \circ T_{B_n}^{i+1})_{0 \leq i \leq d} \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} \left((\tilde{\mathfrak{J}}_\alpha^{(i)}, Y^{(i)})_{0 \leq i \leq d} \right), \quad (4.9)$$

where $\left((\tilde{\mathfrak{J}}_\alpha^{(i)}, Y^{(i)})_{0 \leq i \leq d} \right)$ are i.i.d, $\tilde{\mathfrak{J}}_\alpha^{(1)} \stackrel{\text{law}}{=} \tilde{\mathfrak{J}}_\alpha$, $Y^{(1)} \stackrel{\text{law}}{=} \mathbb{Q}_k$ and $\tilde{\mathfrak{J}}_\alpha^{(1)}$ and $Y^{(1)}$ are independent.

As we did for the point 1/2 in Proposition 4.8 and building up again on [PSZ17, (xii)-(xiii)], we prove the result by induction. Proposition 4.13 together with Lemma 4.10 gives

$$(\gamma(\mu(B_n)) r_{B_n}, \phi \circ T_{B_n}) \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} (\tilde{\mathfrak{J}}_\alpha, Y), \quad (4.10)$$

where $Y \stackrel{\text{law}}{=} \mathbb{Q}$ and $\tilde{\mathfrak{J}}_\alpha$ and Y are independent, proving the result for $d = 0$.

Now, we do the inductive step from d to $d + 1$. By invariance of the measure, for all $k \geq 1$, we have

$$(\gamma(\mu(B_n)) r_{B_n} \circ T_{E_n}^k, \phi \circ T_{B_n}^{k+1}) \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} (\tilde{\mathfrak{J}}_\alpha, Y),$$

using that $T_{B_n} \circ T_{E_n}^i = T_{B_n}^{i+1}$ on E_n for all $i \geq 0$. Let $t_0, \dots, t_d > 0$ and $z^{(0)}, \dots, z^{(d)} \in T^{-k}\{1/2\}$. Let

$$M_n := E_n \cap \bigcap_{i=0}^d \{\gamma(\mu(B_n)) r_{B_n} \circ T_{E_n}^i \leq t_i\} \cap \{\phi \circ T_{B_n}^{i+1} = z^{(i)}\}.$$

Let $t > 0$ and $z^{(d+1)} \in T^{-k}\{1/2\}$. We show

$$\mu_{M_n} \left(\{\gamma(\mu(B_n)) r_{B_n} \circ T_{E_n}^{d+1} \leq t\} \cap \{\phi \circ T_{B_n}^{d+2} = z^{(d+1)}\} \right) \xrightarrow[n \rightarrow +\infty]{} \mathbb{P}(\tilde{\mathfrak{J}}_\alpha \leq t) \mathbb{Q}_k(z^{(d+1)}).$$

Because E_n is an union of 1-cylinders, we can give a partition η_n in cylinders of M_n and a partition η'_n of $N_n := M_n \cap \{\gamma(\mu(B_n)) r_{B_n} \circ T_{E_n}^{d+1} \leq t\} \cap \{\phi \circ T_{B_n}^{d+2} = z^{(d+1)}\}$. Furthermore, associated to the partition η_n , we set \mathcal{I}_n such that

$$\eta_n = \{V := [0a_0^{m-1}0(\geq n)] \mid (a_0^{m-1}) \in \mathcal{I}_n\}.$$

This yields

$$\eta'_n = \left\{ [0a_0^{m-1}0b_0^{j-1}0(\geq n)] \mid a_0^{m-1} \in \mathcal{I}_n, b_0 \geq n, \right. \\ \left. \exists! i b_i = n, j \leq t/\gamma(\mu(B_n)), b_{j-1-k}^{j-1} = (z^{(d+1)})_0^{k-1} \right\}.$$

Thus,

$$\begin{aligned} \mu(N_n) &= \sum_{[0a_0^{m-1}0b_0^{j-1}0(\geq n)] \in \eta'_n} \mu[0a_0^{m-1}0b_0^{j-1}0(\geq n)] \\ &= (1 \pm C \text{diam}(E_n)) \sum_{[0a_0^{m-1}0b_0^{j-1}0(\geq n)] \in \eta'_n} \mu[0a_0^{m-1}0(\geq n)] \frac{\mu[0b_0^{j-1}0(\geq n)]}{\mu(E_n)} \\ &= (1 \pm C \text{diam}(E_n)) \mu_{E_n}(\{\gamma(\mu(B_n)) r_{B_n} \leq t\} \cap \{\phi \circ T_{B_n} = z^{(d+1)}\}) \mu(M_n), \end{aligned}$$

where we used Corollary 4.5, meaning that

$$\mu_{M_n}(N_n) \xrightarrow[n \rightarrow +\infty]{} \mathbb{P}(\tilde{\mathfrak{J}}_\alpha \leq t) \mathbb{Q}_k(z^{(d+1)}).$$

Since $r_{B_n} \circ T_{E_n}^i = r_{B_n} \circ T_{B_n}^i - k$ for $i \geq 1$ on E_n and $\gamma(\mu(B_n))k \xrightarrow[n \rightarrow +\infty]{} 0$, it is sufficient to get (4.9) and it concludes the proof of Theorem 4.1. \square

4.3.4 Proof of Theorems 2.7 and 2.9

Proof (of Theorem 2.7). By Theorem 4.1, we have

$$\psi_n^{\gamma, k} = \sum_{j \geq 1} \delta_{(\gamma(\mu(B_n)) r_{B_n}^{(j)}, \phi \circ T_{B_n}^{j+1})} \xrightarrow[n \rightarrow +\infty]{\mu_{E_n}} P_k.$$

Furthermore, for all $z \in T^{-k}\{1/2\}$, we have

$$\left\| \widehat{T}^k \left(\frac{\mathbf{1}_{B_n, z}}{\mu(B_n, z)} \right) - \frac{1}{\mu(E_n)} \mathbf{1}_{E_n} \right\|_{L^\infty(\mu_{E_n})} \xrightarrow[n \rightarrow +\infty]{} 0.$$

Together with $k\gamma(\mu(B_n)) \xrightarrow[n \rightarrow +\infty]{} 0$ and $r_{B_n} \circ T^k = r_{B_n} - k$ on B_n , this is enough to ensure

$$\Psi_n^{\gamma, k} \xrightarrow[n \rightarrow +\infty]{\mu_{B_n, z}} P_k.$$

Thus, we get

$$\sum_{j \geq 1} \delta_{\gamma(\mu(B_n)) r_{B_n, z}^{(j)}} = \sum_{j \geq 1} \delta_{(\gamma(\mu(B_n)) r_{B_n, z}^{(j)}, \phi \circ T_{B_n}^{j+1})} \mathbf{1}_{\{\phi \circ T_{B_n}^{j+1} = z\}} \xrightarrow[n \rightarrow +\infty]{\mu_{B_n, z}} P_k(\cdot \times \{z\}) \stackrel{\text{law}}{=} \text{RPP}(\tilde{\mathfrak{J}}_\alpha)^{(\mathbb{Q}_k(z), 1)}$$

Finally, using that

$$\lim_{n \rightarrow +\infty} \gamma(\mu(B_{n, z})) / \gamma(\mu(B_n)) = \lim_{n \rightarrow +\infty} (\mu(B_{n, z}) / \mu(B_n))^{1/\alpha} = \mathbb{Q}_k(z)^{1/\alpha},$$

we get, for every $k \geq 1$ and $z \in T^{-k}\{1/2\}$,

$$N_{B_n, z}^\gamma \xrightarrow[n \rightarrow +\infty]{\mu_{B_n, z}} \text{RPP}(\tilde{\mathfrak{J}}_\alpha)^{(\mathbb{Q}_k(z), \mathbb{Q}_k(z)^{1/\alpha})}.$$

Therefore, by Corollary 2.1, it also gives

$$N_{B_n, z}^\gamma \xrightarrow[n \rightarrow +\infty]{\mathcal{L}(\mu)} \text{DRPP}(\mathfrak{J}_\alpha, \tilde{\mathfrak{J}}_\alpha)^{(\mathbb{Q}_k(z), \mathbb{Q}_k(z)^{1/\alpha})}.$$

□

Proof (of Theorem 2.9). The proof of Theorem 2.9 is similar to the proof of Theorem 2.7. The laws \mathfrak{J}_α and $\tilde{\mathfrak{J}}_\alpha$ must be changed to the exponential law \mathcal{E} but, apart from that, the proof stays identical and especially within the key Theorem 4.1. Indeed, from Theorem 2.8 and Lemma 2.8, the same proof as for Proposition 4.8, gives this time a renewal point process of exponential waiting times, *i.e.* the homogeneous Poisson process. Building up on it, an equivalent of Theorem 4.1 can be proven the same way. Finally, since the Poisson point process is stable through thinning and rescaling by the same parameter, *i.e.* if N is a Poisson point process and $\tau > 0$, $N^{(\tau, \tau)}$ is again a Poisson process, and since $\alpha = 1$, the limit for the preimages of $1/2$ are again Poisson point processes. □

4.3.5 Proof of Proposition 2.7

Proof (of Proposition 2.7). Recall the formula (2.12) for the Laplace transform of the law $\tilde{\mathfrak{J}}_\alpha$.

$$\mathbb{E} \left[e^{-s \tilde{\mathfrak{J}}_\alpha} \right] = 1 - \frac{s^\alpha}{\Gamma(1 + \alpha)} \left(e^{-sd_\alpha} + sd_\alpha \int_0^1 y^{-\alpha} e^{-d_\alpha s y} dy \right)^{-1}, \quad s \geq 0.$$

In particular, we have

$$\mathbb{E} \left[e^{-s \tilde{\mathfrak{J}}_\alpha} \right] = 1 - \frac{s^\alpha}{\Gamma(1 + \alpha)} + o(s^\alpha).$$

However, for a renewal process $\text{RPP}(X)$ with $\mathbb{E}[e^{-sX}] = 1 - \lambda s^\alpha + o(s^\alpha)$, we have

$$\text{RPP}(X)^{(\tau, \tau^\alpha)} \xrightarrow[\tau \rightarrow 0]{} \text{FPP}_\alpha(\lambda^{-1}).$$

This can be found for example in [GKMR20, §10.4.4 p.355]. Hence, it remains to show that for every sequence $(x_k)_{k \geq 0}$ with $x_k \in T^{-k}\{1/2\}$, $\mathbb{Q}_k(x_k) \xrightarrow[k \rightarrow +\infty]{} 0$ which is equivalent to $\rho(x_k) / (T^k)'(x_k) \xrightarrow[k \rightarrow +\infty]{} 0$. Since on every interval on the form $[\varepsilon, 1]$, ρ is bounded away from $+\infty$ and $T' \geq c(\varepsilon) > 1$, this is enough to show it when $x_k \xrightarrow[k \rightarrow +\infty]{} 0$. We consider the case $x_k = T_1^{-k} 1/2 = c_{k+1}$, the other cases can be considered similarly. Since the return map Y is full-branched Gibbs-Markov, for each branch $T^{k+1} : [\delta_{k+1}, \delta_k]$ is such that there exists $y \in [\delta_{k+1}, \delta_k]$ with $(T^{k+1})'(y) = (2(\delta_k - \delta_{k+1}))^{-1} = (c_k - c_{k+1})^{-1}$ by the mean value theorem. By bounded distortion, it yields $(T^{k+1})'(\delta_{k+1}) \asymp (c_k - c_{k+1})^{-1}$ and since $T'(\delta_{k+1}) = 1/2$, we obtain $(T^k)'(c_{k+1}) \asymp (c_k - c_{k+1})^{-1}$. On the other side, by (2.11), we have $\rho(c_k) \asymp c_{k+1}(c_{k+1} - c_{k+2})^{-1}$ and implying $\rho(c_k) / (T^k)'(c_k) \xrightarrow[k \rightarrow +\infty]{} 0$. □

5 Fractional Poisson processes in infinite ergodic theory: a concluding perspective

In the recent article [PS24] ([Yas18] for the first return), the authors showed a convergence for a recurrence REPP in the infinite measure preserving context. They consider a \mathbb{Z} -extension over a one-sided subshift of finite type, defined as follows: let (Ω, σ, ν) be a topologically mixing subshift of finite type endowed with a Gibbs measure ν and let $h : \Omega \rightarrow \mathbb{Z}$ be a centered integrable Hölder observable. On the phase space $X := \Omega \times \mathbb{Z}$, we define the dynamics T by

$$T : (x, a) \mapsto (\sigma(x), a + h(x)), \quad x \in \Omega, a \in \mathbb{Z}.$$

The σ -finite measure $\mu := \nu \otimes (\sum_{z \in \mathbb{Z}} \delta_z)$ is preserved by T .

Theorem 5.1. [PS24, Theorem 3.10] For all $r > 0$, let N_r be the point process defined as follows

$$N_r(x) := \sum_{k \geq 1} \delta_{\nu(B(x,r))^2 r_{B(x,r) \times \{0\}}^{(k)}} \quad \forall x \in \mathcal{X}.$$

Then, for the vague convergence,

$$N_r \xrightarrow[r \rightarrow 0]{\mu} N \circ L_0 \quad \text{and} \quad N_r \xrightarrow[r \rightarrow 0]{\mu_{B(x,r) \times \{0\}}} \tilde{N} \circ \tilde{L}_0,$$

where N, \tilde{N} are standard Poisson process and L_0, \tilde{L}_0 is local times at 0 of the standard Brownian motion, independent of N, \tilde{N} .

Here, the target set depends on the point x considered. However, the techniques developed could have also given the convergence of the hitting REPP for generic points x and shrinking balls $B_n = B(x, \eta_n)$ for some $\eta_n \xrightarrow[n \rightarrow +\infty]{} 0$. We justify that both results are linked and we think combining the two approaches could provide fruitful insights for future developments.

For simplicity, consider the subshift of finite type on $\Omega := \{-b, \dots, b\}^{\mathbb{N}}$ for some $b > 0$, $\nu = m^{\otimes \mathbb{N}}$ a Bernoulli measure on Ω with m a probability of mean 0 and variance $\sigma^2 = 1$ and $h := \text{pr}_0$ the projection onto the first coordinate. Then the \mathbb{Z} -extension (X, T) of (Ω, σ) models the usual random walk on \mathbb{Z} and we assume furthermore that it is aperiodic. By inducing on $Y := \Omega \times \{0\}$ and using [Aar97, Lemma 3.7.4], it is possible to show that the system is PDE. Furthermore, using classical estimates for random walks, we easily an equivalent of the wandering rate $w_n(Y) := \mu(Y) \mathbb{E}_{\mu_Y} [r_Y \wedge n] = \sum_{k=0}^{n-1} \mu(Y \cap \{r_Y > n\})$ when $n \rightarrow +\infty$.

Lemma 5.1. For $Y := \Omega \times \{0\}$, we have

$$w_n(Y) \underset{n \rightarrow +\infty}{\sim} \frac{2\sqrt{2}}{\sqrt{\pi}} \sqrt{n}.$$

Proof (of Lemma 5.1). Using classical local limit theorems for random walks, we know that $\mu_Y(r_Y > k) = \mathbb{P}_0(r_0 > k) \underset{n \rightarrow +\infty}{\sim} \sqrt{\frac{2}{\pi k}}$. Since it is not summable, we get

$$w_n(Y) = \mu(Y) \sum_{k=0}^{n-1} \mu_Y(r_Y > k) \underset{n \rightarrow +\infty}{\sim} \sum_{k=0}^{n-1} \sqrt{\frac{2}{\pi k}} \underset{n \rightarrow +\infty}{\sim} \frac{2\sqrt{2}}{\sqrt{\pi}} \sqrt{n}.$$

□

Then, by [Aar97, Proposition 3.8.7 p.137], we can identify the normalizing sequence $(a_n)_{n \geq 0}$ as we have

$$a_n \underset{n \rightarrow +\infty}{\sim} \frac{1}{\Gamma(2-\alpha)\Gamma(1+\alpha)} \frac{n}{w_n(Y)}.$$

Since $w_n(Y) \in \text{RV}(1/2)$ we have $\alpha = 1/2$ and thus, using Lemma 5.1, we have

$$a_n \underset{n \rightarrow +\infty}{\sim} \frac{\sqrt{\pi}}{2\sqrt{2}\Gamma(3/2)^2} \sqrt{n} = \frac{4}{\pi} \frac{\sqrt{\pi}}{2\sqrt{2}} \sqrt{n} = \sqrt{\frac{2}{\pi n}},$$

using that $\Gamma(3/2) = \Gamma(1/2)/2$ and $\Gamma(1/2)^2 = \pi$. Now, by definition (2.2) of γ , $\gamma : s \mapsto 2s^2/\pi$ is a suitable scaling. Thus, Theorem 5.1 implies the following, for x generic in $\Omega \times \{0\}$ and $B_n := B(x, r_n) \times \{0\}$ with $r_n \xrightarrow[n \rightarrow +\infty]{} 0$,

$$\sum_{k \geq 1} \delta_{\gamma(\mu(B_n)) r_{B_n}^{(k)}} = \sum_{k \geq 1} \delta_{\frac{2}{\pi} \mu(B_n)^2 r_{B_n}^{(k)}} \xrightarrow[n \rightarrow +\infty]{\mu} N \circ L_0 \left(\frac{2}{\pi} \cdot \right).$$

It turns out this point process is in fact equal to the limit process we expect with the methods developed in the previous sections.

Lemma 5.2. Let N be a standard Poisson point process and L_0 be the local time at 0 of a standard Brownian motion. Then,

$$N \circ L_0 \left(\frac{2}{\pi} \cdot \right) \stackrel{\text{law}}{=} \text{FPP}_{1/2}(\Gamma(1+1/2)).$$

In Definition 2.6, the fractional Poisson process is introduced as a specific type of renewal point process. However, this is not the only perspective; alternative definitions have been shown to be equivalent. One such approach characterizes it as a classical Poisson process where the time parameter is randomly rescaled, as we now explain. Let $\alpha \in (0, 1]$ and let D_α be the standard α -stable subordinator, that is to say an increasing Lévy process such that for all $s, t \geq 0$, $\mathbb{E}[e^{-sD_\alpha(t)}] = \exp(-ts^\alpha)$ (see [Ber96] for more details on Lévy processes and subordinators). Let E^α be the inverse stable subordinator defined as the generalized inverse of D_α , i.e.

$$E^\alpha(t) = D_\alpha^\leftarrow(t) := \inf\{u > 0 \mid D_\alpha(u) > t\} \quad \text{for } t \geq 0.$$

This process E^α is also non decreasing and is the correct random time scaling to define the fractional Poisson process.

Remark 5.1. *In infinite ergodic theory, this process is also defined as the Mittag-Leffler process and appears as a functional limit for averages of integrable observables for infinite CEMPT [OS15, Ser20].*

Proposition 5.3. [MNV11, Theorem 2.2] *Let N be a standard Poisson point process of parameter $\lambda > 0$ and E^α an inverse stable subordinator independent of N . Then, $N \circ E^\alpha$ is a fractional Poisson process of parameter α and λ .*

In particular, for $\alpha = 1/2$, $E^{1/2}$ has the same law as $\sqrt{2}\bar{B}$ where \bar{B} is the supremum of the Brownian motion [App09, Theorem 2.2.9 p.95]. We are now able to show Lemma 5.2 showing that the point process obtained in [PS24, Theorem 3.10] is the fractional Poisson process of parameter $1/2$.

Proof (of Lemma 5.2). We have

$$\begin{aligned} N \circ L_0 \left(\frac{2}{\pi} \cdot \right) &\stackrel{\text{law}}{=} N \circ \bar{B} \left(\frac{2}{\pi} \cdot \right) \stackrel{\text{law}}{=} N \circ \sqrt{\frac{\pi}{2}} \bar{B} \stackrel{\text{law}}{=} N_{\frac{\sqrt{\pi}}{2}} \circ \sqrt{2} \bar{B} \\ &\stackrel{\text{law}}{=} N_{\Gamma(3/2)} \circ E^{1/2} \stackrel{\text{law}}{=} \text{FPP}_{1/2}(\Gamma(1 + 1/2)), \end{aligned}$$

where N_λ is a Poisson point process of parameter λ and every processes and point processes are assumed to be independent. \square

This closes the gap between the two approaches when $\alpha = 1/2$, the right scale when considering a \mathbb{Z} -extension. However, as we have seen, the construction of the fractional Poisson point process $N \circ E^\alpha$ could provide a useful approach for $\alpha \neq 1/2$ in future works.

Funding

DBT is partially supported by CMUP, which is financed by national funds through FCT – Fundação para a Ciência e a Tecnologia, I.P., under the project with reference UIDB/00144/2020. He is also financed by CNRS and École Polytechnique.

References

- [Aar97] Jon Aaronson, *An introduction to infinite ergodic theory*, Mathematical Surveys and Monographs, vol. 50, American Mathematical Society, Providence, RI, 1997. MR 1450400
- [Alv20] José F. Alves, *Nonuniformly hyperbolic attractors—geometric and probabilistic aspects*, Springer Monographs in Mathematics, Springer, Cham, [2020] ©2020. MR 4226156
- [App09] David Applebaum, *Lévy processes and stochastic calculus*, second ed., Cambridge Studies in Advanced Mathematics, vol. 116, Cambridge University Press, Cambridge, 2009. MR 2512800
- [Ber96] Jean Bertoin, *Lévy processes*, vol. 121, Cambridge university press Cambridge, 1996.
- [BGT89] Nicholas H. Bingham, Charles M. Goldie, and Jef L. Teugels, *Regular variation*, no. 27, Cambridge university press, 1989.
- [BTF23] Dylan Bansard-Tresse and Jorge Milhazes Freitas, *Inducing techniques for quantitative recurrence and applications to misiurewicz maps and doubly intermittent maps*, to appear in Annales de l’Institut Henri Poincaré Probabilités Statistiques (2023).
- [BZ01] Xavier Bressaud and Roland Zweimüller, *Non exponential law of entrance times in asymptotically rare events for intermittent maps with infinite invariant measure*, Ann. Henri Poincaré **2** (2001), no. 3, 501–512. MR 1846853

- [CC13] J.-R. Chazottes and P. Collet, *Poisson approximation for the number of visits to balls in non-uniformly hyperbolic dynamical systems*, Ergodic Theory Dynam. Systems **33** (2013), no. 1, 49–80. MR 3009103
- [CG93] Pierre Collet and Antonio Galves, *Statistics of close visits to the indifferent fixed point of an interval map*, J. Statist. Phys. **72** (1993), no. 3-4, 459–478. MR 1239564
- [CI95] Massimo Campanino and Stefano Isola, *Statistical properties of long return times in type I intermittency*, Forum Math. **7** (1995), no. 3, 331–348. MR 1325560
- [DT23] Mark Demers and Mike Todd, *A trichotomy for hitting times and escape rates for a class of unimodal maps*, arXiv preprint arXiv:2309.09624 (2023).
- [FFTV16] Ana Cristina Moreira Freitas, Jorge Milhazes Freitas, Mike Todd, and Sandro Vaienti, *Rare events for the Manneville-Pomeau map*, Stochastic Process. Appl. **126** (2016), no. 11, 3463–3479. MR 3549714
- [GKMR20] Rudolf Gorenflo, Anatoly A. Kilbas, Francesco Mainardi, and Sergei Rogosin, *Mittag-Leffler functions, related topics and applications*, Springer Monographs in Mathematics, Springer, Berlin, 2020, Second edition. MR 4179587
- [Gou04] Sébastien Gouëzel, *Vitesse de décorrélation et théorèmes limites pour les applications non uniformément dilatantes*, Ph.D. thesis, Ph. D. Thesis, Ecole Normale Supérieure, 2004.
- [Hay13] N. T. A. Haydn, *Entry and return times distribution*, Dyn. Syst. **28** (2013), no. 3, 333–353. MR 3170620
- [HLV05] N. Haydn, Y. Lacroix, and S. Vaienti, *Hitting and return times in ergodic dynamical systems*, Ann. Probab. **33** (2005), no. 5, 2043–2050. MR 2165587
- [Kal02] Olav Kallenberg, *Foundations of modern probability*, second ed., Probability and its Applications (New York), Springer-Verlag, New York, 2002. MR 1876169
- [Las03] Nick Laskin, *Fractional Poisson process*, vol. 8, 2003, Chaotic transport and complexity in classical and quantum dynamics, pp. 201–213. MR 2007003
- [LFF⁺16] Valerio Lucarini, Davide Faranda, Ana Cristina Freitas, Jorge Miguel Milhazes Freitas, Mark Holland, Tobias Kuna, Matthew Nicol, Mike Todd, and Sandro Vaienti, *Extremes and recurrence in dynamical systems*, Pure and Applied Mathematics (Hoboken), John Wiley & Sons, Inc., Hoboken, NJ, 2016. MR 3558780
- [LP18] Günter Last and Mathew Penrose, *Lectures on the Poisson process*, Institute of Mathematical Statistics Textbooks, vol. 7, Cambridge University Press, Cambridge, 2018. MR 3791470
- [LSV99] Carlangelo Liverani, Benoît Saussol, and Sandro Vaienti, *A probabilistic approach to intermittency*, Ergodic Theory Dynam. Systems **19** (1999), no. 3, 671–685. MR 1695915
- [Mar17] Jens Marklof, *Entry and return times for semi-flows*, Nonlinearity **30** (2017), no. 2, 810–824. MR 3604363
- [MNV11] Mark M. Meerschaert, Erkan Nane, and P. Vellaisamy, *The fractional Poisson process and the inverse stable subordinator*, Electron. J. Probab. **16** (2011), no. 59, 1600–1620. MR 2835248
- [MS19] Mark M. Meerschaert and Alla Sikorskii, *Stochastic models for fractional calculus*, second ed., De Gruyter Studies in Mathematics, vol. 43, De Gruyter, Berlin, 2019. MR 3971272
- [OS15] Takashi Owada and Gennady Samorodnitsky, *Functional central limit theorem for heavy tailed stationary infinitely divisible processes generated by conservative flows*, Ann. Probab. **43** (2015), no. 1, 240–285. MR 3298473
- [PS10] Françoise Pène and Benoît Saussol, *Back to balls in billiards*, Comm. Math. Phys. **293** (2010), no. 3, 837–866. MR 2566164
- [PS24] ———, *Quantitative recurrence for t, t^{-1} transformation*, Probability Theory and Related Fields (2024).
- [PSZ13] Françoise Pène, Benoît Saussol, and Roland Zweimüller, *Recurrence rates and hitting-time distributions for random walks on the line*, Ann. Probab. **41** (2013), no. 2, 619–635. MR 3077520

- [PSZ17] ———, *Return- and hitting-time limits for rare events of null-recurrent Markov maps*, Ergodic Theory Dynam. Systems **37** (2017), no. 1, 244–276. MR 3590502
- [RZ20] Simon Rechberger and Roland Zweimüller, *Return- and hitting-time distributions of small sets in infinite measure preserving systems*, Ergodic Theory Dynam. Systems **40** (2020), no. 8, 2239–2273. MR 4120779
- [Sar01] Omri M. Sarig, *Thermodynamic formalism for null recurrent potentials*, Israel J. Math. **121** (2001), 285–311. MR 1818392
- [Sau09] Benoît Saussol, *An introduction to quantitative Poincaré recurrence in dynamical systems*, Rev. Math. Phys. **21** (2009), no. 8, 949–979. MR 2568049
- [SB22] Yaofeng Su and Leonid A. Bunimovich, *Poisson approximations and convergence rates for hyperbolic dynamical systems*, Comm. Math. Phys. **390** (2022), no. 1, 113–168. MR 4381186
- [Ser20] Toru Sera, *Functional limit theorem for occupation time processes of intermittent maps*, Nonlinearity **33** (2020), no. 3, 1183–1217. MR 4063962
- [Tha80] Maximilian Thaler, *Estimates of the invariant densities of endomorphisms with indifferent fixed points*, Israel J. Math. **37** (1980), no. 4, 303–314. MR 599464
- [Tha83] ———, *Transformations on $[0, 1]$ with infinite invariant measures*, Israel J. Math. **46** (1983), no. 1–2, 67–96. MR 727023
- [Yas18] Nasab Yassine, *Quantitative recurrence of some dynamical systems preserving an infinite measure in dimension one*, Discrete and Continuous Dynamical Systems **38** (2018), 343–361.
- [Yas24] ———, *Quantitative recurrence for \mathbb{Z} -extension of three-dimensional axiom A flows*, preprint, 2024.
- [You99] Lai-Sang Young, *Recurrence times and rates of mixing*, Israel J. Math. **110** (1999), 153–188. MR 1750438
- [Zwe00] Roland Zweimüller, *Ergodic properties of infinite measure-preserving interval maps with indifferent fixed points*, Ergodic Theory Dynam. Systems **20** (2000), no. 5, 1519–1549. MR 1786727
- [Zwe07a] ———, *Infinite measure preserving transformations with compact first regeneration*, J. Anal. Math. **103** (2007), 93–131. MR 2373265
- [Zwe07b] ———, *Mixing limit theorems for ergodic transformations*, J. Theoret. Probab. **20** (2007), no. 4, 1059–1071. MR 2359068
- [Zwe08] ———, *Waiting for long excursions and close visits to neutral fixed points of null-recurrent ergodic maps*, Fund. Math. **198** (2008), no. 2, 125–138. MR 2369126
- [Zwe16] ———, *The general asymptotic return-time process*, Israel J. Math. **212** (2016), no. 1, 1–36. MR 3504316
- [Zwe19] ———, *Hitting-time limits for some exceptional rare events of ergodic maps*, Stochastic Process. Appl. **129** (2019), no. 5, 1556–1567. MR 3944776
- [Zwe22] ———, *Hitting times and positions in rare events*, Ann. H. Lebesgue **5** (2022), 1361–1415. MR 4526257

A Additional proofs

A.1 Proof of Corollary 2.6

Corollary 2.6 can be easily deduced from Theorem 2.6 and the following Lemma A.1.

Lemma A.1. *Let $(B_n)_{n \in \mathbb{N}}$ and $(\eta_n)_{n \in \mathbb{N}}$ be defined as in Theorem 2.6. Let $Q(B_n) := B_n \cap T^{-1}(B_n^c)$. Then we have*

$$\gamma(\mu(T_2^{-1}B_n)) = \gamma(\mu(Q(B_n))) \underset{n \rightarrow +\infty}{\sim} \left(\frac{1}{\Gamma(1+\alpha)\Gamma(1-\alpha)} \right)^{1/\alpha} \frac{1}{I(\eta_n)}. \quad (\text{A.1})$$

Proof (of Lemma A.1). The first equality follows directly from the T -invariance of μ . Recall (2.11) stating that the density $\rho := d\mu/d\text{Leb}$ is of the special form

$$\rho(x) = h_0(x) \times \frac{x}{x - T_1^{-1}x}, \quad \forall x \in (0, 1],$$

where $h_0 : [0, 1] \rightarrow \mathbb{R}$ is a positive continuous function (see for example [Tha83, Lemma 4]). Furthermore, recall that $\gamma : s \mapsto 1/b(1/s)$ where b is the asymptotic inverse of a . Since we are studying a special AFN-map we know that (see [Zwe00, Theorem 4]):

$$a_n \underset{n \rightarrow +\infty}{\sim} \frac{2(1-\alpha)}{h_0(0)\Gamma(1+\alpha)\Gamma(2-\alpha)\alpha^\alpha} n^\alpha.$$

Hence, we can choose

$$\gamma(s) = \left(\frac{2(1-\alpha)}{h_0(0)\Gamma(1+\alpha)\Gamma(2-\alpha)\alpha^\alpha} \right)^{1/\alpha} s^{1/\alpha}, \quad \forall s \geq 0.$$

On the other part, we have

$$\begin{aligned} \mu(Q(B_n)) &= \int_{T_1^{-1}\eta_n}^{\eta_n} \rho(x) dx \underset{n \rightarrow +\infty}{\sim} h_0(0) \int_{T_1^{-1}\eta_n}^{\eta_n} \frac{x}{x - T_1^{-1}x} dx = h_0(0)\xi_n \frac{\eta_n - T_1^{-1}\eta_n}{\xi_n - T_1^{-1}\xi_n} \quad \text{for some } \xi_n \in [T_1^{-1}\eta_n, \eta_n] \\ &\underset{n \rightarrow +\infty}{\sim} h_0(0)\xi_n \underset{n \rightarrow +\infty}{\sim} h_0(0)\eta_n. \end{aligned}$$

Hence,

$$\gamma(\mu(Q(B_n))) \underset{n \rightarrow +\infty}{\sim} \left(\frac{2(1-\alpha)}{\Gamma(1+\alpha)\Gamma(2-\alpha)\alpha^\alpha} \right)^{1/\alpha} \eta_n^{1/\alpha}. \quad (\text{A.2})$$

On the other side, we have

$$I(\eta_n) = \int_{\eta_n}^1 \frac{dx}{x - T_1^{-1}x} \underset{n \rightarrow +\infty}{\sim} \int_{\eta_n}^1 \frac{dx}{2^{1/\alpha}x^{1/\alpha+1}} = \frac{\alpha}{2^{1/\alpha}} \left(\frac{1}{\eta_n^{1/\alpha}} - 1 \right) \underset{n \rightarrow +\infty}{\sim} \frac{\alpha}{2^{1/\alpha}\eta_n^{1/\alpha}}, \quad (\text{A.3})$$

where we use the fact that $x - T_1^{-1}x \underset{x \rightarrow 0^+}{\sim} 2^{1/\alpha}x^{1/\alpha+1}$ since $T_1^{-1}x = x - 2^{1/\alpha}x^{1/\alpha+1} + o(x^{1/\alpha})$. Putting together (A.2) and (A.3) gives (A.1) using that $\Gamma(2-\alpha) = (1-\alpha)\Gamma(1-\alpha)$. \square

A.2 Proof of Lemma 2.8

Recall that $B_n = [0, c_n]$ and let $E_n := T_2^{-1}B_n = Y \cap \{r_Y > n\} = [1/2, \delta_n]$. Recall that we have chosen the renormalization of μ so that $\mu(Y) = 1$. We know by [Zwe00, Theorem 4] that, in the barely infinite case, we have $a_n \sim \frac{2n}{h_0(0)\log(n)}$ and thus $b_n := \frac{h_0(0)}{2}n \log(n)$ is an asymptotic inverse of $(a_n)_{n \geq 0}$ (recall that $\gamma(s) := b(s^{-1})^{-1}$).

Fix some $p \geq 0$ and assume $n \geq p$ (this is no issue as we are interested by the limit on n). Let $\tilde{r}_{B_n}^{(p)}$ be the random variable defined as follows:

$$\tilde{r}_{B_n}^{(p)} := h_{E_n} \circ T_{E_p} \mathbf{1}_{E_n},$$

where $h_A(x) := \inf\{n \geq 0 \mid T^n(x) \in A\}$.

Lemma A.2. *There exist a constant C_p , with $C_p \xrightarrow{p \rightarrow +\infty} 1$ such that for all $n \geq p$,*

$$C_p^{-1} \mu(E_n) \leq \frac{\mathbb{E}_{\mu_Y}[\tilde{r}_{B_n}^{(p)}]}{\mathbb{E}_{\mu_{E_p}}[r_{B_n}]} \leq C_p \mu(E_n).$$

Proof (of Lemma A.2). As usual, we use the bounded distortion result Corollary 4.5 and work branch by branch. Let

$$A_n := \{[0a_0^{j-1}b_0^k], a_0 > n, \exists! i a_i = p, b_0 = 0, b_1 > p, b_{k-1} = 0, b_k > n, \forall 0 \leq \ell \leq k-1 b_\ell \leq n\}$$

be a partition in branches of $\mathbf{1}_{E_n}$ for $\tilde{r}_{B_n}^{(p)}$. Then, we have

$$\begin{aligned}
\mathbb{E}_{\mu_Y}[\tilde{r}_{B_n}^{(p)}] &= \sum_{[0a_0^{j-1}b_0^k] \in A_n} (k-1)\mu[0a_0^{j-1}b_0^k] \\
&\leq (1 + C \operatorname{diam}(E_p)) \sum_{[0a_0^{j-1}b_0^k] \in A_n} (k-1) \frac{\mu[0a_0^{j-1}0(>p)]\mu[b_0^k]}{\mu[0(>p)]} \\
&\leq C'_p \frac{\mu(E_n)}{\mu(E_p)} \mathbb{E}_{\mu_Y}[h_{E_n} \mathbf{1}_{E_p}] \\
&\leq C'_p \frac{\mu(E_n)}{\mu(E_p)} \mathbb{E}_{\mu_Y}[(r_{B_n} - 1)\mathbf{1}_{E_p}] \\
&\leq C_p \frac{\mu(E_n)}{\mu(E_p)} \mathbb{E}_{\mu_Y}[r_{B_n} \mathbf{1}_{E_p}].
\end{aligned}$$

The lower bound is obtained similarly. Since $\operatorname{diam}(E_p) \xrightarrow{p \rightarrow +\infty} 0$, we have $C_p \xrightarrow{p \rightarrow +\infty} 1$. \square

Lemma A.3. *For a fixed $p \geq 0$, we have the following asymptotic result:*

$$\mathbb{E}_{\mu_Y}[\tilde{r}_{B_n}^{(p)}] \sim \int (r_Y \wedge n) d\mu_Y = w_n(Y).$$

Proof (of Lemma A.3). We look at the Birkhoff's sums for the induced map on Y and consider the map $\ell_n := r_Y \mathbf{1}_{E_n^c} = r_Y \mathbf{1}_{\{r_Y \leq n\}}$. Let $j \geq 1$, $x \in Y$ and assume for simplicity that $T_Y^j x \in E_n$. Write $k(j) := \sup\{i \geq 0 \mid r_{E_n}^{Y,(i)} < j\}$ (in particular, with the hypothesis we made, we have $j = r_{E_n}^{Y,(k(i)+1)}$). We have

$$\begin{aligned}
\frac{1}{j} S_j^Y \ell_n(x) &= \frac{1}{j} \sum_{k=0}^{j-1} r_Y(T_Y^k x) \mathbf{1}_{E_n^c}(T_Y^k x) \\
&= \frac{1}{j} \sum_{k=1}^{k(j)} r_{E_n}^{Y \circ T_{E_n}^k(x)-1} \sum_{i=1}^{k(j)} r_Y(T_Y^i \circ T_{E_n}^k x) + \frac{1}{j} h_{E_n} \\
&= \frac{1}{j} \sum_{k=1}^{k(j)} \left(\sum_{i=r_{E_p}^Y \circ T_{E_n}^k(x)}^{r_{E_n}^Y \circ T_{E_n}^k(x)-1} r_Y(T_Y^i \circ T_{E_n}^k x) + \sum_{i=1}^{r_{E_p}^Y \circ T_{E_n}^k(x)-1} r_Y(T_Y^i \circ T_{E_n}^k x) \right) + \frac{1}{j} h_{E_n}.
\end{aligned}$$

On one side, we have

$$\begin{aligned}
\frac{1}{j} \sum_{k=1}^{k(j)} r_{E_p}^{Y \circ T_{E_n}^k(x)-1} \sum_{i=1}^{k(j)} r_Y(T_Y^i \circ T_{E_n}^k x) &= \frac{1}{j} \sum_{k=1}^{k(j)} h_{E_p} \circ T_Y \circ T_{E_n}^k(x) \\
&= \frac{1}{j} \sum_{k=1}^{j-1} (\mathbf{1}_{E_n} h_{E_p} \circ T_Y) \circ T_Y^k(x) \\
&\sim \frac{1}{j} S_j^Y (\mathbf{1}_{E_n} h_{E_p} \circ T_Y) \xrightarrow{j \rightarrow +\infty} \mathbb{E}_{\mu_Y}[\mathbf{1}_{E_n} h_{E_p} \circ T_Y]
\end{aligned}$$

On the other side, we have

$$\frac{1}{j} \sum_{k=1}^{k(j)} \sum_{i=r_{E_p}^Y \circ T_{E_n}^k(x)}^{r_{E_n}^Y \circ T_{E_n}^k(x)-1} r_Y \circ T_Y^i \circ T_{E_n}^k(x) = \frac{1}{j} \sum_{k=1}^{k(j)} h_{E_n} \circ T_{E_p} \circ T_{E_n}^k(x) \quad \text{using } E_n \subset E_p$$

The h_{E_n} instead of r_{E_n} comes from the possibility that the sum is empty if $r_{E_p}^Y = r_{E_n}^Y$ and the 0 we get is exactly h_{E_n} in that case.

For $\tilde{r}_{B_n}^{(p)}$, we have

$$\frac{1}{j} S_j^Y \tilde{r}_{B_n}^{(p)}(x) = \frac{1}{j} \sum_{k=0}^{j-1} \tilde{r}_{B_n}^{(p)} \circ T_Y^k(x)$$

$$= \frac{1}{j} \sum_{k=1}^{k(j)} h_{E_n} \circ T_{E_p} \circ T_{E_n}^k(x) \quad \text{by definition of } \tilde{r}_{B_n}^{(p)}.$$

Putting everything together, we get by Birkhoff's theorem,

$$\mathbb{E}_{\mu_Y}[\tilde{r}_{B_n}^{(p)}] = \mathbb{E}_{\mu_Y}[\ell_n] - \mathbb{E}_{\mu_Y}[\mathbf{1}_{E_n} h_{E_p} \circ T_Y].$$

In particular, when n goes to $+\infty$, we have the asymptotic equivalence

$$\mathbb{E}_{\mu_Y}[\tilde{r}_{B_n}^{(p)}] \underset{n \rightarrow +\infty}{\sim} \mathbb{E}_{\mu_Y}[\ell_n].$$

Finally, we have

$$\mathbb{E}_{\mu_Y}[\ell_n] = \int_Y r_Y \mathbf{1}_{\{r_Y \leq n\}} d\mu_Y = w_n(Y) - n\mu_Y(r_Y > n).$$

However, in the particular barely infinite case, we have $\mu_Y(r_Y > n) \asymp 1/n$ whereas $w_n(Y) \asymp \log(n)$. Thus,

$$\mathbb{E}_{\mu_Y}[\ell_n] \underset{n \rightarrow +\infty}{\sim} w_n(Y).$$

This yields the wanted result $\mathbb{E}_{\mu_Y}[\tilde{r}_{B_n}^{(p)}] \underset{n \rightarrow +\infty}{\sim} w_n(Y)$. □

Proof (of Lemma 2.8). With Lemma A.2 and A.3, we obtain

$$\begin{aligned} C_p^{-1} \mu(E_n) &\lesssim \frac{w_n(Y)}{\mathbb{E}_{\mu_{E_p}}[r_{B_n}]} \lesssim C_p \mu(E_n) \\ C_p^{-1} &\lesssim \frac{\mathbb{E}_{\mu_Y}[r_{B_n}]^{-1} w_n(Y) / \mu(E_n)}{\mathbb{E}_{\mu_Y}[r_{B_n}]^{-1} \mathbb{E}_{\mu_{E_p}}[r_{B_n}]} \lesssim C_p, \end{aligned}$$

where $u_n \lesssim v_n$ means that $\limsup_{n \rightarrow +\infty} u_n/v_n \leq 1$.

Taking the limit in n (lim sup and lim inf) and since the convergence in Theorem 2.8 is true $\mu_{E_p} \in \mathcal{L}(\mu)$, we get, taking finally the limit $p \rightarrow +\infty$,

$$C_p^{-1} \lesssim \frac{w_n(Y)}{\mu(E_n)} \mathbb{E}_{\mu_Y}[r_{B_n}]^{-1} \lesssim C_p.$$

Taking afterwards $p \rightarrow +\infty$, it yields

$$\frac{w_n(Y)}{\mu(E_n)} \underset{n \rightarrow +\infty}{\sim} \mathbb{E}_{\mu_Y}[r_{B_n}].$$

However,

$$\begin{aligned} \gamma(\mu(E_n))^{-1} &= b(\mu(E_n))^{-1} \underset{n \rightarrow +\infty}{\sim} \frac{h_0(0)}{2\mu(E_n)} \log(\mu(E_n))^{-1} \\ &\underset{n \rightarrow +\infty}{\sim} \frac{h_0(0)}{2\mu(E_n)} \log(n) \underset{n \rightarrow +\infty}{\sim} w_n(Y) / \mu(E_n). \end{aligned}$$

□

Remark A.1. In fact, in Lemma A.3, we actually showed that $\mathbb{E}_{\mu_Y}[\tilde{r}_{B_n}^{(p)}] \underset{n \rightarrow +\infty}{\sim} w_n(Y) - n\mu_Y(Y \cap \{r_Y > n\})$, whenever $p > 1$. In the case $p > 1$, both terms are of same order. Since we also have

$$w_n(Y) = \sum_{k=0}^{n-1} \mu(Y \cap \{r_Y > k\})$$

and we know that $\mu(Y \cap \{r_Y > n\}) \underset{n \rightarrow +\infty}{\sim} Cn^{-\alpha}$ for some constant α , we obtain

$$\mathbb{E}_{\mu_Y}[r_{B_n}] \underset{n \rightarrow +\infty}{\sim} \frac{\alpha}{1-\alpha} n,$$

which can also be proven with Theorem 2.6 which implies

$$\frac{1}{n} r_{B_n} \xrightarrow[n \rightarrow +\infty]{\mu_Y} \mathcal{J}_\alpha \tag{A.4}$$

In particular, since $\mathbb{E}[\mathcal{J}_\alpha] = \alpha/(1-\alpha)$, it also gives $\mathbb{E}_{\mu_Y}[r_{B_n}] \underset{n \rightarrow +\infty}{\sim} \alpha n/(1-\alpha)$.