

Balloon Occlusion and Stent Graft for Aortic Trauma

Julien Ghelfi, Alice Chopin, Jules Greze, Frederic Thony, Pierre Bouzat

▶ To cite this version:

Julien Ghelfi, Alice Chopin, Jules Greze, Frederic Thony, Pierre Bouzat. Balloon Occlusion and Stent Graft for Aortic Trauma. JVIR: Journal of Vascular and Interventional Radiology, 2021, 32 (9), pp.1402. 10.1016/j.jvir.2021.05.031. hal-04821106

HAL Id: hal-04821106 https://hal.science/hal-04821106v1

Submitted on 19 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



TITLE PAGE

Title:

Balloon occlusion and stent-graft for aortic trauma

Author Names:

Julien Ghelfi 1,2

Alice Chopin ³

Jules Greze³

Frederic Thony ²

Pierre Bouzat ^{1,3}

- 1. Grenoble Alps University, F-38000 Grenoble, France
- 2. Department of radiology, Grenoble University Hospital, F-38000, Grenoble, France
- 3. Grenoble Alps Trauma center, Department of anesthesiology and intensive care medicine, Grenoble University Hospital, F-38000, Grenoble, France

Contact Information / Corresponding author:

Julien GHELFI

Address: CHU Grenoble Alpes, Boulevard de la Chantourne, 38700 La Tronche, France

Telephone number: +33 4 76 76 89 09

Fax: +33 4 76 76 88 30

Email: jghelfi@chu-grenoble.fr

Conflict of interest:

None

Balloon occlusion and stent-graft for aortic trauma

A 27-year-old skier fell from a cliff and landed on a frozen river. On admission, he had no traumatic brain injury or penetrating trauma, but focused assessment sonography for trauma showed a large retroperitoneal hematoma. Despite resuscitation and transfusion of 14 units of red blood cells, arterial blood pressure remained 50/30 mm Hg. A 20mm occlusion balloon (Equalizer; Boston Scientific, Natick, Massachusetts) was inflated in the supra celiac aorta, restoring his pressure to 90/40 mm Hg. Aortography showed extravasation from the infra-renal aorta (Figure 1a). The most readily available stent graft, a 14mm x 80mm Wallgraft (Boston Scientific) was deployed (Figure 1b). Postprocedural computed tomography revealed an anterior wall fracture of the second lumbar vertebra (Figure 2). After decompressive laparotomy, the patient was extubated on day 7 and was discharged from the hospital 2 months later, without any balloon-related complications.

Figure legends

Figure 1. (a) after partial deflation of the occlusion balloon (arrowhead), aortography showed a large infra renal aortic injury with rapid extravasation (arrow). (b) after deployment of a covered stent, repeat aortography confirmed successful treatment of the injury, and patency of renal arteries.

Figure 2. Sagittal reconstruction of postprocedural computed tomography angiography revealed fracture of the second lumbar vertebra responsible for the aortic injury (star), and satisfactory placement of the covered stent (arrowhead).





