



HAL
open science

The buckling and clamped plate problems on differential forms

Fida El Chami, Nicolas Ginoux, Georges Habib, Ola Makhoul, Simon Raulot

► **To cite this version:**

Fida El Chami, Nicolas Ginoux, Georges Habib, Ola Makhoul, Simon Raulot. The buckling and clamped plate problems on differential forms. 2024. hal-04823801

HAL Id: hal-04823801

<https://hal.science/hal-04823801v1>

Preprint submitted on 6 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

The buckling and clamped plate problems on differential forms

Fida El Chami^{*1}, Nicolas Ginoux^{†2}, Georges Habib^{‡1,2}, Ola Makhoul^{§1}, and Simon Raulot^{¶3}

¹Lebanese University, Faculty of Sciences II, Department of Mathematics, P.O. Box 90656 Fanar-Matn, Lebanon

²Université de Lorraine, CNRS, IECL, F-57000 Metz

³Univ Rouen Normandie, CNRS, Normandie Univ, LMRS UMR 6085, F-76000 Rouen, France

December 6, 2024

Abstract. We extend the buckling and clamped-plate problems to the context of differential forms on compact Riemannian manifolds with smooth boundary. We characterize their smallest eigenvalues and prove that, in the case of Euclidean domains, their spectra on forms coincide with the spectra of the corresponding problems. We obtain various estimates involving the first eigenvalues of the mentioned problems and the ones of the Hodge Laplacian with respect to Dirichlet and absolute boundary conditions on forms. These estimates generalize previous ones in the case of functions.

1 Introduction

Let (M, g) be an n -dimensional compact Riemannian manifold with smooth boundary ∂M and let ν be the inward unit vector field normal to ∂M . For a smooth function f on M , we consider the following two problems

$$\begin{cases} \Delta^2 f = \Gamma f & \text{on } M \\ f = 0 & \text{on } \partial M \\ \frac{\partial f}{\partial \nu} = 0 & \text{on } \partial M \end{cases} \quad (1)$$

and

$$\begin{cases} \Delta^2 f = \Lambda \Delta f & \text{on } M \\ f = 0 & \text{on } \partial M \\ \frac{\partial f}{\partial \nu} = 0 & \text{on } \partial M \end{cases} \quad (2)$$

called the clamped plate and the buckling problem respectively. Note that $\Delta f = -\text{tr}(\nabla^2 f)$ is the Laplace operator of f and Δ^2 its square, which is sometimes called the bi-Laplace operator. It is well known that these two problems have discrete spectra consisting of eigenvalues of finite multiplicities

$$0 < \Gamma_1 \leq \Gamma_2 \leq \dots \leq \Gamma_k \leq \dots \longrightarrow \infty$$

and

$$0 < \Lambda_1 \leq \Lambda_2 \leq \dots \leq \Lambda_k \leq \dots \longrightarrow \infty,$$

*fchami@ul.edu.lb

†nicolas.ginoux@univ-lorraine.fr

‡ghabib@ul.edu.lb

§ola.makhoul@ul.edu.lb

¶simon.raulot@univ-rouen.fr

where each eigenvalue is repeated according to its multiplicity. Physically, problem (1) describes the vibrations of a clamped plate, whereas problem (2) describes the critical buckling load of a clamped plate subjected to a uniform compressive force around its boundary.

These two problems were studied by numerous authors. In 1955, Payne [13] proved that, if M is a planar domain, then

$$\Lambda_1 \geq \lambda_2,$$

where λ_2 is the second eigenvalue of the Dirichlet problem on M (see also [8] for a corrected proof). In 1996, Ashbaugh and Laugesen showed, in their work in [2], that whenever M is a bounded and connected open subset of the Euclidean space \mathbb{R}^n ,

$$\Lambda_1^2 \geq \Gamma_1 \geq \Lambda_1 \lambda_1 > \lambda_1^2, \quad (3)$$

where λ_1 is the first eigenvalue of the Dirichlet problem on M . In [4], Chen, Cheng, Wang and Xia proved that, if the Ricci curvature is bounded below by $n - 1$, then

$$\Gamma_1 > n\lambda_1 \quad \text{and} \quad \Lambda_1 > n. \quad (4)$$

Ilias and Shouman gave in [12] an estimate relating the first eigenvalue Λ_1 of the buckling problem to μ_1 , the first nonzero eigenvalue of the Neumann problem:

$$\mu_1 < \Lambda_1. \quad (5)$$

In this paper, we first generalize problems (1) and (2) to the context of differential forms on the manifold M . We prove that each problem has a discrete spectrum consisting of a nondecreasing sequence of real eigenvalues of finite multiplicities and the corresponding eigenforms form a Hilbert basis of L^2 -integrable p -forms on M . We also characterize the first eigenvalue of each problem (see Theorems 2.2 and 2.4).

In Section 3, we prove that if (M, g) is a compact domain of the Euclidean space \mathbb{R}^n , the spectra without multiplicities of both problems on p -forms, for $p = 1, \dots, n$, and on functions on M coincide (see Proposition 3.1). This allows, for example, to determine the first eigenvalues of both of the problems on p -forms for the Euclidean ball of arbitrary radius. In the same section, namely in Theorem 3.3, we establish a relationship between the first eigenvalues of the buckling and clamped-plate problems on an arbitrary compact Riemannian manifold (M, g) with smooth boundary. In the same context, we consider the following problem, called the Dirichlet problem on forms,

$$\begin{cases} \Delta\omega &= \lambda\omega & \text{in } M \\ \omega &= 0 & \text{on } \partial M \end{cases} \quad (6)$$

and give estimates relating the first eigenvalues of the buckling, clamped-plate and Dirichlet problems. The estimates that we obtain generalize inequalities (3) to differential forms on M . We prove as well that there exists a connection with the first eigenvalue of the absolute boundary value problem on forms

$$\begin{cases} \Delta\omega &= \mu\omega & \text{on } M \\ \nu \lrcorner \omega &= 0 & \text{on } \partial M \\ \nu \lrcorner d\omega &= 0 & \text{on } \partial M \end{cases} \quad (7)$$

given in Theorem 3.11. This connection allows to extend (5) to differential forms.

Further in the same section, we show that, under the condition that the Weitzenböck curvature operator is bounded below by a positive constant γ , the first eigenvalue $\lambda_{1,p}$ of problem (6) is also bounded below by a quantity depending on γ , see Theorem 3.16. This gives new estimates of the first eigenvalues of the buckling and clamped-plate problems under the same conditions, see Corollary 3.18.

Finally, we end Section 3 by considering domains M of the unit sphere \mathbb{S}^n and derive inequalities relating the first eigenvalues of the buckling and clamped plate problems on forms of different degrees (see Theorem 3.20).

Acknowledgments: This collaboration is supported by the International Emerging Action ‘‘HOPF’’ (Higher Order boundary value Problems for differential Forms) of the French CNRS, which the authors would like to thank. We also acknowledge the support of Universit  de Rouen Normandie, Universit  de Lorraine, and Universit  Libanaise.

2 The buckling and clamped plate problems

In order to make this work self-contained, we collect here some classical and useful formulae in the study of p -forms on manifolds with boundary. In the following, (M^n, g) denotes an n -dimensional compact Riemannian manifold with smooth boundary and ν the inner unit normal to ∂M . Then we recall from [16, Lemma 18] that for any p -form ω on M , we have

$$\iota^*(\nabla_\nu \omega) = \nu \lrcorner d\omega + d(\nu \lrcorner \omega) + S^{[p]}(\iota^* \omega) \quad (8)$$

and

$$\nu \lrcorner \nabla_\nu \omega = \delta^{\partial M}(\iota^* \omega) - \iota^*(\delta \omega) - S^{[p-1]}(\nu \lrcorner \omega) + (n-1)H\nu \lrcorner \omega. \quad (9)$$

Here $\delta^{\partial M}$ denotes the codifferential on ∂M , $S^{[p]}$ is the natural extension as an endomorphism of $\Omega^p(\partial M)$ of the shape operator $S := -\nabla \nu$ of the embedding ι of ∂M in M and $H = \frac{1}{n-1} \text{tr}(S)$ is its mean curvature, see e.g. [16, p. 624].

We are especially interested in p -forms $\omega \in \Omega^p(M)$ which satisfy the boundary condition $\omega|_{\partial M} = 0$. Then, using (8) and (9), it is not difficult to compute that, along ∂M , we have $\iota^* \nabla_\nu \omega = \nu \lrcorner d\omega$ and $\nu \lrcorner \nabla_\nu \omega = -\iota^* \delta \omega$. So it is straightforward to see that

$$\begin{cases} \omega|_{\partial M} = 0 \\ \nabla_\nu \omega|_{\partial M} = 0 \end{cases} \iff \begin{cases} \iota^* \omega = 0 \\ \nu \lrcorner \omega = 0 \\ \iota^* \delta \omega = 0 \\ \nu \lrcorner d\omega = 0. \end{cases} \quad (10)$$

In the following, we will denote by Z the vector space of smooth p -forms on M which satisfy these boundary conditions that is

$$Z := \{\omega \in \Omega^p(M) \mid \omega|_{\partial M} = 0 \text{ and } \nabla_\nu \omega|_{\partial M} = 0\}. \quad (11)$$

In this work, we will also often use integration by parts formulae which are cumbersome to write down in the general frame work of p -forms on manifolds with boundary. However, as we restrict our attention to elements in Z , it may be very useful to observe that in this context they become very simple. In fact, for any $\omega, \omega' \in \Omega^p(M)$, it holds that:

$$\int_M \langle \Delta \omega, \omega' \rangle d\mu = \int_M (\langle d\omega, d\omega' \rangle + \langle \delta \omega, \delta \omega' \rangle) d\mu + \int_{\partial M} (\langle \nu \lrcorner d\omega, \iota^* \omega' \rangle - \langle \iota^* \delta \omega, \nu \lrcorner \omega' \rangle) d\sigma \quad (12)$$

$$= \int_M \langle \omega, \Delta \omega' \rangle d\mu + \int_{\partial M} (\langle \nu \lrcorner d\omega, \iota^* \omega' \rangle - \langle \iota^* \omega, \nu \lrcorner d\omega' \rangle + \langle \nu \lrcorner \omega, \iota^* \delta \omega' \rangle - \langle \iota^* \delta \omega, \nu \lrcorner \omega' \rangle) d\sigma \quad (13)$$

and so we immediately deduce from (10) that if $\omega, \omega' \in Z$ then

$$\int_M \langle \Delta \omega, \omega' \rangle d\mu = \int_M (\langle d\omega, d\omega' \rangle + \langle \delta \omega, \delta \omega' \rangle) d\mu = \int_M \langle \omega, \Delta \omega' \rangle d\mu. \quad (14)$$

Here $d\mu$ (resp. $d\sigma$) denotes the Riemannian measure density of (M^n, g) (resp. ∂M endowed with the induced metric). Now by replacing ω by $\Delta\omega$ in (13), we obtain

$$\begin{aligned} \int_M \langle \Delta^2\omega, \omega' \rangle d\mu &= \int_M \langle \Delta\omega, \Delta\omega' \rangle d\mu + \int_{\partial M} \left(\langle \nu \lrcorner d\Delta\omega, \iota^* \omega' \rangle - \langle \iota^* \Delta\omega, \nu \lrcorner d\omega' \rangle \right) d\sigma \\ &\quad + \int_{\partial M} \left(\langle \nu \lrcorner \Delta\omega, \iota^* \delta\omega' \rangle - \langle \iota^* \delta\Delta\omega, \nu \lrcorner \omega' \rangle \right) d\sigma \end{aligned}$$

and so if $\omega, \omega' \in Z$, we get

$$\int_M \langle \Delta^2\omega, \omega' \rangle d\mu = \int_M \langle \Delta\omega, \Delta\omega' \rangle d\mu. \quad (15)$$

In the following, we will also denote by $(\cdot, \cdot)_{L^2(M)}$ the L^2 -scalar product on $\Omega^p(M)$ and $\|\cdot\|_{L^2(M)}$ its associated norm. We finally notice that the boundary conditions studied here turn out to be elliptic in the sense of Lopatinskiĭ-Shapiro (see [18, Def. 1.6.1]). This was proved in a more general setting by the first four authors (see [6, Lemma 6.1]) and we restate this result in our context.

Lemma 2.1 *Let (M^n, g) be a compact Riemannian manifold with smooth boundary ∂M and let ν be the inward unit normal vector field to the boundary. The following boundary value problem*

$$\begin{cases} \Delta^2\omega &= f & \text{on } M \\ \omega &= \omega_1 & \text{on } \partial M \\ \iota^*\delta\omega &= \omega_2 & \text{on } \partial M \\ \nu \lrcorner d\omega &= \omega_3 & \text{on } \partial M \end{cases}$$

for given $f \in \Omega^p(M)$, $\omega_1 \in \Omega^p(M)|_{\partial M}$, $\omega_2 \in \Omega^{p-1}(\partial M)$, $\omega_3 \in \Omega^p(\partial M)$, is elliptic in the sense of Lopatinskiĭ-Shapiro.

2.1 The buckling problem

The buckling eigenvalue problem on differential forms is

$$\begin{cases} \Delta^2\omega &= \Lambda\Delta\omega & \text{on } M \\ \omega &= 0 & \text{on } \partial M \\ \nabla_\nu\omega|_{\partial M} &= 0 & \text{on } \partial M \end{cases} \quad (16)$$

for some real constant Λ . Let us begin with the following existence result.

Theorem 2.2 *There exists a Hilbert basis of the space of L^2 -integrable p -forms on (M^n, g) consisting of eigenforms solutions of the problem (16) associated to an unbounded and positive sequence of eigenvalues $(\Lambda_{i,p})_{i \geq 1}$. Moreover, each eigenspace has a finite multiplicity and the corresponding eigenforms are smooth. Finally, the first eigenvalue $\Lambda_{1,p}$ is characterized by*

$$\Lambda_{1,p} = \inf \left\{ \frac{\|\Delta\omega\|_{L^2(M)}^2}{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}, \omega \in \Omega^p(M) \setminus \{0\}, \omega|_{\partial M} = 0 \text{ and } \nabla_\nu\omega|_{\partial M} = 0 \right\}. \quad (17)$$

Equality holds if and only if ω is an eigenform associated to the first eigenvalue.

Proof. The proof is classical and so we only recall the main steps. For all p -forms ω, ω' , the two bilinear forms

$$(\omega, \omega')_V := \int_M \langle \Delta\omega, \Delta\omega' \rangle d\mu \quad \text{and} \quad (\omega, \omega')_W := \int_M \left(\langle d\omega, d\omega' \rangle + \langle \delta\omega, \delta\omega' \rangle \right) d\mu$$

define scalar products on Z whose associated norms will be denoted by $\|\cdot\|_V$ and $\|\cdot\|_W$. We will also denote by V and W the completions of Z with respect to these norms. Then one can easily show that there exists a positive constant C such that $\|\cdot\|_W \leq C\|\cdot\|_V$ on Z so that there is a natural bounded linear operator $\mathcal{I}: V \rightarrow W$ extending the identity map on Z . Since by [1] – see [3] for a corrected proof – any closed and co-closed p -form vanishing along ∂M must vanish identically on M , the operator \mathcal{I} is actually injective.

Now let $\mathcal{K}: V \rightarrow V$ be the linear operator defined by

$$(\mathcal{K}\omega, \omega')_V = (\mathcal{I}\omega, \mathcal{I}\omega')_W$$

for all $(\omega, \omega') \in V^2$. By definition, the operator \mathcal{K} is self-adjoint and positive-definite. On the other hand, since from standard elliptic estimates both norms $\|\cdot\|_V$ and $\|\cdot\|_{H^2(M)}$ are equivalent on Z , the Rellich theorem ensures that \mathcal{I} is compact and so is \mathcal{K} . The spectral theorem for positive compact self-adjoint operators applies and yields the existence of a countable Hilbert orthonormal basis $(\omega_i)_{i \geq 1}$ of V associated to a monotonously nonincreasing positive real sequence of eigenvalues of finite multiplicities $(\alpha_{i,p})_{i \geq 1}$ going to 0 such that $\mathcal{K}\omega_i = \alpha_{i,p}\omega_i$ for all $i \geq 1$. Now fixing $i \geq 1$ and using the definition of \mathcal{K} as well as the integration by part formula (14), it can be computed that, for every $\omega \in Z$,

$$\alpha_{i,p}(\Delta\omega_i, \Delta\omega)_{L^2(M)} = (\mathcal{K}\omega_i, \omega)_V = (\omega_i, \omega)_W = (\Delta\omega_i, \omega)_{L^2(M)}.$$

At the same time, we also have by (15) that

$$(\Delta\omega_i, \Delta\omega)_{L^2(M)} = (\Delta^2\omega_i, \omega)_{L^2(M)}$$

for every $\omega \in Z$ and so the previous equality now reads as

$$(\Delta^2\omega_i - \Lambda_{i,p}\Delta\omega_i, \omega)_{L^2(M)} = 0$$

where we let $\Lambda_{i,p} = \frac{1}{\alpha_{i,p}}$. It follows that ω_i is a weak solution of the eigenvalue problem (16) which, by ellipticity (see in Lemma 2.1), is in fact smooth. Thus the form ω_i becomes a smooth eigenform to problem (16) associated with the eigenvalue $\Lambda_{i,p} = \frac{1}{\alpha_{i,p}}$ which is of finite multiplicity, since $\alpha_{i,p}$ is.

Conversely, observe that if there exists a nontrivial solution ω to (16) for a certain $\Lambda \in \mathbb{R}$, then by (15), we have

$$(\omega, \omega')_V = \Lambda (\omega, \omega')_W$$

for every $\omega' \in Z$. Note that $\Lambda > 0$, since otherwise $\Delta^2\omega = 0$ which from (15) implies that $\Delta\omega = 0$ and then $\omega = 0$ by [1] since $\omega|_{\partial M} = 0$. By definition of \mathcal{K} , we have $(\omega, \omega')_V = \Lambda (\mathcal{K}\omega, \omega')_V$ for all $\omega' \in Z$ and hence in V , therefore $\mathcal{K}\omega = \frac{1}{\Lambda}\omega$. This shows that ω is an eigenform of \mathcal{K} associated to the eigenvalue $\alpha = \frac{1}{\Lambda}$.

Finally, given any eigenform ω associated to a positive eigenvalue Λ of (16), we have by formula (15) that

$$\Lambda \int_M \langle \Delta\omega, \omega \rangle d\mu = \int_M |\Delta\omega|^2 d\mu.$$

Applying (14) to the left-hand side of this equality ensures that

$$\Lambda_{1,p} \leq \frac{\|\Delta\omega\|_{L^2(M)}^2}{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}$$

for every such eigenform, with equality for ω associated to $\Lambda_{1,p}$. Finally, if $\omega \in V$, one may write its decomposition in the Hilbert basis $(\omega_i)_{i \geq 1}$ so that

$$\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 = (\mathcal{K}\omega, \omega)_V = \sum_{i \geq 1} \frac{1}{\Lambda_{i,p}} |(\omega, \omega_i)_V|^2 \leq \frac{1}{\Lambda_{1,p}} \sum_{i \geq 1} |(\omega, \omega_i)_V|^2 = \frac{1}{\Lambda_{1,p}} \|\Delta\omega\|_{L^2(M)}^2.$$

This prove the characterization (17) since Z is dense in V . \square

Remark 2.3 *The Hodge \star operator is an isometry commuting with the Laplacian and preserving the boundary conditions in the buckling problem so that $\Lambda_{i,p} = \Lambda_{i,n-p}$ for any $i \geq 1$ and $1 \leq p \leq n$.*

2.2 The clamped plate problem

The clamped plate eigenvalue problem on differential forms is

$$\begin{cases} \Delta^2 \omega & = \Gamma \omega & \text{on } M \\ \omega & = 0 & \text{on } \partial M \\ \nabla_\nu \omega|_{\partial M} & = 0 & \text{on } \partial M \end{cases} \quad (18)$$

for some real constant Γ . As previously, we immediately get the following existence result.

Theorem 2.4 *There exists a Hilbert basis of the space of L^2 -integrable p -forms on (M^n, g) consisting of eigenforms solutions of the problem (18) associated to an unbounded and positive sequence of eigenvalues $(\Gamma_{i,p})_{i \geq 1}$. Moreover, each eigenspace has a finite multiplicity and the corresponding eigenforms are smooth. Finally, the first eigenvalue $\Gamma_{1,p}$ is characterized by*

$$\Gamma_{1,p} = \inf \left\{ \frac{\|\Delta \omega\|_{L^2(M)}^2}{\|\omega\|_{L^2(M)}^2}, \omega \in \Omega^p(M) \setminus \{0\}, \omega|_{\partial M} = 0 \text{ and } \nabla_\nu \omega|_{\partial M} = 0 \right\}. \quad (19)$$

Equality holds if and only if ω is an eigenform associated to the first eigenvalue.

Proof. It is enough to take $(\cdot, \cdot)_W$ to be the $L^2(M)$ -scalar product on $\Omega^p(M)$ in the proof of Theorem 2.2 and then the proof goes the same. Note that if $\omega \in \Omega^p(M)$ is an eigenform associated to $\Gamma_{1,p}$, it follows from (15) that $\Gamma_{1,p} \geq 0$. Moreover if $\Gamma_{1,p} = 0$ then any associated eigenform ω has to be harmonic with $\omega|_{\partial M} = 0$ and so $\omega = 0$ by [1]. In particular, $\Gamma_{1,p} > 0$. \square

Remark 2.5 *As for the buckling problem, the Hodge \star operator preserves the clamped plate problem so that $\Gamma_{i,p} = \Gamma_{i,n-p}$ for any $i \geq 1$ and $1 \leq p \leq n$.*

3 Eigenvalues of the buckling and clamped plate operators

3.1 Eigenvalues for Euclidean domains

In this section, we completely describe the spectrum of the buckling and clamped plate problems for compact domains in the Euclidean space. More precisely, we prove the following characterization.

Proposition 3.1 *Let (M^n, g) be a compact domain in the Euclidean space \mathbb{R}^n . Then the spectrum of the buckling problem on p -forms on (M^n, g) coincide with the spectrum of the buckling problem on functions that is $\Lambda_{i,p} = \Lambda_{i,0}$ for all $i \geq 1$ and $p \in \{1, \dots, n\}$. The same holds for the clamped plate problem that is $\Gamma_{i,p} = \Gamma_{i,0}$ for all $i \geq 1$ and $p \in \{1, \dots, n\}$.*

Proof. First recall that on \mathbb{R}^n , there exists for each $p \in \{1, \dots, n\}$ a maximal number of parallel p -forms.

Fix $p \in \{1, \dots, n\}$ and denote by ω_0 a nontrivial parallel p -form on M . Then note that for any smooth function f on M with $f|_{\partial M} = 0$ and $\frac{\partial f}{\partial \nu}|_{\partial M} = 0$, the p -form $\omega_f := f\omega_0$ satisfies

$$\omega_f|_{\partial M} = 0 \quad \text{and} \quad \nabla_\nu \omega_f|_{\partial M} = 0. \quad (20)$$

On the other hand, since ω_0 is parallel, we have $d\omega_f = df \wedge \omega_0$ and $\delta\omega_f = -df \lrcorner \omega_0$ and therefore $\Delta\omega_f = (\Delta f)\omega_0$. Applying twice this formula leads to

$$\Delta^2\omega_f = \Delta((\Delta f)\omega_0) = (\Delta^2 f)\omega_0. \quad (21)$$

Now if we take f_1 (resp. f_2) to be an eigenfunction for the buckling (resp. clamped plate) problem (2) (resp. (1)) associated with the eigenvalue Λ (resp. Γ), we conclude combining (20) and (21) that ω_{f_1} (resp. ω_{f_2}) is a p -eigenform for (16) (resp. (18)) associated with the eigenvalue Λ (resp. Γ).

Conversely, first note that if $f \in C^\infty(M)$ is a smooth nontrivial function such that $f = \langle \omega, \omega_0 \rangle$ where ω is a smooth p -form and ω_0 is a smooth parallel p -form we have

$$\Delta f = \langle \nabla^* \nabla \omega, \omega_0 \rangle = \langle \Delta \omega, \omega_0 \rangle \quad (22)$$

where the last equality follows from the Bochner formula (see (32) below) and the fact that M is Euclidean. Note also that if ω satisfies the boundary condition (10) then f and $\partial f / \partial \nu$ vanish on ∂M . Now if ω_1 and ω_2 denote respectively p -eigenforms to the problems (16) and (18) associated to the eigenvalues Λ and Γ , then there exist two parallel p -forms ω_0^i on \mathbb{R}^n such that $f_i := \langle \omega_i, \omega_0^i \rangle$ are smooth nontrivial functions for $i = 1, 2$. Therefore we easily deduce from (22) that these functions are smooth eigenfunctions of (2) and (1) respectively associated to Λ and Γ . \square

Remarks 3.2

1. Proposition 3.1 gives immediately the value of the first eigenvalues of the buckling and clamped plate problems on p -forms for the Euclidean ball of radius $R = 1/H_0$. More precisely, it follows from [2, Sec. 1] that

$$\Lambda_{1,p} = j_{\frac{n}{2},1}^2 H_0^2 \quad \text{and} \quad \Gamma_{1,p} = k_{\frac{n}{2}-1,1}^4 H_0^4,$$

where $j_{\frac{n}{2},1}$ is the first positive zero of the Bessel function $J_{\frac{n}{2}}$ and $k_{\frac{n}{2}-1,1}$ is the first positive zero of $J_{\frac{n}{2}-1} I_{\frac{n}{2}} + J_{\frac{n}{2}} I_{\frac{n}{2}-1}$, with I_ℓ being the corresponding modified Bessel function of the first kind.

2. If (M^n, g) is a compact Riemannian manifold carrying a nontrivial parallel p -forms for a certain $p \in \{1, \dots, n\}$, we can mimic the first part of the proof of Proposition 3.1 to ensure that if $\text{Spec}_{\Lambda,p}(M)$ and $\text{Spec}_{\Gamma,p}(M)$ denote respectively the buckling and clamped plate spectra on p -forms for $p \in \{0, \dots, n\}$ then we have

$$\text{Spec}_{\Lambda,0}(M) \subset \text{Spec}_{\Lambda,p}(M) \quad \text{and} \quad \text{Spec}_{\Gamma,0}(M) \subset \text{Spec}_{\Gamma,p}(M)$$

3.2 General estimates

In this section, we prove general estimates between the first eigenvalues of the buckling and the clamped pate problems and other classical problems. These estimates hold on (M^n, g) without any extra assumption, except being a compact Riemannian manifold with smooth boundary.

The first result of this part establishes a direct link between the buckling and the clamped plate problems.

Theorem 3.3 *On an n -dimensional compact Riemannian manifold (M^n, g) with smooth boundary, we have $\Gamma_{1,p} < \Lambda_{1,p}^2$ for all $p \in \{0, \dots, n\}$.*

Before giving the proof of this result, we first note that if ω is a p -form satisfying $\omega|_{\partial M} = \nabla_\nu \omega|_{\partial M} = 0$ then it follows from the identity $\Delta = d\delta + \delta d$ and from an integration by parts that

$$\|\Delta\omega\|_{L^2(M)}^2 = \|\delta d\omega\|_{L^2(M)}^2 + \|d\delta\omega\|_{L^2(M)}^2$$

and from (14) that

$$(\Delta\omega, \omega)_{L^2(M)} = \|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2. \quad (23)$$

Proof. Let ω be a smooth nonzero p -form on M with $\omega|_{\partial M} = 0$ and $\nabla_\nu \omega|_{\partial M} = 0$. By (23) and the Cauchy-Schwarz inequality, we get

$$\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 \leq \|\Delta\omega\|_{L^2(M)} \|\omega\|_{L^2(M)}$$

and this implies that

$$\frac{\|\Delta\omega\|_{L^2(M)}}{\|\omega\|_{L^2(M)}} \leq \frac{\|\Delta\omega\|_{L^2(M)}^2}{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}.$$

Taking ω to be an eigenform of the buckling problem associated to $\Lambda_{1,p}$ gives the desired inequality. Moreover, if equality holds, then for any eigenform ω of the buckling problem associated to $\Lambda_{1,p}$, the p -form $\Delta\omega$ is pointwise proportional to ω , therefore $\Delta\omega = \Lambda_{1,p}\omega$ on M and thus ω is an eigenform of the Dirichlet problem associated to the eigenvalue $\Lambda_{1,p}$. But because ω satisfies $\nabla_\nu \omega|_{\partial M} = 0$ along ∂M , the unique continuation theorem for elliptic second-order linear operators (see e.g. [17, Theorem 1.4] and [11, chap. VIII]) implies that $\omega = 0$ on M . This shows that the inequality must actually be strict. \square

Now we consider $\lambda_{1,p}$ the smallest eigenvalue of the Hodge Laplacian of M with Dirichlet boundary condition. It is well-known that there exists a Hilbert basis of the space of L^2 -integrable p -forms on (M^n, g) consisting of smooth eigenforms solutions of the problem

$$\begin{cases} \Delta\omega &= \lambda\omega & \text{on } M \\ \omega &= 0 & \text{on } \partial M \end{cases} \quad (24)$$

associated to an unbounded and positive sequence of eigenvalues $(\lambda_{i,p})_{i \geq 1}$. Moreover the first eigenvalue is characterized by

$$\lambda_{1,p} = \inf \left\{ \frac{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}{\|\omega\|_{L^2(M)}^2}, \omega \in \Omega^p(M) \setminus \{0\}, \omega|_{\partial M} = 0 \right\}$$

and has to be positive because of [1].

Remarks 3.4

1. The Hodge \star operator preserves the Dirichlet problem so that $\lambda_{i,p} = \lambda_{i,n-p}$ for any $i \geq 1$ and $1 \leq p \leq n$.
2. If (M^n, g) is a compact domain in the Euclidean space, one can show, reasoning as in the proof of Proposition 3.1, that the Dirichlet eigenvalues on p -forms do not depend on p and correspond to the Laplacian Dirichlet eigenvalues on functions.

We are now in position to give explicit estimates between the first eigenvalues of the three previous problems, restricting our attention to $1 \leq p \leq \lfloor \frac{n}{2} \rfloor$ by Hodge symmetry of these spectra. Namely we get:

Theorem 3.5 *On an n -dimensional compact Riemannian manifold (M^n, g) with smooth boundary, the following inequalities hold:*

1. $\Lambda_{1,p}\lambda_{1,p} < \Gamma_{1,p}$ for $0 \leq p \leq n$;
2. $\inf(\lambda_{1,p+1}, \lambda_{1,p-1}) \leq \Lambda_{1,p}$ for $1 \leq p \leq n$;

3. $\lambda_{1,1} \leq \Lambda_{1,0}$.

Remarks 3.6

1. The inequalities of Theorems 3.3 and 3.5 give the analogue of some well-known results in the case of functions (see [2] for example).
2. In general, it is not clear whether equality can occur in the second and third inequalities of Theorem 3.5. We will see that this cannot occur for Euclidean domains.

Proof. To prove the first inequality, we consider ω an eigenform of the clamped plate problem (18) associated with $\Gamma_{1,p}$. Then the form ω can be considered as a test-form for the first eigenvalue of the buckling problem as well as for the first eigenvalue of the Dirichlet problem on differential forms. Therefore we first get

$$\Lambda_{1,p} \left(\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 \right) \leq \|\Delta\omega\|_{L^2(M)}^2.$$

which can be rewritten as

$$\Lambda_{1,p} \left(\frac{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}{\|\omega\|_{L^2(M)}^2} \right) \leq \frac{\|\Delta\omega\|_{L^2(M)}^2}{\|\omega\|_{L^2(M)}^2},$$

from which the estimate follows. If the inequality is an equality, then any eigenform ω of the clamped plate problem must be an eigenform of the Dirichlet problem for the Hodge Laplacian on M . As above, the unique continuation theorem implies that $\omega = 0$. Therefore the inequality is strict.

For the second estimate, we consider ω an eigenform of the buckling eigenvalue problem (16) associated with the first eigenvalue $\Lambda_{1,p}$ for $1 \leq p \leq [\frac{n}{2}]$. We first notice that the two differential forms $\omega_1 = d\omega \in \Omega^{p+1}(M)$ and $\omega_2 = \delta\omega \in \Omega^{p-1}(M)$ cannot be both trivial otherwise ω would be a harmonic p -form which vanishes on the boundary which is impossible by [1]. Moreover, it follows from (10) that

$$\iota^*\omega_1 = d\iota^*\omega = 0, \quad \nu \lrcorner \omega_1 = \nu \lrcorner d\omega = 0$$

and

$$\iota^*\omega_2 = \iota^*(\delta\omega) = 0, \quad \nu \lrcorner \omega_2 = -\delta^{\partial M}(\nu \lrcorner \omega) = 0$$

which imply that both ω_1 and ω_2 vanish along the boundary. So one can respectively take these forms as test-forms in (24) leading to

$$\lambda_{1,p+1} \|d\omega\|_{L^2(M)}^2 \leq \|\delta d\omega\|_{L^2(M)}^2 \quad \text{and} \quad \lambda_{1,p-1} \|\delta\omega\|_{L^2(M)}^2 \leq \|d\delta\omega\|_{L^2(M)}^2.$$

By adding these two inequalities, we get

$$\inf(\lambda_{1,p+1}, \lambda_{1,p-1}) \left(\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 \right) \leq \|\delta d\omega\|_{L^2(M)}^2 + \|d\delta\omega\|_{L^2(M)}^2$$

and the variational characterization (17) allows to conclude.

For the last one, take a smooth eigenfunction f for the buckling problem (2) on functions associated to $\Lambda_{1,0}$ and let $\omega = df \in \Omega^1(M)$. Then as above we observe that ω vanishes on ∂M so that it can be used as a test-form in the variational characterization of $\lambda_{1,1}$ leading to the inequality

$$\lambda_{1,1} \leq \frac{\int_M (\Delta f)^2 d\mu}{\int_M |df|^2 d\mu} = \Lambda_{1,0}.$$

This proves the third inequality in the broad sense. □

A direct consequence of Theorem 3.3 and the first inequality in Theorem 3.5 is the following estimate.

Corollary 3.7 *Let (M^n, g) be a compact Riemannian manifold with boundary then*

$$\lambda_{1,p} < \sqrt{\Gamma_{1,p}} < \Lambda_{1,p} \quad (25)$$

for all $0 \leq p \leq n$.

Since for domains in Euclidean space, the eigenvalues $\lambda_{1,p}$ and $\Lambda_{1,p}$ do not depend on p , it follows from the previous corollary with $p = 0$ that equality cannot occur in the second and third inequalities of Theorem 3.5.

Remark 3.8 *In [5], the Robin problem is defined as*

$$\begin{cases} \Delta\omega &= \lambda\omega & \text{on } M \\ \nu \lrcorner \omega &= 0 & \text{on } \partial M \\ \nu \lrcorner d\omega &= \tau \iota^* \omega & \text{on } \partial M \end{cases}$$

and it is proven that the first eigenvalue satisfies $\lambda_{1,p}(\tau) \leq \lambda_{1,p}$ for any parameter $\tau > 0$. Hence, it follows from (25) that $\lambda_{1,p}(\tau) < \sqrt{\Gamma_{1,p}} < \Lambda_{1,p}$.

Remark 3.9 *It is in fact not difficult to show that $\lambda_{j,p} \leq \Lambda_{j,p}$ for all $j \geq 1$ and $p = 0, \dots, n$, where the inequality is strict for $j = 1$ by Corollary 3.7. Indeed, it can be checked that the eigenvalue problem (24) is equivalent to the problem*

$$\begin{cases} \Delta^2\omega &= \lambda\Delta\omega & \text{on } M \\ \omega &= 0 & \text{on } \partial M \\ \Delta\omega &= 0 & \text{on } \partial M \end{cases}$$

in such a way that the Dirichlet eigenvalues are determined by applying the min-max formula to the functional

$$\frac{\int_M |\Delta\omega|^2 d\mu}{\int_M |d\omega|^2 + |\delta\omega|^2 d\mu}$$

on subspaces of p -forms which are in $(H^2 \cap H_0^1)(M)$. Namely, any critical point $\omega \in (H^2 \cap H_0^1)(M)$ of that functional associated with a critical value ℓ must satisfy

$$\int_M \langle \Delta\omega, \Delta\omega' \rangle d\mu = \ell \cdot \int_M (\langle d\omega, d\omega' \rangle + \langle \delta\omega, \delta\omega' \rangle) d\mu$$

for all $\omega' \in (H^2 \cap H_0^1)(M)$, that is,

$$\int_M \langle \Delta^2\omega, \omega' \rangle d\mu + \int_{\partial M} (\langle \iota^* \Delta\omega, \nu \lrcorner d\omega' \rangle - \langle \nu \lrcorner \Delta\omega, \iota^* \delta\omega' \rangle) d\mu = \ell \cdot \left(\int_M \langle \Delta\omega, \omega' \rangle \right),$$

which reduces to

$$\int_M \langle \Delta^2\omega - \ell\Delta\omega, \omega' \rangle d\mu = \int_{\partial M} (\langle \nu \lrcorner \Delta\omega, \iota^* \delta\omega' \rangle - \langle \iota^* \Delta\omega, \nu \lrcorner d\omega' \rangle) d\mu.$$

Here we have used that ω' vanishes along ∂M . It can be deduced in the usual way that $\Delta^2\omega - \ell\Delta\omega = 0$ on M as well as $\nu \lrcorner \Delta\omega = 0$ and $\iota^* \Delta\omega = 0$ along ∂M , using the fact that, for $\omega' \in H_0^1(M)$, we have $\iota^* \delta\omega' = -\nu \lrcorner \nabla_\nu \omega'$ and $\nu \lrcorner d\omega' = \iota^* \nabla_\nu \omega'$, and both the tangential and normal components of $\nabla_\nu \omega'$ can be prescribed independently. Thus $\Delta\omega = 0$ along ∂M , which yields the above equivalent problem.

Comparing now the above characterization to that of $\Lambda_{j,p}$, where the same quotient is considered on the smaller space $H_0^2(M)$, allows to conclude. Mind that it cannot be deduced for $j \geq 2$ whether the equality can be attained or not.

Now for $p \in \{0, \dots, n\}$, we consider the Hodge Laplacian with respect to the absolute boundary condition which satisfies the eigenvalue problem

$$\begin{cases} \Delta\omega & = \mu\omega & \text{on } M \\ \nu \lrcorner \omega & = 0 & \text{on } \partial M \\ \nu \lrcorner d\omega & = 0 & \text{on } \partial M. \end{cases} \quad (26)$$

We will denote by $\mu_{1,p}$ the first *positive* eigenvalue which is given by

$$\mu_{1,p} = \inf_{\omega \in \Omega^p(M) \setminus \{0\}} \left\{ \frac{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}{\|\omega\|_{L^2(M)}^2}, \nu \lrcorner \omega = 0 \text{ and } \omega \in H_A^p(M)^\perp \right\}. \quad (27)$$

The eigenspace associated with the zero eigenvalue, if non empty, corresponds to the absolute de Rham cohomology group in degree p defined by

$$H_A^p(M) = \{\omega \in \Omega^p(M) \mid d\omega = \delta\omega = 0 \text{ on } M \text{ and } \nu \lrcorner \omega = 0\}.$$

With these notations, it is clear that $\mu_{1,0}$ corresponds to the first nonzero eigenvalue of the Laplace operator under the Neumann boundary condition. Note also that the dual eigenvalue problem is the relative one whose zero eigenvalue reflects the relative de Rham cohomology group

$$H_R^p(M) = \{\omega \in \Omega^p(M) \mid d\omega = \delta\omega = 0 \text{ on } M \text{ and } \iota^*\omega = 0\}$$

and is related to the absolute boundary condition by the Hodge star operator which induces an isomorphism $H_A^p(M) \simeq H_R^{n-p}(M)$ for all $p = 0, \dots, n$. The Hodge star operator maps (26) onto the boundary problem

$$\begin{cases} \Delta\omega & = \kappa\omega & \text{on } M \\ \iota^*\omega & = 0 & \text{on } \partial M \\ \iota^*(\delta\omega) & = 0 & \text{on } \partial M \end{cases}$$

where κ is the corresponding eigenvalue. As usual, up to the possible eigenvalue 0, the (monotonously nondecreasing) sequence of real eigenvalues is denoted by $(\kappa_{i,p})_{i \geq 1}$, where $\kappa_{1,p}$ is the smallest positive one. It is a straightforward consequence of that correspondence via the Hodge star operator that, for all $i \geq 1$ and $p \in \{0, \dots, n\}$, the identity $\mu_{i,p} = \kappa_{i,n-p}$ holds. Moreover, $\mu_{1,n} = \kappa_{1,0} > 0$ holds because of $H_R^0(M) = \{0\}$. By definition, the identity $\kappa_{1,0} = \lambda_{1,0}$ also holds.

Remark 3.10 *If $H_A^p(M) = \{0\}$, an eigenform for the Dirichlet problem (24) can be taken as a test-form in the variational characterization of $\mu_{1,p}$ leading to the estimate $\mu_{1,p} \leq \lambda_{1,p}$ for all $p = 1, \dots, n$. Note that this inequality does not hold in general for $p = 0$. However, it is a well-known result of Pólya [14] (see also [7]) that $\mu_{1,0} < \lambda_{1,0}$ for Euclidean domains. On the other hand, for $p = 0, \dots, [\frac{n}{2}]$, Guerini and Savo proved in [10, Theorem 2.6, b)] that $\mu_{1,p} \leq \kappa_{1,p}$ for a convex domain M in the Euclidean space. Even more, if M is strictly convex, this inequality is strict for $0 \leq p < [\frac{n}{2}]$ (see [10, Theorem 2.6, c)]). Therefore, for a convex Euclidean domain with $H_R^p(M) = \{0\}$ for some $p = 0, \dots, [\frac{n}{2}]$, we have*

$$\mu_{1,p} \leq \kappa_{1,p} = \mu_{1,n-p} \leq \lambda_{1,n-p} = \lambda_{1,p}$$

This is the case for example for strictly convex domains in \mathbb{R}^n and in addition the first inequality is strict for $0 \leq p < [\frac{n}{2}]$ by the above discussion.

It turns out that the first nonzero eigenvalues of the Hodge Laplacian under the absolute boundary condition can be related with the first eigenvalue of the buckling problem, as is stated below.

Theorem 3.11 *On an n -dimensional compact Riemannian manifold (M^n, g) with smooth boundary, we have*

$$\max(\mu_{1,p}, \mu_{1,n-p}) \leq \Lambda_{1,p} \quad (28)$$

for all $p = 0, \dots, n$. Moreover, equality occurs if and only if there exist $k \in \{p, n-p\}$, $\omega \in \Omega^k(M)$ and $\omega_0 \in H_A^k(M) \setminus \{0\}$ satisfying the overdetermined boundary value problem

$$\begin{cases} \Delta\omega &= \mu_{1,k}\omega & \text{on } M \\ \omega &= \omega_0 & \text{on } \partial M \\ \iota^*(\delta\omega) &= 0, \nu \lrcorner d\omega = 0 & \text{on } \partial M. \end{cases} \quad (29)$$

Proof. Let $\omega_1 \in \Omega^p(M)$ be an eigenform of the buckling problem associated to the eigenvalue $\Lambda_{1,p}$. We denote by ω_0 the $L^2(M)$ -orthogonal projection of ω_1 onto $H_A^p(M)$ and we set $\omega := \omega_1 - \omega_0 \in \Omega^p(M)$. Then $d\omega = d\omega_1$ and $\delta\omega = \delta\omega_1$ on M , and in particular $\Delta\omega = \Delta\omega_1$, as well as $\nu \lrcorner \omega = 0$ and $\nu \lrcorner d\omega = 0$ along ∂M . For this form ω , using the first equality in (14) and the Cauchy-Schwarz inequality leads to

$$\left(\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 \right)^2 = (\Delta\omega, \omega)_{L^2(M)}^2 \leq \|\Delta\omega\|_{L^2(M)}^2 \|\omega\|_{L^2(M)}^2,$$

so that

$$\frac{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}{\|\omega\|_{L^2(M)}^2} \leq \frac{\|\Delta\omega\|_{L^2(M)}^2}{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}.$$

Note that $\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 = \|d\omega_1\|_{L^2(M)}^2 + \|\delta\omega_1\|_{L^2(M)}^2 > 0$. Moreover since $\Delta\omega = \Delta\omega_1$ and ω_1 is an eigenform associated with $\Lambda_{1,p}$, we deduce that

$$\frac{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2}{\|\omega\|_{L^2(M)}^2} \leq \frac{\|\Delta\omega_1\|_{L^2(M)}^2}{\|d\omega_1\|_{L^2(M)}^2 + \|\delta\omega_1\|_{L^2(M)}^2} = \Lambda_{1,p}.$$

But $\nu \lrcorner \omega = 0$ and $(\omega, \omega')_{L^2(M)} = 0$ for all $\omega' \in H_A^p(M)$ so that the p -form ω can be taken as a test-form in (27) and therefore $\mu_{1,p} \leq \Lambda_{1,p}$. The same reasoning ensures that $\mu_{1,n-p} \leq \Lambda_{1,n-p}$ and the result follows since $\Lambda_{1,n-p} = \Lambda_{1,p}$ as noticed in Remark 2.3.

Now if equality occurs one can assume without loss of generality that $k = p$ (recall that $k \in \{p, n-p\}$ by assumption). Then the p -form ω is an eigenform with respect to the absolute boundary condition for the eigenvalue $\mu_{1,p}$, so that it satisfies the problem (27). On the other hand, it is clear that since $\omega = \omega_1 - \omega_0$ with ω_1 an eigenform for the buckling problem and $\omega_0 \in H_A^p(M)$ we have $\iota^*\omega = -\iota^*\omega_0$ and $\iota^*(\delta\omega) = 0$ which gives the first part of the equality case. Conversely, assume that there exists $\omega \in \Omega^p(M)$ satisfying (29) with $\omega_0 \in H_A^p(M) \setminus \{0\}$. Now if we let $\omega_1 := \omega - \omega_0$, it is not difficult to check using (10) that

$$\omega_1|_{\partial M} = 0 \quad \text{and} \quad \nabla_\nu \omega_1|_{\partial M} = 0$$

so that ω_1 can be taken as a test-form in the variational characterization of $\Lambda_{1,p}$. On the other hand, since ω_0 is an harmonic field we have $d\omega_1 = d\omega$ and $\delta\omega_1 = \delta\omega$ and then

$$\Lambda_{1,p} \leq \frac{\|\Delta\omega_1\|_{L^2(M)}^2}{\|d\omega_1\|_{L^2(M)}^2 + \|\delta\omega_1\|_{L^2(M)}^2} = \frac{\|\Delta\omega\|_{L^2(M)}^2}{\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2} = \mu_{1,p}.$$

This implies that equality occurs in our eigenvalue estimate. The same holds if we replace p by $n-p$. \square

Note that problem (29) would not be elliptic without the boundary condition $\nu \lrcorner d\omega = 0$.

Remark 3.12 *In other words, Theorem 3.11 states that*

$$\max(\mu_{1,p}, \kappa_{1,p}) \leq \Lambda_{1,p} \quad (30)$$

for all $p = 0, \dots, n$. Moreover, equality occurs if and only if there exist either $\omega \in \Omega^p(M)$ and $\omega_0 \in H_A^p(M) \setminus \{0\}$ satisfying (29), or $\tilde{\omega} \in \Omega^p(M)$ and $\tilde{\omega}_0 \in H_R^p(M) \setminus \{0\}$ such that

$$\begin{cases} \Delta \tilde{\omega} &= \kappa_{1,p} \tilde{\omega} & \text{on } M \\ \tilde{\omega} &= \tilde{\omega}_0 & \text{on } \partial M \\ \iota^*(\delta \tilde{\omega}) &= 0, \nu \lrcorner d\tilde{\omega} = 0 & \text{on } \partial M. \end{cases} \quad (31)$$

As a direct consequence of Theorem 3.11, we get:

Corollary 3.13 *Let (M^n, g) be an n -dimensional compact Riemannian manifold with boundary, then the following hold:*

1. *if $H_A^p(M) = \{0\}$ then $\mu_{1,p} < \Lambda_{1,p}$ for $p = 0, \dots, n-1$;*
2. *if $H_A^p(M) = \{0\}$ and $H_R^p(M) = \{0\}$ then Inequality (28) is strict for $p = 0, \dots, [\frac{n}{2}]$.*

Note that, when $H_A^p(M) = \{0\}$, the inequality $\mu_{1,p} < \Lambda_{1,p}$ is a straightforward consequence of Corollary 3.7 because of $\max(\mu_{1,p}, \mu_{1,n-p}) \leq \lambda_{1,p}$ by [5, Prop. 5.4].

Remarks 3.14

1. *Since $H_A^n(M) = \{0\}$ it follows from Theorem 3.11 that $\mu_{1,n} < \Lambda_{1,n}$ which is just another formulation of the inequality $\lambda_{1,0} < \Lambda_{1,0}$.*
2. *Let $\omega_i \in \Omega^p(M)$, $i = 1, 2$, be two p -forms on M such that $\omega_1|_{\partial M} = \omega_2|_{\partial M}$. Then it follows directly from (8) and (9) that:*

$$\left(\iota^*(\delta \omega_1) = \iota^*(\delta \omega_2) \quad \text{and} \quad \nu \lrcorner d\omega_1 = \nu \lrcorner d\omega_2 \right) \iff \nabla_\nu \omega_1|_{\partial M} = \nabla_\nu \omega_2|_{\partial M}.$$

Using this characterization, it is not difficult to show that $\omega \in \Omega^p(M)$ and $\omega_0 \in H_A^p(M)$ satisfy (29) if and only if

$$\begin{cases} \Delta \omega &= \mu_{1,p} \omega & \text{on } M \\ \omega &= \omega_0 & \text{on } \partial M \\ \nabla_\nu \omega|_{\partial M} &= \nabla_\nu \omega_0|_{\partial M} & \text{on } \partial M. \end{cases}$$

Obviously, the same holds for the eigenvalue boundary problem (31) with $\tilde{\omega} \in \Omega^p(M)$ and $\tilde{\omega}_0 \in H_R^p(M)$.

From Corollary 3.7, the estimate (30) for $p = 0$ reads

$$\mu_{1,0} \leq \Lambda_{1,0} \quad \text{and} \quad \lambda_{1,0} < \Lambda_{1,0}.$$

The first inequality was proved in [12] and states more precisely that $\mu_{1,0} < \Lambda_{1,0}$ on any compact Riemannian manifold with boundary. However the arguments given in the proof of [12, Lemma 3.1] which establishes that the inequality is strict are not clear. Indeed if M is assumed to be connected, we have $H_A^0(M) \simeq \mathbb{R}$ and so, as indicated in Theorem 3.11, the equality $\mu_{1,0} = \Lambda_{1,0}$ ensures the existence of a smooth function f on M satisfying the overdetermined problem

$$\begin{cases} \Delta f &= \mu_{1,0} f & \text{on } M \\ f &= \frac{1}{\text{Vol}(M, g)} \int_M f d\mu & \text{on } \partial M \\ \frac{\partial f}{\partial \nu} &= 0 & \text{on } \partial M \end{cases}$$

which is a particular case of the so-called Schiffer conjecture (see for example [15] and the references therein). However the fact that the inequality is strict can be proved at least for Euclidean domains.

Corollary 3.15 *Let (M^n, g) be a compact domain in the Euclidean space. Then $\mu_{1,0} < \Lambda_{1,0}$.*

Proof. As recall in Remark 3.10, we have $\mu_{1,0} < \lambda_{1,0}$ for any domains in the Euclidean space and so the result follows from Corollary 3.7. \square

On a convex Euclidean domain with $H_R^p(M) = \{0\}$, since the spectrum of the Dirichlet and the buckling problems do not depend on p , we deduce from Remark 3.10 that

$$\mu_{1,p} \leq \lambda_{1,p} = \lambda_{1,0} < \lambda_{2,0} \leq \Lambda_{1,0} = \Lambda_{1,p}$$

for $p = 0, \dots, [\frac{n}{2}]$. The last inequality is due to Payne [13] (with a gap in the proof filled by Friedlander [8]).

3.3 Manifolds with Weitzenböck curvature operator bounded from below

In this section, we give a lower bound for the first eigenvalue $\lambda_{1,p}$ of the Dirichlet Hodge Laplacian under the condition that the Weitzenböck curvature operator of the Riemannian manifold (M^n, g) is bounded from below by a positive constant $\gamma > 0$. Combined with some results of the former part, we obtain estimates for the first eigenvalues of the buckling and clamped plate problems in this context. These last results generalize previous results by Chen, Cheng, Wang and Xia [4].

For this, we first recall that the Weitzenböck formula for p -forms states that

$$\Delta\omega = \nabla^*\nabla\omega + W^{[p]}\omega \tag{32}$$

for any $\omega \in \Omega^p(M)$ where ∇ (resp. ∇^*) is the Levi-Civita connection (resp. its L^2 -adjoint) on forms and $W^{[p]}$ is the curvature term. This last term is usually called the Weitzenböck curvature operator and it defines a self-adjoint endomorphism acting on p -forms. In the following we will say that $W^{[p]}$ is bounded from below by $\gamma p(n-p) \in \mathbb{R}$ for a fixed $p = 1, \dots, n$ if it satisfies

$$\langle W^{[p]}\omega, \omega \rangle \geq \gamma p(n-p)|\omega|^2 \tag{33}$$

for all $\omega \in \Omega^p(M)$. For $p = 1$, this is equivalent to the fact that the Ricci curvature satisfies $\text{Ric} \geq (n-1)\gamma g$. From [9], this condition is satisfied for all p if the curvature operator is bounded from below by $\gamma \in \mathbb{R}$.

Now integrating (32) on M leads to the so-called *Reilly formula* on p -forms [16, Theorem 3] which writes as

$$\int_M (|d\omega|^2 + |\delta\omega|^2) d\mu = \int_M (|\nabla\omega|^2 + \langle W^{[p]}\omega, \omega \rangle) d\mu + \int_{\partial M} \mathcal{B}(\omega, \omega) d\sigma \tag{34}$$

for all $\omega \in \Omega^p(M)$ and where

$$\mathcal{B}(\omega, \omega) = 2\langle \nu \lrcorner \omega, \delta^{\partial M}(\iota^*\omega) \rangle + \langle S^{[p]}(\iota^*\omega), \iota^*\omega \rangle + (n-1)H|\nu \lrcorner \omega|^2 - \langle S^{[p-1]}(\nu \lrcorner \omega), \nu \lrcorner \omega \rangle.$$

As a direct consequence of this formula, we get the following estimate on $\lambda_{1,p}$.

Theorem 3.16 *Let (M^n, g) be a compact Riemannian manifold with boundary whose Weitzenböck curvature operator $W^{[p]}$ is bounded from below by a positive constant $\gamma p(n-p)$ for some $1 \leq p \leq [\frac{n}{2}]$. Then we have $\lambda_{1,p} > \gamma p(n-p+1)$.*

Proof. We first observe that if ω is a p -form on M satisfying $\omega|_{\partial M} = 0$, the boundary term in the Reilly formula (34) vanishes and then, from (33), we get

$$\int_M (|d\omega|^2 + |\delta\omega|^2) d\mu \geq \int_M (|\nabla\omega|^2 + \gamma p(n-p)|\omega|^2) d\mu.$$

On the other hand, with the help of the pointwise inequality

$$|\nabla\alpha|^2 \geq \frac{1}{p+1}|d\alpha|^2 + \frac{1}{n-p+1}|\delta\alpha|^2$$

which is true for any p -form α (see [9, Lemme 6.8]), we obtain

$$\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 \geq \frac{1}{n-p+1}(\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2) + \gamma p(n-p)\|\omega\|_{L^2(M)}^2$$

since $1 \leq p \leq [\frac{n}{2}]$. Note that equality occurs if ω is a conformal Killing form (see Lemma 3.17 below for a precise definition and [19] for more general properties) and if $n = 2p$ or if $1 \leq p < [\frac{n}{2}]$ and $d\omega = 0$. We finally get

$$\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 \geq \gamma p(n-p+1)\|\omega\|_{L^2(M)}^2 \quad (35)$$

which, by taking ω an eigenform associated to $\lambda_{1,p}$, leads to the desired estimate in a broad sense. Assume now that equality holds so that ω is a conformal Killing p -form which vanishes on the boundary. Then Lemma 3.17 ensures that $\nabla_\nu\omega|_{\partial M} = 0$ and this is impossible since ω is also a p -eigenform for the Dirichlet Hodge Laplacian. \square

In the previous proof, we considered conformal Killing p -forms ω , that is p -forms satisfying the following equation

$$\nabla_X\omega = \frac{1}{p+1}X \lrcorner d\omega - \frac{1}{n-p+1}X^\flat \wedge \delta\omega \quad (36)$$

for all $X \in \Gamma(TM)$ and used the following result that we prove now.

Lemma 3.17 *On a (not necessarily compact) Riemannian manifold (M^n, g) with boundary ∂M , a conformal Killing p -form with $1 \leq p \leq n-1$ which vanishes on ∂M satisfies $\nabla_\nu\omega|_{\partial M} = 0$.*

Proof. It is a direct consequence of (36) that

$$\iota^*(\nabla_\nu\omega) = \frac{1}{p+1}\nu \lrcorner d\omega \quad \text{and} \quad \nu \lrcorner \nabla_\nu\omega = -\frac{1}{n-p+1}\iota^*(\delta\omega).$$

On the other hand, since $\omega \in \Omega^p(M)$ vanishes on ∂M it is straightforward from (8) and (9) to compute that

$$\iota^*(\nabla_\nu\omega) = \nu \lrcorner d\omega \quad \text{and} \quad \nu \lrcorner \nabla_\nu\omega = -\iota^*(\delta\omega).$$

Putting these relations together imply that

$$\nu \lrcorner d\omega = 0 \quad \text{and} \quad \frac{n-p}{n-p+1}\iota^*(\delta\omega) = 0$$

which, with the help of (10), allows to conclude. \square

Combining Theorem 3.16 with (25) leads to the following estimates for the buckling and clamped plate first eigenvalues.

Corollary 3.18 *Let (M^n, g) be a compact Riemannian manifold with boundary whose Weitzenböck curvature operator $W^{[p]}$ is bounded from below by $\gamma p(n-p) > 0$ for some $1 \leq p \leq [\frac{n}{2}]$. Then we have*

$$\Lambda_{1,p} > \gamma p(n-p+1) \quad \text{and} \quad \Gamma_{1,p} > \gamma^2 p^2(n-p+1)^2.$$

Remarks 3.19

1. From these estimates, known lower bounds for the first eigenvalues of the buckling and clamped plate problems on functions [4] can be deduced. Indeed, for $p = 1$, the fact that $W^{[1]}$ is bounded from below by $n - 1$ is equivalent to the fact that the Ricci tensor of (M^n, g) satisfies $\text{Ric} \geq (n - 1)g$. Then the first estimate in Corollary 3.18 reads $\Lambda_{1,1} > n$ under this curvature assumption.
2. Independently, the third inequality of Theorem 3.5, which reads $\Lambda_{1,0} \geq \lambda_{1,1}$, combined with the inequality $\lambda_{1,1} > n$ from Theorem 3.16, yields $\Lambda_{1,0} > n$. This is exactly [4, Theorem 1.6]. Now putting together this estimate with the first inequality in Theorem 3.5 gives $\Gamma_{1,0} > n\lambda_{1,0}$, which is precisely [4, Theorem 1.5].

3.4 Domains in the unit sphere

In this section, we derive an inequality which relates the first eigenvalues of the clamped plated problem for different degrees for domains of the n -dimensional unit sphere \mathbb{S}^n . This result is a consequence of a more general estimate for submanifolds isometrically immersed in the Euclidean space \mathbb{R}^{n+m} which can be obtained using [6, Lemma 3.8]. However, it is in general difficult to control all the terms which appear in these estimates so that we shall restrict ourselves to the case when M is a domain of \mathbb{S}^n . We also obtain a similar result for the first buckling eigenvalue. More precisely, we obtain:

Theorem 3.20 *Let (M^n, g) be a compact domain in the unit sphere \mathbb{S}^n .*

1. For $1 \leq p \leq [\frac{n}{2}]$, we have

$$p\Gamma_{1,p-1} + (n - p)\Gamma_{1,p+1} < C_{n,p}\Gamma_{1,p}, \quad (37)$$

and

$$\min(p\Lambda_{1,p-1}, (n - p)\Lambda_{1,p+1}) < \frac{C_{n,p}}{2} \cdot \Lambda_{1,p}, \quad (38)$$

$$\text{where } C_{n,p} = n + \frac{4 + 2(n - 2p)^2}{p(n - p + 1)} + \frac{n(n - 2p)^2}{p^2(n - p + 1)^2}.$$

2. For $n = 2p$, we have

$$\Lambda_{1, \frac{n}{2}-1} < \left(1 + \frac{16}{n^2(n + 2)}\right) \cdot \Lambda_{1, \frac{n}{2}}. \quad (39)$$

Proof. Let ω be a smooth p -form on Ω which vanishes on ∂M and whose normal covariant derivative is also zero on the boundary. For any $1 \leq i \leq n + 1$, we consider the $(p - 1)$ -form $\partial_{x_i}^T \lrcorner \omega$ on M where $\partial_{x_i}^T$ denotes the tangential part in TM of the unit parallel vector field ∂_{x_i} of \mathbb{R}^{n+1} . It is not difficult to check that

$$(\partial_{x_i}^T \lrcorner \omega)|_{\partial M} = 0 \quad \text{and} \quad \nabla_\nu(\partial_{x_i}^T \lrcorner \omega)|_{\partial M} = 0$$

so that it can be used as a test-form in the variational characterizations of $\Gamma_{1,p-1}$ and $\Lambda_{1,p-1}$.

1. From (19), we get

$$\Gamma_{1,p-1} \int_M |\partial_{x_i}^T \lrcorner \omega|^2 d\mu \leq \int_M |\Delta(\partial_{x_i}^T \lrcorner \omega)|^2 d\mu.$$

Summing that inequality over i and using the pointwise identity $\sum_{i=1}^{n+1} |\partial_{x_i}^T \lrcorner \omega|^2 = p|\omega|^2$ (see e.g. [6, Eq. (22)]), we obtain

$$p\Gamma_{1,p-1} \int_M |\omega|^2 d\mu \leq \int_M \sum_{i=1}^{n+1} |\Delta(\partial_{x_i}^T \lrcorner \omega)|^2 d\mu.$$

Now the right-hand side of that inequality was computed for every p -form ω in the right-hand side of [6, Eq. (31) and (33)] and we obtain

$$p\Gamma_{1,p-1}\|\omega\|_{L^2(M)}^2 \leq 4\|\delta\omega\|_{L^2(M)}^2 + p\|\Delta\omega + (2p-n)\omega\|_{L^2(M)}^2. \quad (40)$$

Since Inequality (40) is true for any p -eigenform ω , we can apply it to the $(n-p)$ -eigenform $\star\omega$ to get

$$(n-p)\Gamma_{1,n-p-1}\|\omega\|_{L^2(M)}^2 \leq 4\|d\omega\|_{L^2(M)}^2 + (n-p)\|\Delta\omega + (n-2p)\omega\|_{L^2(M)}^2.$$

Summing both inequalities and using the fact that $\Gamma_{1,n-p-1} = \Gamma_{1,p+1}$ yields

$$\begin{aligned} (p\Gamma_{1,p-1} + (n-p)\Gamma_{1,p+1})\|\omega\|_{L^2(M)}^2 &\leq n\|\Delta\omega\|_{L^2(M)}^2 + n(2p-n)^2\|\omega\|_{L^2(M)}^2 \\ &\quad + (4 + 2(n-2p)^2)(\Delta\omega, \omega)_{L^2(M)} \\ &\leq n\|\Delta\omega\|_{L^2(M)}^2 + n(2p-n)^2\|\omega\|_{L^2(M)}^2 \\ &\quad + (4 + 2(n-2p)^2)\|\Delta\omega\|_{L^2(M)}\|\omega\|_{L^2(M)} \end{aligned}$$

where we used the Cauchy-Schwarz inequality in the last inequality. On the other hand, since M is in the unit sphere, the estimate (35) holds with $\gamma = 1$, and therefore we get

$$(p\Gamma_{1,p-1} + (n-p)\Gamma_{1,p+1})\|\omega\|_{L^2(M)}^2 \leq C_{n,p}\|\Delta\omega\|_{L^2(M)}^2$$

which allows to deduce (37) in the broad sense by taking ω to be an eigenform associated to $\Gamma_{1,p}$. If (37) were an equality, then for any clamped-plate-eigenform ω associated to $\Gamma_{1,p}$, the p -form $\Delta\omega$ would be pointwise proportional to ω because of the equality in the Cauchy-Schwarz inequality. But, because of $\Delta^2\omega = \Gamma_{1,p}\omega$ on M , we would have $\Delta\omega = \sqrt{\Gamma_{1,p}\omega}$ on M , therefore ω would be a Dirichlet eigenform associated to the eigenvalue $\sqrt{\Gamma_{1,p}}$. Again, because ω satisfies $\nabla_\nu\omega|_{\partial M} = 0$ along ∂M , the unique continuation property for elliptic second-order linear operators would imply that $\omega = 0$ on M . This would lead to a contradiction and shows that (37) must be strict.

Now, we prove (38). Using the variational characterization (17) of $\Lambda_{1,p-1}$ for $p = 1, \dots, n$ gives

$$\Lambda_{1,p-1} \int_M \left(|d(\partial_{x_i}^T \lrcorner \omega)|^2 + |\delta(\partial_{x_i}^T \lrcorner \omega)|^2 \right) d\mu \leq \int_M |\Delta(\partial_{x_i}^T \lrcorner \omega)|^2 d\mu$$

and summing as above on i from 1 to $n+1$ leads to

$$\Lambda_{1,p-1} \sum_{i=1}^{n+1} \int_M \left(|d(\partial_{x_i}^T \lrcorner \omega)|^2 + |\delta(\partial_{x_i}^T \lrcorner \omega)|^2 \right) d\mu \leq 4\|\delta\omega\|_{L^2(M)}^2 + p\|\Delta\omega + (2p-n)\omega\|_{L^2(M)}^2.$$

By the Cartan identity and [10, id. (4.3)] (see also [6, id. (20)]), we have, for every $1 \leq i \leq n+1$,

$$\begin{aligned} d(\partial_{x_i}^T \lrcorner \omega) &= \mathcal{L}_{\partial_{x_i}^T} \omega - \partial_{x_i}^T \lrcorner d\omega \\ &= \nabla_{\partial_{x_i}^T} \omega + \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega - \partial_{x_i}^T \lrcorner d\omega, \end{aligned}$$

where $\mathcal{L}_X\omega$ is the Lie derivative of ω in the X -direction, $\mathbb{I}_{\partial_{x_i}^\perp}^{[p]}$ is the natural extension onto $\Lambda^p T^*M$ of the pointwise endomorphism field $\mathbb{I}_{\partial_{x_i}^\perp}$ of TM defined for all $X, Y \in TM$ by $\langle \mathbb{I}_{\partial_{x_i}^\perp}(X), Y \rangle = \langle \mathbb{I}(X, Y), \partial_{x_i}^\perp \rangle$ and \mathbb{I} is the second fundamental form of M in \mathbb{R}^{n+1} . Here $\partial_{x_i}^\perp$ denotes the normal component of ∂_{x_i} that is, $\partial_{x_i}^\perp = \partial_{x_i} - \partial_{x_i}^T$. Note that, because $\mathbb{I}_x = -g_x \otimes x$

for every $x \in \mathbb{S}^n$ and hence $x \in M$, we have $\mathbb{I}_{\partial_{x_i}^\perp} = -x_i \cdot \text{Id}$ at $x = (x_1, \dots, x_{n+1})$, so that $\mathbb{I}_{\partial_{x_i}^\perp}^{[p]} = -px_i \cdot \text{Id}$ at x . It can be deduced that

$$\begin{aligned} |d(\partial_{x_i}^T \lrcorner \omega)|^2 &= \left| \nabla_{\partial_{x_i}^T} \omega + \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega - \partial_{x_i}^T \lrcorner d\omega \right|^2 \\ &= \left| \nabla_{\partial_{x_i}^T} \omega \right|^2 + \left| \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega \right|^2 + \left| \partial_{x_i}^T \lrcorner d\omega \right|^2 \\ &\quad + 2\langle \nabla_{\partial_{x_i}^T} \omega, \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega \rangle - 2\langle \nabla_{\partial_{x_i}^T} \omega, \partial_{x_i}^T \lrcorner d\omega \rangle - 2\langle \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega, \partial_{x_i}^T \lrcorner d\omega \rangle. \end{aligned}$$

Choosing any orthonormal basis $(e_j)_{1 \leq j \leq n}$ of $T_x M$ for some $x \in M$, we have

$$\begin{aligned} \sum_{i=1}^n \left| \nabla_{\partial_{x_i}^T} \omega \right|^2 &= \sum_{i=1}^{n+1} \sum_{j,k=1}^n \langle \partial_{x_i}^T, e_j \rangle \langle \partial_{x_i}^T, e_k \rangle \langle \nabla_{e_j} \omega, \nabla_{e_k} \omega \rangle \\ &= \sum_{i=1}^{n+1} \sum_{j,k=1}^n \langle \partial_{x_i}^T, e_j \rangle \langle \partial_{x_i}^T, e_k \rangle \langle \nabla_{e_j} \omega, \nabla_{e_k} \omega \rangle \\ &= \sum_{j,k=1}^n \left(\sum_{i=1}^{n+1} \langle \partial_{x_i}^T, e_j \rangle \langle \partial_{x_i}^T, e_k \rangle \right) \cdot \langle \nabla_{e_j} \omega, \nabla_{e_k} \omega \rangle \\ &= \sum_{j,k=1}^n \langle e_j, e_k \rangle \langle \nabla_{e_j} \omega, \nabla_{e_k} \omega \rangle \\ &= \sum_{j=1}^n \left| \nabla_{e_j} \omega \right|^2 \\ &= \left| \nabla \omega \right|^2. \end{aligned}$$

As a second step, at every $x \in M$, because of $|x| = 1$,

$$\sum_{i=1}^{n+1} \left| \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega \right|^2 = p^2 \sum_{i=1}^{n+1} x_i^2 |\omega|^2 = p^2 |x|^2 |\omega|^2 = p^2 |\omega|^2.$$

By [6, eq. (22)],

$$\sum_{i=1}^{n+1} \left| \partial_{x_i}^T \lrcorner d\omega \right|^2 = (p+1) |\omega|^2.$$

Moreover,

$$\begin{aligned} \sum_{i=1}^{n+1} 2\langle \nabla_{\partial_{x_i}^T} \omega, \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega \rangle &= p \sum_{i=1}^{n+1} 2x_i \langle \nabla_{\partial_{x_i}^T} \omega, \omega \rangle \\ &= p \sum_{i=1}^{n+1} x_i \partial_{x_i}^T (|\omega|^2) \\ &= px^T (|\omega|^2) \\ &= 0 \end{aligned}$$

because of $x^T = 0$ for every $x \in \mathbb{S}^n$. For the same reason,

$$\begin{aligned} \sum_{i=1}^{n+1} \langle \mathbb{I}_{\partial_{x_i}^\perp}^{[p]} \omega, \partial_{x_i}^T \lrcorner d\omega \rangle &= p \sum_{i=1}^{n+1} x_i \langle \omega, \partial_{x_i}^T \lrcorner d\omega \rangle \\ &= p \langle \omega, x^T \lrcorner d\omega \rangle \\ &= 0. \end{aligned}$$

Applying the same computational method as above, we also have, in any pointwise orthonormal basis $(e_j)_{1 \leq j \leq n}$ of TM ,

$$\begin{aligned} \sum_{i=1}^{n+1} \langle \nabla_{\partial_{x_i}^T} \omega, \partial_{x_i}^T \lrcorner d\omega \rangle &= \sum_{j=1}^n \langle \nabla_{e_j} \omega, e_j \lrcorner d\omega \rangle \\ &= \sum_{j=1}^n \langle e_j^\flat \wedge \nabla_{e_j} \omega, d\omega \rangle \\ &= |d\omega|^2. \end{aligned}$$

On the whole, we obtain

$$\begin{aligned} \sum_{i=1}^{n+1} |d(\partial_{x_i}^T \lrcorner \omega)|^2 &= |\nabla \omega|^2 + p^2 |\omega|^2 + (p+1) |d\omega|^2 - 2 |d\omega|^2 \\ &= |\nabla \omega|^2 + p^2 |\omega|^2 + (p-1) |d\omega|^2. \end{aligned}$$

Therefore

$$\sum_{i=1}^{n+1} \|d(\partial_{x_i}^T \lrcorner \omega)\|_{L^2(M)}^2 = \|\nabla \omega\|_{L^2(M)}^2 + p^2 \|\omega\|_{L^2(M)}^2 + (p-1) \|d\omega\|_{L^2(M)}^2.$$

Using the Weitzenböck formula (32) with $W^{[p]} = p(n-p) \cdot \text{Id}$ and that $\omega|_{\partial M} = \nabla_\nu \omega|_{\partial M} = 0$ leads to

$$\begin{aligned} \sum_{i=1}^{n+1} \|d(\partial_{x_i}^T \lrcorner \omega)\|_{L^2(M)}^2 &= (\nabla^* \nabla \omega, \omega)_{L^2(M)} + p^2 \|\omega\|_{L^2(M)}^2 + (p-1) \|d\omega\|_{L^2(M)}^2 \\ &= (\Delta \omega, \omega)_{L^2(M)} - (W^{[p]} \omega, \omega)_{L^2(M)} + p^2 \|\omega\|_{L^2(M)}^2 + (p-1) \|d\omega\|_{L^2(M)}^2 \\ &= (\Delta \omega, \omega)_{L^2(M)} + (p^2 - p(n-p)) \|\omega\|_{L^2(M)}^2 + (p-1) \|d\omega\|_{L^2(M)}^2 \\ &= p(2p-n) \|\omega\|_{L^2(M)}^2 + (p-1) \|d\omega\|_{L^2(M)}^2 + (\Delta \omega, \omega)_{L^2(M)}. \end{aligned}$$

On the other hand, for every $i \in \{1, \dots, n+1\}$, we can compute, using a local orthonormal basis $(e_j)_{1 \leq j \leq n}$ of TM as well as $\nabla_X \partial_{x_i}^T = -x_i X$ for every $X \in T_x M$,

$$\begin{aligned} \delta(\partial_{x_i}^T \lrcorner \omega) &= - \sum_{j=1}^n e_j \lrcorner \nabla_{e_j} (\partial_{x_i}^T \lrcorner \omega) \\ &= - \sum_{j=1}^n e_j \lrcorner ((\nabla_{e_j} \partial_{x_i}^T) \lrcorner \omega + \partial_{x_i}^T \lrcorner \nabla_{e_j} \omega) \\ &= - \sum_{j,k=1}^n \langle \nabla_{e_j} \partial_{x_i}^T, e_k \rangle e_j \lrcorner e_k \lrcorner \omega - \partial_{x_i}^T \lrcorner \delta \omega \\ &= \underbrace{\sum_{j,k=1}^n x_i \langle e_j, e_k \rangle e_j \lrcorner e_k \lrcorner \omega}_{0} - \partial_{x_i}^T \lrcorner \delta \omega \\ &= -\partial_{x_i}^T \lrcorner \delta \omega, \end{aligned}$$

where we used the skew-symmetry of $\langle e_j, e_k \rangle e_j \lrcorner e_k \lrcorner \omega$ in (j, k) . As a consequence, again by [6, eq. (22)],

$$\begin{aligned} \sum_{i=1}^{n+1} |\delta(\partial_{x_i}^T \lrcorner \omega)|^2 &= \sum_{i=1}^{n+1} |\partial_{x_i}^T \lrcorner \delta \omega|^2 \\ &= (p-1) |\delta \omega|^2, \end{aligned}$$

from which $\sum_{i=1}^{n+1} \|\delta(\partial_{x_i}^T \lrcorner \omega)\|_{L^2(M)}^2 = (p-1)\|\delta\omega\|_{L^2(M)}^2$ follows. Finally, we deduce that

$$\begin{aligned} \Lambda_{1,p-1} & \left(p(2p-n)\|\omega\|_{L^2(M)}^2 + (p-1)(\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2) + (\Delta\omega, \omega)_{L^2(M)} \right) \\ & \leq 4\|\delta\omega\|_{L^2(M)}^2 + p\|\Delta\omega + (2p-n)\omega\|_{L^2(M)}^2, \end{aligned}$$

that is, using $\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 = (\Delta\omega, \omega)_{L^2(M)}$,

$$p\Lambda_{1,p-1} \left((\Delta\omega, \omega)_{L^2(M)} + (2p-n)\|\omega\|_{L^2(M)}^2 \right) \leq 4\|\delta\omega\|_{L^2(M)}^2 + p\|\Delta\omega + (2p-n)\omega\|_{L^2(M)}^2.$$

Notice that the l.h.s. of that last identity must be positive since it is actually $\sum_{i=1}^{n+1} \|d(\partial_{x_i}^T \lrcorner \omega)\|_{L^2(M)}^2 + \|\delta(\partial_{x_i}^T \lrcorner \omega)\|_{L^2(M)}^2$. Replacing ω by $\star\omega$ and using the Hodge symmetry of the buckling eigenvalues ensure that

$$\begin{aligned} (n-p)\Lambda_{1,p+1} & \left((\Delta\omega, \omega)_{L^2(M)} + (n-2p)\|\omega\|_{L^2(M)}^2 \right) \\ & \leq 4\|d\omega\|_{L^2(M)}^2 + (n-p)\|\Delta\omega + (n-2p)\omega\|_{L^2(M)}^2. \end{aligned}$$

Adding those two inequalities, we obtain

$$\begin{aligned} 2 \min(p\Lambda_{1,p-1}, (n-p)\Lambda_{1,p+1}) \cdot (\Delta\omega, \omega)_{L^2(M)} & \leq 4(\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2) \\ & \quad + p\|\Delta\omega + (2p-n)\omega\|_{L^2(M)}^2 \\ & \quad + (n-p)\|\Delta\omega + (n-2p)\omega\|_{L^2(M)}^2 \\ & = 4(\Delta\omega, \omega)_{L^2(M)} + n\|\Delta\omega\|_{L^2(M)}^2 \\ & \quad + 2(p(2p-n) + (n-p)(n-2p))(\Delta\omega, \omega)_{L^2(M)} \\ & \quad + (p(2p-n)^2 + (n-p)(n-2p)^2)\|\omega\|_{L^2(M)}^2 \\ & = n\|\Delta\omega\|_{L^2(M)}^2 + n(n-2p)^2\|\omega\|_{L^2(M)}^2 \\ & \quad + (4 + 2(n-2p)^2)(\Delta\omega, \omega)_{L^2(M)}. \end{aligned} \tag{41}$$

But (35) with $\gamma = 1$ gives

$$\|\omega\|_{L^2(M)} \leq \frac{1}{p(n-p+1)} \|\Delta\omega\|_{L^2(M)}.$$

Substituting that estimate in (41) yields the following inequality:

$$2 \min(p\Lambda_{1,p-1}, (n-p)\Lambda_{1,p+1}) \cdot \left(\|d\omega\|_{L^2(M)}^2 + \|\delta\omega\|_{L^2(M)}^2 \right) \leq C_{n,p} \|\Delta\omega\|_{L^2(M)}^2,$$

where $C_{n,p}$ is the constant defined in the first statement of Theorem 3.20. We deduce inequality (38) by taking ω as an eigenform of the buckling eigenvalue problem associated to $\Lambda_{1,p}$. Similarly to the first case, the equality cannot occur in (38).

2. If $n = 2p$, then (38) becomes, because of $\Lambda_{1,p-1} = \Lambda_{1,n-p+1} = \Lambda_{1,p+1}$,

$$\Lambda_{1, \frac{n}{2}-1} < \frac{C_{n, \frac{n}{2}}}{n} \cdot \Lambda_{1, \frac{n}{2}},$$

which is (39). □

Notice that, by the assumption $p \leq n-p$, Inequality (38) implies that

$$\min(\Lambda_{1,p-1}, \Lambda_{1,p+1}) < \frac{C_{n,p}}{2p} \cdot \Lambda_{1,p}.$$

References

- [1] C. Anné, *Principe de Dirichlet pour les formes différentielles*, Bull. Soc. Math. France **117** (1989), no. 4, 445–450.
- [2] M.S. Ashbaugh and R.S. Laugesen, *Fundamental tones and buckling loads of clamped plates*, Ann. Sc. Norm. Super. Pisa (4) **23** (1996), 383–402.
- [3] T. Chakradhar, K. Gittins, G. Habib and N. Peyerimhoff, *Magnetic Steklov operator on differential forms*, in progress.
- [4] D. Chen, Q.-M. Cheng, Q. Wang and C. Xia, *On eigenvalues of a system of elliptic equations and of the biharmonic operator*, J. Math. Anal. Appl. **387** (2012), no. 2, 1146–1159.
- [5] F. El Chami, N. Ginoux and G. Habib, *New eigenvalue estimates involving Bessel functions*, Publ. Mat. **65** (2021), 681–726.
- [6] F. El Chami, N. Ginoux, G. Habib and O. Makhoul, *Biharmonic Steklov operator on differential forms*, to appear in Ann. Math. Blaise Pascal.
- [7] L. Friedlander, *Some inequalities between Dirichlet and Neumann eigenvalues*, Arch. Ration. Mech. Anal. **116**, No. 2, 153–160 (1991).
- [8] L. Friedlander, *Remarks on the membrane and buckling eigenvalues for planar domains*, Mosc. Math. J **4** (2004), no. 2, 369–375.
- [9] S. Gallot and D. Meyer, *Opérateur de courbure et laplacien des formes différentielles d’une variété riemannienne*, J. Math. Pures Appl. **54** (1975), 259–284.
- [10] P. Guerini and A. Savo, *Eigenvalue and gap estimates for the Laplacian acting on p -forms*, Trans. Amer. Math. Soc. **356** (2004), no. 1, 319–344.
- [11] L. Hörmander, *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*, Springer, 2003.
- [12] S. Ilias and A. Shouman, *Inequalities between Dirichlet, Neumann and buckling eigenvalues on Riemannian manifolds*, Calc. Var. Partial Differential Equations **59** (2020), no. 4, Paper No. 127, 15 pp.
- [13] L.E. Payne, *Inequalities for eigenvalues of membranes and plates*, J. Rat. Mech. Anal. **4** (1955), 517–529.
- [14] G. Pólya, *Remarks on the foregoing paper*, J. Math. Phys. **31** (1952), 55–57 .
- [15] L. Provenzano and A. Savo, *Isoparametric foliations and the Pompeiu property*, Mathematics in Engineering **2** (2023), no. 5, 1–27.
- [16] S. Raulot and A. Savo, *A Reilly formula and eigenvalue estimates for differential forms*, J. Geom. Anal. **21** (2011), no. 3, 620–640.
- [17] M. Salo (ed.), *Unique continuation for elliptic equations*, lecture notes, 2014.
- [18] G. Schwarz, *Hodge decomposition - A method for solving boundary value problems*, Lecture notes in Mathematics, Springer, 1995.
- [19] U. Semmelmann, *Conformal Killing forms on Riemannian manifolds*, Math. Z. **245** (2003), no. 3, 503–527.