



MORPHOLOGICAL VARIABILITY WITHIN THE MITRATE MITROCYSTELLA INCIPIENS (ECHINODERMATA, STYLOPHORA) FROM HIGH-LATITUDE PERI-GONDWANAN REGIONS

Camille Vidal-Marty, Bertrand Lefebvre, Sabrina Renaud

► To cite this version:

Camille Vidal-Marty, Bertrand Lefebvre, Sabrina Renaud. MORPHOLOGICAL VARIABILITY WITHIN THE MITRATE MITROCYSTELLA INCIPIENS (ECHINODERMATA, STYLOPHORA) FROM HIGH-LATITUDE PERI-GONDWANAN REGIONS. 3rd virtual meeting of IGCP 735 Rocks and the Rise of Ordovician Life, Prague, IGCP 735, Nov 2024, Prague (République Tchèque), Czech Republic. hal-04824721

HAL Id: hal-04824721

<https://hal.science/hal-04824721v1>

Submitted on 7 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

3rd virtual meeting of IGCP 735

abstract volume

Prague, 18th – 20th November 2024



MORPHOLOGICAL VARIABILITY WITHIN THE MITRATE *MITROCYSTELLA INCIPiens* (ECHINODERMATA, STYLOPHORA) FROM HIGH-LATITUDE PERI-GONDWANAN REGIONS

Camille Vidal-Marty¹, Bertrand Lefebvre^{1} & Sabrina Renaud²*

*corresponding author: bertrand.lefebvre@univ-lyon1.fr

¹ UMR CNRS 5276 LGLTPE, Université Lyon 1, 69622 Villeurbanne, France

² UMR CNRS 5558 LBBE, Université Lyon 1, 69622 Villeurbanne, France

The mitrate *Mitrocystella incipiens* (Barrande, 1887) is a locally abundant stylophoran echinoderm, with a short stratigraphic range (late Darriwilian, *H. copiosus* graptolite Zone), but a wide palaeobiogeographic distribution in high-latitude peri-Gondwanan regions of the Mediterranean Province: Czech Republic (Prague Basin), France (Armorican Massif), northern Portugal, and Spain (Hesperian Massif). Two regional subspecies were identified by Chauvel (1937), on the basis of the morphology of their posterior thecal margin: slightly curved in Bohemian forms (*M. incipiens incipiens*), straight and truncated in Ibero-Armorican specimens (*M. incipiens miloni*). Therefore, the main objective of this study is to analyze the morphological disparity within the genus *Mitrocystella*, in order to test the existence of the two distinct morphotypes "*incipiens*" and "*miloni*". A representative sample of 40 complete and well-preserved individuals from Armorica, Bohemia and Iberia was selected. The geometric morphometric analysis of their theca confirms the existence of morphological differences between Armorican and Bohemian populations, but it also identifies that the Iberian specimens are morphologically intermediate between those from Brittany and the Prague Basin. These results support the idea that all populations of *Mitrocystella* belong to the same species, *M. incipiens*, which is characterized by a broad phenotypic plasticity. This ecophenotypism likely results from subtle environmental differences between the three regions, possibly related to bathymetry, nutrient availability, and/or volcanic activity (pH). Finally, the first example of teratological thecal plating in *Mitrocystella incipiens* is observed in a specimen from Spain. These results could thus be illustrated during an oral session.

References

- Barrande, J. 1887. *Système Silurien du centre de la Bohême. 1ère Partie. Recherches Paléontologiques. Volume 7. Classe des Echinodermes. Ordre des Cystidées*. Gerhard, Leipzig and Řivnáč, Prague.
 Chauvel, J. 1937. Les hétérostélées armoricaines. *Compte-rendu sommaire des séances de la Société Géologique et Minéralogique de Bretagne* 3, 1–3.