



**HAL**  
open science

## **Challenging the European southern refugium hypothesis: Species-specific structures versus general patterns of genetic diversity and differentiation among small mammals**

Debbi Pedreschi, Oxala García-Rodríguez, Glenn Yannic, Elena Cantarello, Anita Diaz, Duncan Golicher, Amanda Korstjens, Gerald Heckel, Jeremy Searle, Phillipa Gillingham, et al.

### ► **To cite this version:**

Debbi Pedreschi, Oxala García-Rodríguez, Glenn Yannic, Elena Cantarello, Anita Diaz, et al.. Challenging the European southern refugium hypothesis: Species-specific structures versus general patterns of genetic diversity and differentiation among small mammals. *Global Ecology and Biogeography*, 2018, 28 (2), pp.262-274. <10.1111/geb.12828>. <hal-04827439>

**HAL Id: hal-04827439**

**<https://hal.science/hal-04827439v1>**

Submitted on 6 Jan 2025


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

# Challenging the European southern refugium hypothesis: Species-specific structures versus general patterns of genetic diversity and differentiation among small mammals

Debbi Pedreschi<sup>1,2</sup>  | Oxala García-Rodríguez<sup>2</sup> | Glenn Yannic<sup>3</sup> | Elena Cantarello<sup>2</sup> | Anita Diaz<sup>2</sup> | Duncan Golicher<sup>2</sup> | Amanda H. Korstjens<sup>2</sup> | Gerald Heckel<sup>4,5</sup> | Jeremy B. Searle<sup>6</sup> | Phillipa Gillingham<sup>2\*</sup> | Emilie A. Hardouin<sup>2\*</sup> | John R. Stewart<sup>2\*</sup>

<sup>1</sup>Fisheries Ecosystem Advisory Services, Marine Institute, Oranmore, Galway, Ireland

<sup>2</sup>Faculty of Science and Technology, Bournemouth University, Poole, United Kingdom

<sup>3</sup>LECA – Laboratoire d'Ecologie Alpine – UMR CNRS 5553, Université Savoie Mont Blanc, Le Bourget-du-Lac, France

<sup>4</sup>Institute of Ecology and Evolution, University of Bern, Bern, Switzerland

<sup>5</sup>Swiss Institute of Bioinformatics, Lausanne, Switzerland

<sup>6</sup>Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York

## Correspondence

Debbi Pedreschi, Marine Institute, Rinville, Oranmore, Co. Galway, H91 R673, Ireland.  
Email: debbi.pedreschi@marine.ie

## Funding information

Bournemouth University

Editor: Richard Field

## Abstract

**Aim:** In this study, we conduct a quantitative meta-analysis to investigate broad patterns of genetic variation throughout large geographical regions in order to elucidate concordant geographical patterns across species and identify common historical processes to better inform the “cryptic refugia” versus the traditional “southern refugia” hypothesis debate.

**Location:** Europe.

**Time period:** Late Pleistocene to present day.

**Major taxa studied:** Small mammals (Rodentia, Eulipotyphla).

**Methods:** A meta-analysis was performed on large-scale patterns of genetic diversity for 19 species from 59 papers. For each species, haplotype and nucleotide diversity were calculated using the mitochondrial D-loop and compared to the species' range.

**Results:** No consistent patterns were observed between mitochondrial DNA (mtDNA) diversity indices (nucleotide and haplotype diversity) and any of the indicators of distribution examined [latitude and longitude (max, min, centre, range)]. The patterns of genetic diversity observed in all the 19 species studied appear to be species-specific.

**Main conclusions:** In contrast to the traditional southern refugial hypotheses, we found no evidence for a consistent south–north post-glacial expansion. Instead individual species appear to respond to climate oscillations in niche-specific ways. This individual nature of each species' phylogeographical history indicates a complex web of post-glacial recolonization dynamics across Europe.

## KEYWORDS

colonization, Europe, mammals, molecular diversity, mtDNA, phylogeography, refugia

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors Global Ecology and Biogeography Published by John Wiley & Sons Ltd

\*Contributed equally to this work.

## 1 | INTRODUCTION

The investigation of geographical distributions of genetic lineages and genetic diversity has allowed inferences about post-glacial colonization history in recent decades (the phylogeographical approach: Avise, 2000). There has been a particular focus on Europe, with studies on a wide range of plant and animal organisms (Bilton et al., 1998; Hewitt, 1999, 2004; Stewart, Lister, Barnes, & Dalén, 2010; Taberlet, Fumagalli, Wust-Saucy, & Cosson, 1998).

Initially, it was felt the data suggested that the majority of European temperate species recolonized central and northern Europe after the last glaciation from southern refugial areas (Hewitt, 1999, 2004; Seddon, Santucci, Reeve, & Hewitt, 2001). The loss of genetic diversity from southern to northern Europe has long been remarked upon (e.g., Hewitt, 1999, 2001, 2004; Kryštufek & Griffiths, 2002; Taberlet et al., 1998). European studies led to the identification of three glacial refugia in southern Europe, on the Iberian, Italian and Balkan peninsulas, with three main colonization models dominating latitudinal expansion patterns (Hewitt, 1999). Increasingly however, the existence of refugia outside the traditional southern peninsulae (so-called cryptic northern, or extra-Mediterranean, refugia: Bilton et al., 1998; Stewart & Lister, 2001; Schmitt & Varga, 2012 but see Tzedakis, Emerson, & Hewitt, 2013) has been recognized for a wide range of species including small mammals (e.g., central Europe; Deffontaine et al., 2005; Fink, Excoffier, & Heckel, 2004). Boreal and arctic species may be considered to be in refugia today, in comparison to colder periods of our glacial history when they maintained much wider distributions (Stewart & Dalén, 2008). These conflicting reports indicate that there is discordance between competing hypotheses of European post-glacial biogeographical processes.

The loss of both genetic and species diversity from southern to northern Europe has historically been thought to be a result of expansion from few limited refugia, that is, leading edge allele loss (Hewitt, 1996, 1999; Ibrahim, Nichols, & Hewitt, 1996; Taberlet et al., 1998; Waters, Fraser, & Hewitt, 2013), combined with recurrent losses of divergent populations over multiple glacial cycles (Hewitt, 1996; Hofreiter & Stewart, 2009; Seddon et al., 2001; Stewart, 2009; Stewart et al., 2010). Within species, it is generally expected that more northerly populations will be less genetically diverse than their southern counterparts (Hewitt, 2004; Seddon et al., 2001). Although multiple studies have examined patterns within species, few analyses have compared between species. Still fewer have looked at the potential variation in the type of movement (e.g., step-wise versus phalanx-wise; Habel, Ulrich, & Assmann, 2013), or concerned themselves with the lower latitude limit, or "rear edge," of species ranges (Hampe & Petit, 2005).

High-level "overview" hypotheses and trends can encourage understanding at a macro level and enable testing of multiple scenarios. General patterns may not be obvious when individual taxa and their phylogenetic relationships are examined separately (Taberlet et al., 1998). For this reason, syntheses are necessary and useful to provide a consistent overview—it is important to see both the woods *and* the

trees. This paper investigates patterns of genetic diversity by examining various diversity indices across 19 small mammal species to test for trends and commonalities. Small mammals were selected because they comprise many species (Mitchell-Jones et al., 1999; Tougaard & Renvoisé, 2008); have limited dispersal capabilities related to their small size, enabling relatively accurate estimation of range sizes; and have not been affected by man to the same extent as many larger mammals (Kryštufek & Griffiths, 2002) (with the exclusion of commensals such as domestic rats and house mice, which were excluded here). Furthermore, these species have been important in relation to the study of ice-age refugia in Europe (Bilton et al., 1998; Taberlet et al., 1998). Small mammals display short generation times and a fast-evolving genome (Martin & Palumbi, 1993; Tougaard & Renvoisé, 2008) enabling us to detect the genetic variation of interest. A large number of studies from species retaining similar key characteristics were available to make this synthesis possible. Comparison of these species may provide insight into genetic diversity patterns related to colonization pathways in Europe. Here we investigate the previously hypothesized pathways of colonization through the examination of the mitochondrial DNA (mtDNA) control region of a range of small mammal species, and test for congruence of broad-scale genetic interspecific diversity patterns among those species that may infer shared evolutionary history. Further, we conduct an intraspecific analysis of four species with sufficient numerical and geographical coverage in Europe to enable comparison of their individual phylogeographical histories, and use this to further identify whether they support or contradict the "southern refugia" or "cryptic refugia" hypotheses.

## 2 | METHODS

The study area was taken at a broad scale to be political Europe. An initial list of potential species was compiled covering small mammal species from the orders Rodentia and Eulipotyphla. mtDNA was selected due in part to its relatively rapid rate of mutation, and its haploid maternal inheritance mechanism, both of which make it useful for the elucidation of population historical and demographical changes (Avise, 1995; Avise et al., 1987; Moritz, 1994). These features have allowed mtDNA to become one of the most frequently used markers in molecular ecology and phylogeographical studies, particularly in mammals (e.g., Centeno-Cuadros, Delibes, & Godoy, 2009; Centeno-Cuadros & Godoy, 2010; Hewitt, 2004). GenBank ([www.ncbi.nlm.nih.gov/genbank/](http://www.ncbi.nlm.nih.gov/genbank/)), the primary public archive for the deposition of genetic sequences, was searched for mtDNA control region (a.k.a. D-loop) sequences for these species. Where sequences were available, the primary literature and source papers for each study were found through online data searches (e.g., Google Scholar, PubMed). From these papers, frequencies and/or numbers of individuals for each haplotype were obtained (where published), enabling the full reconstruction of the original study data set through manual re-assembly using FASTA files downloaded from GenBank and associated haplotype data. These reconstructed data sets were

subsequently combined for each species to create species-level diversity estimates across the sampled range.

The sequences were aligned using the ClustalW algorithm in MEGA v.6 (Tamura, Stecher, Peterson, Filipki, & Kumar, 2013) and revised manually. The differences in the lengths and/or sections of the mtDNA control region sequenced for each species were too great to enable analysis of the same mtDNA region in the different species. Within each species, sequences were standardized to the minimum overlapping sequence length available for each species. The total number of haplotypes and variable sites observed and the average number of nucleotide differences per sequence were calculated and reported.

Haplotype and nucleotide diversity are frequently used as measures of diversity in population genetics studies (Egeland & Salas, 2008; Goodall-Copestake, Tarling, & Murphy, 2012) and were estimated using DNASP v. 5.10.01 (Librado & Rozas, 2009). The numbers of sequences available per species varied markedly. In an attempt to compensate for this, standardized haplotype and nucleotide diversity estimates were calculated for each species. Standardized haplotype diversity was calculated for each species by random subsampling from the available individuals to match the smallest sample size encountered. Standardized nucleotide diversity was calculated for each species by random subsampling from the available unique haplotypes to match the smallest number of unique haplotypes encountered.

To investigate direction of range expansion, mtDNA diversity indices were related to distribution. Distribution data were downloaded from the International Union for Conservation of Nature (IUCN) terrestrial mammals database (<https://maps.iucnredlist.org>, date of download 27/05/2014). The ranges of our selected species were clipped to the extent of political Europe using country outlines from the Global Administrative Areas database [<http://www.gadm.org>, downloaded 13/05/2014] and maximum and minimum latitude and longitude were calculated for each species. The clipped extents of occurrence were then transformed using the Lambert Azimuthal Equal Area projection for Europe and area (in km<sup>2</sup>) was calculated for each. These variables were then used to test for relationships with absolute and standardized haplotype and nucleotide diversity estimates both across and within species using correlations (Spearman's and Pearson's) and regressions (linear, quadratic and multiple) carried out in the SPSS v. 20 statistical environment (IBM Corp. 2011).

Haplotype sampling coverage (i.e., what proportion of the population/genetic diversity is represented by our samples) was estimated using the methods of Dixon (2006), based on the number of haplotypes and individuals sampled using the Stirling probability distribution and Bayes theorem to obtain a posterior distribution of the total number of haplotypes in the population including those not yet observed. Haplotype accumulation curves were also produced using the R package "spider" (R Core Team, 2017) (Brown et al., 2012).

For a subset of four species, population genetic diversity and distributional trends were investigated. These investigations were limited to those species that had 75% or higher haplotype sampling completeness (according to Dixon's measure, Dixon, 2006),

had large enough distributions to display evidence of post-Last Glacial Maximum (post-LGM) expansion and a minimum of 100 sampled individuals spread reasonably well across their distribution [*Cricetus cricetus*, *Erinaceus europaeus*, *Microtus arvalis*, *Clethrionomys* (= *Myodes*) *glareolus*]. Individuals of these species were pooled into populations (minimum of five individuals per population, Centeno-Cuadros et al., 2009) based upon the reported study sites in the respective papers, and the aforementioned diversity estimates were again calculated. Where this was not possible (location information not provided, or fewer than five individuals per population), individuals were pooled for a country-level estimate. Sampling site information was reported in a number of different ways in the literature (site names/nearest town, latitude and longitude/grid references) and so all were converted to latitudes and longitudes in decimal degrees for analysis. Where only regional location data were provided, central latitude and longitude values for the region were used for the within-species analysis.

### 3 | RESULTS

#### 3.1 | Between-species investigation

The data collection was ceased in May 2015 to allow the meta-analysis to proceed. A total of 113 species were investigated for sequences deposited in GenBank (searched using species name, with D-loop or control region as keywords). Fifty-four species were found to have GenBank data from 123 studies. Authors were contacted for studies in which the required information was not presented in the paper or the supporting materials, or for which a published paper could not be found (22% of studies). Of those contacted 61% agreed to provide the information; however, many of the authors of the older studies were no longer contactable.

Difficulty was met in the re-assembly of data from the primary literature. Similar to Goodall-Copestake et al. (2012), we found a wide range in the behaviours of those depositing sequences in public archives. Of those that did deposit their sequences, some deposited sequences for all individuals examined, some deposited only each unique haplotype encountered, and others deposited only a single haplotype with notes on the edits needed to reconstruct all of the haplotypes from the study (a time-consuming process). Of those that did deposit sequences in GenBank, many papers did not report the frequencies/numbers of individuals for each haplotype, thus making reconstruction impossible and necessitating their exclusion from the study, greatly reducing the number of species/studies that could be included, and highlighting the need for detailed data archiving. Only one study (Tougard, Renvoisé, Petitjean, & Quéré, 2008) had to be excluded from the between-species analysis due to the segments of the mtDNA control region analysed not overlapping with those of the other studies examined. These sequences were retained for the within-species population-level diversity estimates and comparisons. This process resulted in a final count of 19 species with complete/reconstructed useable data from 59 papers (see Table 1).

**TABLE 1** Distributional area and primary habitat type of the study species

Order	Family	Species	Common name	Distributional area (10 <sup>6</sup> km <sup>2</sup> )	Habitat
Eulipotyphla	Erinaceidae	<i>Erinaceus concolor</i>	Eastern European hedgehog	1.35	Urban, suburban and agricultural areas
Eulipotyphla	Erinaceidae	<i>Erinaceus europaeus</i>	Western European hedgehog	3.81	Man-made habitats, agricultural land, deciduous woodland, and grasslands
Eulipotyphla	Erinaceidae	<i>Erinaceus roumanicus</i>	Northern white-breasted hedgehog	6.03	Man-made habitats farmland, scrubby and shrubby habitats
Eulipotyphla	Soricidae	<i>Crocidura russula</i>	Greater white-toothed shrew	1.93	Shrubland, open habitats, forest edges with abundant vegetation, cultivated fields, urban areas
Eulipotyphla	Soricidae	<i>Sorex antinorii</i>	Valais shrew	0.29	Areas with dense vegetation cover
Eulipotyphla	Soricidae	<i>Sorex minutus</i>	Eurasian pygmy shrew	14.29	Damp areas with dense vegetation; swamps, grasslands, heaths, sand dunes, woodland edge, rocky areas, shrubland, and montane forests
Rodentia	Cricetidae	<i>Arvicola sapidus</i>	Southern water vole	0.88	Hydrophilic vegetation and shorelines
Rodentia	Cricetidae	<i>Chionomys nivalis</i>	European snow vole	0.89	Open, rocky areas, typically above the tree line
Rodentia	Cricetidae	<i>Cricetus cricetus</i>	Common hamster	6.36	Man-made habitats, fertile steppe and grassland
Rodentia	Cricetidae	<i>Lemmus lemmus</i>	Norway lemming	0.76	Alpine and subarctic habitats, for example, peat bogs, dwarf shrub heaths, and sparsely vegetated slopes and ridges
Rodentia	Cricetidae	<i>Microtus guentheri</i>	Günther's vole	0.64	Dry grasslands with sparse vegetation on well-drained soil (natural and man-made)
Rodentia	Cricetidae	<i>Microtus arvalis</i>	Common vole	6.89	Open habitats including meadows, steppe, and agricultural areas
Rodentia	Cricetidae	<i>Microtus thomasi</i>	Thomas's pine vole	0.11	Open areas with deep soil; meadows, high mountain pastures and arable farmland
Rodentia	Cricetidae	<i>Clethrionomys (Myodes) glareolus</i>	Bank vole	8.68	All woodland, preferring densely vegetated clearings, woodland edge, and riverbanks. Also found in scrub, parkland, and hedges
Rodentia	Muridae	<i>Apodemus mystacinus</i>	Eastern broad-toothed field mouse	0.80	Forest with rocky areas; sparse cover of grass or shrubs
Rodentia	Muridae	<i>Mus macedonicus</i>	Macedonian mouse	1.39	Cultivated land, road verges, sand dunes, shrubland, and densely vegetated riverbanks. Absent from dense forests, and avoids urban areas. Restricted to areas with > 400 mm of rain per annum
Rodentia	Muridae	<i>Mus spicilegus</i>	Mound-building mouse	0.83	Natural steppe grasslands, pastures, open woodland and clearings
Rodentia	Sciuridae	<i>Sciurus vulgaris</i>	Eurasian red squirrel	19.9	Lowland to subalpine coniferous forests and deciduous woods, along with parks and gardens
Rodentia	Sciuridae	<i>Spermophilus suslicus</i>	Speckled ground squirrel	0.8	Open areas with short grass; steppes, pastures, road verges and occasionally cultivated

Note. Distributional area refers to the area occupied by each species as obtained from the IUCN database to the nearest 10<sup>6</sup> km<sup>2</sup>, which does not take patchiness into account. Habitat is the primary vegetation cover type preference for each species as obtained from IUCN species profiles ([www.iucn-redlist.org](http://www.iucn-redlist.org)). GenBank accession numbers and source papers are available in Supporting Information Appendix S1.

The minimum number of individuals sampled for a single species was 21 for the eastern broad-toothed field mouse (*Apodemus mystacinus*; sampled from Turkey, Syria, Greece, Bulgaria). This figure is close to the recommended minimum sample size of 25 from Goodall-Copestake et al. (2012) for the cytochrome c oxidase subunit I (COX1) mtDNA gene. We realize that this recommendation was for population level rather than species sampling; however, we would highlight that the *cox1* gene is a more conserved region of the mtDNA. This, coupled with the limitations of the available data, led us to accept 21 individuals as the lowest acceptable sample size for this analysis, and thus 21 individuals were randomly selected for each species to calculate the standardized haplotype diversity. The lowest number of haplotypes observed was six haplotypes in 55 individuals of the northern white-breasted hedgehog (*Erinaceus roumanicus*; Czech Republic, Slovak Republic, Poland); accordingly, six randomly selected unique haplotypes were selected from each species for calculating standardized nucleotide diversity values.

Haplotype diversity measures ranged from 0.32 to 0.99 and standardized haplotype diversities from 0.35 to 0.99 (Table 2). Nucleotide diversity measures ranged from 0.002 to 0.048 and standardized nucleotide diversity from 0.006 to 0.069 (Table 2). Although the number of haplotypes increased significantly ( $R^2 = 0.744$ ,  $p < 0.0001$ ) with area, as is expected, no relationship was evident between area and any of the measures of genetic diversity used.

Diversity measures can be influenced by under-sampling (Goodall-Copestake et al., 2012); haplotype diversity generally increases with sample size (Pereira, Cunha, & Amorim, 2004). Haplotype sampling coverage estimates (i.e., the proportion of the population genetic diversity represented by our samples) ranged from 19 to 100% complete with the majority (15 out of 19) estimated at over 80% complete (Supporting Information Table S2.1, Appendix S2). However, caution must also be employed when interpreting the level of sampling completeness: a report of 90% coverage of the haplotypes that occur in one region may only be a fraction of the haplotypes that occur within the species' whole range. As such, completeness values of 100% are unlikely to indicate that every haplotype that exists has been sampled but instead indicate a reasonable sample of the population. Estimates of haplotype sampling completeness (Dixon's measure: Dixon, 2006) were well supported by the haplotype accumulation curves (Supporting Information Figure S2.1, Appendix S2), another method for estimating sampling coverage.

Species mtDNA diversity indices (outlined in Table 2) were not found to consistently correlate with any of the indicators of distribution examined [latitude and longitude (max, min, centre, range), and area (Figure 1)], either when tested individually or in multiple regressions.

### 3.2 | Within-species investigation

For a subset of four species, within-species diversity and population distributional trends were investigated. These investigations were limited to those species that had 75% or higher haplotype

sampling completeness, had large enough distributions to display evidence of post-LGM expansion and a minimum of 100 sampled individuals spread reasonably well across their distribution (*C. cricetus*, *E. europaeus*, *M. arvalis*, *C. glareolus*). No consistent patterns were observed between any diversity index and distribution (Supporting Information Appendix S2): patterns of diversity varied in species-specific ways (Figure 2).

## 4 | DISCUSSION

Overall, the patterns of genetic diversity observed appear species-specific as individual species responded to climate oscillations in niche-specific ways. Each of the taxa that have been sufficiently studied to allow within-species analysis illustrate complex and divergent phylogeographical histories. It has been recognized that most species are not likely to "track" their preferred habitat back to the refugial areas as climate changes (Dalén et al., 2007; Lagerholm et al., 2014). As such, for each glaciation that occurred, a substantial proportion of each species' lineage was lost, and with that its genetic heritage, complicating the interpretation of colonization patterns. The replacement of one phylogroup with another in line with climate oscillations has also been observed (Brace et al., 2012, 2016; Martínková et al., 2013; Palkopoulou et al., 2016; Searle et al., 2009) indicating that present-day genetic diversity alone can be insufficient to explain historical patterns (Hofreiter & Stewart, 2009; Searle et al., 2009).

The direction of range expansions can be inferred by genetic diversity gradients (leading-edge founder events, Ibrahim et al., 1996; White, Perkins, Heckel, & Searle, 2013). Many widespread European species dispersed into Europe via western Asia through repeated waves of east-west colonization throughout the Quaternary, thus indicating a decrease in species richness from east to west (an oceanic-continental gradient: Fløjgaard, Normand, Skov, & Svenning, 2011; Stewart et al., 2010). Conversely, many species recolonized after the LGM from the south, creating a south-north genetic diversity gradient (Hewitt, 1996, 1999, 2000). Importantly, this meta-analysis indicates that common genetic signals are missing and instead it is species-specific responses that are of primary importance for reconstructing post-glacial recolonization patterns.

### 4.1 | Within-species analysis

All four species (*C. cricetus*, *E. europaeus*, *M. arvalis*, *C. glareolus*) for which intraspecific investigation was possible highlight a species-specific response to climate change, as each provides its own complex colonization history different to each of the others. Interestingly, these species also highlight the presence of "northern" refugia—a term often used to cover all refugia outside of the traditional "southern" refugia of the European Iberian, Italian and Balkan peninsulas, but that actually refer to a number of regions, such as the Carpathians and areas of central Europe—often around Germany—and are increasingly being viewed not as traditional "refugia" but instead as a continuum of fractured microhabitats (Bhagwat & Willis, 2008).

**TABLE 2** Genetic diversity of the species studied: Species diversity measures; normal and standardized haplotype (HapIDiv, StdHapIDiv) and nucleotide (NucIDiv, StdNucIDiv) diversity measures for the between-species analysis (see Methods)

Species	Individuals included	Haplotypes reported	Haplotypes observed	% Sampling complete	Sequence length	Sequence sites	Variable sites	AvgNucDiff (k)	HapIDiv (SD)	StdHapIDiv (SD)	NucIDiv (SD)	StdNucIDiv (SD)
<i>Apodemus mystacinus</i>	21	21	19	19	1,009	136	136	45.48	0.99 (0.018)	0.99 (0.018)	0.045 (0.009)	0.065 (0.012)
<i>Arvicola sapidus</i>	240	87	76	96	203	61	61	9.47	0.97 (0.005)	0.98 (0.023)	0.047 (0.002)	0.058 (0.007)
<i>Chionomys nivalis</i>	584	19	10	100	804	88	88	24.67	0.76 (0.012)	0.64 (0.095)	0.031 (0.001)	0.004 (0.013)
<i>Cricetus cricetus</i>	397	48	27	100	337	19	19	2.80	0.83 (0.009)	0.86 (0.047)	0.008 (0.0002)	0.011 (0.002)
<i>Crocodyrus russula</i>	388	38	18	100	323	33	33	2.05	0.63 (0.02)	0.48 (0.121)	0.006 (0.004)	0.013 (0.003)
<i>Erinaceus concolor</i>	63	32	17	100	380	23	23	6.96	0.91 (0.019)	0.86 (0.048)	0.018 (0.002)	0.02 (0.069)
<i>Erinaceus europaeus</i>	386	66	56	100	400	47	47	9.23	0.95 (0.005)	0.96 (0.026)	0.023 (0.001)	0.031 (0.005)
<i>Erinaceus roumanicus</i>	55	7	6	100	405	6	6	0.81	0.32 (0.078)	0.35 (0.131)	0.002 (0.001)	0.006 (0.001)
<i>Lemmus lemmus</i>	104	24	17	100	165	17	17	1.31	0.8 (0.037)	0.61 (0.116)	0.008 (0.001)	0.018 (0.003)
<i>Microtus guentheri</i>	33	-	26	52	387	50	50	8.62	0.98 (0.014)	0.99 (0.019)	0.022 (0.002)	0.025 (0.005)
<i>Microtus arvalis</i>	677	114	84	100	273	59	59	4.65	0.88 (0.01)	0.95 (0.031)	0.017 (0.001)	0.037 (0.005)
<i>Microtus thomasi</i>	169	15	84	81	539	96	96	8.32	0.98 (0.004)	0.99 (0.018)	0.015 (0.001)	0.025 (0.004)
<i>Mus macedonicus</i>	71	39	52	49	730	66	66	6.39	0.99 (0.006)	0.99 (0.022)	0.009 (0.001)	0.011 (0.002)
<i>Mus spicilegus</i>	36	21	13	100	429	16	16	4.13	0.88 (0.034)	0.9 (0.047)	0.01 (0.001)	0.01 (0.002)
<i>Clethrionomys (Myodes) glareolus</i>	1,078	104	113	100	258	61	61	3.73	0.86 (0.007)	0.82 (0.06)	0.014 (0.001)	0.036 (0.012)
<i>Sciurus vulgaris</i>	799	225	155	100	225	81	81	4.74	0.95 (0.003)	0.93 (0.033)	0.021 (0.001)	0.034 (0.006)
<i>Sorex antinorii</i>	275	40	82	97	331	41	41	4.79	0.96 (0.005)	0.98 (0.023)	0.014 (0.0003)	0.017 (0.002)
<i>Sorex minutus</i>	250	158	134	76	516	228	228	21.01	0.98 (0.003)	0.995 (0.016)	0.041 (0.001)	0.056 (0.008)
<i>Spermophilus suslicus</i>	75	17	38	81	310	58	58	14.98	0.95 (0.013)	0.94 (0.036)	0.048 (0.002)	0.069 (0.01)

Note. Haplotypes reported refers to the summed number of haplotypes across studies. Haplotypes observed are the number of haplotypes observed when studies were pooled by species. Sequence length refers to the minimum overlapping sequence length available for each species. AvgNucDiff refers to the average number of nucleotide differences.

Evidence from several taxa and multiple studies have suggested the existence of cold-adapted species outside of the Mediterranean peninsulae during glacial periods (e.g., Bilton et al., 1998; Stewart et al., 2010). These glacials in fact represented periods of expansion for cold-adapted species in comparison to modern climate (Hofreiter & Stewart, 2009; Stewart et al., 2010). Taxa such as Norway lemming *Lemmus lemmus*, collared lemming *Dicrostonyx torquatus*, Arctic fox *Alopex lagopus*, rock ptarmigan *Lagopus muta* and mountain avens *Dryas octopetala* were found in mid-latitude Europe during glacial periods (Stewart et al., 2010 and references therein). Some temperate taxa (especially small mammals) may also have been located further north than often suggested (Stewart & Lister, 2001). Furthermore, studies have indicated the presence of a wide range of flora (primarily small-seeded, wind dispersed, coniferous trees) in more northerly locations than previously thought possible (Bhagwat & Willis, 2008; Stewart & Lister, 2001), highlighting that central Europe may have been much more habitable during these periods than previously hypothesized.

#### 4.1.1 | *Cricetus cricetus*

Neumann, Jansman, Kayser, Maak, and Gattermann (2004) and Neumann et al. (2005) revealed the separation of two distinct lineages of *C. cricetus* in Europe, one labelled “North” and one “Pannonian” (based on multiple mtDNA markers and microsatellites). Here, we found that fringe populations of the “North” lineage demonstrate lower diversity than those from the core area centred around Germany (Figure 2a), where the “Central” and “West” clades of the Northern lineage are to be found (Banaszek, Jadwiszczak, Ratkiewicz, & Ziomek, 2009). Although of a similarly high diversity level (Figure 2a), the Czech Republic population shares affinities with the southern “Pannonian” region (Neumann et al., 2004, 2005). Polish populations present a complex pattern with mixed signals from both Russia and Pannonia but with generally low diversity levels reflective of their status as a fringe population, similar to those of the low diversity “West” clade of the North lineage, despite their central location (Banaszek & Ziomek, 2011; Banaszek et al., 2009; Banaszek, Jadwiszczak, Ratkiewicz, Ziomek, & Neumann, 2010). Unfortunately, haplotype information was not available from Neumann et al. (2005); as such we were unable to include their Hungarian samples in this meta-analysis and so the picture presented here is somewhat incomplete in relation to the “Pannonian” lineage. The same main lineages identified above are also identified using cytochrome *b*, 16S ribosomal (r)RNA and microsatellite data (Banaszek et al., 2010; Neumann et al., 2005); however, Neumann et al. (2005) concluded that the contemporary phylogenetic structure observed is due to recolonization from eastern refugia.

#### 4.1.2 | *Clethrionomys* [= *Myodes*] *glareolus*

Despite the availability of over 1,000 mtDNA control region sequences sampled from individuals of *C. glareolus* from multiple studies (see Table 1 and Supporting Information Appendix S2), use of

these sequences has primarily been restricted to investigating the associated *Puumala* virus (Dekonenko et al., 2003; Johansson et al., 2008; Razzauti, Plyusnina, Niemimaa, Henttonen, & Plyusnin, 2012) or the effects of Chernobyl radiation (Matson, Rodgers, Chesser, & Baker, 2000; Meeks et al., 2007; Meeks, Chesser, Rodgers, Gaschak, & Baker, 2009; Wickliffe et al., 2006) on *C. glareolus*. Figure 2b indicates diversity values generally reflective of the aforementioned complexity; however, regionally restricted sampling does not enable accurate phylogeographical inferences to be made using the control region sequences available here. Of note however, even with this low distributional coverage, is the considerable variation in diversity values in Austria and the Ukraine which would seem to indicate an underlying complexity in colonization pattern.

Investigations into colonization patterns of *C. glareolus* have also been carried out using cytochrome *b* data (Colangelo, Aloise, Franchini, Annesi, & Amori, 2012; Deffontaine et al., 2005, 2009; Kotlík et al., 2006), and mitogenome sequencing (Filipi, Marková, Searle, & Kotlík, 2015). These studies have revealed perhaps the most complex phylogeographical pattern of any small mammal studied to date, with up to seven proposed well-defined clades, including some with subclades (Deffontaine et al., 2009; Filipi et al., 2015). Earlier papers provided evidence of multiple refugia outside of the traditional “southern” areas (Deffontaine et al., 2005, 2009; Kotlík et al., 2006); however, full mitogenome analysis (Filipi et al., 2015) has allowed the resolution of both “northern” and “southern” refugia. It appears that post-glacial expansion was likely via multiple routes, from multiple refugia, with the traditional Mediterranean peninsular refugia maintaining long-term intraspecific populations throughout the Pleistocene but without necessarily contributing to the post-glacial recolonization of mainland Europe (Bilton et al., 1998; Colangelo et al., 2012; Deffontaine et al., 2005; Filipi et al., 2015). Kotlík et al. (2006) provided evidence of a Carpathian refugium and subsequent expansion, pre-dating the LGM, supported by fossil evidence, along with plant pollen and macrofossils indicating the presence of suitable habitat. Surprisingly, mitogenomic phylogeny has grouped parts of Britain, Sweden and Norway within the “Carpathian” lineage, which may be due to a two-phase colonization, with the “Western” clade replacing the “Carpathian” clade throughout parts of central Europe and Britain (Filipi et al., 2015). Multiple colonization events are evident throughout the species’ range, resulting in geographically disjointed lineages that arose from multiple northern refugia.


#### 4.1.3 | *Erinaceus europaeus*

The European colonization of *E. europaeus* is perhaps one of the best characterized, and has been held up as a prime example of recolonization from southern refugial areas (Hewitt, 1999, 2000). Three *E. europaeus* clades have been identified, a western group stemming from the Iberian peninsula up through France to the Netherlands, UK and Ireland, a central European group recolonizing from Italy up into Austria, Switzerland, Germany, the Netherlands, Scandinavia and Estonia, and a regionally restricted Sicilian group (Bolfíková & Hulva, 2012; Seddon et al., 2001). These groups are strongly separated









using mtDNA (D-loop, cytochrome *b*) but not discriminated using nuclear microsatellites (Bolfíková & Hulva, 2012). A contact zone with the sister species *E. roumanicus* exists in central Europe (Poland, the Czech Republic, Austria and Italy); however, reproductive isolation appears to be maintained (Bolfíková & Hulva, 2012).

It has been hypothesized that long-distance colonization can cause diversity loss through serial bottlenecks (Hewitt, 1996); however, with the exception of island populations (e.g., UK and Ireland) or those separated by large bodies of water (e.g., Norway) where only single haplotypes were found, diversity was not found to be consistently lower in northern regions compared to southern “refugial” areas (Figure 2c). Indeed, the highest haplotype diversity

values were found in more central European regions (e.g., Germany, France, Switzerland, Netherlands and areas of the Czech Republic all demonstrated higher haplotype diversity than those in Iberia and Italy). This could potentially be due to secondary contact between the two lineages but has also been suggested as possible evidence of a northern refugium for this species (Stewart & Lister, 2001). Seddon et al. (2001) similarly found a distinct group of six mitotypes restricted to southern Germany, northern Switzerland, southern Austria and Poland. Furthermore, the relatively high diversity values across mainland European populations is comparable to levels seen in *M. arvalis* and *C. glareolus*, which are hypothesized to be due to more recent expansion within central Europe. Interestingly, Bhagwat



**FIGURE 1** Mitochondrial DNA (mtDNA) diversity indices of all 19 small mammal species examined in relation to area for (a) haplotype diversity and (b) nucleotide diversity (see Supporting Information Appendix S2 for standardized estimates). The dotted line indicates the mean diversity value and thus an indication for “high” and “low” diversity for the small mammal species investigated. Distributions were further classed into endemic (black closed circles < 1 million km<sup>2</sup>) and widespread areas (open circles between 1–5 million km<sup>2</sup>, or grey circles > 5 million km<sup>2</sup>). Species mtDNA diversity indices were not found to consistently correlate with area (illustrated here), or any of the indicators of distribution examined. Note: axis break and scale change

(a) *Cricetus cricetus*(b) *Clethrionomys glareolus*(c) *Erinaceus europaeus*(d) *Microtus arvalis*

**FIGURE 2** Maps and mitochondrial DNA (mtDNA) control region haplotype diversity estimates of *Cricetus cricetus*, *Clethrionomys (Myodes) glareolus*, *Erinaceus europaeus* and *Microtus arvalis* used for within-species population analyses. Where sufficient information was available (minimum of five individuals from a region/location) individuals were grouped into populations within countries and for diversity estimates. Where this was not possible, individuals were pooled for a country-level estimate. Bar charts indicate the diversity level for each population, bar colours correspond to the map markers and shaded areas map the species' known distribution International Union for Conservation of Nature (IUCN), [www.iucnredlist.org](http://www.iucnredlist.org). "Missing" bars indicate monomorphic populations. No consistent patterns were observed between haplotype diversity and distribution: species diversity patterns varied in species-specific ways [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

and Willis (2008) also suggest that the genetic evidence of a southern refugium for this species does not fit with their analysis, which indicated that *E. europaeus* has the life history traits suited to persistence in northern refugia.

#### 4.1.4 | *Microtus arvalis*

*Microtus arvalis* is a well-known example of a species that does not conform to the southern refugium hypothesis (Fink et al., 2004; Heckel, Burri, Fink, Desmet, & Excoffier, 2005; Tougaard et al., 2008). Multiple genetic markers (cytochrome *b*, control region, microsatellites) indicate that there are at least five main evolutionary lineages of *M. arvalis* in Europe [Western (North and South), Central, Eastern, Italian and Balkan] with further levels of complexity within these lineages (Buzan et al., 2010; Martínková et al., 2013; Stojak, McDevitt, Herman, Searle, & Wójcik, 2015). Furthermore, all estimates of divergence times for these lineages indicate a pre-LGM or LGM (c. 18,000 years ago) split (Fink et al., 2004; Heckel et al., 2005; Lischer, Excoffier, & Heckel, 2013; Stojak et al., 2015). *M. arvalis* was not restricted to the classic southern refugia, and instead persisted in multiple northern refugia, likely in patchy favourable habitats (Buzan et al., 2010; Martínková et al., 2013; Stojak et al., 2015; Tougaard et al., 2008). The Eastern lineage is hypothesized to have arisen in the northern Balkans and/or the Carpathian basin, with the Central lineage arising north of the Alps (Braaker & Heckel, 2009; Buzan et al., 2010; Beysard & Heckel, 2014; Stojak et al., 2015; Stojak et al., 2016) although more detailed sampling is required to confirm this. Switzerland has been found to present an interesting case of contact and limited admixture (Italian, Western and Central lineages), as a result of comparatively recent Alpine colonization and hybridization (Beysard & Heckel, 2014; Braaker & Heckel, 2009; Sutter, Beysard, & Heckel, 2013).

Although patterns are not immediately evident from Figure 2d, closer inspection of the control region data reveals support for the above reported results. High genetic diversity is evident throughout the range of *M. arvalis*, with geographical partitioning in keeping with the lineages described above. Furthermore, the region with the most variable diversity values is Switzerland, the location identified as a multi-lineage contact zone (Beysard & Heckel, 2014; Braaker & Heckel, 2009; Stojak et al., 2015; Sutter et al., 2013). In contrast to Heckel et al. (2005) we did not detect a significant relationship between haplotype diversity and latitude or longitude.

## 5 | CONCLUSION

Meta-analyses, such as this, allow the examination and testing of general relationships that enable us to ask ecological questions on a larger scale than usually possible at a single study level (Kaiser et al., 2006). They do however come with a set of caveats that must be highlighted. Firstly, they are subject to publication bias (Kaiser et al., 2006), and importantly for this study, accuracy of species distribution data is a concern (Kryštufek & Griffiths, 2002; Taberlet et al., 1998). Not only do substantial differences in mapping efforts occur between countries, but also ranges such as those extracted from the

IUCN (used here) necessarily ignore the fact that realized ranges are mosaics rather than continuous. However, at the continental scale of this analysis and in the absence of fitting to environmental variables, these effects should be minimal.

The results presented here, along with the recognition of the implications of recurrent periods of glaciation (Fink et al., 2004; Stewart et al., 2010), the primarily individualistic nature of retreat (Graham & Grimm, 1990; Huntley, 1991; Stewart 2008, 2009), and evidence of refugia within refugia (Centeno-Cuadros et al., 2009; Colangelo et al., 2012), have highlighted how the acceptance of broad patterns can conceal a deeper complexity. Despite our use of a much tighter taxonomic grouping than Taberlet et al.'s (1998) use of 10 taxa from diverse clades across mammals, amphibians, arthropods and plants, our findings reflect theirs; "congruence on a continental scale may be the exception rather than the rule", and the southern refugia highlighted by many (Hewitt, 1996, 1999, 2000; Seddon et al., 2001; Taberlet et al., 1998) may better represent a "hotspot" of endemism rather than the sole recolonization sources for Europe (Bilton et al., 1998; Kryštufek & Griffiths, 2002; Stewart et al., 2010; Tougaard et al., 2008). It seems that perhaps unsurprisingly, distribution was then, and is now, primarily dictated by species' biological traits (tolerance ranges, adaptive capacity, plasticity) in combination with competition rather than simply regional restriction, and that northerly regions were neither as hostile nor barren as previously assumed. Further investigations using a nestedness approach (e.g., Habel et al., 2013) may provide further insight into these patterns, and help to identify the type of movement that led to the recolonization of these species.

## ACKNOWLEDGMENTS

This study was fully funded through grant assistance generously provided by the Department of Life and Environmental Science at Bournemouth University and Richard Stillman in particular is thanked. The authors are grateful to all the original authors that deposited their data in GenBank, and those that answered the requests, and provided additional information where available (S. Alasaad, J. R. Michaux, V. Fedorov, F. Suchentrunk, R.J. Baker, C.W. Matson, A. Borkowska, S. Simpson, O.A. Ermakov, P. Wandeler, V.A. Matrosova, D. Ehrich and J.E. Stacy). Without those studies, syntheses like these would not be possible.

## CONFLICT OF INTEREST

The authors confirm that no conflict of interest exists.

## DATA ACCESSIBILITY

As detailed in the Methods, all data are sourced from published sources freely available online. Sequences were obtained through GenBank (see Supporting Information Appendix S1 for the source papers and GenBank accession numbers). Area occupied by each species and primary habitat/vegetation preferences were obtained from IUCN species profiles ([www.iucnredlist.org](http://www.iucnredlist.org)).

## ORCID

Debbi Pedreschi  <http://orcid.org/0000-0001-6881-1051>

## REFERENCES

- Avise, J. C. (1995). Mitochondrial DNA polymorphism and a connection between genetics and demography of relevance to conservation. *Conservation Biology*, 9, 686–690. <https://doi.org/10.1046/j.1523-1739.1995.09030686.x>
- Avise, J. C. (2000). *Phylogeography: The history and formation of species*. Cambridge, Massachusetts and London: Harvard University Press.
- Avise, J. C., Arnold, J., Ball, R. M., Bermingham, E., Lamb, T., Neigel, J. E., & Saunders, N. C. (1987). Intraspecific phylogeography: The mitochondrial DNA bridge between population genetics and systematics. *Annual Review of Ecology and Systematics*, 18, 489–522. <https://doi.org/10.1146/annurev.es.18.110187.002421>
- Banaszek, A., Jadwyszczak, K. A., Ratkiewicz, M., & Ziomek, J. (2009). Low genetic diversity and significant structuring of the common hamster populations *Cricetus cricetus* in Poland revealed by the mtDNA control region sequence variation. *Acta Theriologica*, 54, 289–295. <https://doi.org/10.4098/j.at.0001-7051.021.2009>
- Banaszek, A., Jadwyszczak, K. A., Ratkiewicz, M., Ziomek, J., & Neumann, K. (2010). Population structure, colonization processes and barriers for dispersal in Polish common hamsters (*Cricetus cricetus*). *Journal of Zoological Systematics and Evolutionary Research*, 48, 151–158.
- Banaszek, A., & Ziomek, J. (2011). The common hamster, *Cricetus cricetus* (L.) populations in the Lower San River Valley. *Zoologica Poloniae*, 56, 49–58. <https://doi.org/10.2478/v10049-011-0007-5>
- Beysard, M., & Heckel, G. (2014). Structure and dynamics of hybrid zones at different stages of speciation in the common vole (*Microtus arvalis*). *Molecular Ecology*, 23, 673–687.
- Bhagwat, S. A., & Willis, K. J. (2008). Species persistence in northerly glacial refugia of Europe: A matter of chance or biogeographical traits? *Journal of Biogeography*, 35, 464–482. <https://doi.org/10.1111/j.1365-2699.2007.01861.x>
- Bilton, D. T., Mirol, P. M., Mascheretti, S., Fredga, K., Zima, J., & Searle, J. B. (1998). Mediterranean Europe as an area of endemism for small mammals rather than a source for northwards postglacial colonization. *Proceedings of the Royal Society of London B: Biological Sciences*, 265, 1219–1226. <https://doi.org/10.1098/rspb.1998.0423>
- Bolfíková, B., & Hulva, P. (2012). Microevolution of sympatry: Landscape genetics of hedgehogs *Erinaceus europaeus* and *E. roumanicus* in Central Europe. *Heredity*, 108, 248–255. <https://doi.org/10.1038/hdy.2011.67>
- Braaker, S., & Heckel, G. (2009). Transalpine colonisation and partial phylogeographic erosion by dispersal in the common vole (*Microtus arvalis*). *Molecular Ecology*, 18, 2518–2531.
- Brace, S., Palkopoulou, E., Dalén, L., Lister, A. M., Miller, R., Otte, M., ... Barnes, I. (2012). Serial population extinctions in a small mammal indicate Late Pleistocene ecosystem instability. *Proceedings of the National Academy of Sciences USA*, 109, 20532–20536. <https://doi.org/10.1073/pnas.1213322109>
- Brace, S., Ruddy, M., Miller, R., Schreve, D. C., Stewart, J. R., & Barnes, I. (2016). The colonization history of British water vole (*Arvicola amphibius* (Linnaeus, 1758)): Origins and development of the Celtic fringe. *Proceedings of the Royal Society of London B: Biological Sciences*, 283, 20160130.
- Brown, S. D., Collins, R. A., Boyer, S., Lefort, M. C., Malumbres-Olarte, J., Vink, C. J., & Cruickshank, R. H. (2012). Spider: An R package for the analysis of species identity and evolution, with particular reference to DNA barcoding. *Molecular Ecology Resources*, 12, 562–565. <https://doi.org/10.1111/j.1755-0998.2011.03108.x>
- Buzan, E. V., Forster, D. W., Searle, J. B., & Krystufek, B. (2010). A new cytochrome b phylogroup of the common vole (*Microtus arvalis*) endemic to the Balkans and its implications for the evolutionary history of the species. *Biological Journal of the Linnean Society*, 100, 788–796. <https://doi.org/10.1111/j.1095-8312.2010.01451.x>
- Centeno-Cuadros, A., Delibes, M., & Godoy, J. A. (2009). Phylogeography of Southern Water Vole (*Arvicola sapidus*): Evidence for refugia within the Iberian glacial refugium? *Molecular Ecology*, 18, 3652–3667.
- Centeno-Cuadros, A., & Godoy, J. A. (2010). Structure, organization and nucleotide diversity of the mitochondrial control region and cytochrome b of southern water vole (*Arvicola sapidus*). *Mitochondrial DNA*, 21, 48–53.
- Colangelo, P., Aloise, G., Franchini, P., Annesi, F., & Amori, G. (2012). Mitochondrial DNA reveals hidden diversity and an ancestral lineage of the bank vole in the Italian peninsula. *Journal of Zoology*, 287, 41–52. <https://doi.org/10.1111/j.1469-7998.2011.00884.x>
- Dalén, L., Nyström, V., Valdiosera, C., Germonpré, M., Sablin, M., & Turner, E. ... Götherström, A. (2007). Ancient DNA reveals lack of postglacial habitat tracking in the arctic fox. *Proceedings of the National Academy of Sciences*, 104, 6726–6729. <https://doi.org/10.1073/pnas.0701341104>
- Deffontaine, V., Ledevin, R., Fontaine, M. C., Quéré, J. P., Renaud, S., Libois, R., & Michaux, J. R. (2009). A relict bank vole lineage highlights the biogeographic history of the Pyrenean region in Europe. *Molecular Ecology*, 18, 2489–2502. <https://doi.org/10.1111/j.1365-294X.2009.04162.x>
- Deffontaine, V., Libois, R., Kotlík, P., Sommer, R., Nieberding, C., Paradis, E., ... Michaux, J. R. (2005). Beyond the Mediterranean peninsulas: evidence of central European glacial refugia for a temperate forest mammal species, the bank vole (*Clethrionomys glareolus*). *Molecular Ecology*, 14, 1727–1739. <https://doi.org/10.1111/j.1365-294X.2005.02506.x>
- Dekonenko, A., Yakimenko, V., Ivanov, A., Morozov, V., Nikitin, P., Khasanova, S., ... Schmaljohnn, C. (2003). Genetic similarity of Puumala viruses found in Finland and western Siberia and of the mitochondrial DNA of their rodent hosts suggests a common evolutionary origin. *Infection, Genetics and Evolution*, 3, 245–257. [https://doi.org/10.1016/S1567-1348\(03\)00088-1](https://doi.org/10.1016/S1567-1348(03)00088-1)
- Dixon, C. J. (2006). A means of estimating the completeness of haplotype sampling using the Stirling probability distribution. *Molecular Ecology Notes*, 6, 650–652. <https://doi.org/10.1111/j.1471-8286.2006.01411.x>
- Egeland, T., & Salas, A. (2008). Estimating haplotype frequency and coverage of databases. *PLoS ONE*, 3, e3988. <https://doi.org/10.1371/journal.pone.0003988>
- Filipi, K., Marková, S., Searle, J. B., & Kotlík, P. (2015). Mitogenomic phylogenetics of the bank vole *Clethrionomys glareolus*, a model system for studying end-glacial colonization of Europe. *Molecular Phylogenetics and Evolution*, 82, 245–257. <https://doi.org/10.1016/j.ympev.2014.10.016>
- Fink, S., Excoffier, L., & Heckel, G. (2004). Mitochondrial gene diversity in the common vole *Microtus arvalis* shaped by historical divergence and local adaptations. *Molecular Ecology*, 13, 3501–3514. <https://doi.org/10.1111/j.1365-294X.2004.02351.x>
- Fløjgaard, C., Normand, S., Skov, F., & Svenning, J. C. (2011). Deconstructing the mammal species richness pattern in Europe – Towards an understanding of the relative importance of climate, biogeographic history, habitat heterogeneity and humans. *Global Ecology and Biogeography*, 20, 218–230. <https://doi.org/10.1111/j.1466-8238.2010.00604.x>
- Goodall-Copestake, W. P., Tarling, G. A., & Murphy, E. J. (2012). On the comparison of population-level estimates of haplotype and nucleotide diversity: A case study using the gene *cox1* in animals. *Heredity*, 109, 50–56. <https://doi.org/10.1038/hdy.2012.12>
- Graham, R. W., & Grimm, E. C. (1990). Effects of global climate change on the patterns of terrestrial biological communities. *Trends in Ecology and Evolution*, 5, 289–292. [https://doi.org/10.1016/0169-5347\(90\)90083-P](https://doi.org/10.1016/0169-5347(90)90083-P)

- Habel, J. C., Ulrich, W., & Assmann, T. (2013). Allele elimination recalculated: Nested subset analyses for molecular biogeographical data. *Journal of Biogeography*, 40, 769–777. <https://doi.org/10.1111/jbi.12054>
- Hampe, A., & Petit, R. J. (2005). Conserving biodiversity under climate change: The rear edge matters. *Ecology Letters*, 8, 461–467. <https://doi.org/10.1111/j.1461-0248.2005.00739.x>
- Heckel, G., Burri, R., Fink, S., Desmet, J. F., & Excoffier, L. (2005). Genetic structure and colonization processes in European populations of the common vole, *Microtus arvalis*. *Evolution*, 59, 2231–2242. <https://doi.org/10.1111/j.0014-3820.2005.tb00931.x>
- Hewitt, G. M. (1996). Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnean Society*, 58, 247–276. <https://doi.org/10.1111/j.1095-8312.1996.tb01434.x>
- Hewitt, G. M. (1999). Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society*, 68, 87–112. <https://doi.org/10.1111/j.1095-8312.1999.tb01160.x>
- Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, 405, 907–913. <https://doi.org/10.1038/35016000>
- Hewitt, G. M. (2001). Speciation, hybrid zones and phylogeography—Or seeing genes in space and time. *Molecular Ecology*, 10, 537–549. <https://doi.org/10.1046/j.1365-294x.2001.01202.x>
- Hewitt, G. M. (2004). The structure of biodiversity—insights from molecular phylogeography. *Frontiers in Zoology*, 1, 4–16. <https://doi.org/10.1186/1742-9994-1-4>
- Hofreiter, M., & Stewart, J. (2009). Ecological change, range fluctuations and population dynamics during the Pleistocene. *Current Biology*, 19, R584–R594. <https://doi.org/10.1016/j.cub.2009.06.030>
- Huntley, B. (1991). How plants respond to climate change: Migration rates, individualism and the consequences for plant communities. *Annals of Botany*, 67, 15–22. <https://doi.org/10.1093/oxfordjournals.aob.a088205>
- IBM Corp. (2011). *IBM SPSS Statistics for Windows, Version 20.0*. Armonk, NY: Author.
- Ibrahim, K. M., Nichols, R. A., & Hewitt, G. M. (1996). Spatial patterns of genetic variation generated by different forms of dispersal. *Heredity*, 77, 282–291.
- Johansson, P., Olsson, G. E., Low, H. T., Bucht, G., Ahlm, C., Juto, P., & Elgh, F. (2008). Puumala hantavirus genetic variability in an endemic region (Northern Sweden). *Infection, Genetics and Evolution*, 8, 286–296. <https://doi.org/10.1016/j.meegid.2008.01.003>
- Kaiser, M. J., Clarke, K. R., Hinz, H., Austen, M. C. V., Somerfield, P. J., & Karakassis, I. (2006). Global analysis of response and recovery of benthic biota to fishing. *Marine Ecology Progress Series*, 311, 1–14. <https://doi.org/10.3354/meps311001>
- Kotlík, P., Deffontaine, V., Mascheretti, S., Zima, J., Michaux, J. R., & Searle, J. B. (2006). A northern glacial refugium for bank voles (*Clethrionomys glareolus*). *Proceedings of the National Academy of Sciences USA*, 103, 14860–14864. <https://doi.org/10.1073/pnas.0603237103>
- Kryštufek, B., & Griffiths, H. I. (2002). Species richness and rarity in European rodents. *Ecography*, 25, 120–128. <https://doi.org/10.1034/j.1600-0587.2002.250114.x>
- Lagerholm, V. K., Sandoval-Castellanos, E., Ehrlich, D., Abramson, N. I., Nadachowski, A., Kalthoff, D. C., ... Dalén, L. (2014). On the origin of the Norwegian lemming. *Molecular Ecology*, 23, 2060–2071. <https://doi.org/10.1111/mec.12698>
- Librado, P., & Rozas, J. (2009). DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25, 1451–1452. <https://doi.org/10.1093/bioinformatics/btp187>
- Lischer, H. E., Excoffier, L., & Heckel, G. (2013). Ignoring heterozygous sites biases phylogenomic estimates of divergence times: Implications for the evolutionary history of *Microtus* voles. *Molecular Biology and Evolution*, 31, 817–831. <https://doi.org/10.1093/molbev/mst271>
- Martin, A. P., & Palumbi, S. R. (1993). Body size, metabolic rate, generation time, and the molecular clock. *Proceedings of the National Academy of Sciences USA*, 90, 4087–4091. <https://doi.org/10.1073/pnas.90.9.4087>
- Martínková, N., Barnett, R., Cucchi, T., Struchen, R., Pascal, M., Pascal, M., ... Searle, J. B. (2013). Divergent evolutionary processes associated with colonization of offshore islands. *Molecular Ecology*, 22, 5205–5220. <https://doi.org/10.1111/mec.12462>
- Matson, C. W., Rodgers, B. E., Chesser, R. K., & Baker, R. J. (2000). Genetic diversity of *Clethrionomys glareolus* populations from highly contaminated sites in the Chernobyl region, Ukraine. *Environmental Toxicology and Chemistry*, 19, 2130–2135. [https://doi.org/10.1897/1551-5028\(2000\)019aabb2130:GDOCGPaabb2.3.CO;2](https://doi.org/10.1897/1551-5028(2000)019aabb2130:GDOCGPaabb2.3.CO;2)
- Meeks, H. N., Chesser, R. K., Rodgers, B. E., Gaschak, S., & Baker, R. J. (2009). Understanding the genetic consequences of environmental toxicant exposure: Chernobyl as a model system. *Environmental Toxicology and Chemistry*, 28, 1982–1994. <https://doi.org/10.1897/08-578.1>
- Meeks, H. N., Wickliffe, J. K., Hooper, S. R., Chesser, R. K., Rodgers, B. E., & Baker, R. J. (2007). Mitochondrial control region variation in bank voles (*Clethrionomys glareolus*) is not related to Chernobyl radiation exposure. *Environmental Toxicology and Chemistry*, 26, 361–369. <https://doi.org/10.1897/06-346R.1>
- Mitchell-Jones, A. J., Amori, G., Bogdanowicz, W., Krystufek, B., Reijnders, P. J. H., Spitzenberger, F., ... Zima, J. (1999). *The atlas of European mammals* (Vol. 3). London: Academic Press. ISBN-13: 978-0856611308
- Moritz, C. (1994). Applications of mitochondrial DNA analysis in conservation: a critical review. *Molecular Ecology*, 3, 401–411. <https://doi.org/10.1111/j.1365-294X.1994.tb00080.x>
- Neumann, K., Jansman, H., Kayser, A., Maak, S., & Gattermann, R. (2004). Multiple bottlenecks in threatened western European populations of the common hamster *Cricetus cricetus* (L.). *Conservation Genetics*, 5, 181–193. <https://doi.org/10.1023/B:COGE.0000030002.01948.b3>
- Neumann, K., Michaux, J. R., Maak, S., Jansman, H. A. H., Kayser, A., Mundt, G., & Gattermann, R. (2005). Genetic spatial structure of European common hamsters (*Cricetus cricetus*)—A result of repeated range expansion and demographic bottlenecks. *Molecular Ecology*, 14, 1473–1483. <https://doi.org/10.1111/j.1365-294X.2005.02519.x>
- Palkopoulou, E., Baca, M., Abramson, N. I., Sablin, M., Socha, P., Nadachowski, A., ... Dalén, L. (2016). Synchronous genetic turnovers across Western Eurasia in Late Pleistocene collared lemmings. *Global Change Biology*, 22, 1710–1721. <https://doi.org/10.1111/gcb.13214>
- Pereira, L., Cunha, C., & Amorim, A. (2004). Predicting sampling saturation of mtDNA haplotypes: An application to an enlarged Portuguese database. *International Journal of Legal Medicine*, 118, 132–136. <https://doi.org/10.1007/s00414-003-0424-1>
- R Core Team. (2017). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>
- Razzauti, M., Plyusnina, A., Niemimaa, J., Henttonen, H., & Plyusnin, A. (2012). Co-circulation of two Puumala hantavirus lineages in Latvia: A Russian lineage described previously and a novel Latvian lineage. *Journal of Medical Virology*, 84, 314–318. <https://doi.org/10.1002/jmv.22263>
- Schmitt, T., & Varga, Z. (2012). Extra-Mediterranean refugia: The rule and not the exception? *Frontiers in Zoology*, 9, 22. <https://doi.org/10.1186/1742-9994-9-22>
- Searle, J. B., Kotlík, P., Rambau, R. V., Marková, S., Herman, J. S., & McDevitt, A. D. (2009). The Celtic fringe of Britain: Insights from small mammal phylogeography. *Proceedings of the Royal Society of London B: Biological Sciences*, 276, 4287–4294.
- Seddon, J. M., Santucci, F., Reeve, N. J., & Hewitt, G. M. (2001). DNA footprints of European hedgehogs, *Erinaceus europaeus* and *E.*

- concolor: Pleistocene refugia, postglacial expansion and colonization routes. *Molecular Ecology*, 10, 2187–2198. <https://doi.org/10.1046/j.0962-1083.2001.01357.x>
- Stewart, J. R. (2008). The progressive effect of the individualistic response of species to Quaternary climate change: an analysis of British mammalian faunas. *Quaternary Science Reviews*, 27, 2499–2508. <https://doi.org/10.1016/j.quascirev.2008.08.003>
- Stewart, J. R. (2009). The evolutionary consequence of the individualistic response to climate change. *Journal of Evolutionary Biology*, 22, 2363–2375. <https://doi.org/10.1111/j.1420-9101.2009.01859.x>
- Stewart, J. R., & Dalén, L. (2008). Is the glacial refugium concept relevant for northern species? A comment on Pruett and Winker 2005. *Climatic Change*, 86, 19–22. <https://doi.org/10.1007/s10584-007-9366-9>
- Stewart, J. R., & Lister, A. M. (2001). Cryptic northern refugia and the origins of the modern biota. *Trends in Ecology and Evolution*, 16, 608–613. [https://doi.org/10.1016/S0169-5347\(01\)02338-2](https://doi.org/10.1016/S0169-5347(01)02338-2)
- Stewart, J. R., Lister, A. M., Barnes, I., & Dalén, L. (2010). Refugia revisited: Individualistic responses of species in space and time. *Proceedings of the Royal Society of London B: Biological Sciences*, 277, 661–671.
- Stojak, J., McDevitt, A. D., Herman, J. S., Kryštufek, B., Uhlíková, J., Purger, J. J., ... Wójcik, J. M. (2016). Between the Balkans and the Baltic: Phylogeography of a common vole mitochondrial DNA lineage limited to Central Europe. *PLoS ONE*, 11, e0168621. <https://doi.org/10.1371/journal.pone.0168621>
- Stojak, J., McDevitt, A. D., Herman, J. S., Searle, J. B., & Wójcik, J. M. (2015). Post-glacial colonization of eastern Europe from the Carpathian refugium: Evidence from mitochondrial DNA of the common vole *Microtus arvalis*. *Biological Journal of the Linnean Society*, 115, 927–939.
- Sutter, A., Beysard, M., & Heckel, G. (2013). Sex-specific clines support incipient speciation in a common European mammal. *Heredity*, 110, 398–404. <https://doi.org/10.1038/hdy.2012.124>
- Taberlet, P., Fumagalli, L., Wust-Saucy, A. G., & Cosson, J. F. (1998). Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology*, 7, 453–464. <https://doi.org/10.1046/j.1365-294x.1998.00289.x>
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. (2013). MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*, 30, 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Tougaard, C., & Renvoisé, E. (2008). Rodents and palaeogenetics: New perspectives. *Comptes Rendus Palevol*, 7, 125–134. <https://doi.org/10.1016/j.crpv.2007.12.002>
- Tougaard, C., Renvoisé, E., Petitjean, A., & Quéré, J. P. (2008). New insight into the colonization processes of common voles: Inferences from molecular and fossil evidence. *PLoS ONE*, 3, e3532. <https://doi.org/10.1371/journal.pone.0003532>
- Tzedakis, P. C., Emerson, B. C., & Hewitt, G. M. (2013). Cryptic or mystic? Glacial tree refugia in northern Europe. *Trends in Ecology and Evolution*, 28, 696–704. <https://doi.org/10.1016/j.tree.2013.09.001>
- Waters, J. M., Fraser, C. I., & Hewitt, G. M. (2013). Founder takes all: Density-dependent processes structure biodiversity. *Trends in Ecology and Evolution*, 28, 78–85. <https://doi.org/10.1016/j.tree.2012.08.024>
- White, T. A., Perkins, S. E., Heckel, G., & Searle, J. B. (2013). Adaptive evolution during an ongoing range expansion: The invasive bank vole (*Myodes glareolus*) in Ireland. *Molecular Ecology*, 22, 2971–2985.
- Wickliffe, J. K., Dunina-Barkovskaya, Y. V., Gaschak, S. P., Rodgers, B. E., Chesser, R. K., Bondarkov, M., & Baker, R. J. (2006). Variation in mitochondrial DNA control region haplotypes in populations of the bank vole, *Clethrionomys glareolus*, living in the Chernobyl environment, Ukraine. *Environmental Toxicology and Chemistry*, 25(2), 503–508. <https://doi.org/10.1897/05-327R.1>

## BIOSKETCH

**DEBBI PEDRESCHI** comes from a varied background employing multidisciplinary methods to inform management practices, including through improving phylogeographical understanding. Her interests centre around understanding anthropogenic pressures, ecosystem management, and investigating diversity at a range of levels. The team's research has focused on the examination of genetic variation in a range of species to reconstruct post-glacial geographical range change and contribute to understanding the role of ice-age refugia.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Pedreschi D, García-Rodríguez O, Yannic G, et al. Challenging the European southern refugium hypothesis: Species-specific structures versus general patterns of genetic diversity and differentiation among small mammals. *Global Ecol Biogeogr*. 2019;28:262–274. <https://doi.org/10.1111/geb.12828>