

Self-assembling proteins for bio-electronics

Julien Hurtaud, Joffrey Champavert, Suhail Usmani, Olivier Constantin, Cécile Delacour, Patrice Rannou, Vincent Forge

▶ To cite this version:

Julien Hurtaud, Joffrey Champavert, Suhail Usmani, Olivier Constantin, Cécile Delacour, et al.. Selfassembling proteins for bio-electronics. MRS Fall conference - Symposium SB12-Conductive Biological Materials, Dec 2024, Boston MA, United States. hal-04829073

HAL Id: hal-04829073 https://hal.science/hal-04829073v1

Submitted on 10 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

Symposium SB12-Conductive Biological Materials

Self-Assembling Proteins for Bio-Electronics

Julien Hurtaud¹, Joffrey Champavert¹, Suhail Usmani², Olivier Constantin², Cécile Delacour³, Patrice Rannou⁴ and Vincent Forge¹

1. Univ. Grenoble Alpes, CNRS, CEA, IRIG, Laboratoire de Chimie et Biologie des Métaux, F-38000, Grenoble, France

2. Univ. Grenoble Alpes, CEA, LETI, Département des Technologies pour l'Innovation en Santé, F-38000, Grenoble, France

3. Institut Néel, Univ. Grenoble Alpes, CNRS, Grenoble INP, 38000 Grenoble, France

4. Univ. Grenoble Alpes, Univ. Savoie Mont-Blanc, CNRS, Grenoble INP, LEPMI, 38000 Grenoble, France

Bioelectronics is a fast-growing field of research aimed at controlling the interface between biological systems and electronic devices, which requires the conversion of biological signals into electrical signals usable by conventional electronic devices. Here we demonstrate the potential of protein nanowires resulting from the self-assembly of the prion domain of a filamentous fungus protein. These nanowires are in fact functional, non-pathological amyloid fibres; they perform a biological function, namely recognition of the self and non-self of a filamentous fungus. With a diameter of 5 nm and an average length of 12 μ m, these nanowires, whose structure is known on an atomic scale, have a high aspect ratio (1). What's more, hydrogels can be obtained in an easily controllable way (2,3). Like all amyloid fibres, these nanowires have specific optical properties (4), as well as very good intrinsic ionic/protonic conductivity (2). Finally, the addition of a redox protein domain (2) or a redox chemical compound (3) on the prion domain gives them electronic conductivity. After self-assembly of the prion domains, these redox domains are aligned along the nanowire axes, enabling electronic conductivity by electron hopping.

Measurements of ionic and electronic conductivities will be described from the macroscopic scale (film) to the nanoscopic scale (single nanowire). Applications that take advantage of the various properties of these nanowires will be illustrated. For the first application, the ability to form a hydrogel and the ionic conductivity of nanowires make it possible to produce a biocompatible electrode coating for the primary culture of neurons over very long periods and to detect neuronal electrical activity. For a second application, in the field of sensors, electronically conductive nanowires, which can transport electrons over long distances, can be used as redox mediators between enzymes and glassy carbon electrodes (2,3,5).

These results represent a first step towards innovative all-in-one materials for bioelectronics. Overall, we have observed that our nanowires formed by the self-assembly of a functional, non-pathological prion domain combine all the properties needed to bridge the gap between the world of biology and that of technology, particularly electronics.

(1) Doussineau et al. Mass Determination of Entire Amyloid Fibrils by Using Mass Spectrometry. *Angew Chem Int Ed Engl.* 55: 2340-2344 (2015)

(2) Altamura et al. A synthetic redox biofilm made from metalloprotein-prion domain chimera nanowires. *NATURE CHEMISTRY* 9: 157-163 (2017)

(3) Duraffourg et al. Hybrid Amyloid-Based Redox Hydrogel for Bioelectrocatalytic H2 Oxidation. *Angew Chem Int Ed Engl. 60*: 14488-14497 (2021)

(4) Pansieri et al. Ultraviolet-visible-near-infrared optical properties of amyloid fibrils shed light on amyloidogenesis. *NATURE PHOTONICS* 13: 473-479 (2019)

(5) Rengaraj et al. Interprotein Electron Transfer between FeS-Protein Nanowires and Oxygen-Tolerant NiFe Hydrogenase. *Angew Chem Int Ed Engl.* **56**: 7774-7778 (2017)