

A finite volume scheme accurate at low Mach number on quadrangular mesh by space velocity enrichment

Jonathan Jung, Vincent Perrier

▶ To cite this version:

Jonathan Jung, Vincent Perrier. A finite volume scheme accurate at low Mach number on quadrangular mesh by space velocity enrichment. ECCOMAS 2024 - 9th European Congress on Computational Methods in Applied Sciences and Engineering, Jun 2024, Lisboa, Portugal. hal-04836529

HAL Id: hal-04836529 https://hal.science/hal-04836529v1

Submitted on 13 Dec 2024 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

A finite volume scheme accurate at low Mach number on quadrangular mesh by space velocity enrichment

J. Jung¹, V. Perrier²

 ¹ LMA, E2S-UPPA, and Cagire team, Inria Bordeaux Sud-Ouest, Avenue de l'Université, 64013 Pau Cedex, France.
² Cagire team, Inria Bordeaux Sud-Ouest and LMA, E2S-UPPA, Avenue de l'Université, 64013 Pau Cedex, France.

Keywords: Low Mach number flows, Finite volume scheme, Compressible solver, Finite element method, fluid mechanics

Classical finite volume schemes for compressible Euler system are not accurate on quadrangular mesh at low Mach number in the sense that they do not converge to the incompressible limit when the Mach number tends to zero [1]. The spurious mode that jeopardizes the convergence can be identified and corresponds to the long time limit of a first order wave system whose properties and discretization depend on the scheme used for the compressible system [2]. Then, the low Mach number accuracy can be analysed by studying the long time solution of the associated wave system.

In this presentation, we will propose to enrich the velocity space approximation by adding a divergence free element in each cell. Using this new approximation space and a Godunov' numerical flux, the long-time limit of the wave system discretization corresponds to the divergence free component of the initial condition (as expected). This divergence free component will be defined via a discrete Hodge-Helmoltz decomposition.

As a consequence, for Euler equations, using this new approximation space and a numerical flux that degenerates in the low Mach number limit to a Godunov' flux for the wave system, the numerical scheme provides pressure fluctuations of order Mach square and a divergence of the velocity field of order Mach in the low Mach number limit.

This work can be seen as an extension to quadrangles of the properties obtained on triangles [3].

REFERENCES

- [1] H. Guillard and C. Viozat, On the behaviour of upwind schemes in the low Mach number limit. Computers & fluids, 28(1), pp. 63-86, 1999.
- [2] J. Jung and V. Perrier, Steady low Mach number flows: identification of the spurious mode and filtering method. Journal of Computational Physics, 468, 111462, 2022.
- [3] H. Guillard, On the behavior of upwind schemes in the low Mach number limit. IV: P0 approximation on triangular and tetrahedral cells. Computers & fluids, 38(10), pp. 1969-1972, 2009.