



HAL
open science

Variants in UBAP1L lead to autosomal recessive rod-cone and cone-rod dystrophy

Christina Zeitz, Julien Navarro, Leila Azizzadeh Pormehr, Cécile Méjécasse, Luiza M. Neves, Camille Letellier, Christel Condroyer, Shahad Albadri, Andréa Amprou, Aline Antonio, et al.

► **To cite this version:**

Christina Zeitz, Julien Navarro, Leila Azizzadeh Pormehr, Cécile Méjécasse, Luiza M. Neves, et al.. Variants in UBAP1L lead to autosomal recessive rod-cone and cone-rod dystrophy. *Genetics in Medicine*, 2024, 26 (6), pp.101081. 10.1016/j.gim.2024.101081 . hal-04843771

HAL Id: hal-04843771

<https://hal.science/hal-04843771v1>

Submitted on 18 Dec 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genetics in Medicine

Variants in UBAP1L lead to autosomal recessive rod-cone and cone-rod dystrophy --Manuscript Draft--

Manuscript Number:	GENETMED-D-23-00572R4
Full Title:	Variants in UBAP1L lead to autosomal recessive rod-cone and cone-rod dystrophy
Article Type:	Article
Section/Category:	Clinical Genetics and Genomics
Corresponding Author:	Christina Zeitz, PhD Sorbonne University Paris, Ile de France FRANCE
Corresponding Author's Institution:	Sorbonne University
Order of Authors:	Christina Zeitz, PhD Julien Navarro Leila Azizzadeh Pormehr Cécile Méjécasse Luiza M Neves Camille Letellier Christel Condroyer Shahad Albadri Andréa Amprou Aline Antonio Tasnim Ben-Yacoub Juliette Wohlschlegel Camille Andrieu Malo Serafini Lorenzo Bianco Alessio Antropoli Marco Nassisi Said El Shamieh Sandra Chantot-Bastaraud Saddek Mohand-Saïd Vasily Smirnov José-Alain Sahel Filippo Del Bene Isabelle Audo
Manuscript Region of Origin:	FRANCE
Abstract:	Abstract (200 words) Purpose Progressive inherited retinal degenerations (IRDs) affecting rods and cones are clinically and genetically heterogeneous and can lead to blindness with limited therapeutic options. The major gene defects have been identified in Caucasians and Asians with only few reports in North African cases.

Methods

Whole genome, targeted next-generation and Sanger sequencing was applied to cohort of ~4000 IRDs cases. Expression analyses were performed including Chip-seq database analyses, on human-derived retinal organoids (ROs), retinal pigment epithelium (RPE) cells and zebrafish. Variants' pathogenicity was assessed using 3D-modeling and/or ROs.

Results

Here we identified a novel gene defect with three distinct pathogenic variants in UBAP1L in four independent autosomal recessive IRD cases from Tunisia. UBAP1L is expressed in the RPE and retina, specifically in rods and cones, in line with the phenotype. It encodes Ubiquitin-associated protein 1-like, containing a solenoid of overlapping ubiquitin associated (SOUBA) domain, predicted to interact with ubiquitin. In silico and in vitro studies, including 3D-modeling and ROs revealed that the SOUBA domain is truncated and thus ubiquitin binding most likely abolished secondary to all variants identified herein.

Conclusion

Biallelic UBAP1L variants are a novel cause of IRDs, most likely enriched in the North African population.

**Unité Mixte Inserm –
Sorbonne Université Centre
de Recherche 968**



Paris, Janvier 15th, 2024

Dear Editors,

Dr Serge Picaud

Directeur

Pr Isabelle Audo

Directrice adjointe

Manuscript Number: GENETMED-D-23-00572

Dear Dr. Steiner, dear editors,

We are grateful for your reply accepting our article and asking for minor revision of our manuscript “Variants in *UBAPIL* lead to autosomal recessive rod-cone and cone-rod dystrophy” Ref. No.: GENETMED-D-23-00572. Enclosed we return the revised version. Thanks for your patience!

Christina Zeitz and Isabelle Audo

Dr Christina Zeitz

Pr Isabelle Audo

17 rue Moreau

F-75012 PARIS

Tel. : (33) 01 53 46 25 40/42

christina.zeitz@inserm.fr

isabelle.audo@inserm.fr

January 15th, 2024

RE: GENETMED-D-23-00572R, entitled "Variants in UBAP1L lead to autosomal recessive rod-cone and cone-rod dystrophy"

Dear Dr. Steiner, dear editors, dear reviewers,

We are grateful for your reply accepting our manuscript “Variants in *UBAP1L* lead to autosomal recessive rod-cone and cone-rod dystrophy” Ref. No.: GENETMED-D-23-00572. Thanks for your patience!

Sincerely,
Christina Zeitz and Isabelle Audo

Conflict of interest

The authors have no conflict of interest in this study.

1
2
3
4 **Variants in *UBAPIL* lead to autosomal recessive rod-cone and cone-rod dystrophy**

5
6 Christina Zeitz^{1*}, Julien Navarro^{1#}, Leila Azizzadeh Pormehr^{1,2#}, Cécile Méjécase^{1,3#}, Luiza M.
7
8 Neves^{1,4#}, Camille Letellier^{1#}, Christel Condroyer¹, Shahad Albadri¹, Andréa Amprou¹, Aline
9
10 Antonio¹, Tasnim Ben-Yacoub¹, Juliette Wohlschlegel^{1,5} Camille Andrieu⁶, Malo Serafini¹,
11
12 Lorenzo Bianco^{1,7}, Alessio Antropoli^{1,7}, Marco Nassisi^{1,8}, Said El Shamieh^{1,9}, Sandra Chantot-
13
14 Bastaraud¹⁰, Saddek Mohand-Saïd^{1,6}, Vasily Smirnov^{1,11}, José-Alain Sahel^{1,6,12}, Filippo Del Bene¹,
15
16 Isabelle Audo^{1,6*}
17
18
19
20
21

22 *these authors equally contributed to the article and are both corresponding authors. Mail to:
23
24 christina.zeitz@inserm.fr, isabelle.audo@inserm.fr.
25
26

27
28 #these authors equally contributed to the article
29
30

31 ¹Sorbonne Université, INSERM, CNRS, Institut de la Vision, 75012 Paris, France.
32
33

34 ²Mass. Eye and Ear, Ocular Genomics Institute, Berman-Gund Laboratory for the Study of Retinal
35
36 Degenerations, Harvard Medical School, Boston, MA 02114, USA.
37
38

39 ³UCL Institute of Ophthalmology, London EC1V 9EL, UK; The Francis Crick Institute, London
40
41 NW1 1A, UK.
42
43

44 ⁴Instituto Nacional de Saúde da Mulher, da Criança e do Adolescente Fernandes Figueira,
45
46 Fundação Oswaldo Cruz, Rio de Janeiro 22250-020, Brazil.
47
48

49
50 ⁵Department of Biological Structure, University of Washington, Seattle, WA 98195, USA.
51
52

53 ⁶Centre Hospitalier National d'Ophthalmologie des Quinze-Vingts, Centre de Référence Maladies
54
55 Rares REFERET and INSERM-DGOS CIC 1423, 75012 Paris, France.
56
57

58
59 ⁷Department of Ophthalmology, IRCCS San Raffaele Scientific Institute, 20132 Milan, Italy.
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

⁸Department of Clinical Sciences and Community Health, University of Milan, 20122 Milan, Italy.

⁹Molecular Testing Laboratory, Department of Medical Laboratory Technology, Faculty of Health Sciences, Beirut Arab University, 115020 Beirut, Lebanon.

¹⁰APHP, Hôpital Armand-Trousseau, Département de Génétique, UF de Génétique Chromosomique, 75012 Paris, France.

¹¹Exploration de la Vision et Neuro-Ophthalmologie, CHU de Lille, 59037 Lille, France.

¹²Department of Ophthalmology, The University of Pittsburgh School of Medicine, Pittsburg, PA 15213, USA.

1
2
3
4 **Abstract (200 words)**
5
6

7 Purpose
8
9

10 Progressive inherited retinal degenerations (IRDs) affecting rods and cones are clinically and
11 genetically heterogeneous and can lead to blindness with limited therapeutic options. The major
12 gene defects have been identified in subjects of European and Asian decent with only few reports
13 of North African descent.
14
15
16
17
18
19
20

21 Methods
22
23

24 Genome, targeted next-generation and Sanger sequencing was applied to cohort of ~4000 IRDs
25 cases. Expression analyses were performed including Chip-seq database analyses, on human-
26 derived retinal organoids (ROs), retinal pigment epithelium (RPE) cells and zebrafish. Variants'
27 pathogenicity was accessed using 3D-modeling and/or ROs.
28
29
30
31
32
33
34

35 Results
36
37

38 Here we identified a novel gene defect with three distinct pathogenic variants in *UBAPIL* in four
39 independent autosomal recessive IRD cases from Tunisia. *UBAPIL* is expressed in the RPE and
40 retina, specifically in rods and cones, in line with the phenotype. It encodes Ubiquitin-associated
41 protein 1-like, containing a solenoid of overlapping ubiquitin associated (SOUBA) domain,
42 predicted to interact with ubiquitin. *In silico* and *in vitro* studies, including 3D-modeling and ROs
43 revealed that the SOUBA domain is truncated and thus ubiquitin binding most likely abolished
44 secondary to all variants identified herein.
45
46
47
48
49
50
51
52
53
54
55

56 Conclusion
57
58
59
60
61
62
63
64
65

1
2
3
4 Biallelic *UBAPIL* variants are a novel cause of IRDs, most likely enriched in the North African
5
6 population.
7
8
9

10 11 12 **Keywords**

13
14
15 *UBAPIL*, autosomal recessive rod-cone and cone-rod dystrophy, novel gene defect, genome
16
17 sequencing, retinal organoids, retinal pigment epithelium, zebrafish, 3D-modelling
18
19
20
21
22

23 24 **Introduction**

25
26
27 Inherited retinal degenerations (IRDs) are a clinically and genetically heterogeneous group of
28
29 stationary and progressive diseases, affecting vision ¹. Non-syndromic rod-cone dystrophy (RCD),
30
31 also called retinitis pigmentosa (RP), is the most common form of IRD with a prevalence of
32
33 1:4.000. RCD initially starts with night blindness, secondary to rod photoreceptor dysfunction
34
35 followed by peripheral visual field constriction and eventually loss of central vision, secondary to
36
37 cone photoreceptor degeneration ². Non-syndromic cone-rod dystrophy (CRD) or cone dystrophy
38
39 (CD) are less common, with a prevalence of 1:40.000, with primary cone dysfunction and central
40
41 vision loss ³, also leading to blindness in most severe cases ⁴. RCD and CRD are genetically
42
43 heterogeneous diseases and have been associated with variants in more than 80 genes ⁵
44
45 (<https://web.sph.uth.edu/RetNet/sum-dis.htm>). Despite recent advances in high-throughput
46
47 sequencing, the disease-causing variants are still missing for approximately 20%-40% affected
48
49 individuals with IRDs ⁵. These cases may involve variants in novel disease-associated genes ⁶,
50
51 variants in genes previously associated with a different phenotype ^{7,8}, overlooked intronic variants
52
53 ^{9,10}, structural variants ¹¹ and variants in regulatory regions ¹², which require large cohorts and/or
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 functional analyses for validation ¹³. Functional analyses can be based on *in vitro* or *in vivo*
5
6 modeling including expression and protein immunolocalization studies, mini-gene approaches and
7
8 by the characterization of animal models ¹³. More recently, genome sequencing combined with
9
10 investigations on human induced pluripotent stem cell (hiPSC)-lines differentiated into retinal
11
12 pigment epithelium (RPE) cells or retinal organoids (ROs) has proven valuable in identifying
13
14 additional gene defects underlying IRDs ^{11,14}.

15
16
17
18
19 The goal of this work was to identify the gene defect of a subject with autosomal recessive RCD,
20
21 investigate the candidate gene defect in a cohort of ~4.000 index cases with IRDs, validate its
22
23 relevance using 3D-modeling, and perform expression studies in zebrafish and in RPE cells and
24
25 ROs differentiated from human iPSC-lines.
26
27
28
29
30

31 **Material and Methods**

32 **Subjects**

33
34
35 A cohort of ~4.000 cases with IRDs were clinically investigated as previously described ¹⁵. These
36
37 individuals were recruited at the Reference Center for rare diseases, REFERET, Quinze-Vingts
38
39 hospital, Paris. The race and ethnicity of the individuals were self-reported and these French
40
41 subjects were mainly from European and North African ancestry. Written informed consent for
42
43 the genetic analysis was obtained from all individuals. All studies were carried out in accordance
44
45 with the declaration of Helsinki and were approved by a national ethics committee (CPP Ile de
46
47 France V, Project number 06693, N°EUDRACT 2006-A00347-44, 11 December 2006).
48
49
50
51
52
53
54

55 **Genetic Analyses**

56
57 Total genomic DNA was extracted from whole blood. The affected index individual, CIC03225
58
59 of F1362, from a consanguineous Tunisian couple underwent array comparative genomic
60
61
62
63
64
65

1
2
3
4 hybridization (CGH) ⁶, Sanger sequencing in candidate genes ^{16,17}, microarray analysis¹⁸ and
5
6 targeted next generation sequencing (NGS) as previously described ^{19,20}. Exome sequencing (ES)
7
8 was performed on the affected index individual CIC03225, the unaffected father CIC03248, the
9
10 unaffected mother CIC03226 and on one unaffected brother CICC03227 of F1362 (Figure 1A) as
11
12 described before ²¹. Genome sequencing (GS) was done on the affected index individual CIC03225
13
14 and all available family members (Figure 1A). Paired-end sequencing (2 * 150b) was performed
15
16 by a company (Integragen, Evry, France) using a NovaSeq600 with a coverage of 30X. Burrows-
17
18 Wheeler Aligner (BWA) tool was used to map the reads to the Human genome build (hg38). SNVs
19
20 and small indels calling was performed via Broad Institute's GATK Haplotype Caller GVCf tool
21
22 (GATK 3.8.1) and the Ensembl's VEP (Variant Effect Predictor, release VEP 95.1) program
23
24 processed the variants for further annotation. Variants were prioritized on the basis of a minor
25
26 allele frequency (MAF) ≤ 0.0005 in the genome Aggregation Database ([http://www.gnomad-
29
30 sg.org/](http://www.gnomad-
27
28 sg.org/) gnomAD v3.1.1), representing insertions or deletions (InDels), nonsense, missense, splice
31
32 site variants and assessed for their pathogenicity through bioinformatic tools including
33
34 Polymorphism Phenotyping v2 (PolyPhen2, <http://genetics.bwh.harvard.edu/pph2/>), Sorting
35
36 Intolerant From Tolerant (SIFT, <http://sift.jcvi.org/>), MutationTaster
37
38 (<http://www.mutationtaster.org/>), amino acid conservation across species with UCSC Genome
39
40 Browser (<http://ge-nome.ucsc.edu/index.html>; Human GRCh38/hg38 Assembly), SpliceAI ²² and
41
42 Pangolin ²³, similarly as performed before ²¹. Further prioritization was done by keeping variants
43
44 present in regions of homozygosity (ROHs) unique to the affected member. The ROHs were
45
46 established by doing homozygosity mapping using a program (AutoMap 1.2) ²⁴ on all family
47
48 members then discarding regions shared by affected and unaffected members. Direct sanger
49
50 sequencing on genomic DNA from whole blood from all available family members of F1362
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 (Figure 1A) was used to validate putative disease-causing variants or to perform co-segregation
5
6 analyses. To detect putative copy number variants in affected cases with monoallelic variants in
7
8 the candidate gene, quantitative PCR experiments using primers covering all exons and a kit
9
10 (SYBR Green Real-Time PCR Master Mixes, Applied Biosystems, Thermo Fisher, Villebon-sur-
11
12 Yvette, France) was performed similarly as previously ⁶. The conditions are available on request.
13
14
15 Subsequently, the candidate gene was screened in affected individuals from our French cohort
16
17 mainly of European and North African descendants, by direct Sanger sequencing (conditions will
18
19 be delivered on request) and/or by targeted next generation sequencing (NGS), similarly as
20
21 previously described.
22
23
24
25
26
27

28 **Fibroblasts reprogramming into iPSCs and differentiation into RPE cells and into ROs**

29
30 Human dermal fibroblasts from the affected index case CIC03225 and unaffected brother
31
32 CICC03228 of F1362 were obtained after informed consent at the Quinze-Vingts hospital from a
33
34 skin biopsy and expanded as described before ([Ethical approval: Comité de Protection des
35
36 Personnes Ile de France V (2012-A01333-40; P12-02) and the ANSM (B121362-32)] ²⁵.
37
38 Subsequently they were reprogrammed into iPSCs using reprogramming vectors (CytoTune®
39
40 Sendai OCT3/4, SOX2, CMYC, and KLF4, Thermo Fisher Scientific ²⁵. The iPSC colonies were
41
42 obtained and expanded under feeder free conditions ²⁶. Pluripotency was tested as performed
43
44 before, using a Scorecard technique (Thermo Fisher Scientific, Illkirch, France) ²⁷. For genomic
45
46 integrity and concordance with parental fibroblasts the samples were analyzed by Single
47
48 Nucleotide Polymorphism arrays (SNP) using Illumina InfiniumCore-24 technique (Illumina, San
49
50 Diego, USA) at Integragen ²⁷. Subsequently, iPSCs were differentiated in ROs and RPE cells ²⁶.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Expression analysis

The expression of the *UBAPIL* (HGNC:40028) was investigated using publicly and in-house available database (<https://www.proteinatlas.org/search/UBAPIL> and https://singlecell.broadinstitute.org/single_cell/study/SCP484/nucseq-from-human-control-eyes-scrnaseq²⁸) as previously described²¹. Subsequently, its expression was experimentally verified by RT-PCR experiments using commercially available cDNA of different human tissues (Clontech, Saint- Germain-en-Laye, France), cDNA from human fibroblasts, human-induced iPSCs, ROs and RPE cells of an unaffected individual. A detailed protocol can be provided on request.

Splice defect investigation on ROs

The consequence of the homozygous NM_001163692.2:c.910-7G>A on splicing of *UBAPIL* for the affected index individual CIC03225 and his heterozygous unaffected brother CIC03228 of family F1362 was investigated by RT-PCR experiments applied to ROs of the respective individuals. Forward and reverse oligonucleotides in exons 4 and 5, and 6 amplified transcripts including the NM_001163692.2:c.910-7G>A variant present in intron 4 of *UBAPIL*. The respective products were Sanger sequenced. A detailed protocol can be provided on request.

Zebrafish strain and husbandry

Zebrafish (*Danio rerio*) were bred and maintained at the aquarium of the Institut de la Vision, Paris, France as described before²⁹. Zebrafish were maintained at 28°C on a 14-hour light/10-hour dark cycle. All animal procedures were performed in accordance with French and European

1
2
3
4 Union animal welfare guidelines with protocols approved by committee on ethics of animal
5
6 experimentation of Sorbonne Université (APAFIS#21323-2019062416186982).
7
8
9

10 **Zebrafish RNA *in situ* hybridization and hybridization chain reaction (HCR) analyses**

11
12
13 The 5'- CTGCAGGAAACTGAGTATGTGTTC-3' fw and 5'-
14
15 CCAAGCTTCTAATACGACTCACTATAGGGAGATTCATTAAACTGAGCCAGAAGATGC
16
17 -3' T7-rv primer-set was used to amplify *ubap11b* transcript and the digoxigenin-labeled riboprobe
18
19 were prepared as recommended by the manufacturer instructions (Roche, Basel, Switzerland).
20
21 Single whole-mount *in situ* hybridization of the zebrafish *ubap11b* gene (XM_002666995, NCBI
22
23 accession number) was carried out as previously described³⁰. The *ubap11b* riboprobe was labelled
24
25 with digoxigenin-UTP (Roche). Hybridization with the probe was carried overnight at 68°C. Anti-
26
27 DIG primary antibody coupled to alkaline phosphatase (Roche) and NBT-BCIP (Roche) were used
28
29 for signal detection. Embryos, 5 days after fertilization stained with NBT/BCIP were mounted in
30
31 87% glycerol on microscope slides and imaged with a microscope (Leica MZ10F, Nanterre,
32
33 France). For vibratome sections, whole-mount embryos were washed twice in 1x PBS/0.1%
34
35 Tween-20 (PBS-Tw) solution. The samples were embedded in gelatin/albumin with 4% of
36
37 glutaraldehyde and sectioned (20 µm) on a VT1000 S vibrating blade microtome (Leica). Sections
38
39 were mounted in Fluoromount Aqueous Mounting Medium (Sigma, Merck KGaA, Darmstadt,
40
41 Germany).
42
43
44
45
46
47
48
49

50 The HCR procedure was performed following the manufacturer instruction (Molecular
51
52 Instruments, Los Angeles, USA) and the *ubap11b* probe designed. Labelled embryos were
53
54 embedded and mounted on 100 × 15 mm glass bottom dishes in 1% low melting agarose and
55
56 *ubap11b* HCR probe signal was acquired using a confocal microscope (LSM 900, Zeiss, Jena,
57
58
59
60
61
62
63
64
65

1
2
3
4 Germany) employing a 40x water immersion objective. Software were used for image analyses
5
6 (Imaris Viewer 10.0, Oxford Instruments Group, Gometz la Ville, France) and image processing
7
8 (Adobe Photoshop, Adobe Illustrator, Paris, France).
9

10 11 12 13 14 **3D-modeling**

15
16 The wild type 3D structure of UBAP1L (UniprotKB: F5GYI3) was downloaded from the official
17
18 AlphaFold database: <https://alphafold.ebi.ac.uk/>. For the mutated proteins the AlphaFold (version
19
20 2.3.2) protein folding prediction program using the monomer model with default settings was used
21
22
23
24 31.

25 26 27 28 29 30 31 32 **Results:**

33 34 35 **Phenotypic characteristics of affected individuals**

36
37 At presentation, CIC03225, F1362, was a 23-year-old male subject who complained of night
38
39 blindness and progressive visual field constriction since he was 20_years old. He was the oldest
40
41 son from a consanguineous Tunisian couple (Figure 1A). Best corrected visual acuity (BCVA)
42
43 was 20/32 for both eyes (BE) with -7 diopter optical correction. He had a tritan defect on color
44
45 vision testing (Lanthony's 15 desaturated Hue). Kinetic visual field assessment showed the
46
47 presence of paracentral scotomas. Full field electroretinogram (ff-ERG) was undetectable. Retinal
48
49 imaging revealed typical signs of RCD with macular atrophy in BE (Figure 2). CIC06376, F3377
50
51 was a 50-year-old Tunisian male from a consanguineous union (Figure 1A) with a clinical
52
53 diagnosis of CRD. BCVA was 20/200 in the right eye (RE) and 20/50 in the left eye (LE) with no
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 specific optical correction. Kinetic visual field testing revealed the presence of a central scotoma
5
6 in BE. Ff-ERG showed only residual responses under scotopic conditions confirming the severe
7
8 cone-rod dysfunction. Retinal imaging displayed a pale optic disc, narrowed retinal vessels and
9
10 atrophic changes of the posterior pole and the inferior retina in BE (Figure 2). CIC09282, F5326
11
12 was a 55-year-old Tunisian woman, from a consanguineous couple (Figure 1A). She was referred
13
14 to the rare disease center with a presumed diagnosis of Stargardt disease. BCVA was 20/500 with
15
16 $-0.50(-2.25)15^\circ$ in the RE and 20/400 with $-0.50(-1.50)160^\circ$ in the LE. Color vision testing
17
18 revealed complete dyschromatopsia. Kinetic visual field assessment showed severe constriction in
19
20 BE. Ff-ERG revealed only residual responses under scotopic conditions consistent with severe
21
22 cone-rod dysfunction. Retinal imaging revealed a pale optic disc, narrowed retinal vessels and
23
24 atrophic changes predominant at the posterior pole in BE (Figure 2). CIC13094, F7544 was a 50-
25
26 year-old female from a consanguineous Tunisian couple (Figure 1A) when she was first assessed
27
28 at the rare disease center. She has been complaining about night blindness since teenaged. BCVA
29
30 was 20/160 with $-1.25(-2)175^\circ$ in the RE and 20/800 with $-0.50(-2.25)175^\circ$ in the LE. Kinetic
31
32 visual field testing revealed a central scotoma for BE. Again, ff-ERG responses were only
33
34 observed under scotopic conditions, indicating severe cone-rod dysfunction. Retinal imaging
35
36 revealed atrophic changes in the posterior pole and mid periphery in BE (Figure 2). The summary
37
38 of the clinical data of all individuals with *UBAPIL* variants and a figure showing a normal fundus
39
40 upon photography and autofluorescence imaging can be found in the supplementary data.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Identification of three *UBAPIL* variants in four cases from four unrelated families

The genetic causes of RCD for the affected index, CIC03225 (F1362, Figure 1A), from a consanguineous Tunisian couple remained unsolved despite conducting Sanger sequencing of candidate genes implicated in RCD (e.g.^{16,17}), microarray analysis¹⁸, targeted NGS using a panel of 123 genes or candidate genes implicated in IRDs¹⁹ and ES²¹. Similarly, the search for structural variations using ES raw data did not reveal any candidate copy number variation. However, array comparative genomic hybridization (CGH) and homozygosity mapping using WGS data identified three large (>20 Mb) homozygous regions in the affected index with sizes of 27.7 Mb, 20.9 Mb, 25.8 Mb mapping on chromosome 1, 10 and 15, respectively (Figure 1B). Applying stringent filtering, we identified a homozygous variant in chromosome 15: NM_001163692.2:c.910-7G>A (allele frequency in gnomAD: 1.64e-4, with no homozygous appearance) in a gene, *UBAPIL* within the 25.8 Mb homozygous region on chromosome 15q22.31 in this affected individual. Notably, this gene was absent in a database, which associates genes with IRDs: <https://sph.uth.edu/retnet/sym-dis.htm>. This variant co-segregated with the phenotype in this family (Figure 1A) and was predicted to create a new AG acceptor site at position NM_001163692.2:c.910-5, which would lead to an altered longer protein p.(Phe304Profs*92). Alternatively, nonsense mediated mRNA decay could take place. Previous ES analysis had overlooked this variant, since it was not affecting a canonical splice site. A heterozygous *PROM1* (HGNC:9454) variant, NM_006017.3:c.314A>G p.(Tyr105Cys) was considered as not disease causing, since the unaffected mother, CIC03226 and the unaffected brother, CIC03228 were also heterozygous for this variant. Interestingly, *UBAPIL* was one of 19 potential candidate genes previously listed in a genetically unsolved RP case originating from South Asia³². Noteworthy, *UBAPIL* lies within a super enhancer (SE) region, which was identified by chromatin

1
2
3
4 immunoprecipitation sequencing (ChIP-seq) performed on post-mortem adult human retinas^{33,34}.
5
6 This region contains a cluster of retinal *cis*-regulatory elements (CREs). CREs are characterized
7
8 by a high density of histone modifications associated with promoters and active or poised
9
10 enhancers. Specifically, the presence of histone modification H3K27Ac, commonly associated
11
12 with active enhancers is observed in these CREs. In additions, the presence of ATAC-seq signals
13
14 indicate a high chromatin accessibility in these regions (Figure 1C). These elements supported
15
16
17
18 *UBAPIL* as a strong candidate gene underlying IRDs^{33,34}.
19
20

21 Subsequently, Sanger sequencing of our genetically unresolved French cohort mainly of European
22
23 and North African descendants, revealed another affected individual (CIC06376, F3377) from a
24
25 consanguineous Tunisian couple with a presumably homozygous deletion,
26
27 NM_001163692.2:c.634_644del (allele frequency in gnomAD: 2.64e-5, with no homozygous
28
29 appearance) in *UBAPIL*. This deletion was predicted to result in a premature stop codon,
30
31 p.(Ser212Alafs*44) or mRNA decay (Figure 1A). Similarly, an updated targeted NGS, including
32
33 recent candidate genes, identified a third case (CIC13094, F7544) from a consanguineous Tunisian
34
35 couple harboring the same variant (Figure 1A). This patient revealed also a most likely
36
37 homozygous *RPILI* (HGNC:15946) variant, NM_178857.6:c.92C>T p.(Thr31Met), which was
38
39 predicted to be benign. In addition, using this method a presumably homozygous nonsense variant,
40
41 NM_001163692.2:c.859C>T (allele frequency in gnomAD: 3.29e-5, with no homozygous
42
43 appearance), in *UBAPIL*, predicted to lead to a premature stop codon, p.(Arg287*) or nonsense
44
45 mediated mRNA decay was identified in a fourth family (CIC09282, F5326) from another
46
47 consanguineous Tunisian couple (Figure 1A). This patient revealed also a heterozygous *GUCY2D*
48
49 (HGNC:4689) variant, NM_000180.4:c.1984G>A p.(Val662Met), which was predicted to be
50
51 benign.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 All four affected index individuals with *UBAPIL* variants identified herein reported consanguinity
5
6 among their parents and were from Tunisian couples. This was further confirmed by homozygosity
7
8 mapping analysis in F1362, which could be done with available WGS data, showing multiple
9
10 homozygous regions in the affected index case, which were absent in the unaffected family
11
12 members (Figure 1B). For F5326, homozygosity mapping using available targeted NGS data also
13
14 revealed a 35 Mb homozygous region encompassing *UBAPIL*. Furthermore, none of the
15
16 sequenced polymorphisms in the coding regions of *UBAPIL* were found to be heterozygous in
17
18 F3377 and F7544. These findings collectively support the notion that the gene defect in *UBAPIL*
19
20 is indeed located within a homozygous region, consistent with the consanguinity in all families.
21
22
23
24
25
26
27
28
29
30

31 ***UBAPIL* is expressed in the brain, RPE cells, in the retina and more precisely in rod and** 32 **cone photoreceptors**

33 Transcriptomic databases reported that human *UBAPIL* is expressed in the brain, the eye and more
34
35 specifically in RPE cells as well as in rod and cone photoreceptors²⁸ (Figure 3A). These data were
36
37 confirmed by RT-PCR experiments (Figure 3B). *UBAPIL* was found to be expressed in human
38
39 universal embryonal tissue and lung, but more importantly in brain, retina, fibroblasts, hiPSCs,
40
41 ROs and RPE cells differentiated from iPSCs from an unaffected individual (Figure 3B). In human
42
43 universal embryonal tissue and lung other unspecific transcripts appeared (Figure 3B). *In silico*
44
45 analysis revealed two isoforms of *UBAPLI* in the zebrafish, named *ubap1la* and *ubap1lb*. We
46
47 decided to investigate the expression of these two genes during normal zebrafish development.
48
49 Notably, *ubap1lb* exhibited a remarkably distinct expression pattern in zebrafish larvae at 3 and 5
50
51 days after fertilization. *In situ* hybridization and HCR on zebrafish embryos revealed that the
52
53 *ubap1lb* isoform is specifically expressed in the retina, and more specifically in rod and cone
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 photoreceptors (Figure 4). This gave further evidence for a role of UBAP1L during normal retinal
5
6 development and its potential alteration in the pathological retina.
7
8
9

10 11 12 13 14 **The NM_001163692.2:c.910-7G>A variant leads to mis-splicing of *UBAP1L* in ROs**

15
16 To investigate if the NM_001163692.2:c.910-7G>A, located in intron 4 indeed of *UBAP1L* lead
17
18 an altered longer protein p.(Phe304Profs*92) or to nonsense mediated mRNA decay, RT-PCR
19
20 experiments performed on ROs at 150 days of the affected individual CIC03225 and his unaffected
21
22 heterozygous brother, CIC03228 of F1326 were performed. These revealed in ROs of the patient
23
24 the insertion of 5 nucleotides (GCCCA), while in ROs of the unaffected brother, both, the reference
25
26 and mutated mRNA sequence were present, as confirmed by Sanger sequencing (Figure 5). These
27
28 findings confirmed the *in silico* prediction for mis-splicing associated with the
29
30 NM_001163692.2:c.910-7G>A variant in *UBAP1L* leading most likely to a longer protein
31
32 p.(Phe304Profs*92). Other bands occurring in the unaffected heterozygous brother were not
33
34 investigated in detail.
35
36
37
38
39
40
41
42

43 **In Silico predictions and *in vitro* functional characterization of *UBAP1L* variants**

44
45 *UBAP1L* (NM_001163692.2) containing 6 exons codes for Ubiquitin-associated protein 1-like
46
47 comprising 381 amino acids. The open reading frame starts in exon 2 and ends in exon 6 (Figure
48
49 6A). The protein contains a solenoid of overlapping ubiquitin associated (UBA) domains
50
51 (SOUBA), which was shown to interact with ubiquitin in UBAP1³⁵. The SOUBA domain starts
52
53 at amino acid residue Ser269 and ends at the C-terminus. It is composed of 3 overlapping UBAs
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 forming each 3 α -helices (Figure 6B). Thus, the SOUBA domain is formed by 7 α -helices (α 1- α 7),
5
6 as shown in the 3D model (Figure 6C, wild-type).
7

8
9 The three variants in *UBAPIL* identified in this study could all affect the SOUBA domain (Figure
10
11 6). The NM_001163692.2:c.634_644del, located in exon 3 that leads to the frameshift
12
13 p.(Ser212Alafs*44) occurs upstream of the SOUBA domain (Figure 6B). Therefore, none of the
14
15 7 α -helices would remain, meaning there is no UBA left to bind ubiquitin (Figure 6C). Similarly,
16
17 the NM_001163692.2:c.859C>T, located in exon 4 that leads to the premature stop codon
18
19 p.(Arg287*) occurs within the SOUBA domain. Therefore, only α 1 and a small part of α 2 remain
20
21 (Figure 6C), which severely alter the capabilities of the protein to bind ubiquitin. However, both
22
23 variants may most likely cause nonsense mediated mRNA decay leading to complete loss of
24
25 function. Previously another variant in *UBAPIL*, NM_001163692.2:c.121-2A>C (Figure 6A) was
26
27 described as a possible cause for RP³². It was predicted to lead to exon 3 skipping and an in-frame
28
29 deletion, p.(His41_Ala233del) (Figure 6B), which would not affect the SOUBA domain, but delete
30
31 a significant proportion of the coding sequence amino, which will most likely also affect the
32
33 protein function (Figure 6C). The NM_001163692.2:c.910-7G>A, located in intron 4 was
34
35 predicted to affect splicing, with a delta score of 0.99 by SpliceAI and 0.84 by Pangolin leading to
36
37 an altered longer protein, p.(Phe304Profs*92). Indeed, our studies performed on ROs confirmed
38
39 the mis-splicing (Figure 5). The frameshift starts in exon 5, which also codes for the SOUBA
40
41 domain. Therefore, if the variant leads to an altered protein product, only α 1, α 2 and a small part
42
43 of α 3 would remain, severely altering the capabilities of the protein to bind ubiquitin (Figure 6C).
44
45 Therefore, we think that all variants identified herein are predicted to lead to a highly modified
46
47 protein, which will be unable to bind ubiquitin.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Discussion

Variants implicated in progressive non-syndromic RCD or CRD affect genes expressed in photoreceptors and/or RPE, and they code for proteins involved in different cellular functions, including phototransduction, cellular differentiation, structure and maintenance, retinal metabolism and splicing⁴. To date variants in more than 80 genes (<https://web.sph.uth.edu/RetNet/sum-dis.htm>) can lead to these conditions⁵. With the new advent of NGS, novel gene defects were rapidly discovered. It is much likely that all the major gene defects underlying IRDs have now been reported. As shown in our center, up to 80% of the cases can be solved by a combination of Sanger and targeted NGS. However, it remains challenging to determine the genetic cause for the remaining 20% of cases, even when employing comprehensive ES and GS methods. Novel gene defects are nowadays rare and may only be observed within a single family⁶. To enhance the identification of these rare gene defects researchers gather large IRD cohorts, often use collaborative platforms and tools such as GeneMatcher³⁶ (<https://genematcher.org>) and/or engage in collaborations, such as the European Retinal Disease Consortium (ERDC) (<http://www.erd.c.info/>). These resources facilitate the identification of additional cases that provide further evidence supporting the involvement of a newly identified candidate gene. *In silico* predictions as well as *in vitro* and *in vivo* modeling may also be valuable approached to assess the pathogenicity of a novel putative disease-causing gene defect.

Although identifying the gene defect of our initial case (CIC03225) with RCD and macular atrophy was laborious, only WGS in combination with expression data, *in silico* predictions and functional analyses highlighted the candidate gene defect. This one resulted in a homozygous splice site change in *UBAPIL*, most likely leading to an altered protein length, with a truncated SOUBA domain. Interestingly, we identified three additional cases with *UBAPIL* variants through direct

1
2
3
4 Sanger and targeted NGS, in which the *UBAPIL* protein is predicted to have shorter or absent
5
6 SOUBA domains or is not functional. This was only possible while screening a large IRD cohort
7
8 encompassing ~4.000 deeply phenotypically and genetically explored index cases with RCD or
9
10 CRD. Altogether, *UBAPIL* disease causing variants remain rare in our cohort, with an estimated
11
12 prevalence of about 0.1%. Of note, the four cases identified herein are all originating from
13
14 consanguineous Tunisian couples with one recurrent variant (i.e. NM_001163692.2:c.634_644del)
15
16
17 in two presumably unrelated families. The prevalence of Tunisian ancestry cases with progressive
18
19 RCD or CRD in our cohort is unknown. In addition, an *UBAPIL* variant was listed as a potential
20
21 cause of RP originating from South Asia ³². It will be interesting, to determine the prevalence of
22
23 *UBAPIL* variants in Tunisian and other progressive IRD cohorts of different ethnicities.
24
25

26
27
28 Regarding the phenotype of the subjects reported herein, one case reported primary onset of rod-
29
30 related symptoms, consistent with the diagnosis of RCD whereas the other subjects had decreased
31
32 central vision and abnormal light sensitivity at the onset of symptoms more in line with CRD.
33
34 Typically, RCD is characterized by primary rod photoreceptor leading to decreased vision in dim
35
36 light conditions, and secondary cone dysfunction and degeneration, usually starting from the
37
38 peripheral retina, leading to progressive visual field constriction in day light typically preserving
39
40 central vision until late in the course of the disease. On the other hand, CRD usually starts by
41
42 decreased central vision color vision defect, light sensitivity and secondary rod degeneration
43
44 manifesting with secondary night blindness. The distinction between RCD and CRD dystrophy
45
46 rely on patient's visual symptoms, functional tests such as ff-ERG which typically show
47
48 predominant rod dysfunction in case of RCD or cone dysfunction in case of CRD and multimodal
49
50 retinal imaging showing predominant peripheral disease and relatively preserved macula in RCD
51
52
53 whereas CRD may display significant macular changes early on in the course of the disease.
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Nevertheless, the distinction between RCD and CRD may be difficult especially in advanced cases
5
6 where ff-ERG responses are usually undetectable and somewhat artificial some dystrophies where
7
8 both photoreceptor systems may appear concomitantly affected. Noticeably, the RCD case from
9
10 this study (CIC03225) already displayed severe involvement of the central macular region when
11
12 he was 23 years old, unlike typical RCD cases where the macular region is preserved at the time
13
14 of presentation. *UBAPIL*-disease-causing variants may therefore be associated with a phenotypic
15
16 spectrum ranging from RCD with early onset macular involvement to CRD.
17
18
19

20
21 Expression analyses using publicly available databases, including Chip-seq data, as well as
22
23 expression studies on human-derived ROs, RPE cells and zebrafish collected further evidence for
24
25 *UBAPIL* to be an excellent candidate gene underlying rod and cone photoreceptor disease.
26
27

28
29 *UBAPIL*, is a fairly unknown protein. It is predicted to enable ubiquitin binding activity and is
30
31 believed to be involved in the ubiquitin-dependent protein catabolic process through the
32
33 multivesicular body sorting pathway. It is also predicted to be part of endosomal sorting complexes
34
35 required for transport (ESCRT I complex (<https://www.alliancegenome.org/gene/HGNC:40028>)).
36
37
38 *UBAPIL* contains a SOUBA domain, which begins at amino acid residue 269 and extends to the
39
40 C-terminus of the protein. This domain is similar to the one found in *UBAP1*, where it is known
41
42 to interact with ubiquitin ³⁵. The role of the N-terminal part of the protein is unknown. The
43
44 previously reported candidate variant in *UBAPIL*, NM_001163692.2:c.121-2A>C was predicted
45
46 to lead to exon 3 skipping and an in-frame deletion, p.(His41_Ala233del) ³², which would not
47
48 affect the SOUBA domain. However, it is predicted to delete a significant portion of the protein
49
50 and thus leads most likely to a non-functional protein.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 In all affected subjects described so far, the SOUBA domain of UBAP1L may be truncated or the
5
6 protein will be non-functional or absent and thus ubiquitin binding will not correctly occur. This
7
8 implies that the ubiquitin proteasome system (UPS) will be altered. The UPS plays an important
9
10 role in removing abnormal proteins and preventing accumulation of non-functional and harmful
11
12 proteins within the neuron. Different studies highlighted UPS in retinal health and disease
13
14 (reviewed in ³⁷). More specifically, an impairment of the UPS has been associated with pathogenic
15
16 mechanisms of IRD associated with variants in *RHO* (HGNC:10012) ³⁸, *TOPORS* (HGNC:21653)
17
18 ³⁹ or *KLHL7* (HGNC:15646) ⁴⁰ among others. While variants in these genes have been mainly
19
20 associated with autosomal dominant IRD and increased ubiquitination, *UBAP1L* variants
21
22 associated with autosomal recessive IRD leading to truncated SOUBA proteins may induce a
23
24 dysregulation of the UPS resulting in cellular stress and photoreceptor degeneration. Functional
25
26 studies such as those using mutant and CRISPR/Cas9 edited hiPSCs differentiated in ROs and
27
28 RPE cells as well as zebrafish disease modeling will be important to further decipher the
29
30 pathogenic mechanism implicate in *UBAP1L*-associated retinal disorders.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Reference

1. Lindsay S, Inglehearn CF, Curtis A, Bhattacharya S. Molecular genetics of inherited retinal degenerations. *Curr Opin Genet Dev.* 1992;2(3):459-466.
2. Hamel C. Retinitis pigmentosa. *Orphanet J Rare Dis.* 2006;1:40.
3. Roosing S, Thiadens AA, Hoyng CB, Klaver CC, den Hollander AI, Cremers FP. Causes and consequences of inherited cone disorders. *Prog Retin Eye Res.* 2014;42:1-26.
4. Berger W, Kloeckener-Gruissem B, Neidhardt J. The molecular basis of human retinal and vitreoretinal diseases. *Prog Retin Eye Res.* 2010;29(5):335-375.
5. Schneider N, Sundaresan Y, Gopalakrishnan P, et al. Inherited retinal diseases: Linking genes, disease-causing variants, and relevant therapeutic modalities. *Prog Retin Eye Res.* 2021:101029.
6. Zeitz C, Mejecase C, Michiels C, et al. Mutated CCDC51 Coding for a Mitochondrial Protein, MITOK Is a Candidate Gene Defect for Autosomal Recessive Rod-Cone Dystrophy. *Int J Mol Sci.* 2021;22(15).
7. Mejecase C, Hummel A, Mohand-Said S, et al. Whole exome sequencing resolves complex phenotype and identifies CC2D2A mutations underlying non-syndromic rod-cone dystrophy. *Clin Genet.* 2019;95(2):329-333.
8. Solaguren-Beascoa M, Bujakowska KM, Mejecase C, et al. WDR34, a candidate gene for non-syndromic rod-cone dystrophy. *Clin Genet.* 2021;99(2):298-302.
9. Zeitz C, Michiels C, Neuville M, et al. Where are the missing gene defects in inherited retinal disorders? Intronic and synonymous variants contribute at least to 4% of CACNA1F-mediated inherited retinal disorders. *Hum Mutat.* 2019;40(6):765-787.
10. Weissschuh N, Buena-Atienza E, Wissinger B. Splicing mutations in inherited retinal diseases. *Prog Retin Eye Res.* 2021;80:100874.
11. de Bruijn SE, Fiorentino A, Ottaviani D, et al. Structural Variants Create New Topological-Associated Domains and Ectopic Retinal Enhancer-Gene Contact in Dominant Retinitis Pigmentosa. *Am J Hum Genet.* 2020;107(5):802-814.
12. Ueyama H, Li YH, Fu GL, et al. An A-71C substitution in a green gene at the second position in the red/green visual-pigment gene array is associated with deutan color-vision deficiency. *Proc Natl Acad Sci U S A.* 2003;100(6):3357-3362.
13. Cremers FPM, Boon CJF, Bujakowska K, Zeitz C. Special Issue Introduction: Inherited Retinal Disease: Novel Candidate Genes, Genotype-Phenotype Correlations, and Inheritance Models. *Genes (Basel).* 2018;9(4).
14. Biswas P, Borooah S, Matsui H, et al. Detection and validation of novel mutations in MERTK in a simplex case of retinal degeneration using WGS and hiPSC-RPEs model. *Hum Mutat.* 2021;42(2):189-199.
15. Audo I, Friedrich A, Mohand-Said S, et al. An unusual retinal phenotype associated with a novel mutation in RHO. *Arch Ophthalmol.* 2010;128(8):1036-1045.
16. Audo I, Sahel JA, Mohand-Said S, et al. EYS is a major gene for rod-cone dystrophies in France. *Hum Mutat.* 2010;31(5):E1406-1435.

17. Audo I, Lancelot ME, Mohand-Said S, et al. Novel C2orf71 mutations account for approximately 1% of cases in a large French arRP cohort. *Hum Mutat.* 2011;32(4):E2091-2103.
18. Avila-Fernandez A, Cantalapiedra D, Aller E, et al. Mutation analysis of 272 Spanish families affected by autosomal recessive retinitis pigmentosa using a genotyping microarray. *Mol Vis.* 2010;16:2550-2558.
19. Audo I, Bujakowska KM, Leveillard T, et al. Development and application of a next-generation-sequencing (NGS) approach to detect known and novel gene defects underlying retinal diseases. *Orphanet J Rare Dis.* 2012;7(1):8.
20. Boulanger-Scemama E, El Shamieh S, Demontant V, et al. Next-generation sequencing applied to a large French cone and cone-rod dystrophy cohort: mutation spectrum and new genotype-phenotype correlation. *Orphanet J Rare Dis.* 2015;10:85.
21. El Shamieh S, Neuille M, Terray A, et al. Whole-exome sequencing identifies KIZ as a ciliary gene associated with autosomal-recessive rod-cone dystrophy. *Am J Hum Genet.* 2014;94(4):625-633.
22. Jaganathan K, Kyriazopoulou Panagiotopoulou S, McRae JF, et al. Predicting Splicing from Primary Sequence with Deep Learning. *Cell.* 2019;176(3):535-548 e524.
23. Zeng T, Li YI. Predicting RNA splicing from DNA sequence using Pangolin. *Genome Biol.* 2022;23(1):103.
24. Quinodoz M, Peter VG, Bedoni N, et al. AutoMap is a high performance homozygosity mapping tool using next-generation sequencing data. *Nat Commun.* 2021;12(1):518.
25. Terray A, Slembrouck A, Nanteau C, et al. Generation of an induced pluripotent stem cell (iPSC) line from a patient with autosomal dominant retinitis pigmentosa due to a mutation in the NR2E3 gene. *Stem Cell Res.* 2017;24:1-4.
26. Reichman S, Slembrouck A, Gagliardi G, et al. Generation of Storable Retinal Organoids and Retinal Pigmented Epithelium from Adherent Human iPS Cells in Xeno-Free and Feeder-Free Conditions. *Stem Cells.* 2017;35(5):1176-1188.
27. Wohlschlegel J, Letellier C, Liu B, et al. Generation of human induced pluripotent stem cell lines from a patient with ITM2B-related retinal dystrophy and a non mutated brother. *Stem Cell Res.* 2019;41:101625.
28. Orozco LD, Chen HH, Cox C, et al. Integration of eQTL and a Single-Cell Atlas in the Human Eye Identifies Causal Genes for Age-Related Macular Degeneration. *Cell Rep.* 2023;42(3):112298.
29. Rosello M, Vouigny J, Czarny F, et al. Precise base editing for the in vivo study of developmental signaling and human pathologies in zebrafish. *Elife.* 2021;10.
30. Oxtoby E, Jowett T. Cloning of the zebrafish krox-20 gene (krx-20) and its expression during hindbrain development. *Nucleic Acids Res.* 1993;21(5):1087-1095.
31. Jumper J, Evans R, Pritzel A, et al. Highly accurate protein structure prediction with AlphaFold. *Nature.* 2021;596(7873):583-589.
32. Carss KJ, Arno G, Erwood M, et al. Comprehensive Rare Variant Analysis via Whole-Genome Sequencing to Determine the Molecular Pathology of Inherited Retinal Disease. *Am J Hum Genet.* 2017;100(1):75-90.
33. Marchal C, Singh N, Batz Z, et al. High-resolution genome topology of human retina uncovers super enhancer-promoter interactions at tissue-specific and multifactorial disease loci. *Nat Commun.* 2022;13(1):5827.

- 1
- 2
- 3
- 4 34. Cherry TJ, Yang MG, Harmin DA, et al. Mapping the cis-regulatory architecture of the
- 5 human retina reveals noncoding genetic variation in disease. *Proc Natl Acad Sci U S A.*
- 6 2020;117(16):9001-9012.
- 7
- 8 35. Agromayor M, Soler N, Caballe A, et al. The UBAP1 subunit of ESCRT-I interacts with
- 9 ubiquitin via a SOUBA domain. *Structure.* 2012;20(3):414-428.
- 10 36. Sobreira N, Schiettecatte F, Valle D, Hamosh A. GeneMatcher: a matching tool for
- 11 connecting investigators with an interest in the same gene. *Hum Mutat.* 2015;36(10):928-
- 12 930.
- 13
- 14 37. Campello L, Esteve-Rudd J, Cuenca N, Martin-Nieto J. The ubiquitin-proteasome system
- 15 in retinal health and disease. *Mol Neurobiol.* 2013;47(2):790-810.
- 16 38. Illing ME, Rajan RS, Bence NF, Kopito RR. A rhodopsin mutant linked to autosomal
- 17 dominant retinitis pigmentosa is prone to aggregate and interacts with the ubiquitin
- 18 proteasome system. *J Biol Chem.* 2002;277(37):34150-34160.
- 19 39. Czub B, Shah AZ, Alfano G, Kruczek PM, Chakarova CF, Bhattacharya SS. TOPORS, a
- 20 Dual E3 Ubiquitin and Sumo1 Ligase, Interacts with 26 S Protease Regulatory Subunit 4,
- 21 Encoded by the PSMC1 Gene. *PLoS ONE.* 2016;11(2):e0148678.
- 22 40. Kim J, Tsuruta F, Okajima T, Yano S, Sato B, Chiba T. KLHL7 promotes TUT1
- 23 ubiquitination associated with nucleolar integrity: Implications for retinitis pigmentosa.
- 24 *Biochem Biophys Res Commun.* 2017;494(1-2):220-226.
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

1
2
3
4
5
6
7
8 **Figure legends:**
9

10 **Figure 1. Families with autosomal recessive RCD and CRD/CD and pathogenic variants in**
11 ***UBAPIL***
12
13
14

15
16 (A) Pedigrees of the four families (F1362, F3377, F5326 and F7544) with variants in *UBAPIL* co-
17 segregating with the phenotype in all available family members. The arrows indicate the genetic
18 screening performed on each subject. DNA from the family members of family F3377, F5326 and
19 F7544 were unavailable, but due to the reported consanguinity, we presumed that M2 and M3
20 occurred homozygous in the affected index cases.
21
22
23
24
25
26
27
28

29 (B) Homozygosity mapping using WGS data identified three large (>20 Mb) homozygous regions
30 in the affected index individual CIC03225 of F1362 with sizes of 27.7 Mb, 20.9 Mb, 25.8 Mb
31 mapping to chromosome 1, 10 and 15, respectively.
32
33
34
35
36
37

38 (C) Super Enhancer (red), retinal cis-regulatory elements (light green), topologically associated
39 domain (teal), chromatin accessibility (orange), histone modifications (green) and transcription
40 factors residency (blue) for the *UBAPIL* locus. Chromatin accessibility has been defined by
41 ATAC-seq from post-mortem adult human retina. Histone modifications and transcription factor
42 residency have been defined by ChIP-seq from post-mortem adult human retina. Super enhancer
43 and retinal *cis*-regulator elements have been calculated from previously described experimental
44 data^{33,34}.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Figure 2. Retinal images of index cases carrying homozygous variants in *UBAPIL***
5 **(CIC03225, CIC06376, CIC09282 and CIC13094):** color fundus photographs (CFP) and fundus
6
7 autofluorescence imaging (FAF) of the right and left eye (RE and LE respectively).
8
9

10
11
12 **CIC03225**'s fundus examination revealed typical signs of RCD with pale optic discs, narrowed
13 retinal vessels and central macular atrophy; a loss of autofluorescence in the periphery as well as
14
15 in the macular region. **CIC06376** showed central macular atrophy expanding into the mid
16
17 periphery predominantly in the inferior sector; a loss of autofluorescence in the atrophic areas
18
19 surrounded by an area of heterogeneous autofluorescence expanding in the mid periphery.
20
21 **CIC09282** had pale optic discs, narrowed retinal vessels, atrophic changes in the macular area
22
23 with a loss of central autofluorescence surrounded by a ring of increased autofluorescence.
24
25 **CIC13094** had central macular atrophy; speckled appearance of autofluorescence at the posterior
26
27 pole and mid periphery.
28
29
30
31
32
33
34

35 **Figure 3. Expression analysis of *UBAPIL* mRNA in human tissues by RT-PCR experiments**

36
37
38 (A) Expression analyses using databases revealed that the human *UBAPIL* gene is expressed in
39
40 the eye and more specifically in RPE cells and rod and cone photoreceptors with a low expression
41
42 in bipolar cells.
43
44

45
46 (B) *UBAPIL* was found to be expressed in human universal embryonal tissue and lung, but more
47
48 importantly in brain, retina, fibroblasts, hiPSCs, retinal organoids (ROs) and RPE cells
49
50 differentiated from hiPSCs. Other unspecific transcripts appeared in universal tissue and lung.
51
52
53
54

55 **Figure 4. Expression analysis of *ubap1l* in zebrafish**

1
2
3
4 *In situ* hybridization for *ubap1lb* in zebrafish larvae 5 days after fertilization (left). The dorsal view
5
6 is on the top and retina vibratome section is on the bottom. Expression can be detected in the retina
7
8 in the photoreceptor layer. On the right, fluorescent HCR *in situ* hybridization on 3 days after
9
10 fertilization larvae confirms *ubap1lb* mRNA expression in the OS and ONL where photoreceptors
11
12 are located. ONL, outer nuclear layer; INL, inner nuclear layer; GCL, Ganglion cell layer; OS,
13
14 outer segment.
15
16
17
18
19

20 **Figure 5. Mis-splicing in ROs of an affected individual lead to a longer protein**

21
22 RT-PCR experiments performed on ROs at 150 days from the affected individual CIC03225
23
24 revealed a 5-nucleotide insertion, while the unaffected heterozygous brother, CIC03228 revealed
25
26 both, the 5-nucleotide insertion and the reference sequence on transcript level, as confirmed by
27
28 Sanger sequencing. These findings confirmed the *in silico* prediction for mis-splicing of the c.910-
29
30 7G>A in *UBAP1L* leading most likely to a longer abnormal protein p.(Phe304Profs*92).
31
32
33
34

35 **Figure 6. Gene structure, prediction of domains and 3D-modeling of the unaffected and** 36 37 **altered UBAP1L protein**

38
39
40
41 (A) Schematic representation of *UBAP1L* gene structure with the positions of identified variants.
42
43 (B) Schematic representation of UBAP1L protein structure with the positions of identified
44
45 variants. The SOUBA domain structure is detailed showing the composition of UBAs and α -
46
47 helices.
48
49
50 (C) AlphaFold 3D structure predictions of UBAP1L wild type (UniprotKB: F5GYI3),
51
52 p.(His41_Ala233del), p.(Ser212Alafs*44), p.(Arg287*) and p.(Phe304Profs*92). The colour
53
54 scheme is a gradient that goes from N-terminus (blue) to C-terminus (red).
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **Data Availability Statement**
5
6

7 All data and materials will be supplied upon request.
8
9

10
11
12
13
14 **Authors contributions**
15
16

17 Conceptualization: C.Z., I.A.; Writing-original draft: C.Z., I.A.; Writing-review & editing: J.N.,
18 S.A., F.D.B. Data curation: J.N., L.A.P., C.M., L.N., C.C., Ali.A., C.M., C.L., T.BY., J.W., C.A.,
19 L.B., Ale.A., M.N., S.ES., V.S., S.CB., S.MS., I.A., J.AS., M.S., And.A., S.A. Formal analysis:
20 J.N., L.A.P., C.M., L.N., C.C., Ali.A., C.M., C.L., C.A., L.B., Ale.A., M.N., S.ES., V.S., S.CB.,
21 S.MS., I.A., M.S., S.A. Supervision: C.Z., I.A., F.D.B., S.A.; Funding acquisition: C.Z., I.A.
22 F.D.B.
23
24
25
26
27
28
29
30
31

32
33 **Acknowledgments**
34
35

36 The authors are thankful to the patient and family members participating in the study and clinical
37 staff from the Reference Center for rare diseases REFERET at Quinze-Vingts hospital. DNA
38 samples included in this study originate from NeuroSensCol DNA bank (PI: JA Sahel, coPI I
39 Audo, partner with Centre Hospitalier National d’Ophtalmologie des Quinze-Vingts, INSERM
40 and CNRS). We thank Maia Brunstein and Beatrix Meha of the Hearing Institute Bioimaging Core
41 Facility of C2RT/C2RA (Institut de l’Audition-Institut Pasteur, Paris – France) for technical
42 support.
43
44
45
46
47
48
49
50
51
52

53
54 **Funding**
55
56

57 This research was funded by Fondation Voir et Entendre (CZ), LABEX LIFESENSES (reference
58 ANR-10-LABX-65 to I.A. and C.Z.) supported by French state funds managed by the Agence
59
60
61
62
63
64
65

1
2
3
4 Nationale de la Recherche within the Investissements d’Avenir program (ANR-11-IDEX-0004-
5
6 0), IHU FOReSIGHT (ANR-18-IAHU-0001 to I.A. and C.Z) and (ANR-18-IAHU-01 to F.D.B.)
7
8 supported by French state funds managed by the Agence Nationale de la Recherche within the
9
10 Investissements d’Avenir program, Agence Nationale de la Recherche (ANR-18-CE16-0017-01
11
12 IPOGUT, ANR-20-CE17-0020-02 INCEPTION, ANR-22-CE17-028 PEMGeT to F.D.B.) and
13
14 ANR-23-CE17-0014-01 RP_SOLVEANDCURE to C.Z., I.A. and F.D.B.), Fondation pour la
15
16 Recherche Médicale (MND202003011485 to F.D.B.), Retina France (I.A., C.Z. F.D.B.),
17
18 Fondation Fighting Blindness center grant [C-CMM-0907-0428-INSERM04 to I.A. and C.Z.],
19
20 grant (BR-GE-0619-0761-INSERM to I.A. and C.Z.), UNADEV (Union Nationale des Aveugles
21
22 et Déficients Visuels to I.A. and C.Z.) in partnership with ITMO NNP/AVIESAN (alliance
23
24 nationale pour les sciences de la vie et de la santé) for research in visual disorders, and doctoral
25
26 funding from the Ministère de l’Enseignement Supérieur et de la Recherche (MESR) and
27
28 Fondation de France (CM, JW).


29 30 31 32 33 34 35 36 37 **Ethics declaration**

38
39
40 Written informed consent was obtained from all individuals. All studies were carried out in
41
42 accordance with the declaration of Helsinki and were approved by a national ethics committee
43
44 (CPP Ile de France V, Project number 06693, N°EUDRACT 2006-A00347-44, 11 December
45
46 2006).
47
48
49



50 51 52 53 54 **Conflict of interest**

55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Figure 1


[Click here to access/download;Figure;Figure_1_11072023.pptx](#)**B**

Homozygous Regions for CIC03225
Total = 236.88 Mb (autosomes)

**C**

CIC03225 F1362, 23-year old

M1: c.910-7G>A p.(Phe304Profs*92)

CIC06076 F3377, 50-year oldM2: c.634_644del
p.(Ser212Alafs*44)**CIC09282 F5326, 55-year old**

M3: c.859C>T p.(Arg287*)




CIC13094 F7544, 50-year oldM2: c.634_644del
p.(Ser212Alafs*44)


Figure 3


A




UBAP1L expression retina



B









Control

AGTTTCT



CIC03225

AGCCCAAGTTTCT




CIC03228

AGCCCAAGTTTCT


AGTTTCT

Figure 6


A




B





C



**B**

Homozygous Regions for CIC03225
Total = 236.88 Mb (autosomes)


**C**

CIC03225 F1362, 23-year old

M1: c.910-7G>A p.(Phe304Profs*92)

CIC06076 F3377, 50-year old

M2: c.634_644del
p.(Ser212Alafs*44)



CIC09282 F5326, 55-year old

M3: c.859C>T p.(Arg287*)

CIC13094 F7544, 50-year old

M2: c.634_644del
p.(Ser212Alafs*44)







Figure 3




A

UBAP1L expression retina




B






Control

AGTTTCT



CIC03225

AGCCCAAGTTTCT




CIC03228

AGCCCAAGTTTCT


AGTTTCT

Figure 6


A



B




C



Supplementary table: summary of clinical findings in subjects with *UBAP1*-related retinal dystrophy

Individual	Age at examination	Clinical diagnosis at presentation	Symptoms at presentation	BCVA OD/OS	refraction	ff-ERG	Fundus examination	Fundus autofluorescence imaging
CIC03225, F1362	23	RCD	Night blindness and progressive visual field constriction	20/32 BE	-7 d BE	undetectable	pale optic disc, narrowed retinal vessels, pigmentary changes in the periphery, central macular atrophy	loss of autofluorescence in the periphery and in the macular region
CIC06376, F3377	50	CRD	Decreased central vision	20/200 20/50	None	Residual scotopic responses	central macular atrophy extending into the mid periphery predominantly in the inferior sector;	loss of autofluorescence in the atrophic areas ; heterogeneous autofluorescence in the posterior pole
CIC09282, F5326	55	Stargardt disease/CRD	Decreased central vision	20/500 20/400	-0.50(-2.25)15° -0.50(-1.50)160°	Residual scotopic responses	pale optic discs, narrowed retinal vessels, atrophic changes in the macular area	loss of central autofluorescence surrounded by a ring of increased autofluorescence
CIC13094, F7544	50	CRD	Night blindness and decreased central vision	20/160 20/800	-1.25(-2)175° -0.50(-2.25)175°	Residual scotopic responses	central macular atrophy	speckled appearance of autofluorescence at the posterior pole and mid periphery

BCVA : best corrected visual acuity ; OD : oculo dextra; OS: oculo sinistra; BE: both eyes; d: diopter; ff-ERG: full field electroretinogram



Supplementary figure: normal color fundus photograph (upper pictures) and normal fundus autofluorescence imaging (lower pictures) of the right (RE) and left (LE) eye.

January, 16th 2024

GENETMED-D-23-00572R3

Variants in *UBAP1L* lead to autosomal recessive rod-cone and cone-rod dystrophy

Christina Zeitz^{1*}, Julien Navarro^{1#}, Leila Azizzadeh Pormehr^{1,2#}, Cécile Méjécase^{1,3#}, Luiza M. Neves^{1,4#}, Camille Letellier^{1#}, Christel Condroyer¹, Shahad Albadri¹, Andréa Amprou¹, Aline Antonio¹, Tasnim Ben-Yacoub¹, Juliette Wohlschlegel^{1,5}, Camille Andrieu⁶, Malo Serafini¹, Lorenzo Bianco^{1,7}, Alessio Antropoli^{1,7}, Marco Nassisi^{1,8}, Said El Shamieh^{1,9}, Sandra Chantot-Bastaraud¹⁰, Saddek Mohand-Saïd^{1,6}, Vasily Smirnov^{1,11}, José-Alain Sahel^{1,6,12}, Filippo Del Bene¹, Isabelle Audo^{1,6*}

Maia Brunstein



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s). I give permission to be named as a person in the acknowledgments.

Beatrix Meha



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s). I give permission to be named as a person in the acknowledgments.

January, 16th 2024

GENETMED-D-23-00572R3

Variants in *UBAP1L* lead to autosomal recessive rod-cone and cone-rod dystrophy

Christina Zeitz^{1*}, Julien Navarro^{1#}, Leila Azizzadeh Pormehr^{1,2#}, Cécile Méjécase^{1,3#}, Luiza M. Neves^{1,4#}, Camille Letellier^{1#}, Christel Condroyer¹, Shahad Albadri¹, Andréa Amprou¹, Aline Antonio¹, Tasnim Ben-Yacoub¹, Juliette Wohlschlegel^{1,5} Camille Andrieu⁶, Malo Serafini¹, Lorenzo Bianco^{1,7}, Alessio Antropoli^{1,7}, Marco Nassisi^{1,8}, Said El Shamieh^{1,9}, Sandra Chantot-Bastaraud¹⁰, Saddek Mohand-Saïd^{1,6}, Vasily Smirnov^{1,11}, José-Alain Sahel^{1,6,12}, Filippo Del Bene¹, Isabelle Audo^{1,6*}

Christina Zeitz,



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Julien Navarro



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Leila Azizzadeh Pormeh



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional

Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Cécile Méjécase



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Luiza M. Neves



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Camille Letellier




I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Christel Condroyer



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Shahad Albadri



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Andréa Amprou



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Aline Antonio



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Tasnim Ben-Yacoub




I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Juliette Wohlschlegel



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Camille Andrieu



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Malo Serafini




I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Lorenzo Bianco




I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Alessio Antropoli




I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Marco Nassisi



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Said El Shamieh




I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Sandra Chantot-Bastaraud



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Saddek Mohand-Saïd



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Vasily Smirnov



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

José-Alain Sahel



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Filippo Del Bene



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).

Isabelle Audo



I attest that the research included in this report was conducted in a manner consistent with the principles of research ethics, such as those described in the Declaration of Helsinki and/or the Belmont Report. In particular, this research was conducted with the voluntary, informed consent of any research participants, free of coercion or coercive circumstances, and received Institutional Review Board (IRB) or Research Ethics Committee (REC) approval consistent with the principles of research ethics and the legal requirements of the lead authors' jurisdiction(s).